8. INTERNATIONAL MEDITERRANEAN SCIENTIFIC RESEARCH CONGRESS

August 13-15, 2025 / Antalya, Türkiye

FULL TEXTS BOOK

volume-2

Editors

Assoc. Prof. Dr. Ethem İlhan ŞAHİN

Assist Prof. Dr. Özlem ÖZKAN ÖNÜR







ISBN: 979-8-89695-160-5

8. INTERNATIONAL MEDITERRANEAN SCIENTIFIC RESEARCH CONGRESS

August 13-15, 2025 / Antalya, Türkiye

FULL TEXTS BOOK

VOLUME-2

Editors

Assoc. Prof. Dr. Ethem İlhan ŞAHİN
Assist Prof. Dr. Özlem ÖZKAN ÖNÜR

05.09.2025

by Liberty Academic Publishers, New York, USA
ALL RIGHTS RESERVED NO PART OF THIS BOOK MAY BE REPRODUCED IN ANY FORM, BY PHOTOCOPYING OR BY ANY ELECTRONIC OR MECHANICAL MEANS, INCLUDING INFORMATION STORAGE OR RETRIEVAL SYSTEMS, WITHOUT PERMISSION IN WRITING FROM BOTH THE COPYRIGHT OWNER AND THE PUBLISHER OF THIS BOOK.
© Liberty Academic Publishers 2025
The digital PDF version of this title is available Open Access and distributed under the terms of the Creative Commons Attribution-Non Commercial 4.0 license (http://creativecommons. org/licenses/by-nc/4.0/) which permits adaptation, alteration, reproduction and distribution for noncommercial use, without further permission provided the original work is attributed. The derivative works do not need to be licensed on the same terms.
adopted by Mariam Rasulan

ISBN - 979-8-89695-160-5

FULL TEXTS BOOK

CONGRESS ID

TITLE OF CONGRESS

8. INTERNATIONAL MEDITERRANEAN SCIENTIFIC RESEARCH CONGRESS

PARTICIPATION

Face to Face & Online

DATE - PLACE

August 13-15, 2025 / Antalya, Türkiye

ORGANIZATION

İKSAD-Institute of Economic Development and Social Researches

ORGANIZING COMMITTEE MEMBERS

Prof. Dr. Jose Rodolfo Hernandez CARRION - Valencia University

Prof. Dr. Mykola VAS'KIV - Borys Grinchenko University

Assoc. Prof. Dr. Olha BYKOVA, Borys Grinchenko Kyiv University

Dr. Ethem I. ŞAHİN- Adana Alpaslan Türkeş University

Assoc. Prof. Dr. Tuba ERKAYA KOTAN - Ataturk University

Assist. Prof. Dr. Mehmet TATOĞLU - Recep Tayyip Erdoğan University

Dr. Mehtap DEMİR- Adıyaman University

Dr. Mariam S. OLSSON, USA

Dr. Alina Amanzholova - Hodja Akhmet Yassawi University

Dr. Nurlan Akhmetov - Hodja Akhmet Yassawi University

FULL TEXTS BOOK

INTERNATIONAL SCIENTIFIC COMMITTEE

Prof. Ana CAMPINA - University Fernando Pessoa

Prof. Dr. Mykola VAS'KIV - Borys Grinchenko Kyiv University

Prof. Dr. Fahriye ALTINAY - Yakın Doğu Universitesi/Near East University

Prof. Dr. Mohamed RAMMAL - Saint Joseph University

Prof. Dr. Gökmen DAĞLI- Kyrenia University-Girne Üniversitesi

Prof. Dr. Fadi Al-Turjman, Near East University

Prof. Violla MAKHZOUM - Islamic University of Lebanon

Prof. Dr. Robert REPNIK - University of Maribor

Prof. Dr. Ahmed TLILI - Smart Learning Institute of Beijing Normal University (SLIBNU)

Prof. Dr. Maiga CHANG - Athabasca University

Prof. Dr. Zehra ALTINAY– Yakın Doğu Universitesi/Near East University

Prof. Dr. Nilüfer GALIP – Kyrenia-Girne University

Prof. Dr. Ebba Ossiannilsson - Lund Univesity

Prof. Dr. KINSHUK - North Texas University

Prof. Dr. Rustam SHADIEV - Zhejiang University

Prof. Dr. Ramesh SHARMA - Ambedkar University Delhi

Prof. Dr. Divna Vuksanovic - University of Arts in Belgrade

Assoc. Prof. Dr. Olha BYKOVA - Borys Grinchenko Kyiv University

Assoc. Prof. Dr. Tuba ERKAYA KOTAN - Ataturk University

Assoc. Prof. Gökhan OFLUOĞLU - Zonguldak Bülent Ecevit University

Assist. Prof. Dr. Mehmet TATOĞLU - Recep Tayyip Erdoğan University

Assist. Prof. Dr. Katarina Smakic - Universtiy Union – Nikola Tesla

Dr. Elvan CAFAROV, Azerbaijan State Pedagogical University/Azerbaijan

Dr. Alina AMANZHOLOVA - Hodja Akhmet Yassawi University/Kazakhstan

Dr. Nurlan AKHMETOV - Hodja Akhmet Yassawi University/Kazakhstan

CONGRESS COORDINATOR

Assist. Prof. Dr. Alina AMANZHOLOVA - Khoja Akhmet Yassawi International Kazakh-Turkish University, Kazakhstan

PARTICIPATING COUNTRIES (27)

Türkiye, Azerbaijan, TRNC, Pakistan, UK, Romania, Canada, USA, Malaysia, India, Czech Republic, Lebanon, Kazakhstan, Italy, Algeria, Bangladesh, Portugal, Ethiopia, Georgia, Ukraine, Switzerland, Morocco, China, Germany, Nigeria, Croatia, Kosovo

TOTAL ABSTRACTS: 420


The number of abstracts from foreign countries: **214**

The number of abstracts from Türkiye: **206**


LANGUAGES


Turkish, English


PHOTO GALLERY







observer hall-6
Hamza EL KISSA...
Electrochemical sensors also provide fast response times and high sensitivity, allowing for the detection of very low concentrations of 4-NP with minimal sample preparation.
Furthermore, by tailoring the electrode surface with functional materials—such as conductive polymers, nanoparticles, or composite films—the selectivity and detection performance can be significantly enhanced, making electrochemical methods a versatile and powerful tool for environmental monitoring.
In this presentation, the electrochemical determination of 4-NP was performed using a sensor working electrode prepared in two steps.
5



Workplace
Observer Hall-5
HALL 5_Nejat D...
Участники (4)
Observer Ha... (Сопрезентер, я)
Hall-5, Cüneyt YILMAZ
Mehmet Direkli
HALL 5_Nejat DOGAN
Mehmet Direkli



Observer Hall-5
Hall-5 Ananda...
tunen
mohd zeeshan
DEFINITION AND PRINCIPLES OF SMART TOURISM
Smart tourism utilizes technology to improve travel experiences by offering personalized services and efficient operations. Principles include connectivity, data-driven decision making, and user-centered design. By integrating these elements, destinations can elevate their offerings while ensuring that environmental and cultural impacts are minimized.



Observer hall-1
Hall-1, Simge B...
sude kanber
Hemşirelikte Etik ve Malpraktis



OBSERVER H3
Ulya Yürük


Observer Hall-2
OKUL ÖNCESİNDE SOSYAL BECERİLER VE AKRAN İLİŞKİLERİ: BİR BİBLİYOMETRİK ANALİZ
YÜKSEK LİSANS ÖĞRENCİSİ ŞEVVAL ERVA KAYA
DR. ÖĞR. ÜYESİ MİNE KOYUNCU ŞAHİN


Observer Hall-2
Dramaturginin Aurası: Benjamin'den Yapay Zekâya Teknik Yeniden Üretim Çağında Teatral Özgünlük
Dr. Öğr. Üyesi Vecihe Özge Zeren
Çanakkale Onsekiz Mart Üniversitesi
Güzel Sanatlar Fakültesi,



Observer Hall-2
August 13-15, 2025 / Antalya, Türkiye
KARBONATLAŞMANIN MAGNEZYUM ES
YAPAY İRİ AGREGALARIN TANECİK DAY
ÜZERİNDEKİ ETKİLERİ
Enis COŞKUN
Onur ŞAHİN



observer hall-6
Hall-4 Abdelha...
8. INTERNATIONAL MEDITERRANEAN SCIENTIFIC RESEARCH CONGRESS
August 13-15, 2025 / Antalya, Türkiye



Observer Hall-5
Hall 5- Session...
Dr Afshan Nase...
Katılımcılar (6)
Introduction
• Academics plays vital role in developing globalization and Internationalization Trends.
• Along with the opportunities, it contains threats as well.
• International Students play the role of ambassadorship to the host country.
• International students studying in Pakistani universities encounter a myriad of challenges that influence their academic and personal experiences.
• These challenges encompass linguistic barriers, cultural adjustments, financial limitations, visa complexities, and housing concerns.
• Examining these challenges provides insights into the diverse needs of international students and informs the development of targeted support mechanisms to improve their overall educational journey in Pakistan.
UMT



Observer H-2
Buse ALTIN


KANSER
• Kanser, hücrelerin kontrolsüz şekilde bölünmesi ve doku mikroçevresinden bağımsız olarak büyümesiyle karakterize, heterojen yapıda bir hastalıktır.
• Bu heterojenite, tümör içinde farklı özelliklere sahip hücre alt popülasyonlarının bulunmasına ve hastalığın seyrinin değişkenlik göstermesine neden olur.
• GLOBOCAN 2022 verilerine göre, dünya çapında yaklaşık 20 milyon yeni kanser vakası bildirilmiş ve yaklaşık 9,7 milyon kişi kanser nedeniyle yaşamını yitirmiştir
Observer H-3
Hall-3 Aysel Bay...


observer hall-6
chayaraju balas...
INTRODUCTION
Mechanical power transmission is a fundamental aspect of mechanical engineering, facilitating the efficient transfer of energy from one point to another. These categories include mechanical, hydraulic mechanical, hydrostatic (volumetric hydraulic), and electric transmission systems.
i. Mechanical Transmission Systems: They are composed of essential mechanical components such as the clutch, gearbox, driveshaft, and universal joints, which work in concert to effectively transmit engine power.
ii. Hydraulic Mechanical Transmission Systems: These systems utilize fluid, typically in the form of a hydraulic coupling or a hydraulic torque converter, to transfer power.
iii. Electrical Transmission Systems: These systems employ electric motors to either directly drive the wheels or supplement traditional mechanical or hydraulic systems.


observer hall-7
hall -7 murat da...
Mehmet AVŞAR
Hall 7-Ayşe DEMİR


observer hall-6
Dr. Arti Saxena
H


Observer hall-1
1. GİRİŞ
Midyeler, su kirliliği, oksijen seviyeleri, sıcaklık değişiklikleri gibi faktörlere verdikleri tepkilerle çevresel değişimlerin izlenmesinde görev yapmaktadır (Hartmann ve ark., 2016) (Şekil 1).
Midyelerin genomları ve genetik çeşitlilikleri üzerine yapılan çalışmalar, bu organizmaların çevresel faktörlere nasıl yanıt verdiği hakkında derinlemesine bilgi edinmemize olanak tanımaktadır (Tanguy ve ark., 2008).

8. INTERNATIONAL MEDITERRANEAN SCIENTIFIC RESEARCH CONGRESS

August 13-15, 2025 / Antalya, Türkiye

CONGRESS PROGRAM

Meeting ID: 862 1528 3413
Passcode: 131415

Join Zoom Meeting
https://us02web.zoom.us/j/86215283413?pwd=niwCknpRwjg8bV63Cjg789WLFJ7R5x.1

PARTICIPANT COUNTRIES (27):

Türkiye, Azerbaijan, TRNC, Pakistan, UK, Romania, Canada, USA, Malaysia, India, Czech Republic, Lebanon, Kazakhstan, Italy, Algeria, Bangladesh, Portugal, Ethiopia, Georgia, Ukraine, Switzerland, Morocco, China, Germany, Nigeria, Croatia, Kosovo

Before you login to Zoom please indicate your name_surname and HALL number:

Önemli, Dikkatle Okuyunuz Lütfen

- Kongremizde Yazım Kurallarına uygun gönderilmiş ve bilim kurulundan geçen bildiriler için online (video konferans sistemi üzerinden) sunum imkanı sağlanmıştır.
- Online sunum yapabilmek için **https://zoom.us/join** sitesi üzerinden giriş yaparak "Meeting ID or Personal Link Name" yerine ID numarasını girerek oturuma katılabilirsiniz.
- Zoom uygulaması ücretsizdir ve hesap oluşturmaya gerek yoktur.
- Zoom uygulaması kaydolmadan kullanılabilir.
- Uygulama tablet, telefon ve PC'lerde çalışıyor.
- Her oturumdaki sunucular, sunum saatinden 5 dk öncesinde oturuma bağlanmış olmaları gerekmektedir.
- Tüm kongre katılımcıları canlı bağlanarak tüm oturumları dinleyebilir.
- Moderatör – oturumdaki sunum ve bilimsel tartışma (soru-cevap) kısmından sorumludur.

Dikkat Edilmesi Gerekenler- TEKNİK BİLGİLER

- Bilgisayarınızda mikrofon olduğuna ve çalıştığına emin olun.
- Zoom'da ekran paylaşma özelliğine kullanabilmelisiniz.
- Kabul edilen bildiri sahiplerinin mail adreslerine Zoom uygulamasında oluşturduğumuz oturuma ait ID numarası gönderilecektir.
- Katılım belgeleri kongre sonunda tarafınıza pdf olarak gönderilecektir
- Kongre programında yer ve saat değişikliği gibi talepler dikkate alınmayacaktır

IMPORTANT, PLEASE READ CAREFULLY

- To be able to attend a meeting online, login via **https://zoom.us/join** site, enter ID "Meeting ID or Personal Link Name" and solidify the session.
- The Zoom application is free and no need to create an account.
- The Zoom application can be used without registration.
- The application works on tablets, phones and PCs.
- The participant must be connected to the session 5 minutes before the presentation time.
- All congress participants can connect live and listen to all sessions.
- Moderator is responsible for the presentation and scientific discussion (question-answer) section of the session.

Points to Take into Consideration - TECHNICAL INFORMATION

- Make sure your computer has a microphone and is working.
- You should be able to use screen sharing feature in Zoom.
- Attendance certificates will be sent to you as pdf at the end of the congress.
- Requests such as change of place and time will not be taken into consideration in the congress program.

exp. Hall-1, Seda BEYAZ



Session-1, Hall-1
13.08.2025
Moderator: Assoc. Prof. Dr. Çağlar SOYLU
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Assoc. Prof. Alime SELÇUK TOSUN Öznur ALAN	INVESTIGATION OF PERCEIVED HEALTH AND ITS ASSOCIATED FACTORS AMONG ADULTS	Selçuk University, Türkiye
Fatma Betül ÖZKALELİ Şahizer ERAYDIN	ETHICS AND MALPRACTICE IN NURSING	Tokat Gaziosmanpaşa University, Türkiye
Hande SEVEN AVUK Simge Betül AL Ceren AKIN Sude KANBER Elif YAĞCI	THE RELATIONSHIP BETWEEN E-HEALTHY NUTRITION LITERACY, SUSTAINABLE NUTRITION BEHAVIORS, AND ADHERENCE TO THE MEDITERRANEAN DIET AMONG ADULTS	İstanbul Bilgi University, Türkiye
Özgür Timuçin KUTLU	ONE RHABDOMYOLYSIS, ONE METHEMOGLOBINEMIA, ONE DEATH	Isparta Şehir Hastanesi, Isparta, Türkiye
Hale AVŞAR ERDEKLİ Selin KÖSE	ASSESSMENT OF ADHERENCE TO THE MEDITERRANEAN DIET IN ADULTS WITH METABOLIC SYNDROME	Antalya Bilim University, Türkiye
Hale AVŞAR ERDEKLİ Faruk Emre BOZEL	EVALUATION OF ERGOGENIC SUPPLEMENT USE AND EXERCISE ADDICTION IN ADULTS ENGAGED IN EXERCISE	Antalya Bilim University, Türkiye
Çağlar SOYLU	THE ACUTE EFFECTS OF EXERCISE-INDUCED FATIGUE ON HAMSTRING MUSCLE ARCHITECTURE AND CORTICOSPINAL EXCITABILITY: AN EXPERIMENTAL STUDY IN HEALTHY INDIVIDUALS	Health Sciences University, Türkiye Sports Performance, Recovery, Injury & New Technologies (SPRINT) Research Centre
Çağlar SOYLU	PILOT STUDY ON RELATIONSHIPS BETWEEN MUSCLE ARCHITECTURE, CORTICOSPINAL EXCITABILITY, HAMSTRING OUTCOME SCORE, AND ECCENTRIC STRENGTH IN ADULTS WITH PRIOR HAMSTRING INJURY	Health Sciences University, Türkiye Sports Performance, Recovery, Injury & New Technologies (SPRINT) Research Centre

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-1, Hall-2
13.08.2025
Moderator: Assist. Prof. Dr. Murat KONCA
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Selçuk ALBAYRAK Oğuz MERHAN Lale BAŞER	THE ROLE OF COMMON SWIFTS (APUS APUS) IN THE ECOSYSTEM AND THEIR INTERACTION WITH HUMANS: REHABILITATION AND FEEDING PRACTICES	Kafkas University, Türkiye
Selçuk ALBAYRAK Oğuz MERHAN Kadir BOZUKLUHAN	LONG-TERM CASE ANALYSIS OF WILDLIFE REHABILITATION (2018–2025)	Kafkas University, Türkiye
Rezzan UYSAL Ezgi TÜRKAY	THE EFFECT OF PARENTAL FOOD NEOPHOBIA ON CHILD NEOPHOBIA AND EVALUATION OF ANTHROPOMETRIC MEASUREMENTS	Antalya, Türkiye Antalya Bilim University, Türkiye
Su Lal CANSEV Hatice Kübra ARAS	THE EFFECT OF CHANGED DIETARY HABITS DUE TO PREMENSTRUAL SYNDROME ON HEDONIC AND EMOTIONAL EATING BEHAVIOR	Antalya Bilim University, Türkiye
Emine ASLAN Berna Gamze ÜNSAL Busenur AKANSU Nilüfer AKIŞ Mehmet KOCA	DETERMINATION OF UNIVERSITY STUDENTS' GENDER ROLES WITH THEIR KALEIDOSCOPE CAREER CONCEPTION	Malatya Turgut Özal University, Türkiye
Cem Cahit BARIŞIK	CLINICAL, RADIOLOGIC AND PATHOLOGIC REVIEW OF CASTLEMAN DISEASE IN CHILDREN	İstanbul Medipol University, Türkiye
Murat KONCA	THE RELATIONSHIP BETWEEN DEBT AND HEALTH STATUS: THE TURKISH CASE	Çankırı Karatekin University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-1, Hall-3
13.08.2025
Moderator: Assist. Prof. Dr. Murat BİROL
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Cuma AKBAY Hamdi AYYILDIZ	NAVIGATING A CHANGING WORLD: HOW TO OVERCOME CLIMATE GRIEF USING ADAPTIVE CAPACITY?	Kahramanmaras Sutcu Imam University, Türkiye
Cuma AKBAY Hamdi AYYILDIZ	FACILITATING STAKEHOLDER ENGAGEMENT: A MULTIFACETED APPROACH TO POSITIVE SOCIETAL IMPACT OF EMERGING TECHNOLOGIES AGAINST TO CLIMATE CHANGE	Kahramanmaras Sutcu Imam University, Türkiye
Birol ALTUN	COMMUNICATION PROBLEMS EXPERIENCED DURING WORKING FROM HOME DURING THE COVID-19 PANDEMIC	Çalışma Ekonomisi Ve Endüstri İlişkileri, Türkiye
Hazal ZAİMOĞLU Cavit YAVUZ	TARAFSIZLIK MASKESİ ALTINDA DİJİTAL TARAFGİRLİK: GROK VE FİLİSTİN SÖYLEMİ	Ordu University, Türkiye
Murat BİROL Damla Nur TÜRK	DIGITAL PLATFORM SERIES AS A SPACE FOR OBSERVING TOXIC RELATIONSHIPS: CASE STUDIES OF “KİMLER GELDİ KİMLER GEÇTİ” AND “İLK VE SON”	Giresun University, Türkiye Ankara, Türkiye
Harun KILCI Murat Fatih TUNA	ARTIFICIAL INTELLIGENCE VS. REALITY: A COMPARISON OF THE REASONS PREVENTING THE USE OF DIGITAL TWINS IN HOSPITALS WITH THE ACTUAL SITUATION	Sivas Cumhuriyet University, Türkiye
Tuna ŞENER	A COMPARATIVE ANALYSIS OF THE RISK INDICATORS OF THE TURKISH ECONOMY WITH THE GREECE ECONOMY	Trakya University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-1, Hall-4
13.08.2025
Moderator: Dr. Sümeyye ÇITAK
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Aytan HUMMATOVA Ayla EKER SARIBOYACI Onur UYSAL Murat KAYA Huseyn Hasanzade Hüseyin AVCI Hamed GHORBANPOOR	DEVELOPMENT AND OPTIMIZATION OF AN INNOVATIVE MICROFILTRATION SYSTEM FOR CELLULAR SEPARATION IN MICROFLUIDIC CHIPS	Osmangazi University, Türkiye
Timur KOCA Sümeyye ÇITAK Aylin Fidan KORCUM	RADIATION-INDUCED SKIN TOXICITIES IN HEAD AND NECK CANCERS: CLINICAL SIGNIFICANCE AND MANAGEMENT APPROACHES	Akdeniz University, Türkiye
Narmin ADIGOZALOVA Murat KAYA Özlem TOMSUK Onur UYSAL Ayla EKER SARIBOYACI Hüseyin AVCI Hamed GHORBANPOOR	INVESTIGATION OF IN VITRO HEPATOCELLULAR CARCINOMA MODEL WITH MESENCHYMAL STEM CELLS	Eskişehir Osmangazi University, Türkiye
Seljan MAMMADOVA Zineb BENZAİT Ceren ÖZEL Özlem TOMSUK Murat KAYA Merve Nur SOYKAN Burcugül ALTUĞ Hamed GHORBANPOOR Onur UYSAL Ayla EKER SARIBOYACI Hüseyin AVCI	DECELLULARIZATION AND CHARACTERIZATION OF EXTRACELLULAR MATRIX FROM HUMAN UMBILICAL CORD	Eskişehir Osmangazi University, Türkiye Division of Engineering in Medicine, Department of Medicine Brigham and Women's Hospital Harvard Medical School Cambridge, MA 02139, USA. Dokuz Eylül University, Türkiye
V.Martini, M. Cianfriglia, S. Leporale, Roberta Moramarco, C. Marocco, C. Bramucci, F. Sestito, R. Reda, L. Testarelli, T. Testori, G. Perrotti	OPTIMIZING ORTHOGNATHIC OUTCOMES: THE ROLE OF 3D CEPHALOMETRY AND TFA IN VERTICAL SKELETAL ANALYSIS	Sapienza University of Rome, Rome, Italy Fondazione Policlinico Universitario Campus Bio-Medico, Italy IRCCS Istituto Ortopedico Galeazzi, 20161 Milan, Italy; University of Milan, 20126 Milan, Italy University of Michigan, Ann Arbor, MI 48109, USA Lake Como Istitute, 22100, Como, Italy
V.Hariharan, S.Harinisri, M.Farhat yassir, R.Thelshath	HORMONAL CONTROL OF THE REPRODUCTIVE SYSTEM	Bharath Institute Of Higher Education And Research Chennai, India
V.Hariharan, S.Harinisri, M.Farhat yassir, R.Thelshath	MICROSCOPIC STRUCTURE OF HUMAN TISSUES	Bharath Institute Of Higher Education And Research Chennai, India

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-1, Hall-5
13.08.2025
Moderator: Assoc. Prof. Dr. Cüneyt YILMAZ
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Nejat DOGAN	THE EU SECURITY AND DEFENSE ACQUIS: LAW, INSTITUTIONS, AND DYNAMIC LEGAL FUNCTIONALISM	Palacky University Olomouc, Czech Republic
Dr. Mehmet DIREKLI	A REALPOLITIK PERSPECTIVE ON RUSSIA'S INTERVENTION DURING THE SYRIAN CIVIL WAR	Cyprus Aydın University, TRNC
Mr. Satish Kumar Singh	THE ROLE OF LEGAL FRAMEWORKS AND CULTURAL PRACTICES IN SHAPING WOMEN'S HISTORICAL STRUGGLES FOR FOOD SECURITY	Central University of Punjab, India
Cüneyt YILMAZ	SOCIAL AND POLITICAL EFFECTS OF DIGITAL TRANSFORMATION	St.Clements University, Political Science, London, United Kingdom
Cüneyt YILMAZ	UNORGANIZED CROWD BEHAVIOR: A SOCIAL PSYCHOLOGICAL ANALYSIS	St.Clements University, Political Science, London, United Kingdom
Mihaela CUREA	ACCOUNTING FOR INTANGIBLES: STANDARDS, GAPS, AND EMERGING PERSPECTIVES	Alexandru Ioan Cuza University, Faculty of Economics and Business Administration, Iaşi, Romania

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-1, Hall-6
13.08.2025
Moderator: Assist. Prof. Dr. Ziya CAN
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Şener SAĞLAM	ELECTROCHEMICAL DETERMINATION OF 4-NITROPHENOL USING A POLY(4-NITROANILINE) /GOLD NANOPARTICLES MODIFIED GLASSY CARBON ELECTRODE	İstanbul University – Cerrahpaşa, Türkiye
Dr., Andreea Irina Barzic Vasile Bogdan Mardari Dr., Raluca Marinica Albu	THE ROLE OF TiO_2 NANOPARTICLES IN IMPROVING THE OPTICAL AND ELECTRICAL CHARACTERISTICS OF POLYMER NANOCOMPOSITES FOR LED DEVICES	"Petru Poni" Institute of Macromolecular Chemistry, Iasi, Romania "Alexandru Ioan Cuza" University, Faculty of Chemistry, Iasi, Romania
Shaheen Kousar, Shazia Perveen, Sumaira Kanwal	DUAL THERAPEUTIC POTENTIAL OF 2-(2-NITROPHENYL)-1,3-THIAZOLIDINE-4-CARBOXYLIC ACID: HEPATOPROTECTION AND IRON REGULATION IN A ROTENONE-INDUCED PARKINSON'S DISEASE MODEL	The Women University Multan, Pakistan COMSATS University Islamabad, Sahiwal Campus, Pakistan
Kabiri Loubna, Forsal Issam	GREEN CORROSION INHIBITOR OF CARBON STEEL IN ACID MEDIUM: PLANT EXTRACT	Sultan Moulay Slimane University, Morocco.
Hamza EL KISSANY, Hicham EL-BOUKHARY, Hicham EL KISSANI, Adil LAAZIZ, Abdelouahed HAJJAJI	DIVERSITY OF MICROBIAL ENDOPHYTES, THEIR ASSOCIATION WITH PLANTS, AND ROLE IN SECONDARY METABOLITE PRODUCTION: A REVIEW	Sultan Moulay Slimane University, Beni Mellal, Morocco. Sidi Mohamed Ben Abdellah University, Fez, Morocco
ROSSAFI Bouchra, ABCHIR Oussama, ELKOUALI Mhammed, CHTITA Samir	STRUCTURE-BASED VIRTUAL SCREENING AND IN SILICO EVALUATION OF MARINE ALGAE METABOLITES AS POTENTIAL A-GLUCOSIDASE INHIBITORS FOR ANTIDIABETIC DRUG DISCOVERY	Hassan II University of Casablanca, Morocco
Zaher BABETJI Mustafa PEHLİVAN Pınar YUMRUTAŞ Önder YUMRUTAŞ	DETERMINATION OF CYTOTOXIC AND APOPTOTIC OF THYMUS KOTSCHYANUS var. GLABRESCENS-SILVER NITRATE (AGNO3) NANOPARTICLES	Gaziantep University, Türkiye Adıyaman University, Türkiye
Ziya CAN	DETECTION OF NITRIC OXIDE RADICAL USING FUNCTIONALIZED CARBON DOTS	İstanbul University – Cerrahpaşa, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-1
13.08.2025
Moderator: Assist. Prof. Dr. Asuman GÜNAY YILMAZ
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Asuman GÜNAY YILMAZ	TOMATO LEAF DISEASE CLASSIFICATION USING ConvNeXt-T MODEL	Karadeniz Technical University, Türkiye
Veli ÇAPALI Murat KAÇAL Burak DEMİR	INTEGRATION OF UNIVERSITY SOFTWARE APPLICATIONS WITH CENTRAL IDENTITY MANAGEMENT SYSTEM AND INTEGRATED PORTAL DESIGN EXAMPLE, SDUNET	Süleyman Demirel University, Türkiye
Samet GÜRGEN	ARTIFICIAL NEURAL NETWORK MODELS IN SHIP MAIN ENGINES: CURRENT APPLICATIONS AND FUTURE PERSPECTIVES	İskenderun Technical University, Türkiye
Rojhat ERGÜVEN Alp Buğra AYDIN Derya AVCI	SPI PREDICTION USING THE GRADIENT BOOSTING REGRESSOR (GBR) ALGORITHM	Fırat University, Türkiye
Yeliz BASAR	YAPAY ZEKÂNIN İNME GÖRÜNTÜLEMESİNDEKİ ROLÜ: BT TABANLI UYGULAMALARLA TANISAL DOĞRULUK VE İŞ AKIŞI VERİMLİLİĞİNİN ARTIRILMASI	Acibadem Healthcare Group, Türkiye
Sinan KÖSE Yaşar BECERİKLİ	SIGN LANGUAGE RECOGNITION WITH DEEP LEARNING: A CASE STUDY ON THE SIGN LANGUAGE MNIST DATASET	Kocaeli University, Türkiye
Sinan KÖSE Yaşar BECERİKLİ	SAFETY-AWARE MACHINE LEARNING SUPERVISOR FOR CROSS PARTITION MANAGEMENT IN ARINC 653 MULTICORE SYSTEMS	Kocaeli University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-2
13.08.2025
Moderator: Prof. Dr. Işıl Güneş MODIRI DILEK
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Işıl Güneş MODIRI DILEK	EXAMINATION OF FACTORS AFFECTING STUDENTS' PREFERING MUSIC HIGH SCHOOLS: EXAMPLE OF TRABZON PROVINCE	Trabzon University, Türkiye
Aytan MAMMADOVA	THE REASON OF PREVALENCE AND LONGEVITY OF PROVERBS	Sumgayıt State University, Azerbaijan
İrem ALMACI	DETERMINING THE ABILITY OF 4TH-YEAR SCIENCE EDUCATION UNDERGRADUATE STUDENTS TO DEFINE GRAVITATIONAL FORCE AFFECTING MASS AS WEIGHT USING ARGUMENT-BASED LEARNING TECHNIQUES	Kahramanmaraş Sütçü İmam University, Türkiye
Anvar ABBASOV Kamala QULIYEVA	PEDAGOGICAL CRITICISM IN MODERN EDUCATION	Baku Slavic University, Azerbaijan Azerbaijan State Pedagogical University, Azerbaijan
Berkay AKIN Mehmet İNCE	PARENTAL ROLE IN SUPPORTING LANGUAGE AND COMMUNICATION DEVELOPMENT	Abant İzzet Baysal University, Türkiye
Bahadır KILCAN Eyüp Burak KUZPINARI	EXAMINATION OF THE 2024 TÜRKİYE CENTURY EDUCATION MODEL SECONDARY SCHOOL SOCIAL STUDIES COURSE CURRICULUM IN TERMS OF DATA LITERACY	Gazi University, Türkiye
Ahmet KATILMIŞ Ayça ÖZTÜRK	APPLYING THE SYNECTICS METHOD IN SOCIAL STUDIES CLASSES	Marmara University, Türkiye
Esra ZEYNEL	A THEORETICAL DISCUSSION ON THE CONTEMPORARY RELEVANCE OF THE NEED FOR AUTONOMY IN THE WORKPLACE WITHIN THE FRAMEWORK OF SELF-DETERMINATION THEORY	Isparta University of Applied Sciences, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-3
13.08.2025
Moderator: Assist. Prof. Dr. Tuba DOGAN
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Salih Burak PERK Vehbi GÜNEŞ	INVESTIGETION of SEROPREVELANCE OF BOVINE HERPERSVIRUS TYPE 1 (BHV-1) and MANNHEIMIA HAEMOLYTICA IN BEEF AND DAIRY COWS IN KAYSERI PROVINCE	Erciyes University, Türkiye
Ramazan SARIKAYA Mehmet ÇİTİL	INVESTIGATION OF THE PREVALENCE OF BOVINE VIRAL DIARRHEA VIRUS (BVDV) IN NEONATAL CALVES	Erciyes University, Türkiye
Sedat BEHREM Yunus ARZIK	SOME ENVIRONMENTAL FACTORS AFFECTING LAMB GROWTH PERFORMANCE	Aksaray University, Türkiye
Sedat BEHREM Yunus ARZIK	THE STATUS OF WORLD CASHMERE PRODUCTION	Aksaray University, Türkiye
Mustafa Sedat ARSLAN Tuğba Melike PARLAK Ece OKTAY	DETERMINATION OF THE EMBRYOTOXIC EFFECT OF CEFAZOLINE BY IN OVO METHOD	Selçuk University, Türkiye
Mustafa Sedat ARSLAN Tuğba Melike PARLAK Ece OKTAY	DETERMINATION OF THE EMBRYOTOXIC EFFECT OF CEFTRIAXONE BY IN OVO METHOD	Selçuk University, Türkiye
Tuba DOGAN Esra AKTAS SENOCAK	BETA-CARYOPHYLLENE MITIGATES TESTICULAR DAMAGE INDUCED BY THIOACETAMIDE THROUGH ANTIOXIDANT AND ANTI-INFLAMMATORY PATHWAYS	Atatürk University, Türkiye
Hamza YALÇİN	EFFECT OF MISSING DATA COMPLETION METHODS ON PLS SEM PERFORMANCE	Harran University, Türkiye
Metin Can TORUN Davut BAYRAM Hürü Ceyda TÜM	AN ASSESSMENT OF HOUSING, NUTRITIONAL, AND BEHAVIORAL REQUIREMENTS OF WORKING DOGS IN MILITARY AND LAW ENFORCEMENT UNITS AND RECOMMENDATIONS FOR IMPROVEMENT	JAKEM Komutanlığı, Nevşehir, Türkiye Erciyes University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-4
13.08.2025
Moderator: Major Gheorghe GIURGIU
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Rizwana Raheel Humaira Majeed Khan	PROTECTIVE ANTIOXIDANT AND ANTI-INFLAMMATORY EFFECTS OF METHYL FURFURAL AGAINST DRUG-INDUCED INJURY IN RATS	Lahore College for Women University, Lahore, Pakistan. / Minhaj University, Lahore, Pakistan.
Major Gheorghe GIURGIU, Prof. dr. med. Manole COJOCARU	MICROBIOTA MODULATION AS THERAPEUTIC APPROACH IN THE NEUROPATHIC PAIN IN DOG WITH SPINAL CORD INJURY: IMPACT OF POLENOPLASMIN	Deniplant-Aide Sante Medical Center, Biomedicine, Bucharest, Romania Academy of Romanian Scientists Titu Maiorescu University, Faculty of Medicine, Bucharest, Romania
V.Hariharan,S.Harinisri,M.Farhat yassir,R.Thelshath	CARDIOVASCULAR ADAPTATION TO EXERCISE	Bharath Institute Of Higher Education And Research, Chennai, India
Saraswathi.V,R. Vidhyalakshmi, Kowsika.M, Dhanush kumar.S, Sastha.K, Prashanth.S, Dr.W.Helen, Dr.R.Srinivasam	ROLE OF COFACTORS AND COENZYMES IN METABOLIC PATHWAYS	Bharath Institute of Higher Education and Research, Chennai, Tamilnadu India.
V.Ranjani, S.Deen Ansar, V.Varalakshmi, E.Velmurugan, Dr.W.Helen	HISTORICAL DEVELOPMENT OF PUBLIC HEALTH PROGRAMS IN INDIA	Bharath Institute of Higher Education and Research, Chennai, Tamilnadu, India.
USMAN SAMA'ILA IBRAHIM SAMAILA AISHA ABUBAKAR ABDU	THE APPLICATION OF NANOPARTICLES IN DIAGNOSIS OF INFECTIOUS DISEASE: A REVIEW	Umaru Musa Yaradua Unıversity, Nigeria
ARRAJI Maryem, CHAHBOUNE Mohamed	PREVALENCE AND DETERMINANTS OF MEDICATION ADHERENCE IN MOROCCAN PATIENTS WITH TYPE 2 DIABETES: A MULTICENTER CROSS-SECTIONAL STUDY	Hassan First University of Settat
Adegbola Oluwatosin Oluwabukola, Fakeye Adeola Olajumoke	MOLECULAR DOCKING OF BETACARYOPHELLENE AND MIFEPRISTONE AGAINST COLON CANCER-RELATED PROTEINS IN CHEMOPREVENTION AND CHEMOTHERAPY COLON CANCER	Federal Polytechnic, Ilaro, Nigeria

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-5
13.08.2025
Moderator: Dr. G. I. Onwuka
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Dr. Gerald Onwuka, Dr. Babayemi A.W, KENNETH Ijere	ANALYSING FACTORS INFLUENCING HEPATITIS B INFECTIVITY A LOGISTIC REGRESSION-BASED MODELS	Kebbi State University of Science and Technology, Aliero, Nigeria, Faculty of Science, Department of Mathematics
Shahid Abdullah, Aijaz Ahmad Magray, Neha Choubey	A NOVEL TWO-POINT ITERATIVE METHOD FOR SOLVING NON-LINEAR EQUATIONS WITH MEMORY	VIT Bhopal University, Kothri-Kalan, India JB Institute of Engineering and Tecgnology, India
Amare Worku Demsie, Awoke Andargie Tiruneh, Endalew Getnet Tsega	A FITTED NUMERICAL APPROACH FOR SINGULARLY PERTURBED DELAY DIFFERENTIAL EQUATIONS WITH SHIFT PARAMETERS	Bahir Dar University, P. O. Box: 79, Bahir Dar, Ethiopia Woldia University, P. O. Box: 400, Woldia, Ethiopia
Musa Abdulrahman, Prof. Babayemi, A.W, Dr. G. I. Onwuka, Sumayya Jibril	VOLATILITY OF RICE PRICE DYNAMICS IN KEBBI STATE, NIGERIA	Kebbi State University of Science and Technology, Aliero, Nigeria College of Advance and Legal Studies Yauri
Dr. G. I. Onwuka, Prof. Babayemi A. W., B. Sheh	TWO SAMPLE MULTIVARIATE BEHRENS FISHER PROBLEM FOR HIGH DIMENSIONAL DATA	Kebbi State University of Science and Technology, Aliero, Nigeria Kebbi State Polytechnic Dakingari
Faiza Baghida	FIRST-PRINCIPLES CALCULATIONS BASED ON DFT-PBE WERE EMPLOYED TO INVESTIGATE THE STRUCTURAL, ELECTRONIC, AND THERMOELECTRIC PROPERTIES OF Cu_2Se	Moulay Ismail University Faculty of Sciences, Meknes ,Morocco
Poonam	EFFECT OF ELECTROLYTE ON CHARGE STORAGE PERFORMANCE İN A COMPOST-BASED DEVİCE	Indira Gandhi University, Research Scholar, Department of Physics Rewari, Haryana, India
Chandani Sharma, Ghazala Ansari , Aarti Sharma, Manidipa Roy	ADVANCED DESIGN AND OPTIMIZATION STRATEGIES FOR WEARABLE HEALTH DEVICES FOR CHRONIC DISEASE MANAGEMENT	Deemed to be University, India. SRM Institute of Science and Technology, Modinagar, NCR Campus ABES Engineering College, Ghaziabad ECE Department, ABES Engineering College

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-6
13.08.2025
Moderator: Dr. Dhanraj T. Dhangar
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Aeman Farooq, Dr. Anam Khan, Umm e Eman Syed, Adnan Adil, Neelam Bibi, Dr. Saba Ghayas	RELATIONSHIP BETWEEN LOVE BOMBING AND MENTAL HEALTH AMONG ADULTS: MEDIATING ROLE OF EMOTIONAL ABUSE	Govt. College Women University Sialkot Rawalpindi Institute of Cardiology University of Sargodha
Hajra Bibi, Dr. Anam Khan, Umm e Eman Syed, Neelam Bibi, Dr. Adnan Adil, Dr. Sadia Niazi	MEDIATING ROLE OF COMMUNICATION QUALITY BETWEEN EXCESSIVE INTERNET USAGE AND MARITAL SATISFACTION AMONG MARRIED INDIVIDUALS	Rawalpindi Institute of Cardiology Govt. College Women University Sialkot University of Sargodha
Faiza Shams, Dr. Anam Khan, Mam Umme Aiman Syed, Adnan Adil, Neelam Bibi, Dr. Sadia Niazi	RELATIONSHIP BETWEEN BREADCRUMBING AND RELATIONSHIP SATISFACTION AMONG MARRIED INDIVIDUALS: MEDIATING ROLE OF EMOTIONAL ABUSE	Govt. College Women University Sialkot Rawalpindi Institute of Cardiology University of Sargodha
Kainat Hassan, Dr Anam Khan, Umm e Eman Syed, Neelam Bibi, Dr. Adnan Adil, Dr. Sadia Niazi	MEDIATING ROLE OF EMOTIONAL REGULATION IN COGNITIVE FLEXIBILITY AND DECISION MAKING AMONG FAMILY ADVOCATES	Rawalpindi Women University Govt. College Women University Sialkot University of Sargodha
Syeda Menahil Fatima, Dr. Anam Khan, Umm e Eman Syed, Neelam Bibi, Dr. Adnan Adil, Dr. Sadia Niazi	RELATIONSHIP BETWEEN STRESS AND COMPULSIVE BUYING BEHAVIOR: MODERATING ROLE OF SELF-CONTROL AMONG WOMEN	Rawalpindi Women University Govt. College Women University Sialkot University of Sargodha
Sobia Liaqat, Dr. Anam Khan, Umm Eman Syed, Neelam Bibi, Dr. Saba Ghayas, Anam Yousaf	MODERATING ROLE OF SELF-CONCEPT CLARITY BETWEEN CELEBRITY WORSHIP AND DESIRE FOR FAME AMONG YOUNG ADULTS	Rawalpindi Women University Govt. College Women University Sialkot University of Sargodha
Dr. Dhanraj T. Dhangar	THE SECULAR TADAVI BHIL TRIBE : HISTORY AND CULTURE	MGV's Arts and Commerce College, Yeola Dist.-Nashik, (M.S.) India

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-3, Hall-1
13.08.2025
Moderator: Assoc. Prof. Dr. Öykü BİLGİN
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Öykü BİLGİN	ENRICHMENT OF ASPHALTITE SAMPLE BY CENTRIFUGAL GRAVITY SEPARATION PROCESS	Şırnak University, Türkiye
Tansu ÜNEN Kenan YILDIRIM	THE AFFCET OF THE WOOL FIBER TYPE ON THE YARN TENSILE STRENGTH	Bursa Teknik University, Türkiye
Ali Can YILMAZ Özlem ERDEM YILMAZ	EURO 6 DİZEL MOTORDA 2-ETİLHEKSANOL VE TOLUEN KATKILARININ ISI SALINIMI ve PARÇACIK EMİSYONUNA SİNERJİK ETKİSİ	Çukurova University, Türkiye
Ali Can YILMAZ Özlem ERDEM YILMAZ	LASTİK BASINCININ YAKIT TÜKETİMİNE ve CO_2 EMİSYONLARINA ETKİSİ: ŞEHİR İÇİ VE ŞEHİR DIŞI SÜRÜŞ DÖNGÜLERİNDE DENEYSEL BİR İNCELEME	Çukurova University, Türkiye
Erdi TOSUN Çağrı ADIKTI	COMPARISON OF AIR COOLING CONFIGURATIONS FOR LITHIUM-ION BATTERY THERMAL MANAGEMENT	Çukurova University, Türkiye
İsmail VAROL Fatih KAHRAMAN	EFFECT OF FRICTION STIR PROCESSING ON THE MICROSTRUCTURE AND MECHANICAL PROPERTIES OF METAL MATRIX COMPOSITES	Dokuz Eylül University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-3, Hall-2
13.08.2025
Moderator: Assist. Prof. Dr. Müjgan ELVEREN
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Özcan ÇAĞLAR Harun ALPTEKİN Oğuzhan Meriç ÇAĞLAR	GLOBAL RESEARCH TRENDS IN PRECISION AGRICULTURE IN CEREALS: A BIBLIOMETRIC APPROACH	Iğdır University, Türkiye
Müjgan ELVEREN	POTENTIAL EFFECTS OF TAURINE APPLICATION ON REDUCING SALT STRESS IN BEAN PLANT (*Phaseolus vulgaris* L.)	Erzincan Binali Yıldırım University, Türkiye
Bahar ARGUN KARSLI Taki KARSLI	DETERMINATION OF THE ASSOCIATION BETWEEN THE *STAT3* GENE AND BODY WEIGHT IN ILE DE FRANCE × ANATOLIAN MERINO CROSSBRED SHEEP	Eskişehir Osmangazi University, Türkiye
Pelin YILMAZ SANCAR	MORPHOLOGICAL and PALYNOLOGICAL INVESTIGATIONS of *Polygonum setosum* subsp. *setosum* Jacq.	Fırat University, Türkiye
Muhammed Emre GÖKTAŞ Melih YILAR	POSSIBILITIES OF USING MEDICINAL AND AROMATIC PLANTS AGAINST PLANT PROTECTION AGENTS IN ORGANIC AGRICULTURE	Kırşehir Ahi Evran University, Türkiye
Mehmet Döş Mustafa PEHLİVAN Pınar YUMRUTAŞ Önder YUMRUTAŞ	CYTOTOXIC ACTIVITY AND CHARACTERIZATION OF *SALVIA CASSIA*-DERIVED SILVER NANOPARTICLES ON A549 LUNG CANCER CELLS	Gaziantep University, Türkiye
Ecem KARA Gökhan BAKTEMUR	IN VITRO MORPHOLOGICAL RESPONSES TO SALT STRESS IN CAULIFLOWER PLANT	Sivas Sciense and Technology University, Türkiye
Gökhan BAKTEMUR Ecem KARA	EVALUATION OF IN VİTRO MORPHOLOGICAL RESPONSES TO SALT STRESS IN DILL PLANT	Sivas Sciense and Technology University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-3, Hall-3
13.08.2025
Moderator: Assist. Prof. Dr. Nazlı TARAZ VARIŞLIASLAN
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Kamile Selen ÖZEN Murat TAŞ Nilüfer TAŞ	ARCHITECTURE IN ETHICAL DEADLOCK: THE ROLE OF INVISIBLE ACTORS IN DESIGN DECISION- MAKING	Bursa Uludağ University, Türkiye
Tala Laıth Talal Al-Majali	THE TEMPLE OF ARTEMIS IN GERASA: ARCHITECTURAL ANALYSIS AND COMPARISONS	Selçuk University, Türkiye
Nazlı TARAZ VARIŞLIASLAN	READING SAN CATALDO CEMETERY AS A TEXT: A SEMIOTIC INTERPRETATION OF A FUNERARY COMPLEX	Bursa Technical University, Türkiye
Derya ÇOLAK KESRİKLİOĞLU Murat ŞAHİN	EXAMINATION OF ANATOLIAN SELJUK CARAVANSERAIS USING COMPUTATIONAL METHODS	Fırat University, Türkiye
Pinar MERT CUCE Erdem CUCE Emre ALVUR	BIO-BASED RETROFIT PLASTER INCORPORATING HAZELNUT SHELLS FOR SUSTAINABLE BUILDING ENVELOPES	Recep Tayyip Erdogan University, Türkiye
Pinar MERT CUCE Erdem CUCE Emre ALVUR	THERMAL ENHANCEMENT OF BUILDING FAÇADES USING WASTE-DERIVED MATERIALS: A REVIEW OF SUSTAINABLE RETROFITTING STRATEGIES	Recep Tayyip Erdogan University, Türkiye
Pinar MERT CUCE Erdem CUCE Tamer GUCLU	HYBRID PARABOLIC TROUGH COLLECTOR AND THERMOELECTRIC GENERATOR SYSTEMS FOR MULTIGENERATION	Recep Tayyip Erdogan University, Türkiye
Pinar MERT CUCE Erdem CUCE Tamer GUCLU	SMART ENERGY SYSTEMS FOR SUSTAINABILITY-A REVIEW	Recep Tayyip Erdogan University, Türkiye
Pinar MERT CUCE Erdem CUCE Harun SEN	FROM TRADITIONAL SOLAR CHIMNEY POWER PLANTS TO NEW APPROACHES	Recep Tayyip Erdogan University, Türkiye
Pinar MERT CUCE Erdem CUCE Harun SEN	CFD METHOD OF THE EFFECT OF THE CHIMNEY DESIGN PERFORMANCE IN SOLAR CHIMNEY POWER PLANTS	Recep Tayyip Erdogan University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-3, Hall-4
13.08.2025
Moderator: Dr. Maria Taj
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Aijaz Magray, Shahid Abdullah	ZOONOTIC DISEASES – TRANSMISSION MATHEMATICAL MODELLING, AND IMPACT	JB institute of engineering and technology Hyderabad
Dr. Rüveyde GÜLBAHCE Doç. Dr. Irmak DİK Arş. Gör. Dr. H. Pelin ASLIM Prof. Dr. Sibel YAVRU	DETECTION OF PARVOVIRAL INFECTION IN DOGS WITH IMMUNOCHROMETOGRAPHIC TEST AND ELISA	Selcuk University, Faculty of Veterinary Medicine, Department of Virology, Konya, Turkey
Zohaib Saeed, Arshan Shafqat, Rustam Ali, Qazi Muhammad Awais, Muhammad Rizwan, Rao Zahid Abbas, Muhammad Abdullah Qureshi, Asghar Abbas	FLY BONE INFECTIONS AND INFESTATIONS	Multan University of Science and Technology, Multan, Pakistan. University of Agriculture, Faisalabad, Pakistan Muhammad Nawaz Sharif University of Agriculture Multan, Pakistan
Arshan Shafqat, Zohaib Saeed, Rustam Ali, Qazi Muhammad Awais, Muhammad Rizwan, Rao Zahid Abbas, Muhammad Abdullah Qureshi, Asghar Abbas	PATHOGENICITY OF TYZZER DISEASE	Multan University of Science and Technology, Multan, Pakistan. University of Agriculture, Faisalabad, Pakistan Muhammad Nawaz Sharif University of Agriculture Multan, Pakistan
Fedwa BEGHDADI, El-Hadj DRICHE	ISOLATION AND SCREENING OF ACTINOBACTERIA FROM ALGERIAN SAHARAN SOILS FOR ANTAGONISTIC ACTIVITY AGAINST PHYTOPATHOGENIC BACTERIA AND FUNGI	Genomics and Bioinformatics (LBMGB), Faculty of Natural Sciences and Life (SNV), Hassiba Benbouali University of Chlef, Hay Salem, 02000 Chlef, Algeria
Dr. Maria Taj	CLEAN AND GREEN ENVIRONMENT BY USING AGRICULTURAL WASTE WITH BETTER CLEANING PROPERTIES THAN SYNTHETIC ONES	Jinnah University, Pakistan
Muawuya Ibrahim, Gumi A.M, K. Abdullahi	THE SIGNIFICANCE OF ANTHOCYANINS IN PLANT GROWTH AND DEVELOPMENT: A REVIEW	Kebbi State University of Science and Technology Aliero Usmanu Danfodiyo University Sokoto Federal University Gusau

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Antalya

Session-3, Hall-5
13.08.2025
Moderator: Dr. Gangesh Shah Gondwana
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Dr. Gangesh Shah Gondwana	READING OF INDUS TEXTS IN PRIMEVAL KOYA LANGUAGE, PREVAILING UNIQUE SOCIAL SYSTEM AND CULTURE	Banaras Hindu University, Varanasi, Uttar Pradesh-221005, Republic of India
Hadizat Audu Salihu, Ph.D.	FISHER FOLK, IDENTITY, AND CULTURAL MEMORY: THE HISTORIES OF RIVERINE COMMUNITIES IN THE NIGER-BENUE CONFLUENCE AREA OF NIGERIA	Kwara State University, Malete, Nigeria
Omotosho ISHOLA	A CULTURAL DIPLOMACY OF ILORIN ANNUAL DURBA: AN ASSESSMENT OF ITS IMPACTS ON EXTERNAL ATTRACTIONS	Kwara State University, Malete, Nigeria
Dr James Okolie-Osemene	HOPE FOR REFORM OR CULTURE OF IMPUNITY? INSTITUTIONALIZING ETHICAL CONDUCT IN NIGERIA'S STATE SECURITY ARCHITECTURE	
Basiru Adeyemi ADEBISI	NATURE PARTICIPATION IN SCENE DESIGN IN ADEMOLA ADELAKUN'S OLUDEGUN AND PRODIGY &PENS' LUWO GBAGIDA	Kwara State University, Malete
Zaryab Touseef, Dr. Anam Khan, Umm Eman Syed, Dr Adnan Adil, Dr. Saba Ghayas, Anam Yousaf	MODERATING ROLE OF PERCEIVED SOCIAL SUPPORT BETWEEN CHILDHOOD TRAUMA AND PERSONAL GROWTH INITIATIVE AMONG ADOLESCENTS	Rawalpindi Women University Govt. College Women University Sialkot University of Sargodha
Umm e Aiman, Dr. Anam Khan, Umm e Eman Syed, Dr Adnan Adil, Dr. Sadia Niazi, Anam Yousaf	MODERATING ROLE OF REJECTION SENSITIVITY BETWEEN ROMANTIC GHOSTING EXPERIENCE AND DATE ANXIETY AMONG YOUNG ADULTS	Rawalpindi Women University Govt. College Women University Sialkot University of Sargodha
Aqsa Umer, Dr. Anam Khan, Umm e Eman Syed, Adnan Adil, Neelam Bibi, Dr. Sadia Niazi	RELATIONSHIP BETWEEN CYBERLOAFING AND WORK PERFORMANCE AMONG EMPLOYEES: MODERATING ROLE OF WORK ENVIRONMENT	Rawalpindi Women University Govt. College Women University Sialkot University of Sargodha
Ananda Majumdar	VALLABHBHAI PATEL: ARCHITECT OF INDIAN UNITY AND MODERN GOVERNANCE	Harvard University

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Antalya

Session-3, Hall-6
13.08.2025
Moderator: Dr. Taoufiq KADRI
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Dr. Taoufiq KADRI	CULTURAL DISCOVERY THROUGH LITERARY TEXTS IN FRENCH AS A FOREIGN LANGUAGE: AN INTERCULTURAL PEDAGOGICAL APPROACH	Abdelmalek Essaadi University, Morocco
Rabia Ashraf , Dr. Anam Khan, Umm e Eman Syed, Dr Adnan Adil, Dr. Saba Ghayas, Anam Yousaf	MODERATING ROLE OF MENTORSHIP EFFECTIVENESS BETWEEN PROCRASTINATION AND ACADEMIC STRESS AMONG RESEARCH STUDENTS	Rawalpindi Women University Govt. College Women University Sialkot University of Sargodha
Anisha Parveen, Dr. Anam Khan, Umm e Eman Syed, Adnan Adil, Neelam Bibi, Anam Yousaf	MEDIATING ROLE OF SELF-DOUBT BETWEEN GASLIGHTING AND EMOTIONAL DISTRESS AMONG ADULTS	Govt. College Women University Sialkot Rawalpindi Institute of Cardiology University of Sargodha
Komal Pervaz, Dr. Anam Khan, Umm e Eman Syed, Neelam Bibi, Dr. Adnan Adil, Dr. Saba Ghayas	MEDIATING ROLE OF DIGITAL IDENTITY BETWEEN SOCIAL MEDIA ADDICTION AND SELF-ESTEEM AMONG SOCIAL MEDIA USERS	Rawalpindi Women University Govt. College Women University Sialkot University of Sargodha
Irina-Ana DROBOT	KEYWORDS IN A HAIKU POEM SUGGESTING A LOVE STORY	Technical University of Civil Engineering Bucharest, Faculty of Engineering in Foreign Languages, Department of Foreign Languages and Communication, Bucharest, Romania
Abdulmumuni Fauziyat O., M.S. Maihaja	BREAKING THE BARRIERS: WOMEN IN NIGERIAN POLITICS	Kogi State Judiciary, customary curt of appeal Lokoja Kogi State Nigeria
Assoc. Prof. PhD. Vitalii ANDREIKO Senior lecturer, Anatolii MYSHKO	THE INFLUENCE OF THE ENLIGHTENMENT ON THE DEVELOPMENT OF EDUCATION IN THE USA IN THE EARLY 19TH CENTURY	Uzhhorod National University, Ukraine
Prof. PhD. Oksana PYSMENNA Assoc. Prof. PhD. Svitlana MYSHKO Assoc. Prof. PhD. Nataliya TODOROVA	"SILVER STRINGS" BY BOHDANA FILTS: MUSICAL POETRY OF UKRAINIAN IDENTITY	Mykola Lysenko Lviv National Music Academy, Lviv, Ukraine Uzhhorod National University, Ukraine
Banjo Tolani Christianah	FROM TRADITION TO PREVENTION: CULTURAL DIMENSIONS OF HYGIENE AND HANDWASHING PRACTICES IN NIGERIA	Olabisi Onabanjo University, Ago-Iwoye, Ogun State, Nigeria
Banjo Tolani Christianah	THE CULTURE OF WASTE MANAGEMENT: RETHINKING TRADITIONAL VALUES AND MODERN REALITIES IN NIGERIA	Olabisi Onabanjo University, Ago-Iwoye, Ogun State, Nigeria

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-1, Hall-1
14.08.2025
Moderator: Assist. Prof. Dr. Ayşegül GÜNGÖR ÇELIK
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Didem ALTUN	THE INVESTIGATION OF THE OPTICAL PROPERTIES OF AlGaAs/GaAs MULTIPLE QUANTUM WELLS UNDER THE INFLUENCE OF ELECTRIC AND MAGNETIC FIELDS	Sivas Cumhuriyet University, Türkiye
Mustafa UĞURLU	INVESTIGATION OF THE SECOND PASS EFFECT IN FRICTION STIR PROCESSING OF AA7075-T6 ALUMINUM ALLOY	Çukurova University, Türkiye
Hamza TAŞ	A THEORETICAL AND NUMERICAL INVESTIGATION INTO THE VIBRATION BEHAVIOUR OF HYBRID COMPOSITE PLATES CONSISTING OF CONTINUOUS CARBON FIBER-REINFORCED LAMINAS AND DISCONTINUOUS GLASS FIBER-REINFORCED LAMINAS	Manisa Celal Bayar University, Türkiye
Jale BERRAK Ali KOÇ Özkan KÖSE Yıldız KOÇ	THERMODYNAMIC PERFORMANCE EVALUATION OF THE CO_2/NH_3 MIXTURE IN SEVERAL FRACTIONS	İskenderun Technical University, Türkiye
Serhat AKBABA Zeki KIRAL	DESIGN OF A LINKAGE MECHANISM FOR USE IN INDUSTRIAL FOOD MACHINES	Dokuz Eylül University, Türkiye
Ayşegül GÜNGÖR ÇELIK	ENHANCED EXERGY PERFORMANCE ANALYSIS OF AN AIR SOURCE HEAT PUMP SYSTEM	Manisa Celal Bayar University, Türkiye
Adnan MUTLU Zeki KIRAL	VIBRATION ANALYSIS OF A 3D PRINTER CAUSED BY THE MOVEMENT OF THE PRINTING HEAD	Dokuz Eylül University, Türkiye
Osman ÇIÇEK Hasan MAKARA	COMPARATIVE ANALYSIS OF AI PROGRAMS FOR ELECTRONIC PARAMETER EXTRACTION FROM GAAS P-N DIODE I-V CHARACTERISTICS	Kastamonu University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-1, Hall-2
14.08.2025
Moderator: Prof. Dr. Nida BAYINDIR
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Təranə Nağıyeva	THE INFLUENCE OF RUSSIAN SYNTAX IN THE OFFICIAL STYLE OF THE AZERBAIJANI LANGUAGE	Azerbaijan State Pedagojical University
Şevval Erva KAYA Mine KOYUNCU ŞAHİN	SOCIAL SKILLS AND PEER RELATIONSHIPS IN PRESCHOOL: A BIBLIOMETRIC ANALYSIS	Afyon Kocatepe University, Türkiye
Ramazan Rüçhan KAYA Şevval Erva KAYA	ILLNESS AND HOSPITAL IN THE PRESCHOOL PERIOD: AN EVALUATION BASED ON CHILDREN'S VERBAL AND VISUAL EXPRESSIONS	Uşak University, Türkiye Afyon Kocatepe University, Türkiye
Ceyhune DEMIROVA MAHMUDOVA	COMMUNICATION SKILLS OF FOREIGN LANGUAGE TEACHERS: ANALYSIS OF SOCIAL INTELLIGENCE IN THE CONTEXT OF UNIVERSITY INSTRUCTORS	Azerbaijan Languages University
Mebrure ERDOĞAN Kübra YILMAZ	BIBLIOMETRIC ANALYSIS OF SCIENTIFIC PUBLICATION TRENDS IN THE FIELD OF TISSUE ENGINEERING AND BIOMATERIALS	Kastamonu University, Türkiye
Nida BAYINDIR Mehmet GÖKCE	PRESERVICE TEACHERS' PERCEPTIONS OF THE EDUCATIONAL BENEFITS AND SAFETY RISKS OF YOUTUBE IN PRIMARY EDUCATION	Eskişehir Osmangazi University, Türkiye Kütahya Health Sciences University, Türkiye
Halil İbrahim AKPINAR	SUSTAINABLE BANKING PRACTICES: THE IMPACT OF GREEN CREDIT POLICIES ON THE REAL ECONOMY	İstanbul, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-1, Hall-3
14.08.2025
Moderator: Assist. Prof. Dr. Mehmet DUGAN
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Ercan BARAN	BADI' AL-ZAMAN AL-HAMADHANI AND HIS MAQAMAT (THE MAQAMA OF ADVICE, AL-ASWAD, IRAQ, HAMDAN, AND RUSAFA)	Bingöl University, Türkiye
Ercan BARAN	BADI' AL-ZAMAN AL-HAMADHANI AND HIS MAQAMAT (THE MAQAMA OF MOSUL, MADIRA, THE AMULET, THE ASYLUM, AND HUNGER)	Bingöl University, Türkiye
Sinan YILMAZ	ARABIC-TURKISH AND TURKISH-ARABIC DICTIONARY STUDIES CONDUCTED IN TURKEY	Trabzon University, Türkiye
Zeynep BAŞARAN	PEYGAMBERİMİZİN MUCİZELERİNDEN OLAN NESEBİ DURUMU	Batman University, Türkiye
Mehmet DUGAN	DIGITAL STUDIES AND METHODS ON AVERROES: AN APPLICATION BASED ON THE COMMENTARY ON THE POSTERIOR ANALYTICS	Akdeniz University, Türkiye
Sultanmurat ABZHALOV Kanat BAZARBAYEV Batyrzhan TEMİRKHANOV	THE SIGNIFICANCE OF KHOJA AHMED YASAWI IN THE TURKIC WORLD: THE CASE OF THE KAZAKH NEWSPAPER	Khoja Ahmet Yesevi International Turkish-Kazakh University, Kazakhstan Ankara Hacı Bayram Veli University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-1, Hall-4
14.08.2025
Moderator: Assist. Prof. Dr. Yener ULUS
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Yener ULUS	SPATIAL VARIATION OF HEAVY METALS ALONG A SALINITY GRADIENT IN WINYAH BAY, SOUTH CAROLINA	Auburn University at Montgomery, College of Sciences, Biology and Environmental Science Department, Alabama, USA.
Dr. K. VIGNESH ARSHA G	ROLE OF EFFECTOR PROTEINS IN PATHOGEN VIRULENCE AND PLANT IMMUNE RESPONSES	Palar Agricultural College, Ambur
Dr. K. VIGNESH ARSHA G	FUNCTIONAL GENOMICS OF Fusarium spp. CAUSING WILT DISEASES	Palar Agricultural College, Ambur
Md. Selim Ali	PREVALENCE AND DETECTION OF BACILLUS CEREUS, STAPHYLOCOCCUS AUREUS, ESCHERICHIA COLI, AND TRICHODERMA ON UNWASHED FRUITS SOLD AT ROADSIDE STALLS IN VODRA, RAJSHAHI	University of Rajshahi, Bangladesh
Muhammad Saad Tariq, Wafa Majeed, Ayesha	ROLE OF MORIN IN ATTENUATION OF METABOLIC GLUCOSE AND INSULIN SECRETION IMPAIRMENT DURING HYPERGLYCEMIA	University of Agriculture, Faisalabad, Pakistan
Saira Malik, Syed Mohsan Raza Shah, Uniza Fatima bukhari, Tasawar Fatima bukhari, Iqra	LEAF STRUCTURAL AND FUNCTIONAL MODIFICATION OF IPOMOEA CARNEA JACQ. AN INVASIVE PLANT SPECIES, UNDER DIVERSE SALINITY GRADIENT	University of Education, Lahore, Pakistan
Faisal Nazir	PRODUCTION AND CHARACTERIZATION OF CARBON NANOTUBES FROM BIOCHAR UNDER MICROWAVE IRRADIATION	The University Of Agriculture, Pakistan
Sunnia Afzal, Zara Saeed, Farukh Ameen	FOLIAR APPLICATION OF CHOLINE CHLORIDE AND FERULIC ACID REGULATED OSMOLYTE ACCUMULATION, NUTRIENT ACQUISITION AND ROS METABOLISM TO LESSEN CHROMIUM EFFECTS ON BARLEY (HORDEUM VULGARE)	Government College University, Faisalabad-38000, Pakistan

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-1, Hall-5
14.08.2025
Moderator: H. Nasser
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
H. Nasser, S. Elhadfi, M. Fattah, H. Chadli	FIRST-PRINCIPLES STUDY OF HYDROGEN ADSORPTION ON LITHIUM DOPED CARBON NANOTUBES	Moulay Ismail University, FSM-ESTM, BP 11201, Zitoune, Meknes 50000, Morocco
Abdelillah Taoufik, Hafida Ziouani, Sana Mazouar, El Mostafa Khechoubi, Mahmoud Ettakni	STRUCTURAL AND POLYMORPHIC ANALYSIS OF $[R\text{-}NH_3]_2MCL_4$ BASED MOLECULAR COMPOSITES FOR ADVANCED ENERGY STORAGE	Moulay Ismail University, Meknes, Morocco.
Abdelilah MECHNINE Prof.Ahmed AARAB	A COMPUTATIONAL APPROACH TO PROTEIN ANALYSIS: MODELING, DOCKING, AND MOLECULAR DYNAMICS SIMULATIONS	Abdelmalek Essaadi University, Faculty of Sciences and Techniques of Tangier, Morocco
Owoyemi S.I, Ojo M.O, Popoola F.A., Sanyaolu M.E	CORRELATING THE POTENTIAL INVESTIGATION AND ANALYSIS OF THE IMPACTS OF WATER EFFECTS ON STRUCTURAL FOUNDATION STUDY AND STRUCTURAL FOUNDATION TECHNIQUES OF IGBOKODA AND IGBEKEBO AND ITS ENVIRONS, ONDO SOUTHWESTERN NIGERIA	Adeyemi Federal University of Education Ondo, Nigeria. Glorious Vision University Ogwa, Nigeria Redeemer's University, Ede, Osun State, Nigeria.
Ode Jonathan Acha Dr. Donald Dehiin Hile Agada Felix Ojochegbe	ESTIMATION OF DAILY AND GLOBAL EXTRATERRESTRIAL SOLAR RADIATION AT MAKURDI BENUE STATE	Federal University of Agriculture Makurdi, Benue State. Nigeria
Mohamed Yassine El Brouzi, Wissal Baghdad, Mouloud Lamtai, Abedelhalim Mesfioui	NICKEL NEUROTOXICITY: INVESTIGATION INTO HIPPOCAMPAL LATERALIZATION ON MEMORY PROCESS, ACHE ACTIVITY, NEUROINFLAMMATION AND OXIDATIVE STRESS MARKERS IN WISTAR RATS	Ibn Tofail University, Kenitra, Morocco Hassan II Institute of Agronomy and Veterinary Medicine, Rabat, Morocco Mohammed V Military Instruction Hospital, Rabat, Morocco
Abderrahim Ait Lhaj, Brahim Ouazik, Mohamed El Hasnaoui, Hassan Chaib	DFT STUDY OF THE STRUCTURAL, ELECTRONIC, AND OPTICAL PROPERTIES OF TETRAGONAL SnO2 USING DFT WITH THE BLYP FUNCTIONAL	Ibn Tofail University, Morocco. Ibnou Zohr University, Morocco.

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Antalya

Session-1, Hall-6
14.08.2025
Moderator: Dr.S.Rajeswari
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Noureddine EL OUASTI Adnane AOUIDATE Abdellah OUSAA Moustapha BELMOUDEN	USING CHEMOINFORMATICS AND MACHINE LEARNING TO EXPLORE THE CHEMICAL SPACE OF HERG INHIBITORS	University Ibn Zohr, Morocco
Amadi Oko Amadi, Levi Nnaemeka Osuji	EFFECT OF QUANTUM ANNEALING AND D-WAVE ON THE OPTIMIZATION OF ENHANCED OIL RECOVERY AND PRODUCTION FRAMEWORK	Akanu Ibiam Federal Polytechnic Unwana-Ebonyi state
Amadi Oko Amadi, Levi Nnaemeka Osuji	EFFECT OF QUANTUM ANNEALING AND D-WAVE ON THE OPTIMIZATION OF ENHANCED OIL RECOVERY AND PRODUCTION FRAMEWORK	Akanu Ibiam Federal Polytechnic Unwana-Ebonyi state.
Puniparthi Sunitha M.K. Vijiayalakshmi	REAL-TIME THERAPEUTIC DRUG MONITORING USING WEARABLE DEVICES	Bharath institute of higher education and research
Puniparthi Sunitha M.K. Vijiayalakshmi	THE ROLE OF GUT MICROBIOTA IN CHRONIC DISEASES	Bharath institute of higher education and research
Puniparthi Sunitha M.K. Vijiayalakshmi	3D BIOPRINTING IN ORGAN TRANSPLANTATION: HOPE OR HYPE?	Bharath institute of higher education and research
Rahim Mammadzada	SIMULATION BASED ASSESSMENT OF COOLING FAN ENERGY TRADE-OFFS IN PHOTOVOLTAIC SYSTEMS USING MATLAB	Azerbaijan State Oil and Industry University, Azerbaijan
Rahim Mammadzada	SIMULATION BASED MODELING OF THERMAL AND ELECTRICAL BEHAVIOR IN PHOTOVOLTAIC PANELS USING MATLAB	Azerbaijan State Oil and Industry University, Azerbaijan
Dr.S.Rajeswari	CHANNEL OPTIMIZATION IN WIRELESS MESH NETWORKS USING SUPERVISED RANDOM FOREST ALGORITHM	Saranathan College of Engineering, India

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-2, Hall-1
14.08.2025
Moderator: Assoc. Prof. Dr. Songül KESEN
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
İbrahim YAVUZ Hatice Gözde HOSTA YAVUZ	BIOACTIVE COMPONENTS OF ROYAL JELLY AS A FUNCTIONAL FOOD AND ITS EFFECTS ON FERTILITY	Akdeniz University, Türkiye
Songül KESEN	NEW TEACHING METHODS IN FOOD ENGINEERİNG AND AR/VR APPLICATIONS	Gaziantep University, Türkiye
Songül KESEN	MINOR AND MAJOR COMPONENTS IN OLIVE OIL AND THEIR FUNCTIONS	Gaziantep University, Türkiye
Neslihan Çetin Nisanur Kaynak Can Ali Süzer Sahranaz Kılıç Tuğçe Özlü Karahan	A STUDY ON THE RELATIONSHIP BETWEEN CHRONOTYPE CHARACTERISTICS, ENERGY AVAILABILITY, AND SLEEP BEHAVIOR IN TEAM ATHLETES	İstanbul Bilgi University, Türkiye
Bumin Çağatay AKSU Turgay KARALİNÇ	THE MEDIATING ROLE OF PROFESSIONAL SELF-EFFICACY İN THE EFFECT OF SUSTAINABLE QUALITY PERCEPTIONS OF HEALTHCARE WORKERS ON THEIR SUBJECTIVE WELL-BEING	Istanbul Gelişim University, Türkiye
Zeynep APAYDIN DEMİRCİ	MİNDFUL PARENTING AND DIGITAL TOOL USE IN EARLY CHILDHOOD	Bilecik Şeyh Edebali University, Türkiye
Qaribova Arifa Hajibaba kızı	THE ROLE OF UZEYIR HAJIBAYLI IN THE ESTABLISHMENT OF THE PROFESSIONAL MUSIC EDUCATION SYSTEM IN AZERBAIJAN	Azerbaijan State Pedagogical University

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-2, Hall-2
14.08.2025
Moderator: Assist. Prof. Dr. Mehmet Emin GÖNEN
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Yağmur KARAMAN Bahar SARIBOĞA AKCA	EXAMINATION OF THE FEATURES OF FLUTE FAMILY INSTRUMENTS AND FLUTE HEADJOINTS	Ordu University, Türkiye
Vecihe Özge ZEREN	THE AURA OF DRAMATURGY: FROM BENJAMIN TO ARTIFICIAL INTELLIGENCE – THEATRICAL AUTHENTICITY IN THE AGE OF TECHNICAL REPRODUCTION	Çanakkale Onsekiz Mart University, Türkiye
Meriç DÜZBAŞ	AN ANALYSIS OF THE DEFINITION, STRUCTURE, AND USAGE OF THE DARBEYN USUL IN TURKISH MUSIC	Anadolu University, Türkiye
Lamiye NEBİYEVA	THE SYSTEM OF IMAGES IN ARTISTIC LANGUAGE AND THE STYLIST-ARTISTIC FUNCTION (BASED ON THE CREATIVITY OF NAKHCHIVAN WRITERS)	Naxçıvan State University
Ferhat ACAR	MUSTAFA KEMAL ATATÜRK'S BLACK SEA VISITS (1919-1937)	Kafkas University, Türkiye
Yunus BOR	THE ADMINISTRATION OF THE OTTOMAN PUBLIC DEBT (DÜYUN-U UMUMIYE) IN THE OTTOMAN EMPIRE	Ordu University, Türkiye
Gülnar Hüseynova	20-ci YÜZYIL AZERBAYCAN EDEBİYATI- GENCE EDEBİ ORTAMINDA ŞEHİTLİK KONUSU (ŞEHİT ŞAİR NİZAMİ AYDIN`IN ŞİİRLERİ TEMELİNDE)	Gence State University
Mehmet Emin GÖNEN	THE PSYCHO-SOCIAL REPRESENTATION OF FEMALE CHARACTERS: A COMPARATIVE ANALYSIS THROUGH KUMRU, KEZBAN, AND HATÇE	Yozgat Bozok University, Türkiye
Nəzakət Əbilova	HISTORICAL FORTIFICATIONS OF THE ASTARA PROVINCE OF AZERBAIJAN	Lənkəran State University

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-3
14.08.2025
Moderator: Merve KÜÇÜK
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Ersin YAVUZ Tahsin AVCI	REGIONAL ANALYSIS OF SOCIO-ECONOMIC AND FISCAL PERFORMANCE: AN IMPLEMENTATION ON TÜRKİYE NUTS LEVEL-2	Pamukkale University, Türkiye
Ersin YAVUZ Tahsin AVCI	EXAMINING THE ECONOMIC PERFORMANCE OF OECD COUNTRIES FROM A TAX BURDEN PERSPECTIVE WITH THE PIV METHOD	Pamukkale University, Türkiye
Ekrem YILMAZ Savaş DEMİR	DEPREM ERKEN UYARISININ DAHA KISA ZAMANDA BİLDİRİMİNİN SAĞLANMASI	Multitek Elektronik Research and Development Centre, Istanbul, Turkıye
Ekrem YILMAZ Savaş DEMİR	DEPREM ANINDA KONUM BİLGİLERİNİN TOPLANMASI	Multitek Elektronik Research and Development Centre, Istanbul, Turkıye
Erdal AYDIN	THE EFFECT OF CHILDHOOD SOCIO-ECONOMIC STATUS ON LATER LIFE SATISFACTION	Sabancı University, Türkiye
Barış IŞILDAK Murat Kemal KELEŞ Aşkın ÖZDAĞOĞLU	HELICOPTER SELECTION FOR AERIAL WILDFIRE INTERVENTION: AN ANALYSIS USING LOPCOW, MAUT, WEDBA, MABAC, AND PSI METHODS	Isparta University of Applied Sciences, Türkiye
Murat Kemal KELEŞ Barış IŞILDAK Hasan Hüseyin YİĞİT Aşkın ÖZDAĞOĞLU	COMBINING DIFFERENT MCDM METHODS WITH THE COPELAND METHOD: AN APPLICATION IN AIRLINE SECTOR	Isparta University of Applied Sciences, Türkiye Süleyman Demirel University, Türkiye Dokuz Eylül Ü University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-4
14.08.2025
Moderator: ODU Adejare Samuel
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Sadikhova F.Q.	THE ISSUE OF SOUL AND BODY IN EUROPEAN PHILOSOPHY OF THE NEW AND MODERN ERA	Institute of Philosophy and Sociology of ANAS
Peter Olubunmi Olubiyo Jude Timilehin Olubiyo	EXTENT OF USE OF DIGITAL RESOURCES AMONG PHYSICS AND CHEMISTRY FINAL YEAR UNDERGRADUATE STUDENTS IN ADEYEMI FEDERAL UNIVERSITY OF EDUCATION, ONDO, NIGERIA	University Library, Adeyemi Federal University of Education, Ondo, Nigeria
CHRISTIAN SAMUEL JIMWAN ALIYU ADAMU MULUKU	VALUES REORIENTATION OF SECONDARY SCHOOL STUDENTS THROUGH BIOLOGY EDUCATION FOR SUSTAINABLE DEVELOPMENT IN JOS NORTH, PLATEAU STATE, NIGERIA	University of Jos, Plateau State, Nigeria College of Education Akwanga
Faiza Shams , Dr. Anam Khan, Mam Umme Aiman Syed, Adnan Adil, Neelam Bibi, Dr. Sadia Niazi	RELATIONSHIP BETWEEN BREADCRUMBING AND RELATIONSHIP SATISFACTION AMONG MARRIED INDIVIDUALS: MEDIATING ROLE OF EMOTIONAL ABUSE	Govt. College Women University Sialkot Rawalpindi Institute of Cardiology University of Sargodha
Najma Bi Bi, Dr. Anam Khan, Umm e Eman Syed, Neelam Bibi, Dr. Adnan Adil, Dr. Sadia Niazi	IMPACT OF MINDFULNESS AND SELF- COMPASSION ON PERCEIVED STRESS AMONG MEDICAL PROFESSIONALS	Rawalpindi Institute of Cardiology Govt. College Women University Sialkot University of Sargodha
Quratulain Ashraf, Dr. Anam Khan, Umm e Eman Syed, Neelam Bibi4, Dr. Adnan Adil, Dr. Saba Ghayas	IMPACT OF PARENTAL INVOLVEMENT ON ACADEMIC PERFORMANCE AND SELF EFFICACY AMONG SECONDARY SCHOOL CHILDREN	Rawalpindi Institute of Cardiology Govt. College Women University Sialkot University of Sargodha
ODU Adejare Samuel	TEACHERS' AND STUDENTS' OPINIONS ON LANGUAGE PREFERENCE FOR WRITING AND CONVERSATIONS AMONG YORUBA LANGUAGE SPEAKERS IN NIGERIA	The Federal Polytechnic Ilaro, Ogun State, Nigeria

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-5
14.08.2025
Moderator: Dr Irfan Bashir
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Sadhna Jain, Mamta Sharma, Baldev Gulati	AN EXPLORATORY STUDY ON UNDERGRADUATE LEARNERS' PREPAREDNESS AND PREFERENCES FOR DIFFERENT MODES OF LEARNING IN HIGHER EDUCATION	University of Delhi, Delhi
Fouad Boughanzai, Rochdi Daibi, Anouar Boughanzai, Youssef Ouhassan, Rania Majdoubi, Abdelkader Hadjoudja	CHALLENGES AND CONCERNS ABOUT AI CHATBOT USE IN EDUCATION: A QUALITATIVE ANALYSIS OF TEACHERS' PERSPECTIVES	Ibn Tofail University, Kénitra, Morocco
Glago Frank Jerome Agbenyefia Gladys Mawufemor	DISASTER RISK AWARENESS AMONG COLLEGE STUDENTS: A STUDY OF AKATSI COLLEGE OF EDUCATION.	SDA College of Education, P. O. Box AS 18, Asokore-Koforidua
Dr Irfan Bashir Ms. Fareeha Sherazi Dr Afshan Naseem Dr Yasira Waqar Dr Amna Arif Dr Fariha Gul Dr Almas Shoiab Dr Aisha Sami	PROFESSIONAL DEVELOPMENT OF FACULTY IN HIGHER EDUCATION INSTITUTES OF PAKISTAN: ROLE OF DEPARTMENTAL LEADERSHIP	Dept of Education, University of Management and Technology, Pakistan
Assist. Prof. Dr. Aktolkyn BORANBAYEVA Prof. Dr. Gelişli Yücel Prof. Dr. Assel TASSOVA	FORMATION OF THE FUTURE SPECIALIST'S CREATIVITY	Uzbekali ZhanibekovUniversity, Kazakhstan Gazi University, Türkiye Khoja Akhmet Yassawi International Kazakh-Turkish University, Kazakhstan
Mr. Sunday Samuel OLANREWAJU	GLOBALIZATION, DIGITAL INCLUSION AND REHABILITATION: EXPLORING THE ROLE OF ICT IN SUPPORTING PERSONS WITH SPECIAL NEEDS	Federal College of Education (Special) Oyo, Nigeria

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-2, Hall-6
14.08.2025
Moderator: Dr. G. I. Onwuka
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Abdelhamid Ait M'hid, Guojian Li, Mourad Boughrara, Mohamed Kerouad, Qiang Wang	STRUCTURAL, MAGNETIC, AND ELECTRICAL PROPERTY MODULATION IN ZnO and V- and Cr-doped ZnO THIN FILMS VIA DEFECT ENGINEERING	Northeastern University, Shenyang, 110819, China Moulay Ismail University, Meknes, Morocco
Imane Haji, Loubna Rachidi, Hind Chakchak, Abdelkader Zarrouk, Ghizlan Kaichouh	HETEROGENEOUS ELECTRO-FENTON PROCESS: PERFORMANCE OF A NATURAL BIOWASTE CATALYST FOR THE DEGRADATION OF THE ANTIBIOTIC CFX-Na	Mohammed V University, Rabat, Morocco (CNRST-UATRS), Rabat, Morocco.
Dr. G. I. Onwuka, Prof. Babayemi A. W., B. Sheh	TWO SAMPLE MULTIVARIATE BEHRENS FISHER PROBLEM FOR HIGH DIMENSIONAL DATA	Kebbi State University of Science and Technology Kebbi State Polytechnic Dakingari
Abdallah Lalmi, Badreddine Boumali, Elafri Nedjwa, Hichem Chorfi, Ismail Benteldjoune	LEVERAGING ARTIFICIAL INTELLIGENCE FOR QUALITY CONTROL IN CONSTRUCTION: OPPORTUNITIES, CHALLENGES, AND FUTURE DIRECTIONS	University of Constantine University of Boumerdes
Ikram Mennas, Ahmed Arif, Mohamed Kadour Atouailaa, Abdelkhalk Aboulouard, Mustapha Boulghalat, Mohammed El idrissi	TD-DFT INVESTIGATION OF QUINOXALINE-BASED ACCEPTOR MATERIALS FOR HIGH-PERFORMANCE ORGANIC SOLAR CELLS	Sultan Moulay Slimane University, Beni-Mellal, Morocco Higher School of Education and Training, Beni-Mellal, Morocco National School of Applied Sciences, El Jadida, Morocco
Ahmed Osmane, Khadija Zidan, Moustapha Belmouden	SORGHUM PRODUCTION USING TRICKLING FILTER EFFLUENT: ASSESSING HEALTH AND AGRONOMIC IMPACTS IN SEMI-ARID NORTH AFRICA	Agadir University Ibn Zohr, Morocco University Cadi Ayyad, Marrakech, Morocco
Ahmed Osmane, Khadija Zidan, Moustapha Belmouden	ASSESSING MAIZE HEALTH BENEFITS IN SEMI-ARID REGIONS: IMPACT OF TRICKLING FILTER-TREATED WASTEWATER IRRIGATION	Agadir University Ibn Zohr, Morocco University Cadi Ayyad, Marrakech, Morocco

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-3, Hall-1
14.08.2025
Moderator: Assist. Prof. Dr. Arzu ÖZTÜRK KESEBİR
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Safak YIGIT Handan TANYILDIZI KOKKULUNK	DUAL ASSESSMENT OF SARCOPENIA FOR LUNG CANCER CLASSIFICATION: FUNCTIONAL AND CT-BASED APPROACHES	Istanbul Galata University, Türkiye Altinbas University, Türkiye
Muammer AVCI	CARDIORENAL SYNDROME AFTER DENASUMAB ADMINISTRATION	Isparta Şehir Hastanesi Nefroloj Kliniği
Samed ŞİMŞEK Tuğçe VAROL	LC–MS/MS PROFILING OF PHENOLIC CONSTITUENTS AND IN VITRO ASSESSMENT OF ANTIOXIDANT AND ANTICANCER ACTIVITIES OF HEDYSARUM YILMAZUNALII	Çayırlı Meslek Yüksekokulu, Türkiye Erzincan Binali Yıldırım University, Türkiye
Harun Yıldırım Murtaza Kaya	A RARE CASE OF KOUNIS SYNDROME TYPE II PRESENTING WITH ST-ELEVATION MYOCARDIAL INFARCTION TRIGGERED BY BEE STING	Kütahya Health Sciences University, Türkiye
Elif İrem AKDOĞAN Eftal ŞEHİRLİ	DETECTION AND CLASSIFICATION OF WHITE BLOOD CELLS – A REVIEW	Karabük University, Türkiye
Hilal Kübra GÜÇLÜ EKİNCİ Dilanur GÖKDAĞ	STATURE ESTIMATION BASED ON CALCANEUS LENGTH IN ARCHAEOLOGICAL SKELETONS: THE CASE OF EARLY BYZANTINE PERIOD IN KAYSERİ	Erciyes University, Türkiye Nuh Naci Yazgan University, Türkiye
Arzu ÖZTÜRK KESEBİR	BREAKING CANCER RESISTANCE WITH 6PGD INHIBITION	Ağrı İbrahim Çeçen University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-3, Hall-2
14.08.2025
Moderator: Assist. Prof. Dr. Abdullah Huzeyfe AKCA
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Erdal KESGİN	SWMM-BASED MODELING OF SUSTAINABLE URBAN DRAINAGE PRACTICES ON A UNIVERSITY CAMPUS	Istanbul Technical University, Türkiye
R. İlayda TAN KESGIN	MICROPLASTIC TRANSPORT THROUGH SEDIMENT IN RIVER ENVIRONMENTS	Fatih Sultan Mehmet Vakıf University, Türkiye
İbrahim ÖZ	DONATISIZ YIĞMA YAPILARIN DEPREM PERFORMANSININ SAYISAL YÖNTEMLERLE DEĞERLENDİRİLMESİ	Kırşehir Ahi Evran Ü University, Türkiye
Abdullah Huzeyfe AKCA	ILLUSTRATION OF UPDATES TO THE CONDENSATION CALCULATIONS OF THE THERMAL INSULATION RULES IN BUILDINGS WITH EXAMPLES OF DIFFERENT THERMAL INSULATION DESIGNS	Yıldız Technical University, Türkiye
Enis COŞKUN Onur ŞAHIN Abdullah Huzeyfe AKCA	THE EFFECTS OF CARBONATION ON THE PARTICLE STRENGTH OF MAGNESIUM BASED ARTIFICAL COARSE AGGREAGATE	Yıldız Technical University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-3, Hall-3
14.08.2025
Moderator: Assist. Prof. Dr. Cansu VATANSEVER
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Cansu VATANSEVER	BACTERIAL BIOFILM INFECTIONS AND THERAPEUTIC APPROACHES	Altınbaş University, Türkiye
Fatıma Ulya Aksu Songül Tezcan	EVALUATION OF MEDICATION ADHERENCE IN PATIENTS WITH COMORBIDITIES IN THE CARDIOLOGY CLINIC	Health Sciences University, Türkiye Marmara University, Türkiye
Perçin PAZARCI	CO-MUTATION PROFILE OF PTEN IN GLIOBLASTOMA	Çukurova University, Türkiye
Raviye Özen Koca	AGOMELATINE AND PHYSIOLOGICAL SYSTEMS: AN INTEGRATIVE APPROACH TO CORE MECHANISMS	Necmettin Erbakan University, Türkiye
Zeynep GÜMRÜKÇÜ	ALVEOLAR DEFECT RECONSTRUCTION WITH TITANIUM MESH AND SIMULTANEOUS DENTAL IMPLANT PLACEMENT: A CASE REPORT	Recep Tayyip Erdoğan University, Türkiye
Zeynep GÜMRÜKÇÜ	EVALUATION OF THE BIOMECHANICAL EFFECTS OF QUAD ZYGOMATIC IMPLANT APPLICATIONS IN DIFFERENT MAXILLARY DEFECT TYPES BY FINITE ELEMENT ANALYSIS	Recep Tayyip Erdoğan University, Türkiye
Gamze Akca İmge Madak Şükran Bilge Tanrıseven Merve Gülsen Bal Albayrak Tuğcan Korak	DAIDZEIN INDUCES APOPTOTIC CELL DEATH IN SH-SY5Y NEUROBLASTOMA CELLS	Kocaeli University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-3, Hall-4
14.08.2025
Moderator: Dr.R. Devi
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Robert Karlo Klara Karlo Antonela Karlo Tamara Karlo	GEOGRAPHICAL INEQUALITIES IN ACCESS TO HEALTHCARE: A COMPARISON OF ELDERLY EXPERIENCES IN URBAN AND RURAL AREAS	General Hospital Zadar, Croatia BG Unfallklinik Murnau, Germany Dental Clinic, Zadar, Croatia Dental Clinic, Zadar, Croatia
Mrs.E.Elavarasi,Yuvaraj AR,J.Jayadurka,Dr.R.Srinivasan	DRUG FORMULATION AND DELIVERY SYSTEMS	Bharath Institute of Higher Education and Research,Chennai-73
Mrs.E.Elavarasi,Yuvaraj AR, J.Jayadurka ,Dr.R.Srinivasan	THE ROLE OF CLINICAL PHARMACISTS IN HOSPITAL SETTINGS	Bharath Institute of Higher Education and Research,Chennai-73
Mohd Nadeem, Rahmuddin khan Abul Kalam Najmi	DEVELOPMENT AND OPTIMIZATION OF PIRACETAM AND SHATAVARIN IV LOADED NANOEMULSION FOR ALZHEIMER'S DISEASE THERAPY: IN-SILICO AND EXPERIMENTAL ANALYSIS	School of Pharmaceutical Education and Research, Jamia Hamdard, New Delhi
Dr.R. Devi Dr. V. Dhanapal	TRANSCRANIAL ULTRASOUND COMBINED WITH LIPID NANOPARTICLES ADVANCES BRAIN DELIVERY OF SOD1 ANTISENSE FOR ALS TREATMENT	Sree Sastha Pharmacy College, Tamil Nadu, India
Dr.R. Devi	TARGETED NANOTECHNOLOGY APPROACHES TO BYPASS THE BLOOD–BRAIN BARRIER IN NEURODEGENERATIVE DISORDERS	Sree Sastha Pharmacy College, Tamil Nadu, India
M. Maiss, Y. Ighris, Y. Bouhouchi, M. Alaoui, M. Hssikou, Y. Sadiki, J. Baliti	MESOSCOPIC SIMULATION OF NATURAL CONVECTION IN A CAVITY WITH SINUSOIDAL HEATING USING THE LATTICE BOLTZMANN METHOD	Sultan Moulay Slimane University, Beni Mellal, Morrocco

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-3, Hall-5
14.08.2025
Moderator: Prof. Mariana BĂLAN
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Mohd Zeeshan Wajahat Ali	SUSTAINABLE SMART TOURISM: BALANCING INNOVATION AND RESPONSIBILITY IN THE DIGITAL AGE	Aligarh Muslim University Marwadi University, Rajkot, Gujarat
Prof. Mariana BĂLAN	THE IMPACT OF THE COVID-19 CRISIS ON EMPLOYMENT IN THE CULTURAL AND CREATIVE SECTORS	Lecturer, Scientific Researcher III, Anca CRISTEA, PhD
Professor SADIQ Mohammed Sanusi	RE-IMAGINING AGRICULTURAL POLICY THROUGH BAGONOMICS: TOWARDS INCLUSIVE AND EQUITABLE RURAL DEVELOPMENT IN NIGER STATE	Federal University Dutse, Jigawa State
Olumide Abraham Ajayi (Ph.D.)	IMPACT OF TERRORISM AND BANDITRY ON WOMEN DEVELOPMENT IN NORTHERN NIGERIA	Skyline University Nigeria, Kano
AJIBADE Olalekan Eyitayo OGUNDARE Daniel Olugbenga	EMPLOYEE RETENTION STRATEGIES ON EMPLOYEE TURNOVER IN NIGERIA PUBLIC INSTITUTIONS IN NIGERIA: EVIDENCE FROM THE FEDERAL POLYTECHNIC, ILARO, NIGERIA	Federal Polytechnic, Ilaro, Ogun State, Nigeria
Adesanmi Moses Ademola	PEACE AND THE POLITICS OF EXCLUSION: A PRAGMATIC READING OF EXISTENCE IN NOT MY BUSINESS AND THEY TOO ARE THE EARTH	Federal Polytechnic, Ilaro, Ogun State, Nigeria
Olufemi Timothy Ogunbode PhD. Charles Adedeji Ogidan Saheed Balogun Amusa PhD.	SUSTAINABLE DEVELOPMENT GOALS (SDGS) AND BAPTIST MISSION'S HEALTH FINANCING (HF) MODELS IN YORUBALAND, 1907 - 1986	Ladoke Akintola University of Technology, Ogbomoso, Oyo State Obafemi Awolowo University, Ile-Ife, Osun State

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-3, Hall-6
14.08.2025
Moderator: Taskeen Fatima
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Taskeen Fatima Adnan Sarwar	THE CLINICAL SIGNIFICANCE OF NORMAL PHYSIOLOGICAL BLOOD PARAMETERS IN HEALTH ASSESSMENT	Hajvery University
Taskeen Fatima Mahna Fatima Adnan Sarwar	AN OVERVIEW OF CAD; ETIOLOGICAL FACTORS & SURGICAL MANAGEMENTS	Hajvery University
Md. Rostom Ali Dr. Md. Fakrul Islam Md. Ariful Islam Mst. Surovi Akther	SOCIAL CAPITAL FOR DEALING WITH SOCIO-ENVIRONMENTAL AND BEHAVIORAL IMPACTS OF COVID-19 PANDEMIC ON MIGRANT WORKERS' FAMILY IN NORTHERN BANGLADESH: PREPAREDNESS, RESPONSES, AND RECOVERY MEASURES	University of Rajshahi, Bangladesh Debi Chowdhurani Degree College, Bangladesh
K.R.Padma K.R.Don	HOW TO BALANCE GLOBAL CRISIS MANAGEMENT DURING PANDEMIC	SriPadmavatiMahilaVisvaVidyalayam (Women's University), Tirupati,AP. Bharath University, Chennai, Tamil Nadu, India
Fatima Zahra BOUCHACHI Maryem ARRAJI Mohamed CHAHBOUNE	FACTORS OF CHRONIC PAIN AND PAIN INTERFERENCE IN PATIENTS TREATED BY HEMODIALYSIS: A CROSS-SECTIONAL STUDY	Higher Institute of Health Sciences, Hassan First University of Settat, Morocco
Sodiq Adetunji Adekola, Saheed Ayodeji Adekola, Abdulmalik Abdulkareem Omeiza	EPIDEMIOLOGICAL PROFILE AND SCREENING OUTCOMES OF PARTICIPANTS IN A UNIVERSITY-BASED MEDICAL OUTREACH PROGRAMME (A Case Study of NIMELSSA Health Week 2025, University of Abuja)	University of Abuja, Abuja
Sodiq Adetunji Adekola, Saheed Ayodeji Adekola, Dhikroh Oriyomi Adekola	EVALUATING THE IMPACT OF OCCUPATIONAL STRESS ON MENTAL HEALTH AND JOB PERFORMANCE AMONG HEALTHCARE WORKERS AT THE UNIVERSITY OF ABUJA TEACHING HOSPITAL, GWAGWALADA	University of Abuja, Abuja

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-1, Hall-1
15.08.2025
Moderator: Assoc. Prof. Dr. Suzan CANGÜL
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Sema Nur MUTLU	TECHNICAL ANALYSIS OF UNDERGROUND MINE ACCIDENTS IN TURKEY FROM 1980 TO THE PRESENT	Hitit University, Türkiye
Turgut Ergin Kadir Gök Akil Birkan Selçuk	BIBLIOMETRIC ANALYSIS OF RESEARCH ON OCCUPATIONAL ACCIDENTS; WOS EXAMPLE	Izmir Bakırçay University, Türkiye
Rana KARAKAŞ ÇENGEL Gökcen DOĞAN	YETİŞKİN KADINLARDA DİYET İNFLAMATUAR İNDEKSİN PREMENSTRUAL SENDROM ÜZERİNDE ETKİSİ	Lokman Hekim University, Türkiye
Suzan CANGÜL	ARTIFICIAL INTELLIGENCE IN RESTORATIVE DENTISTRY: A BIBLIOMETRIC ANALYSIS FROM 2020–2025	Dicle University, Türkiye
Habibe ÖZÇELİK Hasan Hüseyin AVCI	EVALUATION OF WHITE CODE CASES EXPERIENCED BY NURSES	Akdeniz University, Türkiye
Gasim RUSTAMLİ Aliakbar EBRAHIMI Özlem TOMSUK Hamed GHORBANPOOR Onur UYSAL Ayla EKER SARIBOYACI Hüseyin AVCI	CONSTRUCTION OF MULTI-CHANNEL MICROFLUIDIC CHIP SYSTEM WITH MESENCHYMAL STEM CELLS	Eskişehir Osmangazi University, Türkiye
Hüsniye Merve DEMİRER	THE ROLE OF TELEREHABILITATION IN THE MANAGEMENT OF PRIMARY DYSMENORRHEA	İstanbul Gelişim University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-1, Hall-2
15.08.2025
Moderator: Eyüp Can İÇLİ
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Eyüp Can İÇLİ	TOPOLOGY OPTIMIZATION ON ENGINE BRACKET: WEIGHT REDUCTION AND MODAL ANALYSIS	Oyak Renault Otomobil Fabrikaları A.Ş
Ahmet Fatih YILMAZ	EFFECT OF INTERLAYER EPOXY–CARBON FIBER POWDER REINFORCEMENT ON THE TENSILE BEHAVIOR OF FFF-PRINTED PETG	Karabük University, Türkiye
Abdurrahman AYDIN Fatih KAHRAMAN	MULTI-AXIS TOOL PATH PLANNING METHODS IN WIRE AND ARC ADDITIVE MANUFACTURING (WAAM) TECHNOLOGIES	Dokuz Eylül University, Türkiye
Buse ALTIN Miraç Tuba ÇELİK Onur ERTİK İsmail Enes PARLAK	ADMET ANALYSIS AND BIOLOGICAL ACTIVITY PREDICTIONS OF NOVEL MOLECULES GENERATED BY A CONDITIONAL TRANSFORMER MODEL	Bursa Technical ürkiye
Sait BAHADIR TEKEREK Rabia KIRKGEÇİT KARAKELLE Muhammet Vahittin KAÇMIŞ	IMPROVEMENT OF MATERIAL PROPERTIES OF CuZnSn Alloy WITH ALUMINUM AND MAGNESIUM ADDITIONS	Gemciler Güven Metal San. ve Tic. A.Ş. , Kahramanmaraş, Türkiye.
Sait BAHADIR TEKEREK Rabia KIRKGEÇİT KARAKELLE Muhammet Vahittin KAÇMIŞ	INVESTIGATION OF MECHANICAL AND PHYSICAL PROPERTIES OF CuNi10FeMn ALLOY WITH TITANIUM ADDITION	Gemciler Güven Metal San. ve Tic. A.Ş. , Kahramanmaraş, Türkiye.
Rabia KIRKGEÇİT KARAKELLE Sait BAHADIR TEKEREK Sefa BİLGİNER Muhammet Vahittin KAÇMIŞ	INVESTIGATION OF TRIBOLOGICAL AND MICROSTRUCTURAL PROPERTIES OF Cu-15Ni-8Sn-0.2X (X: Al_3, Ti_3, Al+Ti) ALLOYS UNDER SEAWATER CONDITIONS	Gemciler Güven Metal San. ve Tic. A.Ş. , Kahramanmaraş, Türkiye.
Rabia KIRKGEÇİT KARAKELLE Sait BAHADIR TEKEREK Sefa BİLGİNER Muhammet Vahittin KAÇMIŞ	INVESTIGATION OF THE EFFECTS OF AL REINFORCEMENT ON THE MECHANICAL, ELECTRICAL, AND TRIBOLOGICAL PERFORMANCE OF CuCrZr Alloy	Gemciler Güven Metal San. ve Tic. A.Ş. , Kahramanmaraş, Türkiye.

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-1, Hall-3
15.08.2025
Moderator: Assist. Prof. Dr. Oğuzhan ÇELİK
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Özlem MADEN Bingül SUBAŞI HARMANCI	INVESTIGATION OF STRESS COPING, PSYCHOLOGICAL WELL-BEING AND ANGER CONTROL IN SPORT AND NON-SPORTS DOERS	Near East University, TRNC
Çiğdem GÖK Bingül SUBAŞI HARMANCI	EXAMINING SOCIAL ANXIETY, ATTACHMENT STYLES AND INTERPERSONAL EMOTION REGULATION IN ADULT INDIVIDUALS	Near East University, TRNC
Ayhan DEMİR Erkan DİKİCİ	PSYCHOSOCIAL SUPPORT INTERVENTION FOR PSYCHOSOCIAL SUPPORT PERSONNEL WORKING IN DISASTER AREAS	Bitlis Eren University, Türkiye
Oğuzhan ÇELİK	THE EFFECT OF THE ENNEAGRAM PERSONALITY MODEL-BASED SOCIAL SKILLS PSYCHOEDUCATION PROGRAMME ON MIDDLE SCHOOL STUDENTS	Erzincan Binali Yıldırım University, Türkiye
Oğuzhan ÇELİK	THE EFFECT OF THE IDENTITY AWARENESS PSYCHOEDUCATION PROGRAMME ON SECONDARY SCHOOL STUDENTS	Erzincan Binali Yıldırım University, Türkiye
All participants must join the conference 10 minutes before the session time. Every presentation should last not longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.		

Session-1, Hall-4
15.08.2025
Moderator: Dr. Saima Abbas
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Motasem Mirza, Dr. Khawer Bilal Baig	THE ROLE OF CULTURAL AND EMOTIONAL INTELLIGENCE IN SHAPING RESILIENT PAKISTANI MINDS IN A GLOBALIZED ERA	Bahria University, Lahore, Pakistan
Dr. Saima Abbas, Dr. Anasuya Jegathevi Jegath	CULTURAL DISSONANCE AND PSYCHOLOGICAL INFLEXIBILITY: AN ACT-BASED ANALYSIS OF AGGRESSION IN URBAN PAKISTANI YOUTH	University of Cyberjaya, Malaysia
Mrs.E.Elavarasi,Yuvaraj AR,J.Jayadurka,DR.R.Srinivasan	NANOTECHNOLOGY-ENABLED DRUG DELIVERY SYSTEMS	Bharath Institute of Higher Education and Research,Chennai-73
Mrs.E.Elavarasi,Yuvaraj AR,J.Jayadurka,DR.R.Srinivasan	BLOCKCHAIN TECHNOLOGY IN PHARMA FOR SUPPLY CHAIN AND CLINICAL TRIALS	Bharath Institute of Higher Education and Research,Chennai-73
Mrs.E.Elavarasi,Yuvaraj AR,J.Jayadurka,DR.R.Srinivasan	PHARMACEUTICAL SUPPLY CHAIN RESILIENCE POST-COVID-19	Bharath Institute of Higher Education and Research,Chennai-73
Mrs.E.Elavarasi,Yuvaraj AR,J.Jayadurka,DR.R.Srinivasan	BIOSIMILARS AND BIOBETTERS: EXPANDING ACCESS TO BIOLOGICS	Bharath Institute of Higher Education and Research,Chennai-73
Mrs.E.Elavarasi,Yuvaraj AR,J.Jayadurka,DR.R.Srinivasan	CELL AND GENE THERAPIES: TRANSFORMING TREATMENT PARADIGMS	Bharath Institute of Higher Education and Research,Chennai-73

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-1, Hall-5
15.08.2025
Moderator: Sharipah Khadijah S.Hashim
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Sharipah Khadijah S.Hashim	INTEGRATING GREEN SKILLS INTO TVET CURRICULUM: A QUALITATIVE STUDY OF STAKEHOLDER PERSPECTIVES IN MALAYSIA	Politeknik Tuanku Sultanah Bahiyah, Malaysia
ZENATI Rania BOUCHERDOUD Ahmed BENDJELLOUL Meriem SEGHIER Abdelkarim AICHOUNI Sara ELANDALOUSSI El Hadj	BIOPOLYMER-BASED ADSORBENTS FROM NATURAL WASTE: A GREEN STRATEGY FOR WASTEWATER PURIFICATION AND RESOURCE VALORIZATION	University of Relizane ,48000, Relizane, Algeria
Bouchra Bourhabi, Latifa Mouhir, Souad Tayane	VALORIZATION OF SEDIMENTS FROM BIN EL OUIDANE DAM (MOROCCO) AS A MATERIAL FOR CONCRETE	Hassan II University, Casablanca, Morocco.
Seloua ESSAIH, Pr. Rekia BELAHSEN, Pr. Belkassem EL AMRAOUI	ECOLOGICAL AND TRADITIONAL HERITAGE PRESERVED BY RURAL MOROCCAN WOMEN: EDIBLE WILD PLANTS BETWEEN APPRECIATION AND NEGLECT	Ibn Zohr University, Morocco
Md. Shadin Ahmed Sojib, Imon Kumar Shikdar, Md. Al-Amin Sarker	DIETARY FATTY ACID MODULATION: EFFECT ON GROWTH, BODY COMPOSITION, AND TISSUE FATTY ACID PROFILE IN RUI Labeo rohita (HAMILTON, 1822)	University of Rajshahi, Rajshahi-6205, Bangladesh
K.R.Padma K.R.Don	UNDERSTANDING HOW BIODIVERSITY UNDERPINS URBAN ECOSYSTEM FUNCTIONS ESSENTIAL FOR HUMAN HEALTH AND WELL-BEING: PLANNING FOR FUTURE OF URBAN BIODIVERSITY	Sri Padmavati Mahila Visvavidyalayam (Women's) University, Tirupati, AP Bharath Institute of Higher Education and Research (BIHER) Bharath University, Chennai, Tamil Nadu, India

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-1, Hall-6
15.08.2025
Moderator: Aniekan Essienubong Ikpe
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Huma Ikram, Hannah Tariq, Muhammad Raza Shah, Darakhshan Jabeen Haleem	POTENTIATION OF COGNITIVE PROFILE BY SYNTHESIZING NANOFORMULATIONS OF APOMORPHINE	University of Karachi, Karachi–75270, Pakistan
Katia BENYAKOUB Mourad KEDDAM Brahim BOUMAALI Michal KULKA	KINETIC MODELLING OF POWDER-PACK BORONIZED 4Cr5MoSiV1 STEEL BY TWO DISTINCT APPROACHES	Laboratoire de Technologie des Matériaux, Faculté G.M. et G.P., USTHB, B.P. No. 32, El-Alia, Bab-Ezzouar, Algiers 16111, Algeria
Youssef Arif, Mohamed Bouhrim, Fatiha Chigr, Ahmed Ait Chaoui, Ahmed Arif , Mustapha Boulghallat	PHYTOCHEMICAL DRUG CANDIDATES FROM ASPARAGUS OFFICINALIS: IN SILICO TARGETING OF B-GLUCOSIDASE, NR4A1, AND GLUT4	Sultan Moulay Slimane University, Béni Mellal. Morocco
Janani Senthil Kamaraj Murugesan	MYCELIUM- BASED PACKAGING: A FUNGAL ALTERNATIVE TO PLASTIC	SRM Institute of Science and Technology, India
Imoh Ime Ekanem Aniekan Essienubong Ikpe Ini-odu Emmanuel Ekanem	INTEGRATION OF MECHANICAL POWER TRANSMISSION IN PNEUMATIC SYSTEMS FOR CONVENTIONAL AUTOMOBILE APPLICATIONS	Akwa Ibom State Polytechnic, Ikot Osurua, Ikot Ekpene, School of Engineering, Department of Mechanical Engineering Technology, Ikot Ekpene, Nigeria
Elmehdi Majdi, Soufiane Zerraf, Mohammed E. Belghiti, Aabdellah Zeroual, Said Belaaouad	MELAMINIUM DIHYDROGEN PYROPHOSPHATE:INCLUSION OF SYNTHESIS, CRYSTAL PACKING, HIRSHFELD SURFACE ANALYSIS, AND REACTIVITY MAPPING-A COMBINED EXPERIMENTAL AND THEORETICAL STUDY	Hassan II University of Casablanca, Casablanca, Morocco Chouaïb Doukkali University, Morocco
Imene Boulhaia, Manuel Bañobre-López, Nadji Moulai-Mostefa	SWELLING PROPERTIES OF FERROGELS BASED ON AGAR GUM AND OCTENYL SUCCINIC ANHYDRIDE (OSA) STARCH	University of Blida 1 Route de Soumaa, Blida 09000, Algeria INL, International Iberian Nanotechnology Laboratory, Portugal University of Medea AinD'heb,Medea 26001, Algeria

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-1, Hall-7
15.08.2025
Moderator: Assoc. Prof. Dr. Suat YILDIZ
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 10:00 – 12:00

Author(s)	Title	Affiliation
Esra ÖZGÖÇMEN TULA Zeynep Ceren ÇELİK Çiğdem ARSLAN	AN ANALYSIS OF DENTAL STUDENTS PERCEPTIONS OF THE CONCEPT OF ORAL HYGIENE THROUGH MIND MAPPING	Bursa Uludağ University, Türkiye
Suat YILDIZ	EFFECT OF LIMIT STABILITY TRAINING ON STRENGTH DEVELOPMENT AND STRENGTH ASYMMETRY IN A PROFESSIONAL FOOTBALL PLAYER	Manisa Celal Bayar University, Türkiye
Murat DAKMANCI Mehmet AVŞAR Yusuf ÖZEL Ayşe DEMİR	FIVE OLYMPICS, FIVE MODELS: THE ECONOMIC ANATOMY OF THE GAMES FROM 2008 TO 2024	Dicle University, Türkiye
Ayşe ÇİFTCİ	SIBLING PRESENCE AS A DETERMINANT OF SOCIAL SKILL ACQUISITION IN EARLY CHILDHOOD	Çankırı Karatekin University, Türkiye
Kadir DİLER	THE IMPACT OF CIRCADIAN RHYTHMS ON ATHLETIC PERFORMANCE AND PERCEPTUAL RESPONSES	Gaziantep University, Türkiye
Kadir DİLER	A COMPARATIVE STUDY OF UNIVERSITY STUDENTS' ATTITUDES TOWARD SPORTS IN TURKEY AND SYRIA: THE ROLE OF DEMOGRAPHIC, SOCIAL, AND PSYCHOLOGICAL VARIABLES	Gaziantep University, Türkiye
Hilal KURAK Güzin Yasemin TUNÇAY	EDUCATION OF CHILDREN WITH SPECIAL NEEDS AND PARENTAL OPINIONS	Çankırı Karatekin University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-2, Hall-1
15.08.2025
Moderator: Assist. Prof. Dr. İlkay ÇORUMLUOĞLU
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Nursel ZORTUL	DRIVER FATIGUE DETECTION AND PREDICTION WIRH MACHINE LEARNING METHODS	Kocaeli University, Türkiye
Iftikhar Chalabi Rahim Oruczade Hemide Halikli	THE TRANSMISSION MECHANISM OF A TRACKED MOBILE ROBOT IN A NEW DESIGN AND THE DEVELOPMENT OF ITS CONTROL SYSTEM	Azerbaijan Technical University, Azerbaijan
Ferhat BOZDUMAN Hatice Kübra KOÇ TOPCUOĞLU İbrahim KÜRTÜL	DESIGN OF A MINIATURE EMG DEVICE FOR USE IN LABORATORY ANIMALS USING ARDUINO AND LABVIEW	Karabük University, Türkiye Bolu Abant İzzet Baysal University, Türkiye
Barış ERGÜL Arzu ALTIN YAVUZ	COMPARATIVE PERFORMANCE ANALYSIS OF ARIMA, SARIMA, AND LSTM MODELS FOR AIR QUALITY INDEX FORECASTING	Eskişehir Osmangazi University, Türkiye
Barış ERGÜL Arzu ALTIN YAVUZ	MACHINE LEARNING-BASED MODELING OF DROUGHT PROCESSES IN TÜRKİYE	Eskişehir Osmangazi University, Türkiye
İlkay ÇORUMLUOĞLU	DETERMINATION OF HISTOLOGICAL FEATURES OF CANCER CELLS USING ARTIFICIAL INTELLIGENCE	Ankara Yıldırım Beyazıt University, Türkiye
Betül YILMAZ Hakan GÜLEN	DESIGN AND SIMULATION OF A LORA-BASED MULTI-SENSOR IoT SYSTEM ARCHITECTURE FOR SMART AGRICULTURE APPLICATIONS	Mersin University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-2, Hall-2
15.08.2025
Moderator: Lect. Müzeyyen YÜCEL
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Sevilay ERMİN Burak ŞEN	CIRCULAR VIBRATING SCREEN DESIGN	SİSTEMAS TEKNOLOJİ SİSTEM ÜRT. TEM. Ve TİC A.Ş.
Göksu GÖZÜKARA Nurdan ÇOLAKOĞLU	EVALUATION OF SALES TEAM PERFORMANCE SYSTEM WITH ANALYTIC HIERARCHY PROCESS (AHP)	İstanbul Arel University, Türkiye
Müzeyyen YÜCEL	ANALYSIS OF DENIM FABRIC IN TERMS OF SUSTAINABILITY	Beykent University, Türkiye
Nazila Ragimova Almammadova Mehriban Hajiyeva Kifayat Kamenieva Nina	DEFINITION OF AN APPROACH TO THE ANALYSIS OF THE SAFETY OF USING SYSTEMS OF ELECTRICAL INTERLOCKING OF SWITCHES AND SIGNALS	Azerbaijan State Oil and Industry University Ukrainian State University of Railway Transport
Miraç Tuba ÇELİK Buse ALTIN İsmail Enes PARLAK Onur ERTİK	SEQUENCE-BASED DEEP LEARNING MODELS FOR DRUG-LIKE MOLECULE GENERATION: PERFORMANCE UNDER DATA CONSTRAINTS	Bursa Technical University, Türkiye
Yavuz KIRIM Özcan BAŞARAN	CAPTURE OF CARBON DIOXIDE FROM WASTE GAS OF INDUSTRIAL ENTERPRISES	Siirt University, Türkiye
Esra MERT Aslan Deniz KARAOĞLAN	DEVELOPMENT OF NATURALLY DYED FUNCTIONAL TEXTILES USING THE TAGUCHI METHOD	Balıkesir University, Türkiye
Gülşah SAYDAN KANBEROĞLU	DEVELOPMENT OF A PVC MEMBRANE POTENTIOMETRIC SENSOR FOR THE DETERMINATION OF DICHROMATE	Van Yüzüncü Yıl University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-2, Hall-3
15.08.2025
Moderator: Assoc. Prof. Dr. Taylan TUĞRUL
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Mustafa NISARI	FORMATION OF PISTACIA VERA SHELL EXTRACT- INCORPORATED NANOFLOWER AND INVESTIGATION OF ITS INTRINSIC PEROXIDASE-MIMIC ACTIVITY	University of Nuh Naci Yazgan, Türkiye
Aysel BAYRAM Sidar BAĞBUDAR Semen ÖNDER	HER2-LOW MEME KANSERİ SIKLIĞI: TEK MERKEZ RETROSPEKTİF ÇALIŞMA	İstanbul University, Türkiye
Zehra VAROL Elif Afra BEŞPARMAK Muhammed Sabit KIYICI Medine DOĞAN SARIKAYA Seçil YILMAZ	ANALYSIS OF CANCER STEM CELL-DRIVEN INVASION IN BREAST CANCER METASTASIS USING ORGAN-ON-A-CHIP TECHNOLOGY	Erciyes University, Türkiye Penn State University, USA Erciyes University, Türkiye
Taylan TUĞRUL	THE EFFECTS OF FULL WIDTH AT HALF MAXIMUM (FWHM) AND ELECTRON MEAN ENERGY VALUES ON PERCENT DEEP DOSE AND LATERAL DOSE PROFILES	Van Yüzüncü Yıl University, Türkiye
İbrahim AYDIN	THE EFFECT OF THE PRESENCE OF MUCINOUS COMPOUNDS ON SURVIVAL IN GASTRICCANCER: A RETROSPECTIVE COHORT STUDY	Health Sciences University, Türkiye Prof.Dr.İlhanVarank Eğitim ve Araştırma Hastanesi
İbrahim AYDIN	COMPARISON OF CONVENTIONAL HEMORRHOIDECTOMY AND HEMORRHOIDAL DEARTERIALIZATIONTECHNIQUES IN THETREATMENT OF GRADE III-IV HEMORRHOIDS: A RETROSPECTIVE ANALYSIS OF 67 PATIENTS	Health Sciences University, Türkiye Prof.Dr.İlhanVarank Eğitim ve Araştırma Hastanesi
Özge ÇETINARSLAN	D-DIMER/ALBUMIN, CRP/ALBUMIN, AND THEIR COMPOSITE INFLAMMATORY-COAGULATION INDEX PREDICT READMISSION AFTER NSTE-ACS	T.C. Demiroğlu Bilim Üniversitesi, Tıp Fakültesi,Kardiyoloji Anabilim Dalı, İstanbul Florence Nightingale Hastanesi
Ünal ÖZTÜRK Nuray ÜREMİŞ Ergül Belge KURUTAŞ	INVESTIGATION OF 8-ISOPROSTAGLANDIN F2A, TAU PROTEIN, AND GPER LEVELS IN PATIENTS WITH SUPRAVENTRICULAR TACHYCARDIA	Kahramanmaraş Sütçü İmam University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-4
15.08.2025
Moderator: Subhashish Dey
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Subhashish Dey	INNOVATIVE STUDY OF STRUCTURAL LIGHT WEIGHT CONCRETE MIX DESIGN(M25) USING EXPANDED CLAY AGGREGATE	Seshadri Rao Gudlavalleru Engineering College, Gudlavalleru, Andhra Pradesh, India
Çağatay Yılmaz	COMPARISON OF AUTOCLAVE AND OUT-OF-AUTOCLAVE COMPOSITE MANUFACTURING METHODS	Abdullah Gül Unıversity, Türkiye
Habibullah UZUN	METAL PARTITIONING AND MOBILITY IN SURFACE WATERS UNDER VARYING FLOW AND LAND USE CONDITIONS	Istanbul Technıcal Unıversıty
Doha doughri, Boubker Mehdaoui, Rachid akhreddine, Abdeslam El Bouari	OPTIMIZING THE STRUCTURE AND DIELECTRIC FEATURES OF SPINEL LITHIUM FOR ENERGY STORAGE IN BATTERY	University Hassan II of Casablanca, Chemistry Department, Casablanca, Morocco
Youssef Arif, Mohamed Bouhrim, Fatiha Chigr , Ahmed Ait Chaoui , Ahmed Arif , Mustapha Boulghallat	PHYTOCHEMICAL DRUG CANDIDATES FROM ASPARAGUS OFFICINALIS: In Silico Targeting of β-Glucosidase, NR4A1, and GLUT4	Sulatan Moulay Slimane University, Béni Mellal. Morocco
N.G. Kalyani Reddy, Chayaraju BalaSai, K.Tharani	ROBOTS IN CONSTRUCTION: ENABLING TECHNOLOGY, APPLICATIONS, AND CHALLENGES	G Pulla Reddy Engineering College, (Autonomous),Kurnool-518002(A.P)
Subhashish Dey	STUDY ON STRENTH PROPERTIES OF CONCRETE BY PARTIAL REPLACEMENT OF CEMENT WITH BLACK CARBON POWDER	Seshadri Rao Gudlavalleru Engineering College, Gudlavalleru, Andhra Pradesh, India
N.G. Kalyani Reddy, Chayaraju BalaSai, K.Tharani	INTEGRATING AUGMENTED AND VIRTUAL REALITY TECHNOLOGIES IN ARCHITECTURAL CIVIL ENGINEERING	G Pulla Reddy Engineering College

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-5
15.08.2025
Moderator: Assistant Professor Dr. Violla Makhzoum
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Dr. Alma Yerzhanova Doç. Dr. Coşkun Fırat Prof. Dr. Mukhtar Saltayevich Yerzhanov	EVALUATING RENEWABLE ENERGY FINANCING MODELS IN EMERGING AND ADVANCED ECONOMIES	University of Turan, Accounting and Auditing Department, Almaty, Kazakhstan Istanbul technical University, Energy Institute, Istanbul, Türkiye
Joy Nneka Uchenye Ejikeme, Emmanuel Chijioke Mmadu	MASQUERADING AS A CULTURAL RESOURCE AND ITS CONTRIBUTIONS TO TOURISM DEVELOPMENT IN IGBO-EZE SOUTH LOCAL GOVERNMENT AREA	University of Nigeria, Nsukka, Enugu State, Nigeria
Assistant Professor Dr. Violla Makhzoum Amal Moussawi Racha Fawaz Sumaya Helbawi	BRIDGING AI AND EDUCATIONAL RESEARCH: OPPORTUNITIES FOR DATA-DRIVEN LEARNING	Modern University for Business and Science (MUBS).Beirut, Lebanon
Omojaro A. O., Taiwo A. A., Taiwo F. R., Babayemi D. I.	RISK MANAGEMENT AND SUPPLY CHAIN RESILIENCE OF OLAM FEED MILL, LAGOS, NIGERIA	The Federal Polytechnic Ilaro, Ogun State, Nigeria.
Dr. Mark Anthony N. Polinar, Atty. Philip Jun A. Pagatpat	DECISION-MAKING ROLES AND WORKPLACE INTERPERSONAL CONFLICT IN AN ORGANIZATIONAL SETTING IN MANDAUE CITY	Cebu Institute of Technology-University
Ebrahim Mohammed Ahmed Professor (Dr) Gurudutta P. Japee	RISK-BASED AUDITING IN PRACTICE: A SYSTEMATIC REVIEW OF IMPLEMENTATION AND IMPACT	Gujarat University, Ahmedabad
Kamar Ismail Shittu Samuel Olatunbosun Binta Abubakar Naimat Kamar Abiola	THEORETICAL PERSPECTIVES ON BUSINESS DEVELOPMENT STRATEGIES AND PERFORMANCE OF SMES: A CONCEPTUAL REVIEW	Federal University Dutsinma Katsina, Nigeria.
Nguyen THİ HANG Tran THI TRANG	OPTIMIZING FINANCIAL ACCOUNTING PROCESSES THROUGH ARTIFICIAL INTELLIGENCE (AI): IMPACTS AND STRATEGIC IMPLICATIONS FOR BUSINESSES IN THE DIGITAL ECONOMY	Thai Nguyen University of Information and Communication Technology University Of Economics

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-6
15.08.2025
Moderator: Dr. R. DEVI
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Dr. R. DEVI	GREEN SYNTHESIS AND CHARACTERIZATION OF SILVER NANOPARTICLES USING EUGENIA ROXBURGHII DC: A SUSTAINABLE APPROACH	Annamalai University, Chidambaram
Mayra ihsan, Wafa Majeed, Ayesha, Muhammad Saad Tariq	EXPLORATION OF ANTI-INFLAMMATORY AND ANTI-OXIDATIVE PROPERTIES OF SALSOLA IMBRICATA IN FREUND'S COMPLETE ADJUVANT INDUCED ARTHRITIS	University of Agriculture, Faisalabad, Pakistan
Hassan Abdulmumini	AGRICULTURE GLOBALIZATION AND HOW IT AFFECTS FOREING RELATION	Federal polytechnic Bali, Taraba state
Babita Gupta	NANOMATERIAL-BASED TRANSDERMAL DRUG DELIVERY FOR SKIN-RELATED DISORDER	Institute of Technology and Management Gida Gorakhpur
ARRAJI Maryem, Khaoula EL BARBI, Fatima Zahra BOUCHACHI, CHAHBOUNE Mohamed	PREVALENCE AND DETERMINANTS OF MEDICATION ADHERENCE IN MOROCCAN PATIENTS WITH TYPE 2 DIABETES: A MULTICENTER CROSS-SECTIONAL STUDY	Hassan First University of Settat
Nabil Ahmed Khan, Ahmad Yusuf Adhami, Mohd Faizan	MULTIOBJECTIVE TRANSPORTATION SURVIVAL COST FUNCTION WITH EXPONENTIAL SUPPLY AND WEIBULL DEMAND: A NEUTROSOPHIC APPROACH	
Wajahat Ali Mohd Khalid	OPTIMIZING HEALTHCARE SYSTEMS UNDER UNCERTAINTY USING FERMATEAN FUZZY AND NEUTROSOPHIC GOAL PROGRAMMING APPROACH	Aligarh Muslim University, Aligarh, India
Vanitha Sri.T, Farhatyassir.M, Elavarasi.E, Dr.R.Srinivasan	BIOREMEDIATION OF PESTICIDE-CONTAMINATED AGRICULTURAL LANDS	Bharath Institute of Higher Education and Research

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-2, Hall-7
15.08.2025
Moderator: Dr. T. O. James
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 12:30 – 14:30

Author(s)	Title	Affiliation
Saraswathi.V,R. Vidhyalakshmi, Kowsika.M, Dhanush kumar.S,Sastha.K, Prashanth.S, Dr.W.Helen, Dr.R.Srinivasam	CLINICAL SIGNIFICANCE OF ELECTROLYTE IMBALANCE IN HUMAN HEALTH	Bharath Institute of Higher Education and Research, Chennai, Tamilnadu India.
Dr. T. O. James, Dr. G. I. Onwuka EKELE, Sunday John	IDENTIFICATION OF SOME RISK FACTORS AND PREVALENCE OF OBESITY ON CHILDREN BETWEEN FIVE TO TEN YEARS	Kebbi State University of Science and Technology, Aliero, Nigeria
Muqaddas Ishfaq, Dr. Anam Khan, Umm e Eman Syed, Neelam Bibi, Dr. Saba Ghayas, Sadia Niazi	IMPACT OF SCREEN ADDICTION AND HEADACHE ON IRRITABILITY AMONG MIGRAINE PATIENTS	Rawalpindi Institute of Cardiology University of Sargodha
Idris Saratu, Dr Abubakar N. Muhammed, Idris Ibrahim	THE IMPACT OF ADMINISTRATIVE CONSTRAINTS IN SPORTS DEVELOPMENT IN NIGERIA, A CASE STUDY OF THE NATIONAL SPORTS COMMISSION	Prince Abubakar Audu University Ayangba, Kogi State. Technical Education Departmrnt Kogi State College Of Education Technical Mopa
Anitha P, Vardhana Janakiraman	FUNGAL DIVERSITY FROM AGRO-INDUSTRIAL OILSEED CAKES: ISOLATION, CHARACTERIZATION, AND BIOTECHNOLOGICAL POTENTIAL OF LIPOLYTIC SPECIES	Vels Institute of Science, Technology and Advanced Studies, Pallavaram, Chennai-600117
Neg Imane, Bouda Said, Zaggoumi Hasna, Mardoume Ibtissam, Haddioui Abdelmajid	COMPARATIVE ANALYSIS OF METHANOL EXTRACTS FROM JUNIPERUS OXYCEDRUS ACROSS ELEVEN MOROCCAN POPULATIONS	Sultan Moulay Slimane University, Beni Mellal, Morocco
Adekola Sodiq Adetunji, Adekola Saheed Ayodeji, Dhikroh Oriyomi Adekola, Fridaus Muhammad Lawal, Suliyat Wahab	THE IMPACTS OF FIBROIDS ON WOMEN'S QUALITY OF LIFE	University of Abuja, Abuja

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-3, Hall-1
15.08.2025
Moderator: Assoc. Prof. Dr. Özcan ÇAĞLAR
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Hülya ŞEREFLİŞAN Ahmet ALKAYA	INNOVATIVE APPROACHES TO WATER QUALITY MANAGEMENT THROUGH THE GENETIC POTENTIAL OF FRESHWATER MUSSELS	İskenderun Technical University, Türkiye
Hülya ŞEREFLİŞAN Ahmet ALKAYA	THE EFFICACY AND APPLICATION POTENTIAL OF MUSSEL AND SNAIL SHELLS IN HEAVY METAL REMOVAL: A BIOREMEDIATION APPROACH	İskenderun Technical University, Türkiye
Mustafa HÜSREVOĞLU Aslı BOZDAĞ	TECH-ENABLED PARTICIPATORY PLANNING IN URBAN TRANSFORMATION	Niğde Ömer Halisdemir University, Türkiye
Utku TOKSUN Yusuf SAYAN	STRESS ANALYSIS OF COOLING WATER LINES FOR UREA-FORMALDEHYDE CONCENTRATE (UFC-85) PRODUCTION PLANT	Rönesans Endüstri Tesisleri Tasarım Merkezi, İstanbul, Türkiye
Büşran SUNYAR Mesude Figen DÖNMEZ	BIBLIOMETRIC ANALYSIS OF SCIENTIFIC PUBLICATIONS ON BACTERIAL DISEASES OF TOMATO	Iğdır University, Türkiye
Büşran SUNYAR Mesude Figen DÖNMEZ	BIBLIOMETRIC PROFILE OF STUDIES ON BACTERIAL CANCER IN STONE FRUIT TREES	Iğdır University, Türkiye
Özcan ÇAĞLAR Harun ALPTEKİN Oğuzhan Meriç ÇAĞLAR	BIBLIOMETRIC ANALYSIS OF ACADEMIC PUBLICATIONS ON NITROGEN AND WATER USE EFFICIENCY IN CEREALS. AN ASSESSMENT FROM AN AGRICULTURAL SUSTAINABILITY PERSPECTIVE	Iğdır University, Türkiye
Mehmet Ali KARGICAK Engin ERTAN	EFFECTS OF ABIOTIC STRESS FACTORS IN FRUIT GROWING	İncir Araştırma Enstitüsü Müdürlüğü, Aydın, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-3, Hall-2
15.08.2025
Moderator: Assoc. Prof. Dr. Semra TETİK
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Sabit MENTEŞE Eda Burcu YILMAZ	INVESTIGATION OF WORK ACCIDENTS WITHIN THE SCOPE OF ARTICLE 13 OF LAW NO. 5510	Munzur University, Türkiye
İlyas GÜNEŞ Olgun UKŞAL	ESCAPE FROM THE SIEGE OF THE UNIPOLAR ORDER: THE PRIMAKOV DOCTRINE IN RUSSIAN FOREIGN POLICY	Marmara University, Türkiye Kafkas University, Türkiye
Olgun UKŞAL	A SOCIAL RESPONSE TO HUMAN RIGHTS VIOLATIONS: THE MOTHERS OF THE PLAZA DE MAYO MOVEMENT IN ARGENTINA	Kafkas University, Türkiye
Burcu KAYA SAYARI Hazel Andrews	FRAMING A CREATIVE TOURISM SYSTEM IN HERITAGE-LED DESTINATIONS	Liverpool John Moores University, Liverpool, UK Anadolu University, Türkiye
Ummühani USLU	DISCUSSION OF SOCIO-ECONOMIC PROBLEMS OF ELDERLY WOMEN IN THE CONTEXT OF SOCIAL WORK	Alanya Alaaddin Keykubat University, Türkiye
D. Burcu Eğilmez-Gürsoy	ENVY, BLAME, AND NEOLIBERAL COMMON SENSE: PUBLIC REACTION TO THE 2024 İZMIR MUNICIPAL WORKERS' STRIKE	İzmir Economics University, Türkiye
Semra TETİK	THE RELATIONSHIP BETWEEN CAREER STRESS AND HOPELESSNESS: A STUDY ON UNIVERSITY STUDENTS	Manisa Celal Bayar University, Türkiye
Semra TETİK	A RESEARCH ON THE RELATIONSHIP BETWEEN SOCIAL INTELLIGENCE AND ENTREPRENEURSHIP TENDENCY	Manisa Celal Bayar University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Session-3, Hall-3
15.08.2025
Moderator: Assoc. Prof. Dr. Tayfun ACARER
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Sinem DEMİROĞLU	THE EFFECT OF SOCIAL ASSISTANCE ON INTERPROVINCIAL MIGRATION: THE TÜRKİYE EXAMPLE	Türkiye
Cem ÖZKURT	THE SPONTANEOUS DIALECTICS OF SOCIAL FORMATION: MIND, SYMBOLIC ORDER, AND THE WEB OF MEANING	Trabzon University, Türkiye
Zeynep YAVUZ Zeynep Nur SANDIKCI	EFFECTIVE PERSONAL FINANCE MANAGEMENT WITH SMART ASSISTANT	Sipay Electronic Money and Payment Services Inc., Istanbul, Turkiye
Emine ARDUÇ KARA	WATCHING THE POWER: A NEW METHOD OF CONTROL IN JOURNALISM FROM PANOPTIC SURVEILLANCE TO SOUSVEILLANCE	Süleyman Demirel University, Türkiye
Tayfun ACARER	DIFFERENCES CAUSED BY RECENT TECHNOLOGICAL DEVELOPMENTS IN THE INFORMATION INDUSTRY IN THE WORKING CONDITIONS OF SEAMEN WORKING IN THE MARITIME SECTOR AND THE FUNCTIONS OF SHIPS	Piri Reis University, Türkiye

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Antalya

Session-3, Hall-4
15.08.2025
Moderator: Asst. Prof. Dr. Elza Bitsadze
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Khalid Boujdi	EFFICIENT SYNTHESIS, STRUCTURAL ELUCIDATION, ANTIBACTERIAL SCREENING, AND IN SILICO EVALUATION OF NOVEL ESTER-FUNCTIONALIZED ISOXAZOLES	University Hassan II, B.P. 146, Mohammedia 28800, Morocco
P C Praveen Kumar, B.Uday Kumar Reddy, Gandla Roopeshwar	5G ANTENNA GAIN ENHANCEMENT USING A NOVEL METASURFACE	G.Pulla Reddy Engineering College(Autonomous):Kurnool,AP, India-518003
P C Praveen Kumar, B.Koushik Sai, Gorrepati Charan Kumar	TUNABLE COMPACT METAMATERIAL-BASED RESONATOR FOR 6G THZ WIRELESS APPLICATIONS	G.Pulla Reddy Engineering College(Autonomous):Kurnool,AP, India-518003
P C Praveen Kumar, N K Deekshith, Pappaka Ganesh	THz MIMO RECONFIGURABLE ANTENNA WITH A MULTI PORT INPUT	G.Pulla Reddy Engineering College(Autonomous):Kurnool,AP, India-518003
Asst. Prof. Dr. Elza Bitsadze	SMART SENSORS, DATA ANALYTICS, AND GEOGRAPHIC INFORMATION SYSTEMS (GIS) FOR COASTAL ECOSYSTEMS MONITORING	Akaki Tsereteli State University, Georgia
Alpha Friday Ebole Akeem Olanrewaju Ajani Khadijah Kubura Agbalaya	DESIGN AND IMPLEMENT OF AN IOT-BASED SYSTEM FOR REAL-TIME MONITORING OF LIVESTOCK HEALTH AND GPS LOCATION TRACKING USING GSM TECHNOLOGY	Lagos State University of Science and Technology, Ikorodu, Lagos, Nigeria
Halliru Sani Musbahu Salisu	HYPERTENSION RISK PREDICTION SYSTEM FOR EARLY DETECTION: A MACHINE LEARNING APPROACH USING SUPERVISED LEARNING	Al-Qalam University Katsina, Nigeria
Anshul Negi, Sushma Bahuguna, Nupur Dang, Hemwati Nandan Bahuguna	FROM DATA TO DECISION: EXPLORING DEFORESTATION TRENDS, INSIGHTS, AND REPORTING	Tecnia Institute of Advanced Studies CDL, Delhi, India

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Antalya

Session-3, Hall-5
15.08.2025
Moderator: Dr. Deepali Tomar
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
KEWULERE Olayimika Wasiu ALABI Oladimeji Muhammad	HISTORICIZING THE UPS AND DOWNS OF ECONOMY IN MORO DISTRICTS, ILORIN EMIRATE, NIGERIA	Kwara State University, Malete, Kwara State, Nigeria
Md Amirul Islam	THE PREDICTION OF MARRIED COUPLES' OBSERVED INTIMACY SKILLS ACROSS THE TRANSITION TO PARENTHOOD IN BANGLADESH: MARITAL SANCTIFICATION AND SPIRITUAL INTIMACY	University of Rajshahi, Rajshahi-6205, Bangladesh Institute of Natural Resources Research and Development, Rajshahi-6206, Bangladesh
Ebrahim Salimikouchi	WALKING THE WORLD TO WRITE THE EARTH : TOWARDS AN ECOPOETICS OF TRAVEL IN IBN BATTUTAT'S RIHLA	University of Neuchâtel, Switzerland
Dr. Deepali Tomar	ASSESSING THE DISRUPTION OF COVID- 19 ON MICRO-ENTREPRENEURS IN UTTARAKHAND	Tecnia Institute of Advanced Studies CDL, Delhi India
Ameen. A. Abdulkadir	HISTORICAL ACCOUNT OF DIPLOMATIC AND CULTURAL EXCHANGES BETWEEN TURKIYE AND NIGERIA FROM THE 16TH CENTURY TO THE 21ST CENTURY	University of Abuja
Dr.B.MAHADEVAN, MANJULA.S, NIVEDHA.S, SHANTHIGA.B	AI IN SKILL DEVELOPMENT AND VOCATIONAL TRAINING	Vellalar College for Women (Autonomous), Thindal (Post), Erode, Tamilnadu, Indi
Anjali Ganesh, Roopesh	HARMONIZING ETHICS AND SPIRITUALITY: A PATHWAY TO ORGANIZATIONAL GROWTH AND EMPLOYEE WELL-BEING	St Joseph Engineering College, Mangalore, Karnataka, India

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-3, Hall-6
15.08.2025
Moderator: Besnik HAJDARI
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Besnik HAJDARI	ARTIFICIAL INTELLIGENCE IN INDUSTRIAL AUTOMATION: TRENDS, TECHNIQUES, AND APPLICATIONS	University "Isa Boletini" Mitrovica
Ms. Hetal Pravinsinh Jadeja Jatinkumar Bhimjibhai Kotadiya	ARTIFICIAL INTELLIGENCE IN ACTION: A COMPARATIVE STUDY OF ML ALGORITHMS FOR TEXT CLASSIFICATION	At Harivandana College, Rajkot, India Atmiya University, Rajkot, India
Jatinkumar Bhimjibhai Kotadiya Dr.Dharmendra Jayantilal Ambani Dr. Amit K. Patel	A COMPARATIVE STUDY OF MACHINE LEARNING ALGORITHMS FOR SENTIMENT ANALYSIS USING ARTIFICIAL INTELLIGENCE TECHNIQUES	Atmiya University, Rajkot, India Harivandana College, Rajkot, India
ChayarajuBalaSai Nadagouda Kalyani K.Tharani	A COMPREHENSIVE ASSESSMENT OF THE APPLICATION OF OPERATIONAL RESEARCH METHODS TO THE DESIGN OF CONCRETE MIXES	G.pulla reddy Engineering college(Autonomous),Kurnool, AP, INDIA
Dr. Sameer Suresh Nanivadekar	BUILDING A GLOBAL CAREER WHILE PURSUING AN ENGINEERING DEGREE: STRATEGIES FOR STUDENTS WORLDWIDE	A P SHAH INSTITUE OF TECHNOLOGY ,THANE ,MAHARSHTRA .INDIA
Arti Saxena	CLASSIFICATION OF VEHICLE INSURANCE CONSUMERS USING MACHINE LEARNING TECHNIQUES	Manav Rachna International Institute of Research and Studies, Faridabad, Haryana, India.
Mohamed BAKIR My Abdelkader Youssefi Rachid DAKIR Mouna EL WAFI Younes El KOUDIA	URBAN AUTONOMOUS NAVIGATION USING MULTI-LIDAR WITH DBSCAN CLUSTERING AND KALMAN FILTER TRACKING	Hassan First University, Faculty of Science and Technology, Laboratory of Engineering, Industrial Management, and Innovation (LIMII), Morocco

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.



Session-3, Hall-7
15.08.2025
Moderator: Dr. Emmanuel Ugochukwu ANASO
Meeting ID: 862 1528 3413/ Passcode: 131415
Ankara Local Time: 15:00 – 17:00

Author(s)	Title	Affiliation
Dr Simran Sikka Dr Mikul	HOW GLOBAL BRANDS INFLUENCE LOCAL CONSUMPTION PATTERNS IN EMERGING ECONOMIES	Manipal University Jaipur Maharishi Markandeshwar (Deemed to be University), Mullana
Nguyen THİ HANG Tran THI TRANG	OPTIMIZING FINANCIAL ACCOUNTING PROCESSES THROUGH ARTIFICIAL INTELLIGENCE (AI): IMPACTS AND STRATEGIC IMPLICATIONS FOR BUSINESSES IN THE DIGITAL ECONOMY	Thai Nguyen University of Information and Communication Technology University Of Economics - Technology For Industries
Grace Charles DIKKI	FINANCIAL RISK MANAGEMENT AND THE VALUE OF LISTED DEPOSIT MONEY BANKS IN WEST AFRICA USING NIGERIA AS A CASE STUDY	Georgia State University, Atlanta, United States
Melik Sami, Khelil Sara, Tallal Abdel Karim Bouzir	ECONOMIC OPTIMIZATION IN ARCHITECTURE: BIM AS A CATALYST FOR COST-EFFICIENT AND AGILE PRACTICE	Mohamed Khider Biskra University, Biskra, Algeria. LACOMOFA Biskra laboratory. Algeria.
Siham NAHLI Abdellatif EDDAKIR	THEORETICAL APPROACHES TO UNDERSTANDING SALARY MANAGEMENT AND SOCIAL PERFORMANCE: A MULTIDISCIPLINARY LITERATURE REVIEW	Sidi Mohammed Ben Abdellah University, FEZ, Morocco
Melik Sami, Khelil Sara, Tallal Abdel Karim Bouzir	DESIGNING FOR DECARBONIZATION: ENVIRONMENTAL BENEFITS OF ZERO-CARBON PUBLIC BUILDINGS	Mohamed Khider Biskra University, Biskra, Algeria. LACOMOFA Biskra laboratory. Algeria.
Dr. Emmanuel Ugochukwu ANASO	STRENGTHENING FEED SECURITY IN WEST AFRICA: UNLEASHING THE STRATEGIC POTENTIAL OF FINANCIAL INSTITUTIONS FOR SUSTAINABLE LIVESTOCK SYSTEMS	Federal University of Agriculture Mubi, Adamawa, Nigeria

All participants must join the conference 10 minutes before the session time.
Every presentation should last not longer than 10-12 minutes.
Kindly keep your cameras on till the end of the session.

Antalya

CONTENT

CONGRESS ID	1
SCIENTIFIC COMMITTEE	2
PHOTO GALLERY	3
PROGRAM	4
CONTENT	5

FULL TEXTS BOOK

Chandani Sharma, Ghazala Ansari , Aarti Sharma, Manidipa Roy ***ADVANCED DESIGN AND OPTIMIZATION STRATEGIES FOR WEARABLE HEALTH DEVICES FOR CHRONIC DISEASE MANAGEMENT***	1
ChayarajuBalaSai, Nadagouda Kalyani, K.Tharani ***A COMPREHENSIVE ASSESSMENT OF THE APPLICATION OF OPERATIONAL RESEARCH METHODS TO THE DESIGN OF CONCRETE MIXES***	7
Selçuk ALBAYRAK, Oğuz MERHAN, Kadir BOZUKLUHAN ***YABAN HAYATI REHABİLİTASYONUNA YÖNELİK UZUN DÖNEMLİ VAKA ANALİZİ (2018–2025)***	29
Murat KONCA ***THE RELATIONSHIP BETWEEN DEBT AND HEALTH STATUS: THE TURKISH CASE***	38
Ramazan Rüçhan KAYA, Şevval Erva KAYA ***ILLNESS AND HOSPITAL IN THE PRESCHOOL PERIOD: AN EVALUATION BASED ON CHILDREN'S VERBAL AND VISUAL EXPRESSIONS***	45
Şevval Erva KAYA, Mine KOYUNCU ŞAHİN ***SOCIAL SKILLS AND PEER RELATIONSHIPS IN PRESCHOOL: A BIBLIOMETRIC ANALYSIS***	51
Yunus BOR ***OSMANLI DEVLETİNDE DÜYUN-I UMUMİYYE İDARESİ***	59
Öznur ALAN, Alime SELÇUK TOSUN ***INVESTIGATION OF PERCEIVED HEALTH AND ITS ASSOCIATED FACTORS AMONG ADULTS***	64
Ahmet Fatih YILMAZ ***EFFECT OF INTERLAYER EPOXY–CARBON FIBER POWDER REINFORCEMENT ON THE TENSILE BEHAVIOR OF FFF-PRINTED PETG***	69
Nida BAYINDIR, Mehmet GÖKCE ***PRESERVICE TEACHERS'PERCEPTIONS OF THE EDUCATIONAL BENEFITS AND SAFETY RISKS OF YOUTUBE IN PRIMARY EDUCATION***	80

Andreea Irina BARZIC, Vasile Bogdan MARDARI, Raluca Marinica ALBU *THE ROLE OF TiO2 NANOPARTICLES IN IMPROVING THE OPTICAL AND ELECTRICAL CHARACTERISTICS OF POLYMER NANOCOMPOSITES FOR LED DEVICES*	95
Timur KOCA, Sümeyye ÇITAK, Aylin Fidan KORCUM *RADIATION-INDUCED SKIN TOXICITIES IN HEAD AND NECK CANCERS: CLINICAL SIGNIFICANCE AND MANAGEMENT APPROACHES*	102
Rahim Mammadzada *SIMULATION BASED ASSESSMENT OF COOLING FAN ENERGY TRADE-OFFS IN PHOTOVOLTAIC SYSTEMS USING MATLAB*	110
Rahim Mammadzada *SIMULATION BASED MODELING OF THERMAL AND ELECTRICAL BEHAVIOR IN PHOTOVOLTAIC PANELS USING MATLAB*	116
Mustafa Sedat ARSLAN, Tuğba Melike PARLAK, Ece OKTAY *DETERMINATION OF THE EMBRYOTOXIC EFFECT OF CEFAZOLIN BY IN OVO METHOD*	122
Cüneyt Yılmaz *SOCIAL AND POLITICAL EFFECTS OF DIGITAL TRANSFORMATION*	132
Emmanuel Ugochukwu ANASO *STRENGTHENING FEED SECURITY IN WEST AFRICA: UNLEASHING THE STRATEGIC POTENTIAL OF FINANCIAL INSTITUTIONS FOR SUSTAINABLE LIVESTOCK SYSTEMS*	143
İsmail VAROL, Fatih KAHRAMAN *EFFECT of FRICTION STIR PROCESSING on MICROSTRUCTURE and MECHANICAL PROPERTIES of METAL MATRIX COMPOSITES*	148
Təranə NAĞIYEVA *THE INFLUENCE OF RUSSIAN SYNTAX IN THE OFFICIAL STYLE OF THE AZERBAIJANI LANGUAGE*	172
Rojhat ERGÜVEN, Alp Buğra AYDIN, Derya AVCI *SPI PREDICTION USING THE GRADIENT BOOSTING REGRESSOR (GBR) ALGORITHM*	183
Mehmet DUGAN *DIGITAL STUDIES AND METHODS ON AVERROES: AN APPLICATION BASED ON THE COMMENTARY ON THE POSTERIOR ANALYTICS*	192
Büşran SUNYAR, Mesude Figen DÖNMEZ *DOMATESİN BAKTERİYEL HASTALIKLARI ÜZERİNE YAPILAN BİLİMSEL YAYINLARIN BİBLİYOMETRİK ANALİZİ*	205
Faruk Emre BOZEL, Hale AVŞAR ERDEKLİ *EVALUATION OF ERGOGENIC SUPPLEMENT USE AND EXERCISE ADDICTION IN ADULTS ENGAGED IN EXERCISE*	219
Elif İrem AKDOĞAN, Eftal ŞEHİRLİ *DETECTION AND CLASSIFICATION OF WHITE BLOOD CELLS – A REVIEW*	231
Jale BERRAK, Ali KOÇ, Özkan KÖSE, Yıldız KOÇ *THERMODYNAMIC PERFORMANCE EVALUATION OF THE CO2/NH3 MİXTURE IN SEVERAL FRACTIONS*	242

Tansu ÜNEN, Kenan YILDIRIM	
THE AFFCET OF THE WOOL FIBER TYPE ON THE YARN TENSILE STRENGTH	**252**
Nəzakət ƏBILOVA	
HISTORICAL FORTIFICATIONS OF THE ASTARA PROVINCE OF AZERBAIJAN	**268**
Samet GÜRGEN	
ARTIFICIAL NEURAL NETWORK MODELS IN SHIP MAIN ENGINES: CURRENT APPLICATIONS AND FUTURE PERSPECTIVES	**276**
Semra TETİK	
THE RELATIONSHIP BETWEEN CAREER STRESS AND HOPELESSNESS: A STUDY ON UNIVERSITY STUDENTS	**287**
Semra TETİK	
A RESEARCH ON THE RELATIONSHIP BETWEEN SOCIAL INTELLIGENCE AND ENTREPRENEURSHIP TENDENCY	**297**
Erdi TOSUN, Çağrı ADIKTI	
COMPARISON OF AIR COOLING CONFIGURATIONS FOR LITHIUM-ION BATTERY THERMAL MANAGEMENT	**308**
Raviye ÖZEN KOCA	
AGOMELATINE AND PHYSIOLOGICAL SYSTEMS: AN INTEGRATIVE APPROACH TO CORE MECHANISMS	**314**
E. CUCE, P. MERT CUCE, T. GUCLU	
HYBRID PARABOLIC TROUGH COLLECTOR AND THERMOELECTRIC GENERATOR SYSTEMS FOR MULTIGENERATION	**319**
Barış ERGÜL, Arzu ALTIN YAVUZ	
COMPARATIVE PERFORMANCE ANALYSIS OF ARIMA, SARIMA, AND LSTM MODELS FOR AIR QUALITY INDEX FORECASTING	**324**
Barış ERGÜL, Arzu ALTIN YAVUZ	
MACHINE LEARNING-BASED MODELING OF DROUGHT PROCESSES IN TÜRKİYE	**335**
E. CUCE, P. MERT CUCE, T. GUCLU	**341**
SMART ENERGY SYSTEMS FOR SUSTAINABILITY-A REVIEW	
Ayça ÖZTÜRK, Ahmet KATILMIŞ	**347**
SOSYAL BİLGİLER DERSİNDE SİNEKTİK TEKNİĞİNİN KULLANIMI	
Cavit Yavuz, Hazal Zaimoğlu	
TARAFSIZLIK MASKESİ ALTINDA DİJİTAL TARAFGİRLİK: GROK VE FİLİSTİN SÖYLEMİ	**360**
Hülya ŞEREFLİŞAN, Ahmet ALKAYA	
INNOVATIVE APPROACHES TO WATER QUALITY MANAGEMENT THROUGH THE GENETIC POTENTIAL OF FRESHWATER MUSSELS	**370**
Olumide Abraham Ajayi	
IMPACT OF TERRORISM AND BANDITRY ON WOMEN DEVELOPMENT IN NORTHERN NIGERIA	**389**
Caglar SOYLU	
THE ACUTE EFFECTS OF EXERCISE-INDUCED FATIGUE ON HAMSTRING MUSCLE ARCHITECTURE AND CORTICOSPINAL EXCITABILITY: AN EXPERIMENTAL STUDY IN HEALTHY INDIVIDUALS	**396**

Caglar SOYLU *THE RELATIONSHIP BETWEEN MUSCLE ARCHITECTURE, CORTICOSPINAL EXCITABILITY, HAMSTRING OUTCOME SCORE, AND ECCENTRIC STRENGTH IN ADULTS WITH PRIOR HAMSTRING INJURY: PILOT STUDY*	405
Dhanraj T. Dhangar *THE SECULAR TADAVI BHIL TRIBE: HISTORY AND CULTURE*	415
Tuna ŞENER *A COMPARATIVE ANALYSIS OF THE RISK INDICATORS OF THE TURKISH ECONOMY WITH THE GREECE ECONOMY*	421
Ummühani USLU *YAŞLI KADINLARIN SOSYO-EKONOMİK SORUNLARININ SOSYAL HİZMET BAĞLAMINDA TARTIŞILMASI*	432
Cüneyt Yılmaz *UNORGANIZED CROWD BEHAVIOR: A SOCIAL PSYCHOLOGICAL ANALYSIS*	439
Ananda Majumdar *VALLABHBHAI PATEL: ARCHITECT OF INDIAN UNITY AND MODERN GOVERNANCE*	448
Banjo, Tolani Christianah *THE CULTURE OF WASTE MANAGEMENT: RETHINKING TRADITIONAL VALUES AND MODERN REALITIES IN NIGERIA*	455
Metin Can TORUN, Davut BAYRAM, Hürü Ceyda TÜM *EVALUATION OF HOUSING, NUTRITION, AND BEHAVIORAL REQUIREMENTS OF WORKING DOGS IN MILITARY AND LAW ENFORCEMENT UNITS: RECOMMENDATIONS FOR IMPROVEMENT*	461
Zeynep GÜMRÜKÇÜ *REPAIR OF ALVEOLAR DEFECT WITH TITANIUM MESH AND SIMULTANEOUS DENTAL IMPLANT PLACEMENT: A CASE REPORT*	470
Aktolkyn BORANBAYEVA, Gelişli, Yücel, Assel TASSOVA *FORMATION OF THE FUTURE SPECIALIST'S CREATIVITY*	477
Rezzan UYSAL, Ezgi TÜRKAY *THE EFFECT OF PARENTAL FOOD NEOPHOBIA ON CHILD NEOPHOBIA AND EVALUATION OF ANTHROPOMETRIC MEASUREMENTS*	485
Özge ÇETİNARSLAN *D-DIMER/ALBUMIN, CRP/ALBUMIN, AND THEIR COMPOSITE INFLAMMATORYCOAGULATION INDEX PREDICT READMISSION AFTER NSTE-ACS*	492
Murat DAKMANCI, Mehmet AVŞAR, Yusuf ÖZEL, Ayşe DEMİR *FIVE OLYMPICS, FIVE MODELS: THE ECONOMIC ANATOMY OF THE GAMES FROM 2008 TO 2024*	498
Vala Nadeşoğlu, Alireza Souri, Nihat İnanç *A RANDOM FOREST-BASED ANOMALY DETECTION APPROACH IN REAL-TIME IOT NETWORKS*	508
Vala Nadeşoğlu, Alireza Souri, Mostafa Habibi *HYBRID INTRUSION DETECTION IN SDN-IOT NETWORKS USING DEEP NEURAL NETWORK AND RANDOM FOREST*	519

Enise Çiçek YILDIRIM, Sinan KÖSE, Salih Hakan YETGİN ***EVALUATION OF TRIBOLOGICAL BEHAVIOURS OF GRAPHITE/COC AND CARBON BLACK/COC COMPOSITES WITH TAGUCHI METHOD***	**530**
Büşran SUNYAR, Mesude Figen DÖNMEZ ***BIBLIOMETRIC PROFILE OF STUDIES ON BACTERIAL CANCER IN STONE FRUIT TREES***	**542**
Su Lal CANSEV, Hatice Kübra ARAS ***THE EFFECT OF CHANGED DIETARY HABITS DUE TO PREMENSTRUAL SYNDROME ON HEDONIC AND EMOTIONAL EATING BEHAVIOR***	**555**
Veli ÇAPALI, Murat KAÇAL, Burak DEMİR ***INTEGRATION OF UNIVERSITY SOFTWARE APPLICATIONS WITH CENTRAL IDENTITY MANAGEMENT SYSTEM AND INTEGRATED PORTAL DESIGN EXAMPLE, SDUNET***	**568**
Lamiye NEBİYEVA ***THE SYSTEM OF IMAGES IN ARTISTIC LANGUAGE AND THE STYLIST-ARTISTIC FUNCTION (BASED ON THE CREATIVITY OF NAKHCHIVAN WRITERS)***	**578**
Harun Yıldırım, Murtaza Kaya ***A RARE CASE OF KOUNIS SYNDROME TYPE II PRESENTING WITH ST-ELEVATION MYOCARDIAL INFARCTION TRIGGERED BY BEE STING***	**586**
Sadikhova F.Q. ***THE ISSUE OF SOUL AND BODY IN EUROPEAN PHILOSOPHY OF THE NEW AND MODERN ERA***	**592**
Zeynep Başaran ***PEYGAMBERİMİZİN MUCİZELERİNDEN OLAN NESEBİ DURUMU***	**596**
Didem ALTUN ***THE INVESTIGATION OF THE OPTICAL PROPERTIES OF AlGaAs/GaAs MULTIPLE QUANTUM WELLS UNDER THE INFLUENCE OF ELECTRIC AND MAGNETIC FIELDS***	**601**
Hetal Pravinsinh Jadeja, Jatinkumar Bhimjibhai Kotadiya ***ARTIFICIAL INTELLIGENCE IN ACTION: A COMPARATIVE STUDY OF ML ALGORI THMS FOR TEXT CLASSIFICATION***	**609**
Jatinkumar Bhimjibhai Kotadiya, Dharmendra Jayantilal Ambani, Amit K. Patel ***A COMPARATIVE STUDY OF MACHINE LEARNING ALGORITHMS FOR SENTIMENT ANALYSIS USING ARTIFICIAL INTELLIGENCE TECHNIQUES***	**618**
Sameer Suresh Nanivadekar ***BUILDING A GLOBAL CAREER WHILE PURSUING AN ENGINEERING DEGREE: STRATEGIES FOR STUDENTS WORLDWIDE***	**628**
E.Elavarasi, Yuvaraj AR, J.Jayadurka, R.Srinivasan ***DRUG FORMULATION AND DELIVERY SYSTEMS***	**632**
R. Devi, V. Dhanapal ***TRANSCRANIAL ULTRASOUND COMBINED WITH LIPID NANOPARTICLES ADVANCES BRAIN DELIVERY OF SOD1 ANTISENSE FOR ALS TREATMENT***	**647**

R. Devi *TARGETED NANOTECHNOLOGY APPROACHES TO BYPASS THE BLOOD–BRAIN BARRIER IN NEURODEGENERATIVE DISORDERS*	652
SADIQ Mohammed Sanusi *RE-IMAGINING AGRICULTURAL POLICY THROUGH BAGONOMICS: TOWARDS INCLUSIVE AND EQUITABLE RURAL DEVELOPMENT IN NIGER STATE*	654
Puniparthi Sunitha, M.K. Vijiayalakshmi *REAL-TIME THERAPEUTIC DRUG MONITORING USING WEARABLE DEVICES*	668
Puniparthi Sunitha, M.K. Vijiayalakshmi *THE ROLE OF GUT MICROBIOTA IN CHRONIC DISEASES*	682
Puniparthi Sunitha, M.K. Vijiayalakshmi *3D BIOPRINTING IN ORGAN TRANSPLANTATION: HOPE OR HYPE?*	696
N.G. Kalyani Reddy, Chayaraju BalaSai, K.Tharani *ROBOTS IN CONSTRUCTION: ENABLING TECHNOLOGY, APPLICATIONS, AND CHALLENGES*	707
Amadi Oko Amadi and Levi Nnaemeka Osuji *EFFECT OF QUANTUM ANNEALING AND D-WAVE ON THE OPTIMIZATION OF ENHANCED OIL RECOVERY AND PRODUCTION FRAMEWORK*	719
N.G. Kalyani, Chayaraju BalaSai, K.Tharani *INTEGRATING AUGMENTED AND VIRTUAL REALITY TECHNOLOGIES IN ARCHITECTURAL CIVIL ENGINEERING*	731
Hassan Abdulmumini *AGRICULTURE GLOBALIZATION AND HOW IT AFFECTS FOREING RELATION*	739
Peter Olubunmi Olubiyo, Jude Timilehin Olubiyo *EXTENT OF USE OF DIGITAL RESOURCES AMONG PHYSICS AND CHEMISTRY FINAL YEAR UNDERGRADUATE STUDENTS IN ADEYEMI FEDERAL UNIVERSITY OF EDUCATION, ONDO, NIGERIA*	744
Zeynep YAVUZ, Zeynep Nur SANDIKCI *EFFECTIVE PERSONAL FINANCE MANAGEMENT WITH SMART ASSISTANT*	753
Ersin YAVUZ, Tahsin AVCI *EXAMINING THE ECONOMIC PERFORMANCE OF OECD COUNTRIES FROM A TAX BURDEN PERSPECTIVE WITH THE PIV METHOD*	759
Ersin YAVUZ, Tahsin AVCI *REGIONAL ANALYSIS OF SOCIO-ECONOMIC AND FISCAL PERFORMANCE: AN IMPLEMENTATION ON TÜRKİYE NUTS LEVEL-2*	773

ADVANCED DESIGN AND OPTIMIZATION STRATEGIES FOR WEARABLE HEALTH DEVICES FOR CHRONIC DISEASE MANAGEMENT

Chandani Sharma, Ghazala Ansari , Aarti Sharma, Manidipa Roy

Associate Professor, Deptt of CSE, MMICTBM (MCA), Maharishi Markandeshwar (Deemed to be University), Mullana-Ambala, Haryana, India.

Assistant Professor , Deptt of ECE, SRM Institute of Science and Technology, Modinagar, NCR Campus

Assistant Professor, ABES Engineering College, Ghaziabad

Professor, ECE Department, ABES Engineering College

ABSTRACT

This paper explores the design and optimization of wearable health devices (WHDs) for chronic disease management, focusing on the intersection of mechanical, electronic, and software systems used in these WHD's. Wearable health technologies, such as continuous glucose monitors, fitness trackers, and smartwatches, are becoming indispensable tools for patients with chronic conditions like diabetes, cardiovascular disease, and respiratory disorders. The paper addresses the critical design components that impact WHD's performance, including sensors, data processing units, communication technologies, and power management systems. Further, Optimization techniques such as model-based design, energy-efficient algorithms, and advanced data fusion methods are examined to highlight the role in enhancing device accuracy, reliability, and user compliance for devices. A detailed approach on how AI and machine learning are integrated into wearable devices to provide real-time health insights and predictive analytics, enabling personalized care and early intervention for chronic disease patients is explored with appreiciable SWOT Analysis.

Keywords: WHD's, sensors, data processing units, communication technologies, and power management systems, predictive analytics, SWOT Analysis

I. Introduction

The integration of wearable health devices (WHDs) into the management of chronic diseases has sparked a revolution in healthcare. Chronic conditions such as diabetes, hypertension, and heart disease require constant monitoring, which is often burdensome and costly. WHDs offer a promising solution by providing real-time data collection, continuous monitoring, and feedback to patients and healthcare providers. These devices, which include continuous glucose monitors (CGMs), smartwatches with heart rate and ECG monitors, and other biofeedback systems, play a vital role in empowering patients to manage their conditions more effectively and prevent complications.

The design of these devices must overcome several challenges, including miniaturization, accuracy, power efficiency, user compliance, and seamless integration with the healthcare ecosystem. This paper outlines key design principles and optimization techniques that improve the functionality and usability of WHDs, with a focus on their application in chronic disease management.

II. Wearable SMART devices availible in Market

Smart wearable devices are gadgets that are worn on the body and are designed to provide various functions such as health tracking, notifications, communication, and entertainment, often with connectivity to smartphones or other devices. AI and machine learning algorithms play a critical role in enhancing the capabilities of smart wearable devices.These devices include smartwatches, fitness trackers, smart glasses, smart clothing, and more, providing a range of functionalities such as health monitoring, activity tracking, and real-time data analysis. These are shown below:

FULL TEXTS BOOK




Fig 1: SMART Wearable Devices a) SMART Watch b) Fitness Tracker c) SMART Glass d) SMART Ring e) SMART Clothing f) SMART eaphones g) SMART Patches

III. Design Principles for Wearable Health Devices

The success of WHDs in chronic disease management depends on their design and how well the individual components function together. Wearable health devices are designed with a focus on comfort, accuracy, power efficiency, and real-time data processing. Key design principles include the use of lightweight, biocompatible materials such as medical-grade silicone or thermoplastic elastomers to ensure long-term wearability.

Sensors play a critical role, with photoplethysmography (PPG) for heart rate monitoring, electrodermal activity (EDA) sensors for stress detection, and accelerometers for activity tracking. Low-power microcontrollers and Bluetooth Low Energy (BLE) connectivity are essential for efficient data transmission while preserving battery life, with rechargeable lithium-polymer (Li-Po) or coin cell batteries often used. Devices such as the Apple Watch and Fitbit leverage AI-driven algorithms to enhance accuracy, filtering out noise from movement artifacts. Moreover, integration with cloud platforms and mobile applications allows for seamless real-time monitoring and predictive analytics, enhancing personalized healthcare interventions. Key design principles include flow chart represented in Fig 2.




Fig 2: Block Diagram of a Wearable Health Device (WHD)

FULL TEXTS BOOK

The description of block diagram is listed below:

1. Sensors and Sensing Technologies

The primary function of any WHD is accurate health monitoring. The choice of sensors is critical to ensure reliable data collection. Common sensors include:

Optical Sensors: These sensors are used in devices like pulse oximeters and smartwatches to measure vital signs such as heart rate, blood oxygen saturation, and pulse.

Electrochemical Sensors: Widely used in CGMs (Continuous Glucose Monitors), sensors detect the levels of glucose in the interstitial fluid, providing continuous feedback for diabetes management.

Bioelectrical Sensors: Used in ECG monitors and smartwatches to track the electrical activity of the heart, these sensors help manage cardiovascular diseases.

Accuracy and sensitivity are paramount for these sensors, as incorrect readings can result in adverse health outcomes. Furthermore, sensors need to be minimally invasive and capable of providing long-term, real-time data to ensure continuous monitoring.

2. Data Processing and Communication Technologies

Data processing units handle the signals collected from sensors, convert them into meaningful insights, and ensure that they are transmitted efficiently. Data must be processed in real-time to enable immediate intervention in case of abnormal readings. Optimization techniques include:

Edge Computing: Moving some data processing to the device itself reduces the load on cloud servers and enables faster response times, which is crucial for chronic disease management.

Low-Power Wireless Communication: Technologies such as Bluetooth Low Energy (BLE) and Zigbee are commonly used to transmit data from the wearable to a smartphone or cloud-based platform for analysis and storage.

Communication technologies must be reliable, ensuring that the data sent from the wearable device is not lost or delayed, which is crucial in cases where timely interventions are needed.

3. Power Management and Miniaturization

Power management is one of the most challenging aspects of wearable health devices. Chronic disease management requires continuous data collection, so energy-efficient algorithms and power-saving technologies must be employed. Design considerations for effective power management include:

Low Power Consumption Sensors: Sensors must be designed to operate efficiently, using as little power as possible while maintaining accuracy.

Energy-Efficient Processors: Processors must be optimized to handle data processing without draining the device's battery.

Long-Lasting Batteries: Batteries need to last for extended periods, ideally for days or weeks, to avoid the need for frequent recharging, which can hinder user compliance.

Miniaturization techniques play a crucial role in ensuring that WHDs are comfortable, lightweight, and unobtrusive for continuous wear. As technology advances, materials such as flexible and stretchable electronics are becoming more common in wearable designs, allowing devices to fit comfortably while maintaining functionality.

IV. Design and Optimization Strategies used

To enhance the performance, accuracy, and user experience of WHDs, several design and optimization techniques are employed. Advanced design and optimization techniques are used in wearable health devices to improve user experience, accuracy, and efficiency. High-precision sensors like photoplethysmography (PPG) for heart rate monitoring and accelerometers for activity tracking are important design factors, as are the use of lightweight, biocompatible materials for comfort and durability. Power efficiency is the main emphasis of optimization strategies, which use Bluetooth Low

FULL TEXTS BOOK

Energy (BLE) connectivity, low-power microcontrollers, and energy-efficient algorithms to prolong battery life.

Cloud integration allows for real-time monitoring and predictive analytics, while AI-driven data processing helps reduce noise and increase measurement accuracy. These techniques, which are found in gadgets like the Fitbit and Apple Watch, guarantee consistent, dependable health tracking, promoting early disease identification and individualized treatment. The strategies could be summarised as under :

1. Model-Based Design

Model-based design (MBD) is a crucial approach in developing WHDs. MBD allows for virtual simulations and modeling of the device components before physical prototypes are created. This reduces design time, improves accuracy, and minimizes manufacturing errors. MBD helps optimize the integration of sensors, data processing units, and communication systems to ensure that the wearable functions optimally within the limitations of size, power, and user comfort.

2. Energy-Efficient Algorithms

Power consumption is a significant challenge for wearable health devices, especially those intended for continuous monitoring. Optimizing algorithms for energy efficiency is critical to extending the device's battery life without compromising data accuracy. Techniques include:

Data Compression: Reducing the amount of data transmitted without losing crucial information.

Adaptive Sampling: Altering the frequency of data collection based on the patient's condition or real-time needs, thus saving power when continuous monitoring is unnecessary.

3. Machine Learning and Predictive Analytics

AI and machine learning algorithms can be integrated into wearable devices to offer real-time health insights and predictive analytics. For example:

Personalized Health Insights: By analyzing past health data, machine learning algorithms can predict trends and offer personalized recommendations for disease management, such as adjusting medication or exercise levels.

Predictive Analytics for Early Intervention: AI can be used to detect early signs of health deterioration, such as abnormal glucose levels or irregular heart rhythms, allowing for timely intervention before complications arise.

Machine learning models can be continuously trained with new data, improving their accuracy and reliability over time. This transforms wearable health devices from simple monitoring tools into proactive health management systems.

V. User-Centered Design and Ergonomics

The long-term success of wearable health devices depends on user compliance. A device that is uncomfortable, difficult to use, or inconvenient will not be used regularly.

To ensure that patients adhere to their health monitoring routines, several design principles must be considered like Comfort and Wearability, ensuring that they can be worn throughout the day without causing discomfort. This includes using lightweight materials, breathable fabrics, and ergonomic shapes. Second factor is Intuitive User Interfaces, essential to ensure that patients can easily interact with the device. For instance, a device for elderly users with diabetes should have large buttons, clear displays, and audible notifications. Thirdly, Data Accessibility and Integration integrated with broader health management systems. This means that data collected by the device should be easily accessible to both patients and healthcare providers. Integration with cloud-based systems or electronic health records (EHRs) allows healthcare professionals to remotely monitor patients' health and intervene when necessary.

To analyze user-centered design and ergonomics in wearable health devices, a numerical analysis is done by comparing different devices based on these parameters and visualize the results with a graph. Based on sample collection and availible market products Table 1 depicts an overview of device details.

FULL TEXTS BOOK

Table 1: Device Details

Device	Weigh t (g)	Battery Life (hrs)	Sensor Accuracy (%)	Comfort Rating (/10)
Apple Watch	38	18	95	8.5
Fitbit Charge	29	96	92	9
Garmin Vivosma rt	24	120	90	9.2

The visualization of above statistics is carried out and reflected in bar chart Fig. 3 that compares key ergonomic factors of different wearable health devices, including weight, battery life, sensor accuracy, and comfort rating.




Fig 3: Comparison of SMART Watches using Ergonomic factors

It can be seen that The Garmin Vivosmart has the longest battery life and highest comfort rating but slightly lower sensor accuracy. The Apple Watch offers the highest accuracy but has a shorter battery life. The Fitbit Charge balances comfort and battery life well. This analysis highlights the trade-offs between weight, battery performance, and user comfort in wearable health device design.

VI. Conclusion and Future Directions

The management of chronic diseases is being revolutionized by wearable health devices, which provide real-time information, individualized therapy, and continuous monitoring. The future of healthcare will be greatly influenced by the continuous optimization of these devices—through improved sensor accuracy, energy-efficient power systems, AI-driven analytics, and user-centered design. WHDs will become increasingly complex as technology develops, providing increased accuracy, smoother integration with medical ecosystems, and easier-to-use user interfaces. Through the use of advancements in software, mechanical, and electronic systems as well as new technologies like flexible electronics and AI-powered diagnostics, WHDs will continue to improve patient outcomes, raise quality of life, and be essential to proactive healthcare management.

REFERENCES

[1] J. Kim, A. S. Campbell, B. E.-F. de Ávila, and J. Wang, "Wearable Biosensors for Healthcare Monitoring," Nature Biotechnology, vol. 37, no. 4, pp. 389–406, Apr. 2019.

FULL TEXTS BOOK

[2] M. Stoppa and A. Chiolerio, "Wearable Electronics and Smart Textiles: A Critical Review," Sensors, vol. 14, no. 7, pp. 11957–11992, Jul. 2014.

[3] S. Patel, H. Park, P. Bonato, L. Chan, and M. Rodgers, "A Review of Wearable Sensors and Systems with Application in Rehabilitation," Journal of NeuroEngineering and Rehabilitation, vol. 9, no. 1, p. 21, Apr. 2012.

[4] A. Pantelopoulos and N. G. Bourbakis, "A Survey on Wearable Sensor-Based Systems for Health Monitoring and Prognosis," IEEE Transactions on Systems, Man, and Cybernetics, Part C (Applications and Reviews), vol. 40, no. 1, pp. 1–12, Jan. 2010.

[5] S. C. Mukhopadhyay, "Wearable Sensors for Human Activity Monitoring: A Review," IEEE Sensors Journal, vol. 15, no. 3, pp. 1321–1330, Mar. 2015.

[6] A. M. Khan, Y.-K. Lee, S. Lee, and T.-S. Kim, "A Triaxial Accelerometer-Based Physical-Activity Recognition via Augmented-Signal Features and a Hierarchical Recognizer," IEEE Transactions on Information Technology in Biomedicine, vol. 14, no. 5, pp. 1166–1172, Sep. 2010.

[7] M. A. Hanson et al., "Body Area Sensor Networks: Challenges and Opportunities," Computer, vol. 42, no. 1, pp. 58–65, Jan. 2009.

[8] C. C. Y. Poon, B. P. L. Lo, M. R. Yuce, A. Alomainy, and Y. Hao, "Body Sensor Networks: In the Era of Big Data and Beyond," IEEE Reviews in Biomedical Engineering, vol. 8, pp. 4–16, 2015.

[9] Chandani Sharma, Anzar Ahmed, Sonika Khandari, Amit Saini “Cloud-Based Smart Cultivation System Using IoT Data Analytics” 5th IEEE International Conference on Computing, Power and Communication Technologies (IC2PCT) on 9-10 February 2024, organized by Galgotia’s University, Greater Noida, India.

[10] Chandani Sharma, Manish Sharma, Shilpy Sharma, “ Creation of an Improved Software Testing Framework for the Cloud Environment”, International Conference on Fourth Industrial Revolution based Technology and Practices ‘(ICFIRTP-2022)" on Nov 23-Nov 24, 2022 in association with IEEE Xplore at Quantum University, Roorkee.

A COMPREHENSIVE ASSESSMENT OF THE APPLICATION OF OPERATIONAL RESEARCH METHODS TO THE DESIGN OF CONCRETE MIXES

***ChayarajuBalaSai*[1]**

ORCID ID: 0009-0003-4055-9773

Assistant Professor Department of Civil Engineering,G.pulla reddy Engineering college(Autonomous),Kurnool, AP, INDIA

***Nadagouda Kalyani*[2]**

Assistant Professor Department of Civil Engineering,G.pulla reddy Engineering college(Autonomous),Kurnool, AP, INDIA

***K.Tharani*[3]**

Assistant Professor Department of Civil Engineering,G.pulla reddy Engineering college(Autonomous),Kurnool, AP, INDIA

ABSTRACT

Conventional approaches to concrete mixture design yield good results, but they do not ensure the best possible composition. As a result, the growing necessity for designers to balance the basic elements of concrete to meet performance criteria such mechanical qualities, chemical properties, workability, sustainability, and cost is driving the use of operational research approaches. Because of this, a lot of writers have been searching for machine learning and mathematical programming ways to forecast the characteristics of concrete mixes and optimize them. The following is a thorough analysis of operational research methods related to concrete mixture design and proportioning, along with a classification system.

Keywords: Concrete Mix Design, Machine learning, Mathematical Optimaization.

1. Introduction

Concrete is one of the most commonly used construction materials in the world. Due to its unique feature combination of work- ability, mechanical properties, durability, and the wide availability of raw materials sources, concrete has become highly attractive for many applications in the civil industry [1]. Generally speaking, concrete can be defined as the mixture of a cement paste and water with fine and coarse aggregates. However, a concrete mix can be way more complex and have a wide range of constituents. It can be combined with materials, such as cement, pozzolans, fly ash, blast-furnace slag, silica, recycled concrete aggregate, polymers, fibers, and superplasticizers. Due to the vast possibility of constituents and the different variations in composition, it is possible to obtain a concrete mixture with different physical and chemical properties. Therefore, varying the components and their quantities allows the properties of both fresh and hardened states to be tailored to achieve the design specifications.

Each concrete application in the construction sector will require a specific mix composition to satisfy the performance re- quirements, such as workability, durability, and compressive strength. Therefore, designing good concrete is utterly vital to guarantee that the mixture will achieve the required specifications. In order to meet the best composition of concrete, designers have adopted concrete mix design to find an optimal concrete mixture. This procedure is defined as selecting suitable raw materials and propor- tioning the quantities of each constituent in order to produce a final product that meets the desired physical and chemical properties [2]. For this reason, promising development in the concrete mix design step can guarantee improvements in physical-mechanical performance, more outstanding durability of structures, and a reduction in production costs.

FULL TEXTS BOOK

Most of the available mix design methods are based on empirical relationships, charts, and graphs developed from experimental investigations. They follow the same principles, and only minor variations exist in the mix design methods in selecting the mix pro- portions [3]. Some traditional methods for proportioning concrete are the American Concrete Institute (ACI) mix design method and the British mix design method [4]. According to DeRousseau [1], the conventional mix proportion classification can be broken down into two main methods: prescriptive specification and performance-based specification. The main difference between the two methods is that the first follows a step-by-step procedure to determine the quantity of each component. On the other hand, the second method does not require strict guidance. According to the design specification, a value of a required property is established; then, the designer can select any amount of cement, water, and aggregates and verify that the mixture achieves the value of the required property.

Although these traditional methods are way useful, the mathematical relationship between variables and the mixture composition can have some drawbacks. The first drawback of current concrete design methods is the lack of contemplation of Supplementary Cementitious Materials' influence (SCMs) on concrete properties. Since SCMs have been used lately to replace a portion of cement, they are prone to dramatically impact mechanical properties such as compressive strength [4]. Another drawback is that the mix design does not capture the true complexity of the relationship between the mixture design and the concrete's properties. The type and quantities of each material used are variable, and the estimation of these properties cannot be quantitatively and precisely evaluated. Lastly, the current concrete design methods do not present optimised values of concrete properties. The methods meet a satisfactory composition of the concrete constituents that reach the desired values of the properties. However, choosing a specific design does not guarantee that it is not an optimal concrete mix solution.

Besides the traditional methods, some research lines encompass the design of experiments to identify a good concrete mix composition. Compared to traditional methods, the application of the design of experiments can exhibit better solutions due to the numerous experiments. Although several studies have been investigating the problems related to concrete mix design, the optimi- sations of experimental design can undergo exponential growth if they increase the number of variables and their values [1]. Furthermore, due to the high degree of non-linearity between the dependent and independent variables, the regression models ob- tained to predict the properties of the concrete can hardly present a precise regression equation [5].

Therefore, both procedures have disadvantages and do not meet an optimised mix composition. For this reason, the computational aid system has been studied over the years in order to solve the problems regarding concrete mix design [6]. It formulates the problem by more accurately modelling the relationships between variables and the output response, leading to an optimal solution [7]. Thus, mathematical programming and machine learning have been prominent in the literature among the operational research techniques used to solve mixture proportioning issues. Both methods can be employed separately or as complementary approaches, frequently used to solve various engineering problems. Considering this fact, many researchers look to predict the properties of concrete in a more accessible, less costly, and less time-consuming way [8]. On the other hand, others seek to develop a more economical and sustainable concrete mix [9].

Several papers in the literature reviewed mathematical models and machine learning techniques in predicting concrete properties or the optimal composition of concrete. Chaabene et al. [10] examined machine learning (ML) models to forecast the mechanical properties of concrete. Another work presented a systematic review of the ML algorithms employed to predict compressive strength [11]. Besides that, Song et al. [12] reviewed the current literature on optimising mixing ratios using ML and metaheuristic algorithms, which offers insight into the continuous development of models in the field of hybrid optimisation. Based on these previous works, this article aims to examine relevant articles that have solved the concrete mix design problem using two types of operational research techniques: machine learning and mathematical programming. The novelty of this article lies in the lack of review articles focusing on these two techniques applied to the concrete mix design, which can bring a holistic understanding of the employment of these tools. Additionally, this article intends to propose a classification framework with the main features of the last works using mathematical programming or machine learning to solve problems with concrete mix design.

FULL TEXTS BOOK

To place things into context, three major questions are addressed in the paper:

1. What are the main machine learning algorithms that have been used in order to come up with the concrete mix design?

2. What are the mathematical optimisation algorithms that have been used in order to help concrete mix design?

3. How is the literature linking both aspects, mathematical optimisation and machine learning, in order to design a concrete mix?

The rest of the paper is organised as follows: Section 2 briefly describes the concrete mix design research background. Next, section 3 describes the adopted study review methodology, presenting a systematic review and proposing a classification framework. The following section, section 4 evaluates the concrete mix design studies. Next, section 5 brings a discussion of the general findings. Finally, the final section contains concluding remarks on the general findings and research proposals.

2.Background

In order to change one or more of the qualities of the concrete in its fresh or hardened condition, additives are substances that are added either before or during the mixing process.Admixtures can have physical or chemical effects. The amounts of admixtures put to concrete are minimal when compared to the amounts of the normal constituents. Admixtures are often added to concrete mixes in amounts of 5% of the mass of cement. The amount of admixtures added to concrete can range from 0.5% to 10% of the mass of cement, depending on how the concrete is applied.Comparing the weight of all the constituents and the overall composition of the concrete, the admixtures' composition is between 0.02% and 0.5%.Dransfield (2003) Admixtures are defined by US Per IS9103:1999 as any material other than water, aggregates, hydraulic cement, and additives (slag, fiber reinforcement, and/or pozzolana). Chemical or mineral admixtures are the two types of additives used in concrete. The fresh state qualities are altered at a minimal cost by mineral admixtures, often known as US additives. The type of additive employed determines how they affect the properties of concrete in its hardened condition (Gandage et al., 2014).Admixtures have been used in concrete since the Before describing the proposed classification framework, some essential aspects of concrete mix design are summarised. Then, Section 2.1 and Section 2.2 outline mathematical optimisation models and machine learning methods adopted as solution strategies for the concrete mix design.

1.1. Overview of mathematical programming techniques

Mathematical models and computational optimisation methods are alternative solutions in concrete mix design to tackle the traditional methods and the time-consuming laboratory experiment optimisations. While the optimisation methods bring an optimal or near-optimal solution with low computational effort, the others only lead the designers to an ideal solution, which cannot be the most improved. These approaches initially establish one or more objectives (e.g., concrete properties) as functions of the decision variables (e.g., concrete constituents). Then, they search for the optimal concrete mixture by adopting optimisation algorithms. Mathematical methods can be classified into meta-heuristic, heuristic, and exact methods. Most researchers have frequently implemented them in the concrete mix design field to deal with mix design issues.

1.1.1. Exact methods

Exact methods have been used for a long time in mathematical optimisations and are very effective in modelling and optimising the properties of concrete. We can elucidate some model examples among the exact methods, such as linear [13], non-linear [14], and mixed programming [15]. Many authors have applied exact methods to estimate the optimal composition or predict other concrete properties. For instance, Jin et al. [15] proposed non-linear and mixed regression models to predict concrete strength based on mixture-design variables and curing age. The proposed model achieved higher accuracy than the linear method using the same var- iables and datasets. Another study that used non-linear programming was

FULL TEXTS BOOK

Habibi and Ghomashi's work [16]. Two mathematical models were developed based on sequential quadratic programming to optimise the self-compacting concrete mix design, minimising cost with compressive strength and slump as constraints. Miller et al. [17] developed a linear method for predicting Global Warming Potential (GWP) from concrete production and compressive strength for concretes containing replacement binder. The work proved the equations were suitable for predicting compressive strength and GWP for mixtures containing replacement binder. Fan and Miller

[18] also took environmental problems into account. They developed a set of mathematical equations to find minimum Greenhouse Gas (GHG) emissions and to dictate the required concrete mix proportions, based on water-to-binder ratio and supplementary material, to achieve optimised GHG emissions. Li et al. [19] proposed a numerical simulation algorithm for calculating the overall elastic properties of ordinary concrete, which involves the finite element method in combination with Monte Carlo simulations. Comparisons with experimental test results of elastic modulus and Poisson's ratio indicated a good agreement with the proposed model.

1.1.2. Heuristic and metaheuristic methods

Heuristic and metaheuristic methods are higher-level procedures aiming to find, develop, or select an algorithm that may provide a sufficiently good solution to an optimisation problem, especially with incomplete information [20]. While many heuristic algorithms are very specific and problem-dependent, a metaheuristic is a high-level problem-independent algorithm. Regarding concrete mix design issues, many studies have been using different heuristic metaheuristic methods to predict and optimise the properties of concrete. For example, Lee et al. [21] applied Harmony Search (HS) algorithm to find the optimum mix proportion satisfying the required strength and slump. The results showed that the HS algorithm could be a potent tool for high-performance concrete mix proportioning compared with other methods. Moayedi et al. [22] investigated the efficiency of the Ant Lion Optimisation (ALO) al- gorithm to predict the concrete slump. Besides, they compared it with Biogeography-Based Optimisation (BBO) and Grasshopper Optimisation Algorithm (GOA). The findings revealed that ALO outperformed both models. Another study [23] applied an Artificial Bee Colony (ABC) algorithm to predict chloride penetration in self-consolidating concretes. Comparisons among ABC and Genetic Algorithm (GA) and Particle Swarm Optimisation (PSO) were conducted and showed higher reliability [23].

Among the metaheuristic methods used to solve concrete mix design problems, many studies applying a genetic algorithm or some variation of swarm optimisation are notorious. For example, to improve the anti-cracking performance of high-strength concrete, Yue et al. [24] used GA to optimise concrete mix proportioning by minimising the cracking risk coefficient. Comparisons between a control concrete and the optimised one showed that crack resistance was considerably improved, and a reduction of 25% of the crack risk coefficient was achieved. Furthermore, Awoyera et al. [25] modelled the compressive strength, split tensile strength, and flexural strength of geopolymer concrete using gene expression programming, which showed a good prediction.

Regarding the swarm optimisations, Mashhadban et al. [7] investigated fibers' effects on self-compacting concrete properties using a hybrid model composed of PSO and a neural network. The model predicted splitting tensile strength, flexural strength, compressive strength, and fracture energy with a high level of accuracy. On the other hand, Kandiri et al. [26] estimated the compressive strength of the ground granulated blast furnace slag concrete by applying a multi-purpose Salp Swarm Algorithm (SSA).

Besides that, some authors have been employing other optimisation tools. For example, Knor et al. [27] applied a numerical so- lution based on the inverse heat transfer problem (IHTP) and direct search optimisation to investigate the thermal properties of hardening concrete, i.e., specific heat and thermal conductivity. Pazouki et al. [28] proposed a hybrid method using a

FULL TEXTS BOOK

neural network and firefly optimisation algorithm to predict and optimise the compressive strength of self-compacting concrete containing fly ash.

1.2. *Overview of machine learning techniques*

Machine learning is considered a branch of artificial intelligence that uses several algorithms to learn from the data. The learning step allows its behaviour to be modified based on its own experience, i.e., the system can predict or classify data based on previous data [29]. This behavioural modification consists of establishing some logical rules according to the input data to explore the potential of hidden patterns in the data used and improve the performance of a specific task. The use of these techniques allows the designers or the decision-makers to predict some desired outputs and choose a cleverer and more assertive decision regarding the properties of concrete [30].

The techniques have an outstanding performance in data processing and a remarkable ability to model the properties of concrete without explicit knowledge of the relationships between the properties of concrete and the variables been extensively used to model various properties of different types of concrete and solve concrete mix design issues. Another factor encouraging machine learning is the development of new types of concrete and the lack of mixing design guidelines, which can be noted in the Portland concrete guidelines that do not consider the new types of concrete developed.

With technological advancement and the search for more sustainable building materials, researchers have been looking for al- ternatives to replace part of the cement and aggregates, aiming to reduce CO_2 emissions and improve the quality of concrete. However, although the works regarding these materials are increasing, guidelines and standards do not cover concrete mixtures with these substitutions for common concrete mixtures. In addition, some types of concrete, such as fly ash-based geopolymer concrete, rub- berised concrete, and recycled aggregate concrete, do not have specific guidelines to guarantee a good concrete mixture.

Machine learning has recently gained prominence in several areas, including the construction sector. Besides that, due to its ac- curacy in property predictions, some researchers have been efficiently applying these algorithms to solve concrete mix issues. The most used machine learning techniques in the literature to solve the concrete mix design issues can be broken down into five broad cate- gories: Artificial Neural Networks (ANN); Support Vector Machine (SVM); Decision Trees (DT); Linear Regression; and Logistic Regression (Fig. 1). The first three categories are the most employed in concrete mix design predictions and optimisations.

1.2.1. *Artificial neural networks*

Artificial neural networks (ANN) are one of the most widespread machine learning techniques [32]. This model is inspired by the human neural system and the learning ability of the human brain. Its schematic representation consists of several layers of inter- connected neurons responsible for letting the information flow from one neuron to another. Although the number of layers and neurons is directly related to the type of problem to be studied, ANN is generally composed of an input layer with the influencing variables, an output layer with the desired predicted variables, and at least one layer hidden between them. Yeh, a pioneer in this field, published some distinct works that influenced the development of the new architecture of ANN [33]. For instance, one of his works showed a higher accuracy of the strength model based on ANN when compared to a regression analysis model [34]. Furthermore, among his recent works, some models are analysing the strength of concrete using the design of experiments [35] and the accuracy of a neural network to predict the slump flow [36].

Different types of ANN are applied to solve concrete mix design problems in the literature. The ANN most used in modelling the prediction of concrete mixture properties are Feed-forward Neural Network (FNN) and the Backpropagation Neural Network (BPNN) approaches. To elucidate some examples, Yaprak et al. [37] applied a Feed-forward Backpropagation Neural Network (FBPNN) al- gorithm to predict the concrete compressive strength using different cement types, ages, and cure conditions as input parameters. Shahmansouri et al. [38] also

FULL TEXTS BOOK

predicted the compressive strength of geopolymer concrete using FNN, varying the specimen's age, NaOH concentration, and contents of neozolite, silica fume, and ground granulated blast-furnace slag. Yue et al. [24] applied FNN to




Fig. 1. Main machine learning techniques used to solve concrete mix design problems

predict concrete slump, compressive strength, tensile strength, and elastic modulus. Regarding BPNN, while Verm et al. [39] predicted the compressive strength by implementing this algorithm for fly-ash blended cement concrete mixes, Kellouche et al. [40] used BPNN to analyse the carbonation depth of fly-ash concrete. Both works indicated good agreement with the experiment results.

Other types of ANN have also been used frequently to predict concrete properties. One model widely applied to problems with concrete mix design is Multilayer Perceptron (MLP). Abbella´n-García [41] applied MLP to predict the 28-day compressive strength of ultra-high-performance concrete using different combinations of supplementary cementitious materials:silica fume, fly ash, slag, glass powder, rice husk ash, fluid catalytic cracking residue, metakaolin, and limestone powder. Sargam et al. [30] compared nine machine learning algorithms, and the MLP model indicated a higher accuracy.

Besides these models, a few works employed other techniques. For example, Zhang et al. [42] established a strength prediction model of rubber fibre concrete using the Extreme Learning Machine (ELM), which showed that the model has high accuracy compared with other models as a conventional neural network. Nunez et al. [29] deployed a novel Recurrent Neural Network (RNN) to predict the compressive strength of recycled aggregate concrete. The model effectively captured the underlying features contributing to the compressive strength of the concrete. Czarnecki et al. [43] compared two ANN, Self-Organizing Feature Map (SOFM) and MLP, for predicting the compressive strength of cementitious composite containing ground granulated blast slag. Al-Mughanam et al. [44] implemented an Adaptive Neuro-Fuzzy Inference System (ANFIS) model to predict the compressive strength of self-compacting concrete containing treated palm oil fuel ash as a partial cement substitution. All of these models indicated high-efficient perfor- mance inpredicting the properties of concrete.

FULL TEXTS BOOK

1.2.2. *Support vector machine*

The Support Vector Machine (SVM) is another machine learning technique highly chosen in concrete properties prediction. SVM is associated with learning algorithms that analyse regression and classification analysis data. According to Suykens and Vandewalle [45], SVM is a powerful and versatile method to deal with linear/non-linear classification, regression, and even outlier detection. The main important objective of the SVM classifier is to separate different classes with the most significant possible margin between two of them. In contrast, Support vector regression (SVR), a subset of SVM, tries to fit as many instances as possible on the street while limiting margin violations [46].

To name a few works applying SVM or SVR, Salimbahrami and Shakeri [47] adopted the SVM method to estimate recycled con- crete's compressive strength. ANN and SVM techniques were compared, and although the results indicated SVM as an effective tool, its predictions are far closer to ANN. Besides, Thilakarathna et al. [48] carried out embodied carbon analysis using a set of machine learning algorithms to minimise the carbon footprint of high-strength and ultra-high-strength concrete without jeopardising the mechanical properties of the concrete. The outcomes found in this work were in good agreement with the ones discovered by Sal- imbahrami and Shakeri. On the other hand, Liu et al. [49] used SVM to predict the autogenous shrinkage of conventional concrete mixtures, and SVM was as accurate as ANN prediction. Other work introduced an advanced version of SVM, least-square SVM, to predict pervious concrete's permeability coefficient and compressive strength [50].

1.2.3. *Decision trees*

The Decision Trees (DT) methods also belong to the supervised learning family and can be applied to solve classification and regression problems. This predictive modelling is represented in a tree structure, where the leaves represent class labels and branches represent the characteristics set that lead to those class labels [48]. Due to its simplicity, some authors use this technique to estimate the properties of concrete mixtures. Random Forest (RF) and M5P trees stand out among the DT algorithms in this research area. Asthe other two methods of ANN and SVM presented previously, DT is also used to predict various properties of concrete.

For example, Gomaa et al. [51] modelled an RF model to predict slump flow and compressive strength as functions of influential parameters of the alkali-activated concrete. The model showed high efficiency in predicting both properties. Han et al. [52] presented an ensemble of ML models formulated by combining RF and SVM to predict the modulus of elasticity of recycled aggregate concrete, which produced more accurate results than those yielded by either standalone ML models. Behnood and Golafshani [53] studied the efficiency of the M5P algorithm in developing predictive models for the mechanical properties of concretes containing waste foundry sand as a partial or total replacement for fine aggregate. This research indicated a successful use of the M5P tree in generating pre- dictive models for compressive strength, modulus of elasticity, flexural strength, and splitting tensile strength.

In addition to the tree-based techniques mentioned above, researchers like Anyaoha et al. [14] and Feng et al. [54] also used other DT techniques to predict the concrete's properties. The former used a boosting ensemble of Smooth Transition Regression Trees (STR-Tree) and a boosting algorithm to create a more robust model. The developed model predicted the compressive strength of high-performance concrete using its constituents and mixture proportions as input variables, which showed its dominance in pre- diction accuracy over the other methods. The latter employed an adaptive boosting algorithm integrated into weak models, such as Classification and Regression Tree (CART), ANN, and SVM. The outcomes indicated CART as the higher performance for the compressive strength prediction of concrete containing slag and fly ash.

3.Review methodology

The systematic method adopted in this review aimed to address the research questions elucidated in the introduction section.

FULL TEXTS BOOK

Therefore, the review methodology can be broken down as follows:

Step 1. The first step is related to the search process for articles involving the concrete mix design topic and selecting the most relevant articles, which fit more into the scope of the questions raised.

Step 2. The second step involves analysing the previously selected articles to specify the general characteristics of the mathematical optimisation modelling and machine learning techniques used in this subject.

Step 3. The third step encompasses the development of a classification framework linking the predicted properties and the methods applied for the prediction and optimisation of concrete mixtures.

Step 4. Finally, the last step involves the evaluation of the material based on the proposed framework.

1.3. Systematic review search

The review search was conducted to find and select the most relevant articles. In order to ensure this, a set of keywords was well- defined. Two main groups of keywords have been used to find the literature related to concrete mix design problems:

Group 1: Words related to concrete mix design: “Concrete Mix Design”; “Concrete Mixture”; “Concrete Proportioning".

Group 2: Words related to mathematical modelling or machine learning: “Mathematical Programming”; “Mathematical Model- ling”; “Optimisation”; “Algorithm”; “Metaheuristic”; “Machine Learning".

Combining the two groups, a general search query was achieved with the mais essential keywords and then employed on academic databases, including Scopus and Science Direct. The following query was applied in the databases for the extraction: “concrete mix design” OR “concrete mixture” OR “concrete mix” OR “concrete proportioning” AND (“mathematical programming” OR

FULL TEXTS BOOK




Fig. 2. The procedure of systematic review search.

"mathematical modelling" OR "mathematical modeling" OR "mathematical model" OR "non-linear model" OR "nonlinear model" OR "particle swarm optimisation" OR "particle swarm optimisation" OR "metaheuristic optimisation" OR "metaheuristic optimisation" OR "machine learning" OR "algorithm" OR "genetic algorithm" OR "neural network" OR "support vector machine" OR "regression tree" OR "random forest" OR "linear regression" OR "polynomial regression" OR "support vector regression" OR "gaussian process regression" OR "boosted tree").

Initially, the search based on the abovementioned databases generated a total of 2040 papers. After the first screening, some constraints, such as conference papers, books, language, and a specific timeline period, were applied to the total of registers, which were reduced to 1087 papers after specifying these constraints After evaluating the title and removing the duplicates, the following screening checked the relevance of the remaining documents, which filtered 310 papers. This step mainly focused on selecting the documents linked to solving concrete mix design problems. Next, further examination was performed to guarantee that the scope of the selected papers was aligned with this review by scrutinising their abstracts, which resulted in a set of 148 remaining documents. Then, they were evaluated by thoroughly reading each paper's introduction and discussion sections. The remaining papers were chosen according to the application of mathematical methods and machine learning techniques to find the best composition of the concrete or any optimised property of this composition. Any article that only focused on experiment outcomes and did not solve an optimisation for concrete mix design

or predict any concrete property based on its constituents was removed. After the last screening, a total of 100 papers were selected to be evaluated in this review. To summarise the systematic review procedure, Fig. 2 describes each step adopted in this review.

1.4. Descriptive analysis

The selected articles were first evaluated according to the year of publication. Then, Fig. 3 was elaborated to reveal the publication rate trend considering the ten years considered in this analysis. It is noticeable that the number of papers related to concrete mix design has continuously risen over the years. Despite being a common topic, this growth can be explained by the different types of concrete studied and several possibilities of the concrete mixture composition aligned with the development of computational techniques. In addition, the search for more sustainable building materials that promote less environmental impact and greater thermal comfort has also stimulated this growth.

Following, the publishing journals most adopted to submit papers regarding concrete mix design were evaluated. Table 1 displays the spread of papers analysed across the publishing journals, bringing 100 documents from 40 different journals. Although there is a wide variety of publishing journals, the Construction and building materials journal contains the most significant publications involving concrete mix design problems, followed by the Journal of Building Engineering and the Journal of Cleaner Production.

Depending on the uses of concrete, a huge variety of concrete can be designed for different objectives, such as high-rise buildings, sustainable buildings, economic buildings, residential buildings, and pavements. As previously described, concrete comprises cement, water, and solid aggregates. However, different materials can be added to this mixture to impart a specific property. Currently, there is a growing demand for building materials with higher performance, more sustainable, and less harmful to the environment. As a result, there is a great deal of adding other materials or partially replacing the cement, which is responsible for emitting significant portions of CO_2 into the atmosphere.

Therefore, the design of concrete mixes is a powerful assistant in specifying the composition of the constituent materials to achieve a specific characteristic. Each concrete mixture design will depend exclusively on the objectives to be achieved in the fresh and hardened states of concrete specified by the designers. Among the papers selected for this review, it was found that there are two well- defined research lines. The first research line is characterised by predicting a specific property of the concrete that one wishes to obtain from the composition of the constituents or a combination of the composition with other properties. While the second research line brings optimisation of the concrete mix design problem, where one or more objectives are selected to be minimised or maximised and thus obtain the optimum concrete mix for each case studied.

Fig. 4 breaks down these two lines of research into three technique classes most applied in the retrieved papers. The first en- compasses machine learning (ML) techniques, the second includes mathematical models (MM), and the third presents a hybrid system (HyS) containing both approaches. Machine learning is the most used method in papers where the properties prediction is sought,

FULL TEXTS BOOK




Fig. 3. Distribution of reference papers by year.

Table 1

Distribution of reference papers by publication.

Journal	Number of studies
Construction and Building Materials	35
Journal of Building Engineering	7
Journal of Cleaner Production	6
Cement and Concrete Research	3
Cement and Concrete Composites	3
Sustainability	3
ACI Materials Journal	3
Applied Sciences	2
Computers and Concrete	2
Frontiers in Materials	2
Frontiers of Structural and Civil Engineering	2
Advances in Civil Engineering	2
European Journal of Environmental and Civil Engineering	2
Materials	2
International Journal of Concrete Structures and Materials	1
Automation in Construction	1
Journal of the International Measurement Confederation	1
Neural Computing and Applications	1
Advanced Engineering Informatics	1
Advances in Engineering Software	1
Advances in Materials Science and Engineering	1
Alexandria Engineering Journal	1
Arabian Journal for Science and Engineering	1
Engineering with Computers	1
Expert Systems with Applications	1
Heat and Mass Transfer	1
Heliyon	1
International Journal of Industrial Engineering: Theory Applications and Practice	1
International Journal of Pavement Research and Technology	1
Journal of Computing in Civil Engineering	1
Journal of Construction Engineering and Management	1

FULL TEXTS BOOK

Journal of Materials Research and Technology	1
KSCE Journal of Civil Engineering	1
Measurement	1
Polymers	1
Renewable and Sustainable Energy Reviews	1
Resources, Conservation and Recycling	1
Science of The Total Environment	1
Soft Computing	1
Structural Concrete	1




Fig. 4. Papers distribution according to the research line.

appearing in 45% of documents. Then comes hybrid systems with 24% and mathematical models with 14%. However, when the main objective is optimisation, there is an inversion of the techniques used. Mathematical models appear as the most applied tool with 12%, followed by hybrid systems with 5%. It is worth mentioning that there is no document in optimisation research presenting machine learning as the unique technique used.

1.5. Literature review classification framework

In this section, a classification framework of the literature review is proposed. The framework classification identifies some cat- egories which characterize the research area examined in the concrete mix design. Besides, each category is mapped to several suitable classes. The main purpose of this framework is to detail the works published in the literature in a consistent and organised way, thus enabling the comprehension of the study and comparisons for future works. Considering the two lines of research found in the literature, the classification of the framework can start from the initial objective of the papers: if it is for the prediction of some property or if it is the optimisation of mixtures with certain minimised or maximised properties. Fig. 5 describes the classification framework broken down into five categories: research line, concrete type, concrete property evaluated, output/objective type of prediction and optimisation cases, and the type of computational aid system used.

The first category of the proposed classification framework is based on the research line adopted in each study. After analysing the selected papers, two specific lines were outlined. One considers the prediction of concrete properties, while the other focuses on optimising the

FULL TEXTS BOOK

design of concrete mixes. After this indication, one can follow the next step and get into the second category to continue the work classification.

The second category examines the main types of concrete investigated in the literature review. It includes the types most used in the




Fig. 5. The proposed classification framework.

construction sector as well as new, more sustainable concretes. Among those evaluated in this review are: conventional concrete, rubberised concrete, recycled aggregate concrete, geopolymer concrete, fly ash concrete, ground granulated blast furnace slag con- crete, silica fume concrete, self-compacting concrete, reinforced concrete, high-performance concrete, ultra-high-performance con- crete, high-strength concrete, plastic concrete, pervious concrete, oil-palm shell concrete, self-consolidating concrete, pozzolanic concrete, and alkali-activated concrete.

According to the research line adopted, there is a wide variety of properties that researchers seek to predict or optimise. The next category allows the specification of which property is evaluated. Among the properties investigated, several properties are related to the concrete's fresh and hardened state. For example, some works consider slump and slump flow regarding the workability's concrete. Furthermore, many predictions and optimisations of compressive strength, flexural strength, tensile strength, and modulus of elasticity regarding mechanical properties are found in the literature. Besides that, some works evaluate other properties such as cost and sustainability-related properties of the material such as embodied carbon, CO_2 emission, carbonation depth, and thermal conductivity. The following classification category characterises the output or objective type from each study. Here, it is evaluated if the pre- diction model output or the optimisation objective considers one or more properties, i.e., single or multi-output/objective. Both research lines present some works predicting or optimising just one property and others considering multiple properties to be predicted

or optimised.

The last category considers the applied method to predict the properties or optimise the objective functions. As previously discussed in the introduction section, mathematical programming and machine learning are widely adopted in concrete mix design problems. The main techniques used are artificial neural networks, decision trees, random forest, support vector machine, ANFIS, linear model, non-linear model, particle swarm optimisation, salp swarm algorithm, firefly algorithm, harmony search, and beetle antennae search. Table 2 summarises the technique acronym used in this review.

4.Evaluation of studies

In this section, studies are evaluated based on the proposed classification framework. In the following subsections, the papers selected in this review are organised according to the categories indicated in Fig. 6.

1.6. Review according to the main objective of the work

The first category relies on the primary purpose of the paper. As previously diagnosed, papers addressing the topic of concrete mix design can be primarily classified into two lines of research: prediction and optimisation. According to Fig. 5, the prediction of concrete properties represents most of the studies found in the literature. It reveals that in the analysis carried out on the papers selected, around 83% of the papers take into account the prediction of properties, and only 17% of the documents aim at optimising the concrete mix design problem, minimising or maximising the objective functions. However, some works adopt both research lines, i.e., the same study considers a property prediction and then performs a concrete mix optimisation.

Both research lines have employed machine learning techniques or mathematical models to characterize and analyse concrete mixtures. For instance, Huang et al. [55] followed the prediction line and developed an ANN model to determine the multiple me- chanical properties of rubberised concrete based on concrete mix composites. On the other hand, Wang et al. [56] aimed to optimise .

FULL TEXTS BOOK

Table 2

List of the techniques acronyms used in this paper.

ABC	Artificial Bee Colony	HS	Harmony Search
ALO	Ant Lion Optimisation	ICA	Imperialist competitive algorithm
ANFIS	Adaptive neuro-fuzzy inference systems	KNN	K-nearest neighbour
ANN	Artificial Neural Network	LCA	Life cycle assessment
BAS	Beetle Antennae Search	LR	Linear regression
BBO	Biogeography-Based Optimisation	LSSVM	Least square support vector machine
BPNN	Back propagation neural network	MARS	Multivariate adaptive regression splines
CART	Classification and regression tree	MEP	Multi-expression programming
CNN	Convolutional neural network	MLP	Multilayer perceptron
DT	Decision Tree	MOLSSVR	Multi-output least squares support vector regression
EA	Evolutionary algorithm	PSO	Particle swarm optimisation
ELM	Extreme Learning Machine	RBFNN	Radial basis function neural network
ESIM	Evolutionary support vector machine inference model	RF	Random forest
FA	Firefly algorithm	RNN	Recurrent neural network
FBPNN	Feed forward back propagation neural network	SLCA	Soccer league competition algorithm
FNN	Feed forward neural network	SOFM	Self-organizing feature map
FRPNN	Feedforward resilient propagation neural network	SQP	Sequential quadratic programming
GA	Genetic algorithm	SSA	Salp swarm algorithm
GEP	Gene expression programming	STR	Smooth Transition Regression
GMDH	Group method of data handling	SVM	Support vector machine
GOA	Grasshopper Optimisation Algorithm	SVR	Support vector regression
GPR	Gaussian process regression	WCA	Water cycle algorithm
GWO	Grey wolves optimisation	wSVM	Weighted support vector machine

concrete mixture of slag-blended concrete with minimum CO2 emission. Zhang et al. [20] employed different ML models to predict concrete objectives first. As a second step, the best prediction model was selected as the objective function for the optimisation procedure.

1.7. Studies based on the concrete type

This section focuses on the types of concrete analysed in the retrieved papers. The documents bring up different combinations of concrete constituents that engender a wide variety of concrete types. Therefore, selecting materials and their quantities allows designing concrete with diverse compositions according to the project requirements. Due to this variety, it is possible to notice that studies have evaluated different types of concrete from 2012 to 2021.

Table 3 presents all types of concrete studied in prediction and optimisation research lines. It is worth mentioning that there is a considerable diversity of concrete in both research lines. However, there is a predominance of conventional concrete, concrete con- taining recycled aggregates, fly ash, silica fume, and slag. These five types of concrete are analysed by 55% and 67% of prediction and optimisation research studies, respectively. Both research lines reveal fly ash concrete as the most studied concrete type. It achieves first place in prediction research and second place in optimisation research.

FULL TEXTS BOOK

Although fly ash-blended concrete appears as the most relevant on this list, other types of concrete are also noteworthy, such as plastic concrete and limestone concrete. Furthermore, with the ongoing search for more sustainable materials and mechanical properties similar to conventional concrete, many authors have been investigating new constituents and, thus, new types of concrete. To name a few, Iqbal et al. [57] modelled the mechanical properties of concrete containing waste foundry sand. Besides that, Chung and Tia [58] investigated the effects of cementitious materials on the compressive strength and surface resistivity of limestone concrete

Table 3

Concrete types evaluated in the literature review.

Predicted			
Concrete type	Qt.	Concrete type	Qt.
Concrete containing fly ash	17	Plastic concrete	3
Recycled aggregate concrete	15	High-strength concrete	2
Conventional concrete	12	Ultra-high-performance concrete	2
High-performance concrete	12	Pozzolanic concrete	2
Concrete containing slag	10	Alkali-activated concrete	1
Concrete containing silica fume	8	Waste foundry sand concrete	1
Reinforced concrete	7	Ultra-high-strength concrete	1
Rubberised concrete	5	Pervious concrete	1
Self-compacting concrete	4	Oil palm shell concrete	1
Self-consolidating concrete	4	Hydraulic concrete	1
Geopolymer concrete	3	Lime-stone concrete	1
Optimised			
Concrete type	**Qt.**	**Concrete type**	**Qt.**
Concrete containing slag	6	High-performance concrete	2
Concrete containing fly ash	5	Reinforced concrete	1
Conventional concrete	3	Plastic concrete	1
Concrete containing silica fume	2	Self-compacting concrete	1
Recycled aggregate concrete	2	Pozzolanic concrete	1
High-strength concrete	2	Lime-stone concrete	1

mixes. Zhang et al. [59] inspected the replacement of natural coarse aggregate with oil palm shell and elucidated the pros and cons of oil palm shell concrete.

1.8. Studies based on concrete property

Designing a concrete mixture relies on specifying the properties of the final product and guaranteeing it will meet the property requirements of the project. For this reason, many researchers and designers focus on experiments and mathematical models that better represent the relationship between concrete composition and its properties. Therefore, the third classification category proposed is the concrete property. Table 4 depicts the distribution of the papers according to the main properties predicted and the objectives optimised. Most of the predictions' outputs are related to the mechanical and chemical properties.

Concerning the predicted properties, several papers evaluate the best technique of machine learning and mathematical models to predict concrete properties. Compressive strength is the most investigated property in all studies, representing 43% of the papers evaluated. The predominance in most studies is justified by its importance in the durability design and predicting the service life of concrete structures. Still, regarding the compressive strength, there are some varieties related to the curing time; however, most works try to predict this property after 28 days. Two other highly evaluated properties are elastic modulus and slump, which appear in 9% and 7% of the works. Elastic modulus is another mechanical property that reflects the ability of concrete to deflect elastically. And the slump, related to the fresh concrete state, is a factor that directly contributes to the workability of the mixture. Forecasting the concrete properties, especially the mechanical properties, helps designers in the early characterization of the concrete, thus saving design time. On the other hand, the papers considering the optimisation process adopt fewer properties as objective functions. Three appear most frequently in single or multi-optimisations among the main properties. 37% of the works look for the optimal concrete mix minimising the cost, 17% look over the compressive strength, and 14% use CO_2 emission as an objective function to minimise. Op- timizations, despite being extremely important for the specification of a more suitable concrete mixture for a project, there is still a lack

of studies focusing on different objective functions. Most studies still focus exclusively on compressive strength and cost.

1.9. Studies based on objective/output type

The following category breaks down the papers into four main classes. Two classes are associated with the predictive research line, characterised as single and multi-output. While the other two are linked to the optimisation research line, defined as single and multi- objective. The distribution of the studies addressing each of these classes is illustrated in Fig. 6. As is seen from Fig. 6, a prediction with a single output occupies the most significant proportion of studies, with 70% of the studies examined. Another interesting point is that the researchers focus more on single-objective optimisation studies than on multi-objective optimisation, representing only 7% of the documents.

A possible reason for many studies in the prediction area is exploring models more effectively for each type of concrete property. Regarding the optimisation of concrete, the studies related to single optimisation are still more significant than those with multiple

FULL TEXTS BOOK

Table 4

Main predicted and optimised properties found in concrete mix design problems.

Predicted			
Properties	Qt.	Properties	Qt.
Compressive strength	66	Permeability coefficient	1
Tensile strength	14	Coefficient of thermal expansion	1
Elastic modulus	13	Modulus of rupture	1
Flexural strength	11	Poisson's ratio	1
Slump	10	Dry shrinkage rate	1
Carbonation depth	4	Shrinkage	1
Mix proportion	4	Porosity	1
Plastic viscosity	3	Transport tortuosity	1
Ultra-pulse velocity	3	Specific heat	1
Thermal conductivity	2	Chloride penetration	1
Slump flow	2	Rapid chloride permeability	1
Interface yield stress	2	Water absorption	1
Autogenous shrinkage	2	GWP	1
Rebound number	1	Fracture energy	1
Embodied carbon	1		
Optimised			
Properties	**Qt.**	**Properties**	**Qt.**
Cost	13	Tensile strength	1
Compressive strength	6	Cracking risk coefficient	1
CO_2 emission	5	Energy consumption	1
Embodied CO_2	2	Resource consumption	1
Slump	2	GHG emission	1
Flexural strength	1	Diffusion coefficient of CO_2	1

4.Discussion

In this discussion section, the papers analysed through the classification framework developed in Section 3 answer the main research questions addressed in this review.

1. What are the main machine learning algorithms that have been used in order to come up with the concrete mix design?

After analysing the selected papers, six main groups are observed when it comes to machine learning techniques applied to concrete mix design: artificial neural networks, decision trees, support vector machines, linear regression, non-linear regression, and logistic

A predominance of the first three groups can be observed in the papers reviewed. In most documents, artificial neural networks have been used in several algorithms (feed-forward neural network, backpropagation neural network, multilayer percep- tron) for property predictions and even used as objective functions for optimisation problems. However, despite the widespread use of this technique for predicting compressive strength at different concrete ages, it is also applied to predict other properties such as tensile strength, elastic modulus, slump, and carbonation depth.

Support Vector Machines (SVM) and Decisions Trees (DT) have also gained prominence in predicting concrete mixture properties, especially support vector regression, random forest, and M5P model trees. However, some studies reveal that in multi-output pre- diction, a model can have higher

FULL TEXTS BOOK

accuracy than others, depending on the property to be predicted. For example, the backpropagation neural network algorithm can achieve superior performance in predicting the compressive strength of concrete. At the same time, RF can provide greater accuracy in predicting properties such as slump [20]. Some comparative studies between the three types of machine learning show that ANN performed considerably better in predicting mechanical properties, such as compressive strength, followed by RF and M5P model tree [79].

Although ANN, SVM, and DT present excellent results in predicting properties, comparative studies between machine learning techniques indicate the superiority of other less explored ones. For example, the Multivariate Adaptive Regression Splines (MARS) technique was compared to the multilayer perceptron and least square support vector machine and performed better compared to these other techniques in predicting compressive strength and ultrasonic pulse velocity [88]. However, notwithstanding the promising results, this technique appears in only five studies.

Besides that, some studies focus on building a computational ensemble model consisting of a set of machine learning algorithms. Some papers compare the individual machine learning techniques with this ensemble approach, revealing a higher accuracy when employing multiple techniques. Thus, new models can be built by integrating different algorithms, allowing vast possibilities to find more accurate models.

2. What are the mathematical optimisation algorithms that have been used in order to help concrete mix design?

As mentioned in the preceding section, mathematical models are often used in concrete mix design to obtain the best concrete composition that meets some objective functions. According to each study, these functions are specified and aim at minimising or maximising some characteristics of the concrete. Different properties are used as objective functions; compressive strength and cost are the most evaluated.

Among the algorithms used in the 2012–2021 period, the Genetic Algorithm (GA) is the optimisation algorithm most widely adopted since the beginning of the period evaluated. However, all the articles selected for this review employed single-objective optimisations to find the best composition varying the objective functions. For instance, Kwon et al. [110] obtained optimum con- crete mix proportions through GA using the diffusion coefficient of CO_2 as an objective function for reinforced concrete structures.

Other algorithms also appear with a particular frequency in concrete proportioning studies. For example, the Firefly Algorithm (FA) has been frequently used in multiobjective optimisations with compressive strength, cost, and CO2 emissions as objective functions [8]. Another example is Gene Expression Programming (GEP) is applied in single-objective optimisation minimising cost [75] and multiobjective optimisation, evaluating compressive strength and slump [96]. Particle Swarm Optimisation (PSO) is also used in both types of optimisation, which is analysed the cost for the single case [29] and compressive strength, cost, and slump for the multi- objective analysis [20].

Another application of mathematical models is optimising other techniques' hyperparameters and increasing the model's accuracy [82]. These four algorithms also appear frequently in optimising hyperparameters; however, GA is the most relevant in this algorithm optimisation application.

3. How is the literature linking both aspects, mathematical optimisation and machine learning, in order to design a concrete mix?

Based on the reviewed papers, a classification framework was proposed, noting the existence of two well-defined lines of research for the concrete mix design project.

The first line of research is linked to the optimisation process of concrete mixtures. Through mathematical models, objective functions that must be maximised or minimised are

FULL TEXTS BOOK

represented to obtain the best composition according to the project objective. The studies analysed showed mathematical models optimising the concrete mix design.

The second line is related to the prediction of concrete properties. In this line, most of the researchers found in the literature are concentrated, and about 83% of the works in this area use machine learning to predict concrete mixtures' characteristics.

Many studies have also adopted hybrid models in optimisation and prediction research lines. The former uses machine learning techniques to represent some properties as an objective function. For example, the application of BPNN represents the compressive strength and slump as objective functions. Then, a particle swarm algorithm is applied to perform a multi-objective optimisation [20]. The latter uses mathematical models to improve the machine learning techniques' parameters and thus optimise the model [50].

5.Conclusions

Operational research techniques for concrete mix design have been used to improve the results achieved by the conventional concrete mix proportioning methods based on many experimental tests. Mathematical programming and machine learning methods applied in concrete mix design have turned attractive for several reasons. To begin with, the different composition combinations due to the diversity of raw materials make the concrete design very challenging to perform all laboratory experiments. In addition, laboratory methods are massive, time-consuming, and do not guarantee an ideal concrete mixing solution. Furthermore, mathematical models and machine learning techniques innovate the concrete design method, ensuring that concrete prediction and optimisation models are carried out quickly and accurately.

This review gathered the most relevant papers on concrete mix design from 2012 to 2021. And then, it was proposed a classification framework that could group the works into different categories. Within the analysis of 100 studies, the framework was broken down into five main categories: type of research line, type of concrete, concrete properties, type of output or objective, and type of computer aid system adopted.

` inclusion of this category would be challenging to group in a single category since the input variables depend on the output to be predicted. For those reasons, the classification framework focused on these five categories. According to the reviewed papers, two research lines were spotted. Around 83% are mainly aimed at predicting the properties of concrete, where 70% developed a prediction model with a single output response, and 13% presented a model with multi-property prediction. The remaining documents, around 17%, look for optimising the concrete mixture, with 10% showing a single objective optimisation and 7% minimising or maximising multi-objective functions.

Among the types of concrete investigated, there is a growing trend of studies in both research lines with concrete containing recycled aggregate and supplementary cementitious material such as fly ash, slag, and silica fume. It can be justified by encouraging the reuse of wasted materials and developing more sustainable materials. Regarding the properties predicted or used as objective functions, mechanical properties are the most evaluated, especially compressive strength. Another objective function often used is cost, which is always considered in optimisation studies. These two properties are presented in 60% and 37% of the examined research lines.

Furthermore, the framework was broken down into three classes to evaluate the computer aid system: mathematical models, machine learning, and hybrid system. Most works applied machine learning to predict some concrete property when it comes to predicting research. About 45% of all documents use at least one machine learning technique. Mathematical models and hybrid systems are used in both research lines, the former accounting for 26% and the latter for 29% of the articles. Although several techniques have been used in both areas, a relevant domain of artificial neural networks, support vector machines, decision trees, and genetic algorithms was noted.

FULL TEXTS BOOK

Although this subject presents many relevant works for predicting properties and optimising concrete mixtures, few focus on multi- objective optimisation. Since several objective functions can show a trade-off relationship, simultaneously optimising them is pretty challenging. Therefore, in order to simplify the mathematical optimisation model, most studies use only a single objective function. However, multi-objective optimisation can adequately represent the concrete mix problem, making it more reliable. Accordingly, future works could focus on multi-objective optimisations, analysing different properties so that the results encompass several concrete characteristics, for example, compressive strength, cost, CO_2 emission, and thermal conductivity. Thus, it can achieve a more resistant, less costly, and more sustainable concrete material.

References

[1] M.A. DeRousseau, J.R. Kasprzyk, W.V. Srubar, Computational design optimization of concrete mixtures: a review, Cement Concr. Res. 109 (April) (2018) 42–53, https://doi.org/10.1016/j.cemconres.2018.04.007.

[2] A.M. Neville, J.J. Brooks, Concrete Technology, second ed., Pearson Education Limited, Harlow, 2010.

[3] M.L. Gambhir, Concrete Technology, fifth ed., McGraw Hill Education, New Delhi, 2013.

[4] M.A. DeRousseau, Concrete Mixture Design Using Machine Learning, Life Cycle Assessment, and Multi-Objective Optimization, University of Colorado Boulder, 2020.

[5] J. Chou, A.-Duc Pham, Enhanced artificial intelligence for ensemble approach to predicting high performance concrete compressive strength, Construct. Build. Mater. 49 (2013) 554–563.

[6] H. Naseri, H. Jahanbakhsh, P. Hosseini, F. Moghadas Nejad, Designing sustainable concrete mixture by developing a new machine learning technique,

J. Clean. Prod. 258 (2020), 120578, https://doi.org/10.1016/j.jclepro.2020.120578.

[7] H. Mashhadban, S.S. Kutanaei, M.A. Sayarinejad, Prediction and modeling of mechanical properties in fiber reinforced self-compacting concrete using particle swarm optimization algorithm and artificial neural network, Construct. Build. Mater. 119 (2016) 277–287, https://doi.org/10.1016/j. conbuildmat.2016.05.034.

[8] J. Zhang, Y. Huang, F. Aslani, G. Ma, B. Nener, A hybrid intelligent system for designing optimal proportions of recycled aggregate concrete, J. Clean. Prod. 273 (2020), 122922, https://doi.org/10.1016/j.jclepro.2020.122922.

[9] H.-S. Lee, S.-M. Lim, X.-Y. Wang, Optimal mixture design of low-CO2 high-volume slag concrete considering climate change and CO2 uptake, Int. J. Concr. Struct. Mater. 13 (1) (2019), https://doi.org/10.1186/s40069-019-0359-7.

[10] W. ben Chaabene, M. Flah, M.L. Nehdi, Machine learning prediction of mechanical properties of concrete: critical review, Nov, Construct. Build. Mater. 260 (2020) 119889, https://doi.org/10.1016/j.conbuildmat.2020.119889.

[11] I. Nunez, A. Marani, M. Flah, M.L. Nehdi, Estimating compressive strength of modern concrete mixtures using computational intelligence: a systematic review, Dec, Construct. Build. Mater. 310 (2021) 125279, https://doi.org/10.1016/j.conbuildmat.2021.125279.

[12] Y. Song, X. Wang, H. Li, Y. He, Z. Zhang, J. Huang, Mixture optimization of cementitious materials using machine learning and metaheuristic algorithms: state of the art and future prospects, Materials 15 (21) (2022) 7830, https://doi.org/10.3390/ma15217830. Nov.

[13] A. Mohammed, S. Rafiq, P. Sihag, R. Kurda, W. Mahmood, Soft computing

FULL TEXTS BOOK

techniques: systematic multiscale models to predict the compressive strength of HVFA concrete based on mix proportions and curing times, J. Build. Eng. 33 (2021), 101851, https://doi.org/10.1016/j.jobe.2020.101851.

[14] U. Anyaoha, A. Zaji, Z. Liu, Soft computing in estimating the compressive strength for high-performance concrete via concrete composition appraisal, Construct. Build. Mater. 257 (2020), 119472, https://doi.org/10.1016/j.conbuildmat.2020.119472.

[15] R. Jin, Q. Chen, A.B.O. Soboyejo, Non-linear and mixed regression models in predicting sustainable concrete strength, Construct. Build. Mater. 170 (2018) 142–152, https://doi.org/10.1016/j.conbuildmat.2018.03.063.

[16] A. Habibi, J. Ghomashi, Development of an optimum mix design method for self-compacting concrete based on experimental results, Construct. Build. Mater. 168 (2018) 113–123, https://doi.org/10.1016/j.conbuildmat.2018.02.113.

[17] S.A. Miller, P.J.M. Monteiro, C.P. Ostertag, A. Horvath, Concrete mixture proportioning for desired strength and reduced global warming potential, Construct. Build. Mater. 128 (2016) 410–421, https://doi.org/10.1016/j.conbuildmat.2016.10.081.

[18] C. Fan, S.A. Miller, Reducing greenhouse gas emissions for prescribed concrete compressive strength, Construct. Build. Mater. 167 (2018) 918–928, https://doi.org/10.1016/j.conbuildmat.2018.02.092.

K.-Q. Li, D.-Q. Li, P.-T. Li, Y. Liu, Meso-mechanical investigations on the overall elastic properties of multi-phase construction materials using finite element method, Construct. Build. Mater. 228 (2019)

YABAN HAYATI REHABİLİTASYONUNA YÖNELİK UZUN DÖNEMLİ VAKA ANALİZİ (2018–2025)

Selçuk ALBAYRAK

Doktora Öğrencisi, Kafkas Üniversitesi, Sağlık Bilimler Enstitüsü, Kars, Türkiye.

ORCID ID: https://orcid.org/0000-0002-0117-1368

Oğuz MERHAN

Prof. Dr., Kafkas Üniversitesi, Veteriner Fakültesi, Biyokimya Anabilim Dalı, Kars, Türkiye.

ORCID ID: https://orcid.org/0000-0002-3399-0667

Kadir BOZUKLUHAN

Prof. Dr., Kafkas Üniversitesi, Veteriner Fakültesi, İç Hastalıkları, Kafkas, Türkiye.

ORCID ID: https://orcid.org/0000-0003-4929-5156

ÖZET

Bu çalışmada, 2018–2025 yılları arasında Erzurum Doğa Koruma ve Milli Parklar (DKMP) Müdürlüğüne teslim edilen yaralı veya bakıma muhtaç yabani hayvanlar temel alınarak, Erzurum'daki yabani hayvan türlerinin dağılımı, yaralanma veya bakıma muhtaç olma nedenleri, yaş grubu özellikleri ile rehabilitasyon süreçlerinin başarı oranları istatistiksel olarak değerlendirilmiştir. Yedi yıllık süreçte, teşhis ve tedavi amacıyla toplam 3386 yabani hayvan Erzurum DKMP Müdürlüğüne teslim edilmiştir. Bu hayvanlar arasında 16 memeli ve 106 kanatlı türü tespit edilmiştir. Tür bazında dağılıma bakıldığında, kendi türleri içerisinde en yüksek orana sahip türler sırasıyla %36,61 ile Kaya Güvercini (*Columba livia*), %32,55 ile Tilki (*Vulpes vulpes*), %22,75 ile Yaban Keçisi (*Capra aegagrus*), %14,90 ile Kirpi (*Erinaceus concolor*) ve %13,12 ile Ebabil (*Apus apus*) olarak belirlenmiştir. Teslim alınan yabani hayvanların %55,58'inin Erzurum DKMP Müdürlüğünde tedavi ve rehabilitasyon süreci başarıyla tamamlanmış ve bu bireyler doğaya yeniden kazandırılmıştır. İncelenen yedi yıllık dönemde teslim alınan hayvan sayılarında yıllar itibarıyla artış eğilimi gözlenmiştir. Hayvanların teslim edilme nedenleri değerlendirildiğinde, %94,98 gibi yüksek bir oranın çevresel faktörlere bağlı doğal zararlar nedeniyle gerçekleştiği belirlenmiştir. Teşhis edilen vakalar arasında ise %42,11 oranı ile bakım ve beslenme eksikliğine bağlı bitkinlik ve %30,71 oranı ile kanat problemleri en sık rastlanan nedenler olarak öne çıkmıştır. Yaş grubu dağılımı değerlendirildiğinde ise teslim alınan yabani hayvanların %67,34'ünün erişkin, %32,66'sının ise yavru olduğu tespit edilmiştir. Elde edilen veriler, Erzurum'daki yaban hayatı rehabilitasyonuna yönelik uzun dönemli çalışmaların kapsamını ve etkinliğini ortaya koymakla birlikte tür bazında müdahale önceliklerinin ve koruma stratejilerinin belirlenmesine de katkı sağlamaktadır.

Anahtar Kelimeler: Erzurum, istatistik, rehabilitasyon, yabani hayvanlar

ABSTRACT

In this study, wild animals that were injured or in need of care and delivered to the Erzurum Directorate of Nature Conservation and National Parks (DKMP) between 2018 and 2025 were evaluated statistically in terms of species distribution, causes of injury or need for care, age group characteristics, and the success rates of rehabilitation processes. Over the seven-year period, a total of 3,386 wild animals were delivered to the Erzurum DKMP Directorate for diagnosis and treatment. Among these animals, 16 mammal species and 106 bird species were identified. When examining species distribution, the species with the highest proportions within their respective groups were determined as Rock Pigeon (*Columba livia*) with 36.61%, Red Fox (*Vulpes vulpes*) with 32.55%, Wild Goat (*Capra aegagrus*) with 22.75%, Hedgehog (*Erinaceus concolor*) with 14.90%, and Common Swift (*Apus apus*) with 13.12%. It was

FULL TEXTS BOOK

found that 55.58% of the wild animals delivered to the Erzurum DKMP Directorate successfully completed the treatment and rehabilitation process and were released back into the wild. Throughout the examined seven-year period, an increasing trend was observed in the number of animals delivered each year. When evaluating the reasons for delivery, it was determined that 94.98% of the cases were due to natural damages caused by environmental factors. Among the diagnosed cases, exhaustion due to lack of care and feeding was the most common condition at a rate of 42.11%, followed by wing injuries at 30.71%. In terms of age group distribution, 67.34% of the delivered wild animals were classified as adult, while 32.66% were classified as juvenile. The data obtained not only reveal the scope and effectiveness of long-term wildlife rehabilitation efforts in Erzurum but also contribute to identifying species-specific intervention priorities and developing conservation strategies.

Keywords: Erzurum, rehabilitation, statistics, wild animals

GİRİŞ

Yaban hayatı rehabilitasyonu, ekosistem sağlığının korunması, türlerin devamlılığının sağlanması ve çevresel tehditlerin izlenmesi açısından kritik bir rol oynamaktadır (Lazarova ve ark. 2024, Engler ve ark. 2025). Günümüzde kentleşme, habitat parçalanması, iklim değişikliği ve artan insan-yaban hayatı etkileşimleri, yabani hayvanların yaralanma ve bakım ihtiyacını önemli ölçüde artırmıştır (Engler ve ark. 2025). Bununla birlikte, rehabilitasyonun başarısı sadece hayvanların serbest bırakılması ile değil, serbest bırakıldıktan sonraki hayatta kalma oranları, üreme başarısı ve ekosisteme uyum gibi uzun vadeli göstergelerle değerlendirilmelidir (Mladonicky ve ark. 2024).

Finlandiya'daki bir çalışmada, rehabilitasyon oranlarının %80 gibi yüksek seviyelerde olduğu bildirilmektedir; ancak kayıt eksiklikleri ve öznel değerlendirmelerin, hayvanların ümitsiz bir vaka olarak kabul edilmesi veya yetersiz koşullarda doğaya salınması gibi etik sorunlara yol açabileceği vurgulanmıştır (White ve ark. 2024).

ABD'de kapsamlı bir literatür incelemesinde, rehabilitasyon sonrası hayatta kalma oranlarının türlere göre büyük farklılıklar olduğu görülmüştür. Özellikle petrol kirliliğine maruz kalan deniz kuşlarında başarının düşük olduğu belirlenmiş; bu da rehabilitasyon stratejilerinin tür bazlı değerlendirilmesini ve her türün biyolojik ve davranışsal özelliklerinin dikkate alınmasını gerekli kılmaktadır (Mladonicky ve ark. 2024).

Yaban hayatı rehabilitasyonunun başarısı yalnızca veteriner hekimlik ve ekolojik bilgiyle değil, aynı zamanda yerel halkın geleneksel bilgi ve tutumlarıyla da doğrudan ilişkilidir. Kenya'daki Maasai topluluğuyla yürütülen bir araştırma, geleneksel bilgiye sahip bireylerin yaban hayatı koruma konusunda daha olumlu tutumlara sahip olduğunu ve bu tutumların koruma davranışlarını doğrudan etkilediğini ortaya koymuştur. Bu bulgu, rehabilitasyon süreçlerinde yerel halkın bilgi sistemlerinin dikkate alınması gerektiğini ortaya koymaktadır (Ochieng ve ark. 021).

Öte yandan, rehabilitasyon uygulamalarında etik boyutlar da göz ardı edilmemelidir. Bununla ilgili Washington Eyaleti Yaban Hayatı Rehabilitasyon Kılavuzu'nda, rehabilitasyonun sadece fiziksel iyileşmeyi değil, aynı zamanda psikolojik sağlığı da gözetmesi gerektiği vurgulanmıştır. Hayvanların evcilleştirilmesi, insanlara alışması (habituasyon) veya yanlış türlere bağlanması (mal-imprinting) gibi durumlar, doğaya salınan bireylerin hayatta kalmasını zorlaştırmakta ve ekosistem için risk oluşturmaktadır. Özellikle yırtıcı kuşlar ve memelilerde başarı oranı, türün doğal davranışlarının korunmasıyla doğrudan ilişkilidir. Hayvanların doğaya salınmadan önce fiziksel, davranışsal ve psikolojik olarak tam iyileşmiş olmaları gerektiği vurgulanmıştır (Thompson 2019). Bu durum, Mladonicky ve ark. (2024) tarafından da desteklenmiştir. Literatür taramasında, rehabilitasyon sürecinde insan teması azaltıldığında hayatta kalma oranlarının arttığı, özellikle kuş türlerinde bu etkinin belirgin olduğu ifade edilmiştir.

Ayrıca, rehabilitasyon sonrası salım için belirlenen kriterler arasında fiziksel iyileşme, davranışsal uygunluk, mevsimsel zamanlama ve salım bölgesinin seçimi gibi unsurlar yer almaktadır. Bu kriterlerin sağlanmadığı durumlarda hayvanların salımı önerilmemektedir. Ayrıca rehabilitasyon sürecinde zoonotik ve epizootik hastalıkların kontrol altına alınmasının hayati önem taşıdığını vurgulamaktadır (Thompson 2019).

FULL TEXTS BOOK

Benzer şekilde, Bulgaristan'daki Green Balkans Yaban Hayatı Rehabilitasyon Merkezi'nde 25 yıllık bir çalışmada, analiz edilen 18.720 bireyin %32'sinin doğal nedenlerden, %18'inin ise antropojenik etkilerden kaynaklı zarar gördüğü raporlanmıştır (Lazarova ve ark. 2024). Almanya'nın Berlin kentinde 20 yıl süren bir araştırmada ise kentleşmenin bina ve araç çarpmaları, evcil hayvan saldırıları ve kötü beslenme gibi faktörlerle rehabilitasyon süreçlerini doğrudan etkilediği gösterilmiştir (Engler ve ark. 2025).

Rehabilitasyon sürecinin şeffaf ve izlenebilir olması için kayıt tutma uygulamaları büyük önem taşımaktadır. Her hayvanın tür bazında kayıt altına alınması gerektiğini ve günlük kayıt defterlerinin düzenli tutulmasının zorunlu olduğunu belirtmektedir. Ayrıca, rehabilitasyon izni almak isteyen kişilerin 1000 saatlik gönüllü çalışma ve veteriner iş birliği gibi kriterleri karşılaması gerekmektedir (Thompson 2019).

ARAŞTIRMA ve BULGULAR

2018–2025 yılları arasında yürütülen uzun dönemli vaka analizi toplanan verilerin analizi sonucunda yapılmıştır. Bu veri analizlerinden yapılan en önemli çıkarımlardan biri rehabilitasyon süresince türler arasında belirgin farklılıklar gösterdiği tespit edilmiştir. Ayrıca bu farklılıkların sadece türler arasında değil yıllara göre, mevsin normellerinde ki değişikliklere, yaş skalalarına ve transfer edildikleri mesafelere görede rehabilitasyon başarı oranlarında da değişiklikler gözlemlenmiştir. Özellikle genç hayvanların kritik dönemlerinde insan teması, geri dönüşü olmayan davranışsal bozulmalara yol açmaktadır.

Yaptığımız veri analizi sonucunda yedi yıllık yapılan 3386 yabani hayvan vaka çalışması sonucunda transferler haricinde başarı oranımızın %55,58 olarak analiz edilmiştir. Ayrıca bu vaka analizlerinden tür dağılımlarını, yıllara göre vaka değişimlerini, vakalardaki zarar kategorilerini, teşhis edilen klinik bulguları, en yoğun vakalarda gözlemlenen kanatlı ve memeli türlerini, kanatlı türlerdeki mevsimles değişimleri ve yaş skalalarını belirledik.

Bu çalışma, 2018–2025 yılları arasında Erzurum'da yürütülen yaban hayatı rehabilitasyon sürecine dair kapsamlı verileri içermektedir. Aşağıda öne çıkan bulgular grafiklerle desteklenmiştir:

1.Yaban Hayvanlarının Türlere Göre Dağılımı: Bu yedi yıllık süreçte teslim alınan yabani hayvanların büyük çoğunluğunu kuş türleri oluşturmuştur. Kanatlı bireyler toplam vakaların yaklaşık %92 (3126 birey)'sini temsil ederken, memeli türler %7 (254) civarında bir orana sahiptir. Sürüngenler ise oldukça düşük bir oranla, yalnızca birkaç birey ile temsil edilmiştir. Bu dağılım, bölgedeki ekolojik yapı ve insan-yaban hayatı etkileşimlerinin en çok kuş türlerini etkilediğini göstermektedir.

2. Rehabilitasyon Sürecinin Başarı Oranları: Erzurum'daki yaban hayatı rehabilitasyon sürecinde sonuçlanan 2573 vakaya dair başarı oranları değerlendirildiğinde, tedavi ve bakım süreci tamamlanan bireylerin yaklaşık %55,58 (1430 birey)'inin doğaya yeniden kazandırıldığı görülmektedir. Buna karşılık, vakaların %44,42 (1143)'si de çeşitli nedenlerle yaşamını yitirmiştir. Ayrıca geriye kalan yabani hayvanlar ise en yakın Yabani Hayvan Rehabilitasyon Merkezine transferi gerçekleştirilmiş ya da transfer sırasında ex olmuştur. Bu oranlar, rehabilitasyon uygulamalarının genel etkinliğini ortaya koyarken, aynı zamanda iyileştirme süreçlerinin daha da geliştirilmesi gerektiğine işaret etmektedir.

3. Yaş Gruplarına Göre Vaka Dağılımı: 2018–2025 yılları arasında Erzurum DKMP Müdürlüğü'ne teslim edilen toplam 3.386 yabani hayvan vakası, yaş gruplarına göre incelendiğinde erişkin bireylerin %67,34 (2280 birey), yavruların ise %32,66 (1106 birey) oranında temsil edildiği görülmektedir.

Yaban hayvanlarında kesin yaş tayini, türler arası fizyolojik farklılıklar, travma sonrası deformasyonlar ve gözlem koşullarının sınırlılığı nedeniyle oldukça güçtür. Bu nedenle, rehabilitasyon sürecinde yaş grupları genellikle "erişkin" ve "yavru" olarak iki temel kategoriye ayrılarak değerlendirilmiştir. Bu sınıflandırma, hem klinik müdahale planlamasında hem de doğaya salım stratejilerinde önemli bir rol oynamaktadır.


Erişkin bireylerin daha yüksek oranda teslim edilmesi, çevresel tehditlere karşı daha fazla maruz kalmalarıyla ilişkilendirilebilirken; yavru bireylerin özellikle ilkbahar ve yaz aylarında yuvadan düşme, bitkinlik ve bakım eksikliği gibi nedenlerle rehabilitasyon merkezlerine ulaştırıldıkları gözlemlenmiştir.

Bu yaş grubu dağılımı, rehabilitasyon merkezlerinin yaşa özgü müdahale protokolleri geliştirmesi

FULL TEXTS BOOK

gerektiğini ortaya koymaktadır. Erişkin bireylerde fiziksel travmalar ve kanat problemleri ön plana çıkarken, yavru bireylerde bakım eksikliği ve yetersiz beslenme gibi nedenler daha yaygındır. Ayrıca, yavru bireylerde insan teması kaynaklı davranışsal bozulmaların daha sık gözlemlenmesi, rehabilitasyon sürecinde dikkatli sosyal izolasyon uygulamalarını zorunlu kılmaktadır.


4. Vaka Sayılarının Yıllara Göre Dağılımı: Yedi yıllık süreçte Erzurum'daki yaban hayatı rehabilitasyon merkezine teslim edilen vaka sayılarında belirgin bir artış eğilimi gözlemlenmiştir. Özellikle 2024 yılı, en yüksek birey sayısının (665 birey) kaydedildiği dönem olarak öne çıkmaktadır. Bu artış, hem çevresel faktörlerin etkisinin arttığını hem de halkın rehabilitasyon merkezlerine yönlendirme konusunda daha bilinçli hale geldiğini göstermektedir.



Grafik 1. 2018–2025 Yılları arasında vaka sayılarının yıllık dağılımı

5. Doğal ve Antropojenik Etkenlere Bağlı Zarar Dağılımı: Erzurum'da 2018–2025 yılları arasında yürütülen yaban hayatı rehabilitasyon sürecinde teslim edilen vakaların zarar teşhisleri incelendiğinde, çoğunluğun %94,98 oranla çevresel faktörlere bağlı doğal etkilerden kaynaklandığı görülmektedir. Bu durum, bölgedeki iklimsel değişimler, habitat bozulmaları ve doğal yaşam alanlarındaki stres faktörlerinin yabani hayvanlar üzerinde ciddi etkiler yarattığını göstermektedir.

Geriye kalan vakalar ise daha düşük oranlarla antropojenik nedenlere dayanmaktadır. Bunlar arasında diğer hayvanlar tarafından gerçekleşen saldırılar (%2,27), trafik kazaları (%1,74), elektrik çarpmaları (%0,56) ve ateşli silah yaralanmaları (%0,44) yer almaktadır. Bu bulgular, hem doğal hem de insan kaynaklı tehditlerin rehabilitasyon sürecinde dikkate alınması gerektiğini ortaya koymaktadır.




Grafik 2. 2018-2025 Yılları arasındaki yabani hayvanlarda gözlemlenen zarar kategorilerinin dağılımı

FULL TEXTS BOOK

6. En Yaygın Klinik Teşhislerin Dağılımı: Erzurum'da 2018–2025 yılları arasında yürütülen yaban hayatı rehabilitasyon sürecinde en sık karşılaşılan klinik teşhisler incelendiğinde, vakaların büyük bir kısmının bitkinlik ve bakım eksikliği nedeniyle gerçekleştiği görülmektedir. Bu teşhis, toplam vakaların %42,11'ini oluşturarak en yaygın neden olarak öne çıkmaktadır. Bunu %30,71 oranıyla kanat problemleri ve %8,27 oranıyla ayak problemleri takip etmektedir.

Daha düşük oranlarla gözlemlenen diğer teşhisler arasında iç kanama (%1,77), yuvadan düşme (%5,43), kafa ve boyun travmaları (%2,51), öksüzlük (%2,63), enfeksiyon çeşitleri (%3,34) ve çeşitli diğer nedenler (%3,22) yer almaktadır. Bu dağılım, rehabilitasyon sürecinde karşılaşılan fizyolojik ve davranışsal sorunların çeşitliliğini ortaya koymakta; özellikle kuş türlerinde kanat ve bakım problemlerinin öncelikli müdahale alanları olduğunu göstermektedir.




Grafik 3. 2018–2025 Yılları arasında en yaygın klinik teşhislerin oransal dağılımı

7. 2018–2025 Yılları Arasında Rehabilitasyon Merkezine Teslim Edilen Memeli Türlerin Oransal Dağılımı: Erzurum'da 2018–2025 yılları arasında yürütülen yaban hayatı rehabilitasyon sürecinde teslim edilen memeli türler incelendiğinde, en yüksek oranla teslim edilen türün Tilki (Vulpes vulpes) olduğu görülmektedir. Toplam memeli vakalarının %32,55'ini oluşturan Tilki'yi, %22,75 ile Yaban Keçisi (*Capra aegagrus*), %14,90 ile Kirpi (*Erinaceus concolor*) ve %7,06 ile Kurt (*Canis lupus*) takip etmektedir.

Daha düşük oranlarla gözlemlenen diğer memeli türler arasında Vaşak, Yaban Tavşanı, Boz Ayı, Yarasa, Sincap, Yediuyur, Alaca Sansar, Kaya Sansarı, Gelengi, Karaca, Porsuk ve Su Samuru yer almaktadır. Bu dağılım, bölgedeki memeli türlerinin rehabilitasyon sürecine dahil olma sıklığını ve tür bazlı müdahale önceliklerini ortaya koymaktadır.


FULL TEXTS BOOK



Grafik 4. Erzurum'da rehabilite edilen memeli türlerin tür bazlı oransal dağılımı

8. Rehabilitasyon Sürecinde En Sık Teslim Edilen İlk 20 Kanatlı Türün Dağılımı: Erzurum'da 2018–2025 yılları arasında yürütülen yaban hayatı rehabilitasyon sürecinde en çok teslim edilen kanatlı türler incelendiğinde, Kaya Güvercini (*Columba livia*) %36,61 oranıyla açık ara en yüksek sayıya ulaşmıştır (1144 birey). Bunu %13,12 ile Ebabil (*Apus apus*), %6,50 ile Kerkenez (*Falco tinnunculus*) ve %5,12 ile Kızıl Şahin (*Buteo rufinus*) takip etmektedir. Küçük Karga, Serçe, Kuzgun ve Leylek gibi türler de %2–5 aralığında önemli oranlara sahiptir.

Daha düşük oranlarla gözlemlenen diğer türler arasında Sığırcık, Saksağan, Kulaklı Orman Baykuşu, Kaya Kartalı, Çobanaldatan, Şahin, Atmaca, Puhu, Kukumav, Çıvgın, Kara Çaylak ve Kumru yer almaktadır. Bu dağılım, rehabilitasyon merkezine en sık getirilen türlerin hem şehirleşme hem de mevsimsel etkilerle ilişkili olduğunu göstermektedir. Özellikle uçuş yeteneği zayıflamış veya yavru döneminde olan bireylerin yoğun şekilde teslim edilmesi, kanatlı türlerin rehabilitasyon sürecindeki öncelikli konumunu ortaya koymaktadır.




Grafik 5. 2018–2025 Yılları arasında en fazla rehabilite edilen kanatlı türlerin ilk 20'sinin dağılımı

FULL TEXTS BOOK

9. 2018–2025 Yılları Arasında En Sık Teslim Edilen 20 Kanatlı Türün Mevsimsel Dağılımı: Erzurum'da 2018–2025 yılları arasında yürütülen yaban hayatı rehabilitasyon sürecinde en çok teslim edilen 20 kanatlı türün mevsimsel dağılımı incelendiğinde, türlerin doğaya teslim edilme zamanlarının belirgin şekilde mevsimlere göre farklılık gösterdiği görülmektedir. Özellikle yaz ve ilkbahar aylarında vaka sayılarında artış gözlemlenmiş; bu dönemler, yavru bireylerin yuvadan düşme, bitkinlik ve uçuş problemleri gibi nedenlerle rehabilitasyon merkezlerine daha sık ulaştırıldığı zamanlar olmuştur.

Kaya Güvercini, Ebabil, Kerkenez ve Kızıl Şahin gibi türler, yaz aylarında en yoğun şekilde teslim edilen türler arasında yer alırken; kış aylarında teslim edilen birey sayısı genel olarak daha düşüktür. Bu durum, hem mevsimsel göç dinamikleri hem de bölgesel iklim koşullarının türlerin sağlık durumları üzerindeki etkisini ortaya koymaktadır.



Grafik 6. En sık teslim edilen 20 kanatlı türün mevsimsel dağılı

TARTIŞMA ve SONUÇ

Bu çalışma, Erzurum'daki yaban hayatı rehabilitasyonuna yönelik uzun dönemli vaka analizinin, yerel türlerin korunması ve uluslararası literatürle karşılaştırmalı değerlendirmeler yapılmasına olanak sağladığını göstermektedir. Elde edilen bulgular, Lazarova ve ark. (2024), Mladonicky ve ark. (2024) ve Engler ve ark. (2025) gibi uluslararası çalışmalarla uyumlu olup Türkiye'de yaban hayatı rehabilitasyonunun bilimsel temellere dayandırılması açısından önemli katkılar sunmaktadır. Bu veriler, rehabilitasyonun yalnızca iyi niyetle değil, bilimsel temellere dayalı karar süreçleriyle yürütülmesi gerektiğini göstermektedir. Bu bulgular, yerel bilgi sistemlerinin modern koruma stratejilerine entegre edilmesinin önemini ortaya koymaktadır.

Yaptığımız bu çalışmadaki Erzurum örneğinde ise, 2018–2025 yılları arasında teslim edilen yabani hayvanların %55,58'i başarılı şekilde doğaya yeniden kazandırılmıştır. Ayrıca, vakaların %94,98'inin çevresel faktörlere bağlı doğal zararlar nedeniyle gerçekleştiği tespit edilmiş, bu bulgu Lazarova ve ark. (2024) tarafından vurgulanan doğal travma nedenleriyle uyumludur. En sık teşhis edilen durumlar arasında %42,11 oranıyla bitkinlik ve %30,71 oranıyla kanat problemleri yer almakta olup benzer sonuçlar Engler ve ark. (2025) çalışmalarında da görülmüştür.

Mladonicky ve ark. (2024) tarafından yapılan kapsamlı literatür taramasında, rehabilitasyon sonrası başarıyı etkileyen en kritik faktörlerin tür özellikleri, yaş grubu, yaralanma tipi ve salım sonrası izleme süresi olduğu vurgulanmıştır. Erzurum örneğinde yaş grubu dağılımında erişkin bireylerin %67,34, yavruların ise %32,66 oranında temsil edilmesi, literatürde özellikle yavru bireylerde insan teması kaynaklı davranışsal bozulmaların uzun vadeli hayatta kalma oranlarını düşürdüğü yönündeki bulgularla örtüşmektedir (Griffiths ve ark. 2010, Thompson 2019).

Ayrıca, Mladonicky ve ark. (2024) çalışmasında vurgulanan bir diğer önemli nokta, rehabilitasyonun dolaylı koruma katkılarıdır. Bunlar arasında hastalık izleme, çevresel toksinlerin tespiti, kamu farkındalığı ve bilimsel veri üretimi yer almaktadır. Erzurum DKMP Müdürlüğü'nün yıllık vaka

FULL TEXTS BOOK

kayıtları, bölgesel ekosistem sağlığına dair önemli bir gösterge niteliği taşımakta olup, bu verilerin dijitalleştirilmesi ve ulusal düzeyde standartlaştırılması, Türkiye'deki rehabilitasyon çalışmalarının bilimsel ve koruma temelli etkisini artıracaktır.

Sonuç olarak, Erzurum'daki yaban hayatı rehabilitasyon süreci, uluslararası literatürdeki örneklerle karşılaştırıldığında hem başarı oranı hem de tür çeşitliliği açısından anlamlı bir düzeyde yer almaktadır. Ancak salım sonrası izleme, davranışsal değerlendirme ve ölüm nedenlerinin belirlenmesi gibi alanlarda daha sistematik ve uzun vadeli çalışmaların yapılması gerekmektedir. Bu bağlamda, Mladonicky ve ark. (2024) tarafından önerilen veri standardizasyonu, izleme araçlarının yaygınlaştırılması ve tür bazlı rehabilitasyon protokollerinin geliştirilmesi gerekmektedir.

Bu çalışmanın bulguları ve uluslararası literatürle yapılan karşılaştırmalar doğrultusunda, yaban hayatı rehabilitasyonunun daha etkin, izlenebilir ve bilimsel temellere dayalı bir yapıya kavuşturulması için çeşitli öneriler geliştirilmiştir.

İlk olarak, veri toplama ve izleme sistemlerinin güçlendirilmesi gerekmektedir. Rehabilitasyon merkezlerinde tutulan kayıtların dijitalleştirilmesi ve tanımların standartlaştırılması, hem ulusal hem de uluslararası düzeyde karşılaştırılabilir veri üretimini mümkün kılacaktır. Salım sonrası izleme araçlarının (GPS, uydu, radyo takip, halkalama gibi) yaygınlaştırılması, rehabilitasyonun gerçek başarısını ölçmek açısından kritik öneme sahiptir. Ayrıca, tedavi süresi ile salım başarısı arasındaki ilişkinin dikkate alınması, uzun süreli tedavilerin başarıyı olumsuz etkileyebileceği gerçeğini göz önünde bulundurmayı gerektirir. Bu bağlamda, ölüm nedenlerinin belirlenmesi için teşhis sistemlerinin güçlendirilmesi de önemli bir ihtiyaçtır.

Etik ve sosyal boyutlar açısından, ağır yaralı hayvanların tedavi edilip edilmemesi gibi kararların bilimsel ve kaynak temelli değerlendirmelerle ele alınması gerekmektedir. Özellikle insanlar tarafından ümitsiz vaka olarak getirilen bireylerin oranının azaltılması için kamu farkındalığının artırılması önemlidir. Bu doğrultuda, eğitim ve bilgilendirme çalışmaları yaygınlaştırılmalı; özellikle yaz aylarında artan vaka sayısına karşı önleyici stratejiler geliştirilmelidir.

Politika ve uygulama düzeyinde ise, Doğa Koruma ve Milli Parklar Genel Müdürlüğünün yabani hayvan potansiyeli yüksek bölgelerde rehabilitasyon merkezi kurma kararı desteklenmeli ve hayata geçirilmelidir. Ayrıca, yerel halkın geleneksel bilgi sistemlerinin modern koruma politikalarına entegre edilmesi, hem ekolojik hem de toplumsal sürdürülebilirlik açısından önem taşımaktadır. Zarar gören topluluklar için tazmin mekanizmalarının geliştirilmesi ve bilimsel-etik temelli, sıkı düzenlemelere dayalı bir rehabilitasyon sisteminin oluşturulması, Türkiye'de yaban hayatı rehabilitasyonunun kurumsal kapasitesini artıracaktır.

Bu öneriler, Erzurum örneğinde elde edilen bulguların daha geniş bir koruma stratejisine dönüştürülmesi açısından yol gösterici niteliktedir.

***Etik Beyan*:** Bu çalışma kapsamında kullanılan tüm veriler, araştırmacı tarafından doğrudan gözlem yöntemiyle elde edilmiştir. Gözlemler rehabilitasyon merkezindeki rutin bakım süreçleri sırasında, hayvanların normal davranışlarını etkilemeden kaydedilmiştir. Herhangi bir kurumun veri tabanından bilgi alınmamış, hayvanlar üzerinde deneysel işlem, müdahale veya örnekleme yapılmamıştır. Gözlemler vakaların teslim edildiği anda gerçekleştirilmiş olup çalışmada yer alan veriler kamuya açık veri kaynaklarında bulunmayan özgün nitelikler taşımaktadır. Bu nedenlerle, etik kurul izni gerektirecek bir durum söz konusu değildir.

KAYNAKÇA

Engler, M., Sens, R., Lundberg, M., Delor, A., Stelter, M., Tschertner, M., Feyer, S., Zein, S., Halter-Gölkel, L., Altenkamp, R., & Müller, K. (2025). Worth the Effort? Rehabilitation Causes and Outcomes and the Assessment of Post-Release Survival for Urban Wild Bird Admissions in a European Metropolis. Animals (Basel), *15*(12), 1746. https://doi.org/10.3390/ani15121746

Griffiths, R., Murn, C., & Clubb, R. (2010). Survivorship of rehabilitated juvenile Tawny Owls (Strix aluco) released without support food, a radio tracking study. Avian Biology Research, 3(1), 1-6.

Lazarova, I., Balieva, G., & Klisurov, I. (2024). Causes of admissions and mortality in patients of a wildlife rehabilitation centre in Bulgaria. Biodiversity Data Journal, 12, e123681.

FULL TEXTS BOOK

https://doi.org/10.3897/BDJ.12.e123681

Mladonicky, J., Sirolli, M., Willette, M., & Ponder, J. (2024). Evaluation of current scientific literature on the impact of wildlife rehabilitation on conservation, documentation of knowledge gaps and identification of future opportunities. University of Minnesota, College of Veterinary Medicine. Prepared for U.S. Fish and Wildlife Service.

Ochieng, C. N., Thenya, T., Shah, P., & Odwe, G. (2021). Awareness of traditional knowledge and attitudes towards wildlife conservation among Maasai communities: The case of Enkusero Sampu Conservancy, Kajiado County in Kenya. African Journal of Ecology, 59(3), 712–723. https://doi.org/10.1111/aje.12872

Thompson, P. (2019). Washington Department of Fish and Wildlife wildlife rehabilitation manual (3rd ed.). Washington Department of Fish and Wildlife.

White, K., Hänninen, L., Sainmaa, S., & Valros, A. (2024). Reasons for admission and rehabilitation rates of various wildlife species in Finland. Frontiers in Veterinary Science, 11, Article 1455632. https://doi.org/10.3389/fvets.2024.1455632

BORÇLANMA VE SAĞLIK STATÜSÜ İLİŞKİSİ: TÜRKİYE ÖRNEĞİ

THE RELATIONSHIP BETWEEN DEBT AND HEALTH STATUS: THE TURKISH CASE

Dr. Öğr.Üyesi Murat KONCA

Çankırı Karatekin Üniversitesi, Sağlık Bilimleri Fakültesi, Sağlık Yönetimi Bölümü, Çankırı.

ÖZET

Bu çalışmada, borçlanmanın sağlık statüsü üzerindeki etkisi Türkiye özelinde ele alınmaktadır. Bu kapsamda, 1.000 canlı doğumda yenidoğan ölüm hızı, gayrisafi yurtiçi hasılaya oranla merkezi hükümet borcu, toplam dış borcun yüzdesi olarak kısa vadeli borçlar ve gayrisafi yurtiçi hasılaya oranla toplam borç servisi değişkenleri arasındaki eş-bütünleşiklik ilişkisi incelenmektedir. Çalışmanın analizleri Türkiye'nin 1985-2023 dönemine ait verileri ile gerçekleştirilmiş olup veriler, Dünya Bankası (2025) veri tabanından alınmıştır. Çalışmada, öncelikle, tanımlayıcı istatistikler ve değişkenler arasındaki korelasyon düzeyleri incelenmiştir. Sonrasında, Augmented Dickey–Fuller Testi ile değişkenlerin hangi seviyede durağan hale geldiği incelenmiş ve tüm değişkenlerin birinci derece farkta durağan hale geldiği görülmüştür. Bu bulgunun bir gereği olarak Johansen eş-bütünleşme testi kullanılmıştır. Johansen eş-bütünleşme testi öncesinde Vektör Otoregresif (VAR) modeli ile uygun gecikme uzunluğuna karar verilmiştir. Eş-bütünleşme testi sonrasında, vektör hata düzeltme modeli kurulmuştur. Johansen eş-bütünleşme testi kapsamında İz (Trace) ve Maksimum Özdeğer (Max-Eigenvalue) test sonuçları elde edilmiştir. İz testi sonuçlarına göre dört serinin tamamı arasında en az bir ve muhtemelen birden fazla eş-bütünleşme ilişkisinin bulunduğu; maksimum özdeğer testi sonuçlarına göre ise en az dört değişkenin bir eş-bütünleşme ilişkisi içinde olduğu görülmüştür. Eş-bütünleşme denklemi sonuçlarına göre, gayrisafi yurtiçi hasılaya oranla toplam borç servisi ve toplam dış borcun yüzdesi olarak kısa vadeli borçlar değişkenlerinin yenidoğan ölüm hızı üzerindeki etkilerinin pozitif ve anlamlı olduğu anlaşılmıştır. Vektör hata düzeltme modeli sonuçları incelendiğinde, hata düzeltme katsayısının sadece toplam dış borcun yüzdesi olarak kısa vadeli borçlar değişkeni için anlamlı olduğu saptanmıştır.

Anahtar Kelimeler: Borçlanma, sağlık statüsü, Türkiye, zaman serisi.

ABSTRACT

This study examines the impact of debt on health status in Türkiye. In this context, the cointegration relationship between the following variables is examined: neonatal mortality rate per 1,000 live births, central government debt as a percentage of gross domestic product, short-term debt as a percentage of total external debt and total debt service as a percentage of gross domestic product. Analyses were conducted using data from Türkiye covering the 1985-2023 period, obtained from the World Bank (2025) database. First, descriptive statistics and correlation levels between variables were examined. The Augmented Dickey–Fuller Test was then used to examine the level at which variables became stationary, and it was found that all variables became stationary at their first-degree differences. As a requirement of this finding, the Johansen cointegration test was used. Before the Johansen cointegration test, the appropriate lag length was determined using the Vector Autoregressive (VAR) model. Following the cointegration test, a vector error correction model was constructed. The results of trace and maximum eigenvalue were obtained within the scope of the Johansen cointegration test. The trace test results indicated that there was at least one, and possibly more than one, cointegration relationship among all four series; the maximum eigenvalue test results indicated that at least four variables had a cointegration relationship. The cointegration equation results indicated that the effects of the variables total debt service as a percentage of gross domestic product and short-term debt as a percentage of total external debt on neonatal mortality rate were positive and significant. When the vector error correction model results were examined, the error correction coefficient was found to be significant only for the variable short-term debt as a percentage of total external debt.

Keywords: Debt, health status, Türkiye, time series.

FULL TEXTS BOOK

GİRİŞ

Borçlanma, akıllıca yapıldığında finansal refahı artırabilse de bilinçsiz ve aşırı borçlanma zararlı etkilere sahip olabilir. Borçlanmanın zararlı etkilerinin görüldüğü konulardan biri de sağlık statüsüdür (Mudrazija ve Butrica, 2023). Borçlanma sonucu ortaya çıkan yüksek maliyetler stresi artırmakta ve bu durum başta mental sağlık olmak üzere birçok sağlık statüsü göstergesini olumsuz etkilemektedir. Borçlanma, ayrıca, başta kabul edilebilir bir düzeyde refahı sağlayacak yaşam şartları ve sağlık hizmetleri için gerekli harcamalarda kesintiye gidilmesine zemin hazırlamaktadır ki bu sebeplerle sağlık statüsünü olumsuz etkilemektedir (Frech vd., 2021).

Borçlanma ve sağlık statüsü ilişkisi, literatürdeki farklı çalışmalarda ele alınmıştır. Bu kapsamda aşağıda, borçlanmanın sağlık statüsü üzerindeki etkisini inceleyen çeşitli çalışmalara yer verilmiştir.

Keese ve Schmitz (2014), Almanya'nın 1999-2009 yılları arasındaki verilerini kullanarak hane halkı borçluluğu ile farklı sağlık sonuçları arasındaki ilişkiyi analiz etmişlerdir. Analiz sonuçlarına göre, borçluluğun; sağlık memnuniyeti, ruh sağlığı ve obezite ile güçlü bir şekilde ilişkili olduğu bulunmuştur.

Clayton ve diğerleri (2015) çalışmasında, 1995-2012 dönemine ve 17 Avrupa ülkesine ait veriler kullanılarak hane halkı borcu ve sağlık sonuçları arasındaki ilişki incelenmiştir. Analiz sonuçlarına göre, hane halkı borcunun sağlık sonuçlarını etkilediği ve etkinin borcun vadesine göre değişkenlik gösterdiği saptanmıştır. Buna göre hem kısa hem de orta vadeli borcun sağlık sonuçları üzerinde olumlu bir etkisi vardır. Uzun vadeli teminatsız borç ve ipotek borcu ise, daha kötü sağlık sonuçlarıyla ilişkili bulunmuştur.

Daoud (2020), 57 düşük ve orta gelirli ülkenin 1990-2015 dönemine ait verilerini kullanarak borçlanma ile çocuk ölümleri arasındaki ilişkiyi ele almıştır. Analiz sonuçlarına göre, borçlanma çocuk ölümlerini anlamlı bir şekilde artırmaktadır.

Ma ve diğerleri (2022), 1991-2019 dönemine ve gelişmekte olan bazı Asya ülkelerine (Bangladeş, Malezya, Filipinler, Tayland, Sri Lanka, Çin ve Hindistan) ait verileri kullanarak bu ülkelerde borcun sağlık sonuçları üzerindeki etkilerini araştırmışlardır. Çalışmanın bulguları, borcun uzun vadede gelişmekte olan Asya ekonomilerinde bebek ölüm hızını artırdığını ve yaşam beklentisini düşürdüğünü ortaya koymuştur.

Aladejare (2023), 1981-2020 dönemine ve Batı Afrika'da bulunan gelişmekte olan ülkelere ait verileri kullanarak dış borcun yaşam beklentisi üzerindeki etkisini değerlendirmiştir. Yazar, dış borcun ve makroekonomik oynaklığın Batı Afrika ülkelerinde yaşam beklentisini uzun vadede kısalttığını göstermiştir.

Dinga ve diğerleri (2024), 2000-2022 dönemine ve 13 Güney Afrika Kalkınma Topluluğu ülkesine ait verileri kullanarak borçlanma ile anne ölüm oranları arasındaki ilişkiyi araştırmışlardır. Sonuçlar, Güney Afrika Kalkınma Topluluğu bölgesi ülkelerinde dış borcun anne ölümlerini azalttığını ortaya koymuştur. Çalışma sonuçları, ayrıca, anne ölümleri ile dış borç arasında çift yönlü bir ilişki olduğunu ortaya koymuştur.

Audi ve Roussel (2024), Suudi Arabistan'ın 1974-2023 dönemine ait verileri kullanarak bu ülkede yaşam beklentisi ile dış borç arasındaki ilişkiyi inceledikleri çalışmalarında, yaşam beklentisi ile dış borç arasında hem kısa hem de uzun vadede anlamlı bir ilişki olduğunu göstermiştelerdir.

Yukarıdaki benzer çalışmalarda da görülebileceği üzere, borçlanma ve sağlık statüsü arasında anlamlı bir ilişki vardır. Bu ilişkiden yola çıkılan mevcut çalışmada, borçlanmanın sağlık statüsü üzerindeki etkisi Türkiye özelinde ele alınmaktadır. Çalışma, bir zaman serisi analizi içermekte olup borçlanmaya ilişkin çeşitli göstergelerin, toplumlar için önemli bir sağlık statüsü göstergesi olan yenidoğan ölüm hızı üzerindeki etkisini incelemektedir. Literatür incelendiğinde, borçlanmanın spesifik bir sağlık göstergesi olan yenidoğan ölüm hızı üzerindeki etkisini Türkiye özelinde inceleyen güncel bir çalışmaya rastlanılmamıştır ki bu durum, mevcut çalışmanın literatüre katkısı olarak değerlendirilmektedir.

FULL TEXTS BOOK

Çalışma kapsamında ilerleyen başlıklarda, öncelikle çalışmanın yöntemi hakkında bilgi verilmektedir. Bunu takiben bulgular sunulmakta ve çalışma, bulgulara dayalı somut önerilerin yer aldığı bölümle sonlandırılmaktadır.

YÖNTEM

Bu çalışmada, 1.000 canlı doğumda yenidoğan ölüm hızı, gayrisafi yurtiçi hasılaya oranla merkezi hükümet borcu, toplam dış borcun yüzdesi olarak kısa vadeli borçlar ve gayrisafi yurtiçi hasılaya oranla toplam borç servisi değişkenleri arasındaki eş-bütünleşiklik ilişkisi ele alınmaktadır. Çalışmanın analizleri Türkiye'nin 1985-2023 dönemine ait verileri ile gerçekleştirilmiş olup veriler, Dünya Bankası (2025) veri tabanından alınmıştır.

Çalışmada, öncelikle, tanımlayıcı istatistikler elde edilmiş ve sonrasında tüm değişkenlerin doğal logaritmaları (LN) alınmıştır. Sonraki tüm analizler LN alınmış değişkenler ile yapılmıştır. Bu kapsamda, ilk olarak, değişkenler arasındaki korelasyon düzeyleri incelenmiştir. Sonrasında, Augmented Dickey–Fuller Testi ile değişkenlerin hangi seviyede durağan hale geldiği ortaya konmuş ve tüm değişkenlerin birinci derece farkta durağan hale geldiği görülmüştür. Bu bulgunun bir gereği olarak Johansen eş-bütünleşme testi kullanılmıştır. Johansen eş-bütünleşme testi öncesinde Vektör Otoregresif (VAR) modeli ile uygun gecikme uzunluğuna karar verilmiştir. Eş-bütünleşme testi sonrasında, vektör hata düzeltme modeli kurulmuştur.

Çalışmanın analizlerinden elde edilen sonuçlar için güven düzeyi %95 olarak belirlenmiştir ve analizler EViews 14 paket programı ile gerçekleştirilmiştir.

BULGULAR

Çalışmada, 1.000 canlı doğumda yenidoğan ölüm hızı, merkezi hükümet borcunun gayrisafi yurtiçi hasılaya (GSYH) oranı, toplam dış borç içindeki kısa vadeli borçların oranı ve toplam borç servisinin GSYH'ye oranı değişkenleri arasında uzun dönemli bir ilişki olup olmadığı Johansen eş-bütünleşme testi ile incelenmiştir. Söz konusu değişkenler Tablo 1'de sunulmaktadır.

Tablo 1. Çalışmanın Değişkenleri

Değişken	Değişken Açıklaması	Kısaltması	LN Dönüşümü	Dönemi	Veri Kaynağı
1.000 canlı doğumda yenidoğan ölüm hızı	Canlı doğan her 1.000 bebekten, doğumdan sonraki ilk 28 gün içinde ölen bebeklerin sayısını ifade etmektedir.	NEOMORT	LNNEOMORT	1985-2023	Dünya Bankası (2025)
Gayrisafi yurtiçi hasılaya oranla merkezi hükümet borcu	Merkezi hükümetin sahip olduğu toplam borç stokunun, gayrisafi yurtiçi hasılaya oranını ifade etmektedir.	GOVDEPT	LNGOVDEPT	1985-2023	Dünya Bankası (2025)
Toplam dış borcun yüzdesi olarak kısa vadeli borçlar	Vadesi 1 yıl veya daha kısa olan dış borçların, toplam dış borç stoku içindeki oranını ifade etmektedir.	STDEPT	LNSTDEPT	1985-2023	Dünya Bankası (2025)
Gayrisafi yurtiçi hasılaya oranla toplam borç servisi	Dış borçlara ilişkin yıllık faiz ödemeleri ve anapara geri ödemelerinin toplamının gayrisafi yurtiçi hasılaya oranını ifade etmektedir.	TDEBTS	LNTDEBTS	1985-2023	Dünya Bankası (2025)

Tablo 2'de çalışmanın değişkenlerine ilişkin tanımlayıcı istatistikler sunulmaktadır. Buna göre, 1.000 canlı doğumda yenidoğan ölüm hızı (NEOMORT) değişkeni, ortalama 16,82 olup geniş bir dağılıma sahiptir; minimum değeri 4,90 iken maksimum değeri 37,90'dır ve standart sapması 10,74'tür. Merkezi

FULL TEXTS BOOK

hükümet borcunun GSYH'ye oranı (GOVDEPT) ortalama %36,84 düzeyindedir ve %28,93 ile %48,47 arasında değişmektedir. Kısa vadeli dış borçların toplam dış borç içindeki oranı (STDEPT) ortalama %23,01 olup, %13,81 ile %35,28 arasında gözlemlenmiştir. Toplam borç servisinin GSYH'ye oranı (TDEBTS) ise ortalama %7,78'dir ve en düşük %4,84, en yüksek %12,12 olarak kaydedilmiştir. Jarque-Bera testine ait p-değerleri serilerin normal dağıldığını göstermektedir (p>0,05).

Tablo 2. Tanımlayıcı İstatistikler

	NEOMORT	GOVDEPT	STDEPT	TDEBTS
Ortalama	16,82	36,84	23,01	7,78
Maksimum	37,90	48,47	35,28	12,12
Minimum	4,90	28,93	13,81	4,84
Standart sapma	10,74	6,14	5,72	1,89
p(Jarque-Bera)	0,14	0,16	0,30	0,12
Gözlem sayısı	39	39	39	39

Tablo 3, çalışmada kullanılan değişkenler arasındaki Pearson korelasyon katsayılarını göstermektedir. Buna göre, LNNEOMORT ile LNSTDEPT arasında pozitif ve istatistiksel olarak anlamlı bir ilişki bulunmaktadır (r =0,65; p<0,05). Benzer şekilde, LNNEOMORT ile LNTDEBTS arasında da anlamlı ve pozitif yönlü bir ilişki tespit edilmiştir (r =0,47; p<0,05). Diğer yandan, LNGOVDEPT ile LNNEOMORT arasında zayıf ve istatistiksel olarak anlamsız bir ilişki mevcuttur (r =-0,15; p>0,05). LNGOVDEPT ile LNTDEBTS arasında pozitif ve anlamlı bir ilişki gözlenmiştir (r = 0,32; p<0,05). LNGOVDEPT ile LNSTDEPT arasındaki zayıf negatif ilişki anlamsızdır (r=-0,01; p>0,05). Son olarak, LNSTDEPT ile LNTDEBTS arasındaki zayıf negatif ilişki de anlamsızdır (r=-0,19; p>0,05).

Tablo 3. Korelasyon Analizi

	(1)	(2)	(3)	(4)
LNNEOMORT (1)	1			
LNGOVDEPT (2)	-0,15	1		
LNSTDEPT (3)	0,65**	-0,01	1	
LNTDEBTS (4)	0,47**	0,32*	-0,19	1

** p<0,01; * p<0,05

Tablo 4'te, serilerin düzeyde (I(0)) veya birinci farkta (I(1)) durağan olup olmadığını belirlemek amacıyla uygulanan Augmented Dickey–Fuller (ADF) test sonuçları gösterilmektedir. Düzey değerlerinde hem sabitli hem de trendli modeller altında yapılan test sonuçları, hiçbir serinin düzeyde durağan olmadığını göstermektedir (p>0,05). Dolayısıyla, serilerin birim kök içerdiği söylenebilir. Ancak birinci farklar alındığında, tüm seriler hem sabitli hem de trendli modellerde durağan hale gelmektedir (p<0,05). Görüldüğü gibi seriler I(1) seviyesinde entegredir ve bu bulgu serilerin Johansen eş-bütünleşme analizine uygun olduğunu göstermektedir.

Tablo 4. Augmented Dickey–Fuller Testi Sonuçları

	Seviyede I(0)				Birinci Farkta I(1)			
	Sabitli		Trendli ve Sabitli		Sabitli		Trendli ve Sabitli	
	t	p	t	p	t	p	t	p
LNNEOMORT	-1,71	0,38	-1,44	0,82	-5,05	0,001**	-4,81	0,001**
LNGOVDEPT	-2,75	0,07	-2,70	0,24	-5,01	0,001**	-4,92	0,001**
LNSTDEPT	2,23	0,19	-3,16	0,11	-4,21	0,001**	-4,16	0,01*
LNTDEBTS	0,32	0,90	-0,51	0,97	-4,68	0,001**	-5,22	0,001**

** p<0,01; * p<0,05

Tablo 5'te, çok değişkenli zaman serisi analizlerinde kullanılan Vektör Otoregresif (VAR) modeline ilişkin uygun gecikme uzunluğunun belirlenmesinde kullanılan farklı bilgi kriterlerini sunmaktadır. Buna göre, en uygun gecikme uzunluğunu 2 olarak belirlemiştir zira farklı kriterler arasında en fazla 2 tercih edilmiştir.

FULL TEXTS BOOK

Tablo 5. VAR Sonuçları

Lag	LogL	LR	FPE	AIC	SC	HQ
0	15,05	NA	6,25e-06	-0,63	-0,45	-0,57
1	186,01	293,05	8,99e-10	-9,48	-8,59*	-9,17
2	210,07	35,76*	5,90e-10*	-9,94*	-8,34	-9,39*
3	222,50	15,63	8,01e-10	-9,74	-7,43	-8,94
4	236,08	13,96	1,13e-09	-9,61	-6,58	-8,56

* kriter tarafından seçilen gecikme sırasını göstermektedir.
LR: sequential modified LR test statistic (each test at 5% level)
FPE: Final prediction error
AIC: Akaike information criterion
SC: Schwarz information criterion
HQ: Hannan-Quinn information criterion

Tablo 6'da, Johansen eş-bütünleşme testi kapsamında İz (Trace) ve Maksimum Özdeğer (Max-Eigenvalue) test sonuçları sunulmaktadır. Bu bulgulardan hareketle, iz testi sonuçlarına göre dört serinin tamamı arasında en az bir ve muhtemelen birden fazla eş-bütünleşme ilişkisinin bulunduğunu; maksimum özdeğer testi sonuçlarına göre ise, en az dört değişkenin bir eş-bütünleşme ilişkisi içinde olduğunu söylemek mümkündür ($p<0,05$).

Tablo 6. İz ve Maksimum Özdeğer Sonuçları

İz (Trace) Testi				
Hipotezler	Özdeğer	İz istatistiği	0,05 Kritik değer	p
Eş-bütünleşme yoktur*	0,46	58,01	47,85	0,004**
En fazla 1 eş-bütünleşme vardır*	0,39	35,04	29,79	0,011*
En fazla 2 eş-bütünleşme vardır*	0,28	16,15	15,49	0,039*
En fazla 3 eş-bütünleşme vardır*	0,10	3,94	3,84	0,047*
Maksimum Özdeğer (Max-Eigenvalue) Testi				
Hipotezler	Özdeğer	Maksimum Özdeğer	0,05 Kritik değer	p
Eşbütünleşme yoktur	0,46	22,95	27,58	0,175
En fazla 1 eş-bütünleşme vardır	0,39	18,89	21,13	0,100
En fazla 2 eş-bütünleşme vardır	0,28	12,21	14,26	0,102
En fazla 3 eş-bütünleşme vardır*	0,10	3,94	3,84	0,047*

** $p<0,01$; * $p<0,05$

Eşitlik (1)'de eş-bütünleşme denklemi sunulmaktadır. Eş-bütünleşme denklemi incelendiğinde, LNSTDEPT değişkenin LNNEOMORT üzerindeki etkisinin pozitif ve anlamlı olduğu (katsayı = 4,28; t = 9,09); benzer şekilde, LNTDEBTS değişkenin de LNNEOMORT üzerindeki etkisinin pozitif ve anlamlı olduğu (katsayı = 1,31; t = 2,92) söylenebilir. LNGOVDEPT değişkenin etkisi ise, anlamsızdır (katsayı = -1,30; t = -1,09).

$$LNNEOMORT_{t-1} = -1{,}30LNGOVDEPT_{t-1}+4{,}28LNSTDEPT_{t-1}+1{,}31LNTDEBTS_{t-1}+13{,}82 \quad \textit{Eşitlik (1)}$$

Tablo 7'de sunulan vektör hata düzeltme modeli (VECM) sonuçları incelendiğinde, hata düzeltme katsayısının (CointEq1); LNNEOMORT, LNGOVDEPT ve LNTDEBTS için anlamlı olmadığı; LNSTDEPT için ise, negatif ve anlamlı olduğu yorumu yapılabilir. Buradan hareketle, LNSTDEPT değişkeninin uzun dönem dengeye dönüş sürecinde önemli bir düzeltici rol üstlendiği söylenebilir.

FULL TEXTS BOOK

Tablo 7. Vektör Hata Düzeltme Modeli Sonuçları

Hata Düzeltme	D(LNNEOMORT)	D(LNGOVDEPT)	D(LNSTDEPT)	D(LNTDEBTS)
CointEq1	0,01	-0,09	-0,24	0,08
	(0,01)	(0,05)	(0,07)	(0,08)
	[0,79]	[-1,80]	[-3,41]	[0,93]
D(LNNEOMORT(-1))	0,56	-0,97	4,76	-1,85
	(0,19)	(2,88)	(4,01)	(4,79)
	[2,84]	[-0,33]	[1,19]	[-0,38]
D(LNNEOMORT(-2))	0,36	4,27	-2,42	1,65
	(0,18)	(2,65)	(3,68)	(4,40)
	[2,01]	[1,60]	[-0,65]	[0,37]
D(LNGOVDEPT(-1))	-0,01	-0,63	-0,26	-0,44
	(0,01)	(0,23)	(0,33)	(0,39)
	[-1,10]	[-2,66]	[-0,80]	[-1,13]
D(LNGOVDEPT(-2))	-0,01	-0,48	-0,36	0,06
	(0,01)	(0,22)	(0,31)	(0,37)
	[-1,12]	[-2,11]	[-1,16]	[0,17]
D(LNSTDEPT(-1))	-0,01	0,311	0,56	-0,16
	(0,01)	(0,19)	(0,27)	(0,32)
	[-0,61]	[1,57]	[2,05]	[-0,51]
D(LNSTDEPT(-2))	0,01	0,32	0,12	0,01
	(0,01)	(0,19)	(0,27)	(0,32)
	[0,14]	[1,63]	[0,45]	[0,01]
D(LNTDEBTS(-1))	-0,01	0,14	0,25	0,07
	(0,01)	(0,18)	(0,25)	(0,30)
	[-0,41]	[0,77]	[1,01]	[0,24]
D(LNTDEBTS(-2))	-0,01	0,310	-0,05	-0,13
	(0,01)	(0,17)	(0,24)	(0,29)
	[-0,47]	[1,77]	[-0,23]	[-0,46]
C	-0,01	0,18	0,14	-0,01
	(0,01)	(0,09)	(0,12)	(0,14)
	[-0,49]	[2,01]	[1,13]	[-0,05]
R^2	0,80	0,33	0,42	0,19
Düzeltilmiş R^2	0,74	0,10	0,24	0,09

( ) içindekiler standart hataları ve [ ] içindekiler t-istatistiklerini göstermektedir.

SONUÇ

Çalışmanın sonuçlarına göre, çalışmanın modelinde yer alan değişkenler arasında uzun dönemli eş-bütünleşik ilişkiler mevcuttur ve özellikle kısa vadeli dış borçların toplam dış borç stoku içindeki ağırlığı hem uzun dönem denge ilişkisi içinde güçlü ve anlamlı bir role sahiptir hem de kısa vadede dengeye dönüş sürecinde etkilidir. Çalışmanın sonuçlarından, ayrıca, gayrisafi yurtiçi hasılaya oranla toplam borç servisinin uzun dönemde anlamlı bir etki gösterdiği ancak yenidoğan ölüm hızının dengeye dönüş sürecinde aktif bir düzeltici rol üstlenmediği, dolayısıyla pasif bir şekilde tepki verdiği görülmektedir.

Görüldüğü gibi ülkelerin kısa vadeli borçlanmaya bağımlılığının sağlık sonuçları üzerinde olumsuz sonuçları mevcuttur. Bununla birlikte, kısa vadeli dış borçların dengeye dönüş sürecinde anlamlı bir düzeltme gücüne sahip olduğu anlaşılmaktadır. Bu bulgu, kısa vadeli borçların yalnızca uzun vadede değil, aynı zamanda sistemin dengeye uyum sürecinde de kritik bir rol oynadığını göstermektedir.

Çalışmanın bulguları doğrultusunda, özellikle kısa vadeli dış borçlanmanın azaltılmasına yönelik mali ve borçlanma politikaları geliştirilmesi gerektiği önerilmektedir. Borçlanma yapısında uzun vadeli ve sürdürülebilir finansman kaynaklarına öncelik verilmesi hem ekonomik istikrar hem de sağlık sonuçlarının iyileştirilmesi açısından önem taşımaktadır. Ayrıca, borç servisi yükünü azaltıcı reformların ve bütçe disiplinine yönelik uygulamaların desteklenmesi, doğrudan insan yaşam kalitesini etkileyen sağlık göstergelerinde iyileşme sağlayabilir. Son olarak, bu tür göstergeler ışığında kamu borçlanma stratejileri, sadece mali sürdürülebilirlik açısından değil, aynı zamanda sosyal kalkınma ve sağlık politikaları açısından da entegre bir biçimde ele alınmalıdır.

FULL TEXTS BOOK

KAYNAKÇA

Aladejare, S. A. (2023). Does external debt promote human longevity in developing countries? Evidence from West African countries. *Fudan Journal of the Humanities and Social Sciences*, *16*(2), 213-237.

Audi, M., & Roussel, Y. (2024). Exploring the Link Between Public Health and External Debt in Saudi Arabia. *Journal of Business and Economic Options*, *7*(4), 1-12.

Clayton, M., Liñares-Zegarra, J., & Wilson, J. O. (2015). Does debt affect health? Cross country evidence on the debt-health nexus. *Social science & medicine*, *130*, 51-58.

Daoud, A. (2020). The wealth of nations and the health of populations: A quasi-experimental design of the impact of sovereign debt crises on child mortality. *arXiv preprint arXiv:2012.14941*.

Dinga, G. D., Mah, G., & Mosikari, T. (2024, October). Government health expenditure and maternal mortality: the moderating role of external debt. *Healthcare*, Vol. 12, No. 20.

Dünya Bankası (2025). Data Bank. https://data.worldbank.org/ 18/07/2025.

Frech, A., Houle, J., & Tumin, D. (2021). Trajectories of unsecured debt and health at midlife. *SSM-Population Health*, *15*, 100846.

Keese, M., & Schmitz, H. (2014). Broke, ill, and obese: Is there an effect of household debt on health? *Review of Income and Wealth*, *60*(3), 525-541.

Ma, Y., Hu, M., & Zafar, Q. (2022). Analysis of the impact of external debt on health in an emerging Asian economy: Does FDI matter?. *Frontiers in Public Health*, *10*, 824073.

Mudrazija, S., & Butrica, B. A. (2023). How does debt shape health outcomes for older Americans? *Social Science & Medicine*, *329*, 116010.

OKUL ÖNCESİ DÖNEMDE HASTALIK VE HASTANE: ÇOCUKLARIN SÖZEL VE GÖRSEL ANLATIMLARI ÜZERİNDEN BİR DEĞERLENDİRME

ILLNESS AND HOSPITAL IN THE PRESCHOOL PERIOD: AN EVALUATION BASED ON CHILDREN'S VERBAL AND VISUAL EXPRESSIONS

Ramazan Rüçhan KAYA

Öğr.Gör. Dr., Uşak Üniversitesi, Sağlık Hizmetleri Meslek Yüksekokulu,

Yönetim ve Organizasyon Bölümü, Uşak, Türkiye.

Şevval Erva KAYA

Yüksek Lisans Öğrencisi, Afyon Kocatepe Üniversitesi, Sosyal Bilimler Enstitüsü,

Temel Eğitim Anabilim Dalı/ Okul Öncesi Programı, Afyon, Türkiye.

ÖZET

Bu araştırma, okul öncesi dönemdeki çocukların hastalık, hastane ve hekimlere ilişkin algılarını incelemeyi amaçlamaktadır. Çalışma, nitel araştırma yöntemi ile kurgulanmış olup, katılımcılar amaçlı örnekleme ve kartopu örnekleme yöntemiyle belirlenmiştir. Araştırma kapsamında okul öncesi eğitimi almakta olan 5 ve 6 yaşlarında, toplam 25 (15 kız, 10 erkek) çocuktan yapılandırılmış görüşme formu aracılığıyla veri toplanmıştır. Elde edilen veriler tematik analiz yöntemiyle çözümlenmiştir. Analiz sonucunda; çocukların hastalığı genellikle göstermiş olduğu belirtileriyle anlamlandırdıkları, sağlıksız olma, mikrop kapma gibi ifadelerle tanımladığı ve ateş, öksürük ve hapşırık gibi belirtilerle açıkladığı tespit edilmiştir. Ayrıca katılımcı çocuklar hastalık ile ilgili değerlendirmelerinde, hasta bireyin özelliklerine değinerek; "fiziksel olarak zayıf", "solgun" ve "yorgun" olarak betimlenmiş; kimi zaman "mutlu" kimi zaman "mutsuz" ifadeleriyle tanımlanmıştır. Hastaneye dair algıların ise mimari unsurlar, çalışanlar, hastaneden beklentiler, hastaneye gidildiğinde hissedilenler ve hastane ortamına yönelik kavramlardan oluştuğu tespit edilmiştir. Okul öncesi dönemde bulunan çocukların hekimlere ilişkin yanıtlarında; hekimin davranışlarına, hekime karşı bireysel tutumlarına, hekimin kıyafetlerine ve hekimin kullandığı aksesuarlara vurgu yaptıkları tespit edilmiştir. Bu kapsamda hekimlerin genellikle "mavi veya beyaz önlük (kazak)" ya da "tişört giyen", "gözlük takan" kişiler olarak değerlendirildiği ayrıca hekimlerin steteskop ile resmedildiği görülmüştür. Ayrıca çocukların hekimlere karşı tutumlarında; kendilerini iyileştirdiği için bir minnet vurgusu olduğu tespit edilmiştir. Bu bulgular, çocukların sağlıkla ilişkili kavramlara dair algılarının gözlemlerine ve kendi gelişim düzeylerine uygun biçimde şekillendiğini göstermektedir. Araştırma, çocukların bakış açısından sağlık kurumlarını anlamaya katkı sağlamakta ve çocuk dostu sağlık ortamlarının geliştirilmesine yönelik öneriler sunmaktadır.

Anahtar Kelimeler: Hekim, Hastane, Hastalık, Çocuklar, Okul Öncesi Eğitimi.

ABSTRACT

This study aims to examine the perceptions of preschool children about illness, hospitals, and doctors. The study was designed with qualitative research method and the participants were determined by purposive sampling and snowball sampling method. Within the scope of the study, data were collected from a total of 25 children (15 girls, 10 boys) aged 5 and 6 who were receiving preschool education through a structured interview form. The obtained data were analyzed using thematic analysis. As a result of the analysis, it was determined that children generally make sense of the illness through its symptoms, define it with terms such as being unhealthy and catching an infection, and explain it with

FULL TEXTS BOOK

symptoms such as fever, cough, and sneezing. In addition, in their evaluations of the illness, the participant children referred to the characteristics of the patient, describing them as “physically weak,” “pale,” and “tired”; they were sometimes described as “happy,” sometimes “unhappy.” It was determined that the perceptions about the hospital consisted of architectural elements, staff, expectations from the hospital, feelings felt when going to the hospital, and concepts related to the hospital environment. In the responses of the preschool children about doctors; It was found that children emphasized the doctor's behavior, their individual attitudes toward the doctor, the doctor's attire, and the accessories they used. In this context, doctors were generally depicted as wearing "blue or white coats (sweaters)" or "T-shirts" or "glasses." Furthermore, doctors were often depicted with stethoscopes. Furthermore, children's attitudes toward doctors emphasized gratitude for their healing. These findings suggest that children's perceptions of health-related concepts are shaped by their observations and their own developmental levels. The research contributes to understanding healthcare institutions from children's perspectives and offers suggestions for developing child-friendly healthcare environments.

Keywords: Physician, Hospital, Illness, Children, Preschool Education.

GİRİŞ

Hastaneler, bireylerin yaşam döngüsü içerisinde sağlık hizmeti almak amacıyla başvurdukları temel kurumlar arasında yer almaktadır. Ancak hastaneler, özellikle çocuklar açısından yalnızca tedavi edici bir kurum değil, aynı zamanda duygusal ve bilişsel açıdan farklı deneyimlerin yaşandığı mekânlar olarak da önem taşır. Çocukluk dönemi, bireyin çevresini algılama, deneyimlerini anlamlandırma ve kavramlara yönelik bilişsel şemalarını oluşturma sürecinin en hızlı ilerlediği gelişimsel evredir (Piaget, 1952; Vygotsky, 1978). Bu süreçte çocukların sağlık kurumlarına, hekimlere ve hastalık kavramına ilişkin edindikleri deneyimler, onların sağlıkla ilgili tutumlarını, tedavi süreçlerine uyumlarını ve sağlık profesyonellerine duydukları güveni doğrudan etkilemektedir (Dalley ve ark., 2019).

Çocukların hastane ve sağlık çalışanlarına ilişkin algıları yalnızca bireysel deneyimlere dayanmaz. Aile tutumları, akran etkisi, medya aracılığıyla edinilen temsiller ve toplumsal kültür, çocukların zihinsel şemalarını şekillendiren önemli unsurlardır (McGrath, 2018). Örneğin, bazı çocuklar için hastane, iyileşmenin, bakımın ve güvenliğin sembolü olurken; bazıları için korku, acı ve yabancılık hissi uyandıran bir mekân olarak algılanabilmektedir (Coyne, 2006). Bu algılar yalnızca kısa vadeli psikolojik etkiler yaratmakla kalmaz, aynı zamanda bireylerin ilerleyen yaşamlarında sağlık hizmetlerinden yararlanma tutumlarına kadar uzanan uzun vadeli sonuçlar doğurabilmektedir (Pelander ve Leino-Kilpi, 2010). Son yıllarda çocukların sağlık kurumlarıyla ilişkilerinde “çocuk dostu hastane” anlayışı önem kazanmıştır. Dünya Sağlık Örgütü (WHO, 2012) ve UNICEF gibi uluslararası kuruluşlar, çocukların sağlık hizmetlerine erişiminde onların psikososyal ihtiyaçlarının da göz önünde bulundurulması gerektiğini vurgulamaktadır. Bu bağlamda çocukların algılarının anlaşılması, hem sağlık profesyonelleri hem de sağlık hizmetlerinin yeniden yapılandırılması açısından kritik bir öneme sahiptir.

Bu araştırmanın temel amacı, çocukların hastane, hekim ve hastalık kavramlarına ilişkin algılarını görsel (çizim vb.) ve sözel (anlatım, açıklama vb.) veriler aracılığıyla ortaya koymaktır. Böylelikle elde edilen bulgular doğrultusunda, sağlık hizmetlerinin çocuk dostu bir yaklaşımla geliştirilmesine yönelik çıkarımlar sunmak hedeflenmektedir.

YÖNTEM

Araştırma kapsamında, geleceğin yetişkinleri olan çocukların sağlık hizmetlerine ilişkin algılarının, sağlık kurumu, sağlık çalışanı ve sağlık durumuna ilişkin değerlendirmeleri aracılığıyla tespit edilmesi amacıyla nitel araştırma yöntemi tercih edilmiş, katılımcılar amaçlı örnekleme ve kartopu örnekleme tekniğiyle belirlenmiştir. Bu kapsamda Afyon ve Isparta illerinde okul öncesi eğitim almakta olan 25 çocuktan veri toplanmıştır. Veri toplama aracı olarak; verilerin daha kolay toplanma, işlenme ve

FULL TEXTS BOOK

düzenlenme kolaylığı sebebiyle yapılandırılmış soru formu kullanılmıştır (Gürbüz ve Şahin, 184-185). Bu minvalde çocukların; hekimler, hastaneler, hastalıklar ve hasta bireyler hakkındaki düşünceleri sorgulanmış ayrıca hekimler, hastaneler ve hasta bireyleri resmederek betimlemeleri istenmiştir. Elde edilen sözel veriler araştırmacılar tarafından not alınarak kayda geçirilmiş, görsellerle birlikte veri tabanı oluşturulmuştur. Elde edilen veriler tematik analiz kapsamında değerlendirilmiştir.

BULGULAR

Elde edilen verilerin analizi sonucunda, okul öncesi dönemdeki çocukların sağlık hizmetlerine yönelik algıları, "hastanelere ilişkin bulgular", "hekimlere ilişkin bulgular" ve "hastalık ve hasta bireye ilişkin bulgular" başlıklarında değerlendirilmiştir.

Tablo 1. Hastanelere ilişkin bulgular

Tema	Alt Tema	Kod
Mimari	Kat	*Çok Büyük, Çok Katlı (2-3-4-5-6-7)*
	Renk	*Mavi, Kırmızı, Beyaz*
Çalışanlar		*Doktor-Hemşire (Yardımcı)*
Beklentiler		*İyileşmek, Ağrının Geçmesi, Ateş Düşmesi*
Hissedilen	Olumlu	*Güzellik, Mutluluk, İyi Hissetme*
	Olumsuz	*Korku, Mutsuzluk, Kötülük*
Hastane Ortamı	Fiziksel Mekanlar	*Yataklar, Odalar, Koltuklar, Asansörler, Merdivenler*
	Hizmet Süreçleri	*Sıra Almak, Muayene Olmak*
	Sağlık Hizmeti	*İlaçlar, Hastalar, Doktorlar, Hemşireler,*
	Diğer	*Temizlik, Ev*

Araştırma kapsamında çocukların hastanelere yönelik olarak kullanmış oldukları ifadeler ve çizmiş oldukları resimler kapsamında; çocukların hastaneleri, mimari özellikleri, hastanelerde çalışanlar, hastanelere yönelik beklentileri, hastaneye gittiklerinde hissettikleri ve hastane ortamına yönelik ifadelerle değerlendirdikleri tespit edilmiştir. Bu kapsamda, çocuklara göre hastaneler genellikle çok büyük ve çok katlı yapılardır ve genellikle kırmızı, mavi veya beyaz renkli binalardır. Bu tespitler çocukların hastaneleri büyük ve karmaşık ortamlar olarak algıladıklarını göstermektedir.

Çocuklar hastane çalışanlarına yönelik olarak genellikle "doktor amca", "hemşire abla" gibi ifadelerle hekim ve hemşirelere yönelik bilgilerini ortaya koymaktadır. Hastanelere yönelik beklentiler kapsamında, çocukların hastanelere gittiklerinde ağrılarının geçmesi ve iyileşmek yönünde beklentileri olduğu tespit edilmiştir. Aynı zamanda hastaneye gittiklerindeki hislerini paylaşan çocukların genellikle hastanelerden korktuğu, mutsuz olduğu görülmekle birlikte, hastanedeki sürecin ardından, iyi hissetme hali ve mutluluğun geleceğine yönelik duygularını ifade ettikleri tespit edilmiştir. Hastane ortamına

FULL TEXTS BOOK

yönelik ifadelerin ise; yataklar, odalar, koltuklar vb. gibi fiziksel ortam, sıra almak muayene olmak gibi hizmet süreçleri, ilaçlar, hastalar, doktorlar, hemşireler gibi sağlık hizmeti ve temiz olan ve ev gibi diğer kavramlardan oluştuğu tespit edilmiştir. Bu bulgular ışığında, çocukların hastanelerin şekil özelliklerine, çalışanlarına yönelik fikir sahibi olduğu, hastane ortamında yaşananlara ve ortamdaki fiziksel değişkenleri tanıdıkları, hastanelerden korkmakla birlikte orada iyileşeceklerinin bilincinde oldukları tespit edilmiştir.

Tablo 2. Hekimlere ilişkin bulgular

Tema	Kod
Hekimin Davranışları	*Muayene Yapmak, Ellerini Yıkama, İğne Yapmak, Güler Yüzlü Olmak, Akıllı Olmak, İlaç Vermek*
Hekime Karşı Tutum	*Minnet, (İyileşme), Sevgi, İyilik, Korku*
Hekimin Kıyafeti	*Mavi Önlük, Beyaz Önlük, Ceket, Tişört, Kazak*
Hekimin Aksesuarları	*Steteskop (Kalp Dineleme Aleti Vb.), Maske, Eldiven, Gözlük*

Araştırma kapsamında, okul öncesi dönemde bulunan çocukların hekimlere ilişkin kullanmış oldukları ifadeler ve oluşturdukları görseller incelendiğinde; çocukların hekimleri tanımlarken, hekimin davranışları, hekime karşı bireysel tutumları, hekimin kıyafetleri ve hekimin kullandığı aksesuarlar üzerinden değerlendirdikleri tespit edilmiştir.

Bu kapsamda, çocuklara göre hekimler, muayene yapmakla görevli, ellerini yıkayarak hijyene dikkat eden, iğne yapan, ilaç veren, akıllı ve zeki aynı zamanda da güler yüzlü bireylerdir. Çocuklar tarafından yapılan resimlerde hekimler genellikle gülen bir yüz ile tasvir edilmiş, mavi veya beyaz kıyafetli olarak tasvir edilmiştir. Aynı zamanda çocuklar hekimlerin kullandığı aksesuar olarak, kalplerini dinlediğini ifade ettikleri steteskopa vurgu yapmıştır. Çocukların hekimlere karşı bireysel tutumlarında ise korku ve minnet vurgusu ön plana çıkmaktadır. Çocuklar her ne kadar hekimlere yönelik korku ile başlayan ifadelerde bulunsalar da onların hastalıklarını iyileştirdiklerinin ve minnet duyulması gereken bir iş yaptıklarının farkında oldukları tespit edilmiştir.

Tablo 3. Hastalıklar ve Hasta bireye ilişkin bulgular

Tema	Alt Tema	Kod
Hastalığı Tanımlama	Hastalığın Belirtileri	*Hapşırık, Öksürük, Mide Bulantısı, Burun Akıntısı, Burun Tıkanıklığı, Ağrı, Ateşlenmek, Baş Ağrısı, Kötü Hissetme*
	Hastalık Türleri	*Alerji, İshal, Grip*
	Hastalık Nedenleri	*Üşütmek, Mikrop, Pislik*
	Fiziksel Durum	*Yorgun (Bitkin), Uykulu, Yüzü Kırmızı*

FULL TEXTS BOOK

Hasta Durumu	**Bireyin**	Hijyen	*Pis*
		Duygusal Durum	*Mutsuz, Kötü, Korkulu*

Çocukların hastalıklara yönelik algılarının temelinde kendi başlarından geçen hastalık deneyimlerinin ve "hapşuruk, öksürük, mide bulantısı, burun akıntısı, burun tıkanıklığı, ağrı, ateşlenmek, baş ağrısı, kötü hissetme" gibi bu hastalıkların belirtilerinin olduğu tespit edilmiştir. Hastalık türlerine ilişkin ise çocukların genellikler "alerji, ishal ve grip" gibi hastalıklara aşina oldukları tespit edilmiştir. Ayrıca hastalıkları ortaya çıkaran sebeplerin ise yine kendi deneyimleri ve ebeveynlerinin ifadeleriyle, "üşütmek, mikroplar, pislik" gibi kavramlarla açıklandığı tespit edilmiştir.

Araştırma kapsamda, hasta bireyleri değerlendiren çocukların, hastaların fiziksel durumlarına, hijyen durumlarına ve duygusal durumlarına vurgu yaptıkları tespit edilmiştir. Bu doğrultuda; çocukların gözünde hasta bireylerin, yorgun, uykulu, hijyene dikkat etmeyen (pis), mutsuz, korkulu ve kötü durumdaki kişiler olduğu tespit edilmiştir.

SONUÇ VE DEĞERLENDİRME

Araştırma bulguları, çocukların hastalık kavramını çoğunlukla sağlıksız olma ve mikrop kapma gibi somut ve günlük yaşam deneyimlerinden beslenen ifadelerle tanımladıklarını ortaya koymuştur. Hastalık belirtilerini ise ateş, öksürük ve hapşırık gibi doğrudan gözlemlenebilir durumlarla açıklamışlardır. Hastane algıları, bir yandan çok büyük, yüksek katlı gibi mimari unsurlara; diğer yandan korku ve mutluluk gibi duygusal tepkilere dayanmaktadır. Çocukların hasta bireylere ilişkin değerlendirmeleri, hem fiziksel görünüm hem de duygusal durumlarına yönelik betimlemeleri içermektedir. Hekim algılarında ise doktorlar genellikle mavi ya da beyaz elbise giyen, gözlük takan, steteskop kullanan kişiler olarak resmedilmiştir. Çocukların doktorlara yönelik tutumları çoğunlukla sevgi ve minnet içerirken, zaman zaman korku hissini de barındırmaktadır. Bu sonuçlar, çocukların sağlıkla ilişkili kavramlara dair algılarının gözlemsel deneyimler, görsel unsurlar ve kendi gelişim düzeyleri çerçevesinde şekillendiğini göstermektedir

Araştırma kapsamında gelecek nesillerin mimarı olacak çocukların sağlık hizmeti unsurlarına ilişkin algıları değerlendirilmiştir. Bu kapsam da çocukları hastanelere, hekimlere ve hasatlık ile hasta bireylere yönelik algıları araştırılmıştır. Çocukların sağlık hizmeti bileşenlerine yönelik algılarının bireysel deneyimler ve ebeveyn yönlendirmeleriyle oluştuğu tespit edilmiştir. Bu doğrultuda çocukların sağlık hizmeti kullanımı ve bu hizmetleri oluşturan unsurlara yönelik gelecek perspektiflerinin olumlu seyri için, öncelikle ebeveynlerin bu hizmet koluna ilişkin olumlu yönlendirmeleri önem arz etmektedir. Aynı zamanda çocukların hizmet aldıkları kurumlarda yaşayacakları olumsuz deneyimlerin, gelecekte ilgili kurumlara ve hizmet sunuculara karşı oluşacak ön yargıya sebep teşkil edebileceği söylenebilir. Dolayısıyla hem ebeveynlerin hem de hizmet sunumunu koordine eden birimlerin çocukların sağlık hizmeti deneyimlerini en kolay ve en az duygusal hasarla atlatmaları için çaba sarf etmeleri gerektiği vurgulanmaktadır.

KAYNAKLAR

Coyne, I. (2006). Children's experiences of hospitalization. *Journal of Child Health Care, 10*(4), 326–336. https://doi.org/10.1177/1367493506067863

FULL TEXTS BOOK

Dalley, J. S., Rahman, A., & Bartram, T. (2019). Children's perceptions of hospitals and health professionals. *Child: Care, Health and Development, 45*(2), 175–185. https://doi.org/10.1111/cch.12618

Gürbüz, S., & Şahin, F. (2018). *Sosyal bilimlerde araştırma yöntemleri* (5.baskı). Seçkin yayıncılık.

McGrath, P. (2018). Hospitalized children and their families: Psychological impact of hospitalization. *Pediatric Nursing, 44*(2), 63–71.

Pelander, T., & Leino-Kilpi, H. (2010). Children's best and worst experiences during hospitalisation. *Scandinavian Journal of Caring Sciences, 24*(4), 726–733. https://doi.org/10.1111/j.1471-6712.2010.00770.x

Piaget, J. (1952). *The origins of intelligence in children*. International Universities Press.

Vygotsky, L. S. (1978). *Mind in society: The development of higher psychological processes*. Harvard University Press.

World Health Organization. (2012). *Standards for child-friendly health care*. WHO Regional Office for Europe.

FULL TEXTS BOOK

OKUL ÖNCESİNDE SOSYAL BECERİLER VE AKRAN İLİŞKİLERİ: BİR BİBLİYOMETRİK ANALİZ

SOCIAL SKILLS AND PEER RELATIONSHIPS IN PRESCHOOL: A BIBLIOMETRIC ANALYSIS

Şevval Erva KAYA

Afyon Kocatepe Üniversitesi, Sosyal Bilimler Enstitüsü,

Temel Eğitim Anabilim Dalı/ Okul Öncesi Programı, Afyon, Türkiye.

Mine KOYUNCU ŞAHİN

Dr. Öğr. Üyesi, Afyon Kocatepe Üniversitesi, Eğitim Fakültesi,

Okul Öncesi Eğitimi Bölümü, Afyon, Türkiye.

ÖZET

Okul öncesi dönemde sosyal beceriler ve akran ilişkileri çocuğun şimdiki ve daha sonraki yaşamında, önemli rol oynamaktadır. Küresel ölçekte tüm bireyler, sözlü ya da sözsüz iletişim biçimlerinden bağımsız olarak, başkalarıyla etkileşim kurma; yaşam sürecinde karşılaşılan çeşitli zorluklar karşısında sosyal çevrelerinden destek alma ve bu doğrultuda psikososyal iyi oluşlarını sürdürebilme amacıyla diğer bireylere güven duyma gereksinimi içerisindedir. Bu araştırmada, okul öncesi dönemdeki çocukların sosyal becerileri ve akran ilişkileri konusunda yapılan çalışmaların bibliyometrik göstergeler yönünden incelenmesi amaçlanmıştır. Araştırmada bibliyometrik analiz tekniği kullanılmıştır. Araştırmaya dahil edilme kriterleri olarak; 2025 yılına kadar hazırlanmış makaleler kullanılmıştır. Veriler, 07.07.2025 tarihinde, Web of Science veri tabanında yer alan yayınlar içerisinde, konu başlığı düzeyinde "Peer Relations" or "Friendship Relations" or "Peer" or "Peer Relationship" or "Friendship" AND "Social Skills" or “Social Development” AND “Preschool Education” or “Early Childhood” anahtar kelimeleri kullanılarak elde edilmiştir. Araştırma 1983-2025 yılları arasında 44 farklı dergide, 49 farklı ülkeden, 130 yazar tarafından yayınlanan 334 makaleyi kapsamaktadır. Yayınlanan makalelerde en çok kullanılan anahtar kelimelerin "social skills", “peer relations”, “social competence” olduğu görülmektedir. 2021 yılında en çok makalenin yayınlandığı, Journal Of Chıld Psychology And Psychıatry ve Clınıcal Psychology Revıew dergilerinin en çok yayın yapan 2 dergi olduğu görülmektedir. Aynı yazar tarafından en çok 2 çalışma gerçekleştirilmiş, 2 yayın yapan 5 yazar saptanmıştır. Dale F. Hay ve Andrew Chadwick bu yazarlardan bazılarına örnektir. En çok yayın yapılan ülkelerin ABD, İngiltere ve Türkiye olduğu tespit edilmiştir. Okul öncesi dönemde sosyal becerilerin geliştirilebilmesi akran ilişkilerinin iyi yöne evrilebilmesi ve günümüz sorunlarından “akran zorbalığı” konusunun önüne geçilebilmesi yönünden büyük önem arz etmektedir.

Anahtar Kelimeler: Okul Öncesi, Sosyal Beceriler, Akran İlişkileri, Bibliyometrik Analiz.

ABSTRACT

Social skills and peer relationships in the preschool period play a crucial role in a child's current and later life. Globally, all individuals, regardless of verbal or nonverbal communication styles, feel the need to interact with others; to receive support from their social environment in the face of various challenges encountered throughout life; and to trust others to maintain their psychosocial well-being. This study aimed to examine studies on the social skills and peer relationships of preschool children in terms of bibliometric indicators. Bibliometric analysis was employed in the study. Inclusion criteria for the study included articles published up to 2025. Data were collected from publications in the Web of Science

FULL TEXTS BOOK

database on July 7, 2025, using the keywords "Peer Relationships" or "Friendship Relationships" or "Peer" or "Peer Relationship" or "Friendship" AND "Social Skills" or "Social Development" AND "Preschool Education" or "Early Childhood" at the subject level. The research covers 334 articles published by 130 authors from 49 different countries in 44 different journals between 1983 and 2025. The most frequently used keywords in the published articles were "social skills," "peer relations," and "social competence." The Journal of Child Psychology and Psychiatry and Clinical Psychology Review, which had the most articles published in 2021, were the two journals with the most publications. The most two studies were conducted by the same author, and five authors published two publications. Dale F. Hay and Andrew Chadwick are examples of some of these authors. The countries with the most publications were identified as the US, the UK, and Turkey. Developing social skills in the preschool period is crucial for improving peer relationships and preventing peer bullying, a contemporary problem.

Keywords: Preschool, Social Skills, Peer Relations, Bibliometric Analysis.

GİRİŞ

Okul öncesi dönem, bireyin yaşamında sosyal gelişimin temellerinin atıldığı ve sosyal becerilerin en yoğun biçimde geliştiği önemli bir evredir. Sosyal beceriler, bireyin sosyal çevresiyle olan sağlıklı iletişim ve etkileşimini sağlayan davranışlar bütününü ifade eder. Özellikle paylaşma, sırayla oynama, iş birliği yapma, duygu tanıma ve empati gibi beceriler, çocukların hem aile hem de akran ortamında olumlu ilişkiler kurmalarına ortam oluşturmaktadır (Çiftçi ve Akgün, 2010).

Akran ilişkileri ise, çocukların sosyal becerilerini uygulama ve geliştirme fırsatı bulduğu en önemli sosyal bağlamlardan biridir. Çocuklar, akranlarıyla etkileşim kurarak sosyal normları öğrenmekte ve benlik algılarını şekillendirmektedir (Rubin ve ark., 2011). Nitelikli akran ilişkileri, sadece sosyal gelişimi değil, diğer gelişim alanları üzerinde de uzun vadeli etkiler göstermektedir (Ladd, 1990). Öte yandan, sosyal beceri alanındaki yetersizlikler akran reddine, sosyal dışlanmaya ve ileriki yıllarda uyum sorunlarına neden olabilmektedir (Denham ve ark., 2003).

Son yıllarda bu alana yönelik bilimsel ilgi giderek artmakta, sosyal beceriler ve akran ilişkileri arasındaki etkileşim çok boyutlu bir şekilde ele alınmaktadır (Gülay ve Erten Sarıkaya, 2014; Öner ve Özbey, 2022; Çakır ve Güntekin Akduman, 2023; Tutkun ve Eskidemir Meral, 2025). Ancak bu çalışmaların genel eğilimlerinin, hangi konulara odaklandıklarının, hangi ülkelerde ve yıllarda yoğunlaştıklarının sistematik olarak ortaya konması, literatürdeki mevcut durumu değerlendirme açısından önemlidir. Bibliyometrik analiz bu noktada devreye girerek, ilgili alandaki yayınların sayısal ve görsel haritalandırmasını sunmakta, araştırma alanlarının zaman içindeki gelişimini ve bilimsel iş birliklerini görünür kılmaktadır (Donthu ve ark., 2021).

Bu bağlamda hazırlanan bu çalışma, okul öncesi dönemde sosyal beceriler ve akran ilişkileri temalı yayınları bibliyometrik yöntemle incelemeyi amaçlamaktadır. Böylece araştırma alanının gelişim seyri, öne çıkan temalar, etkili yayınlar ve yazarlar belirlenerek, hem mevcut literatüre katkı sağlanması hem de gelecekte yapılacak araştırmalara yön verilmesi hedeflenmektedir.

YÖNTEM

Bu araştırma, okul öncesi dönemde sosyal beceriler ve akran ilişkileri konularında yapılan bilimsel çalışmaların, araştırma yoğunluklarını ve bilimsel iş birliklerini ortaya koymak amacıyla yürütülen bir bibliyometrik analiz çalışmasıdır. Literatürde herhangi bir konu başlığında hazırlanmış yayınların sayısal verilerle incelenmesine dayanan bibliyometrik analiz yöntemi, yayınların yapısal özelliklerini sistematik olarak değerlendirmeyi amaçlamaktadır (Broadus, 1987).

FULL TEXTS BOOK

1.1. Araştırmaya Dâhil Edilme Kriterleri,


Araştırma kapsamında, 07.07.2025 tarihinde, Web of Science veri tabanında; konu düzeyinde "Peer Relations" or "Friendship Relations" or "Peer" or "Peer Relationship" or "Friendship" AND "Social Skills" or “Social Development” AND “Preschool Education” or “Early Childhood” anahtar kelimleri aratılmıştır. Bu arama sonucunda 334 yayına ulaşılmış, bibliyometrik analiz gerçekleştirilmiştir.

Verilerin Analizi

Web of Science veri tabanından elde edilen yayın yılı, ülke, dergi, yazar ve atıf sayısına ilişkin veriler, VOSviewer yazılımı aracılığıyla analiz edilmiş ve görselleştirilmiştir.


BULGULAR

Araştırmada yer alan makalelerin yıllara göre dağılımı Şekil 1’de ifade edilmiştir.



Şekil 1. Yıllara Göre Makale Sayısı


Yıl bazında yapılan yayınların dağılımı incelendiğinde, okul öncesi dönemde akran ilişkileri ve sosyal beceriler üzerine yapılan çalışmaların özellikle son 10 yılda artış gösterdiği görülmektedir. Yayın sayısının en yüksek olduğu yıl 2021 olarak tespit edilmiştir. Bu artışa, sosyal-duygusal gelişimin erken çocukluk dönemindeki önemine yönelik farkındalığın artmasının ve pandemi sürecinde çocukların sosyal etkileşimlerinin kısıtlanmasının neden olabileceği yorumu yapılabilmektedir.



Şekil 2. Ülkelere Göre Makale Sayısı (makale sayısı≥5)

FULL TEXTS BOOK

Şekil 2' ye bakıldığında, 50 farklı ülkeden yayın olduğu ve en fazla yayının USA (193), Ingiltere (20) ve Türkiye'de (19) yapıldığı görülmektedir. 2010 yılından sonra bu üç ülkede, eğitim politikaları ve stratejik planlamalar erken çocukluk döneminde sosyal-duygusal gelişimi öncelikli hedefler arasında konumlandırmaktadır. Elde edilen sonuçlar doğrultusunda bilimsel yayınlarında ülke politikasıyla birlikte gelişim gösterdiği düşünülmektedir. Sosyal beceriler ve akran ilişkileri konularında yayın yapan ülkeler arasındaki atıf bağlantıları, şekil 3.'te görselleştirilmiştir.



Şekil 3. Ülkeler Göre Atıf İlişkileri

Tablo 1. Dergilere Göre Makale ve Atıf Sayıları


Sıra No	Dergi adı	Makale Sayısı	Atıf Sayısı
1	Early Chıld Development And Care	3	48
2	Clınıcal Psychology Revıew	2	284
3	Early Chıldhood Research Quarterly	2	184
4	Amerıcan Journal On Mental Retardatıon	2	162
5	Energy Educatıon Scıence And Technology Part B-Socıal And Educatıonal	2	19
6	Journal Of Chıld Psychology And Psychıatry	1	383
7	Chıld Development	1	140
8	Chıld Language Teachıng & Therapy	1	122
9	Journal Of Abnormal Chıld Psychology	1	108
10	Behavıoral Sleep Medıcıne	1	90

FULL TEXTS BOOK

11	Topıcs In Language Dısorders	1	33
12	Behavıor Modıfıcatıon	1	32
13	Infant And Chıld Development	1	32
14	Chıld Psychıatry & Human Development	1	28
15	European Early Chıldhood Educatıon Research Journal	1	24
16	Journal Of Vısual Impaırment & Blındness	1	19
17	Socıal Development	1	19
18	Ajıdd-Amerıcan Journal On Intellectual And Developmental Dısabılıtıes	1	18
19	Mental Retardatıon	1	18
20	Journal Of Research In Personalıty	1	12

Okul öncesinde akran ilişkileri ve sosyal beceriler konularına yönelik araştırmaların yayınlandığı dergiler ve aldıkları atıf sayıları incelendiğinde; "Early Chıld Development And Care", "Clınıcal Psychology Revıew" ve "Early Chıldhood Research Quarterly" dergilerinin öne çıktığı görülmüştür (Tablo 1).

Okul öncesinde akran ilişkileri ve sosyal beceriler konuları üzerine en çok yayın yapan dergiler ve aralarındaki atıf ilişkileri şekil 4.'te görselleştirilmiştir.



Şekil 4. Dergilere Göre Atıf İlişkileri


Okul öncesinde akran ilişkileri ve sosyal beceriler konuları üzerine en çok atıf alan yazarlar tablo 2.'te yer almaktadır.

Tablo 2. En Çok Makale Yayınlayan Yazarlar ve Atıf Sayıları (Makale Sayısı ≥ 1)

Sıra No	Yazarlar (Adı ve Soyadı)	Makale Sayısı	Atıf Sayısı
1	Zarife Seçer	2	10
2	Hülya Gülay Ögelman	2	19
3	Nana Shin	2	109
4	Maria Gladh	2	5
5	Brian E. Vaughn	1	109
6	Andrew Chadwick	1	383
7	Mara Allodi Westling	1	5
8	Sakire Anliak	1	3
9	Steven R. Asher	1	33
10	Pier J M Prins	1	195

Okul öncesinde akran ilişkileri ve sosyal beceriler konuları üzerine çalışan yazarlar içerisinde; Zarife Seçer (2) ve Hülya Gülay Ögelman (2) en çok yayın yapan yazarlardır. Ayrıca Andrew Chadwick (383) ve Pier J M Prins (195) atıfla en çok atıf alan yazarlardır.

Okul öncesinde akran ilişkileri ve sosyal beceriler konuları üzerine yapılan çalışmalarda en çok kullanılan anahtar kelimelere şekil 5'de yer verilmiştir.



Şekil 5. Makalelerde En Çok Kullanılan Anahtar Kelimeler

FULL TEXTS BOOK

Çalışmaların konu sınıflandırmasının yapılmasını sağlayan anahtar kelimelere bakıldığında, en yoğun kullanılan anahtar kelimelerin, "social skills", "peer relations", "preschool education", "friendship" ve "social competence" olduğu tespit edilmiştir.

TARTIŞMA VE SONUÇ

Bu araştırma ile okul öncesinde akran ilişkileri ve sosyal beceriler konularına odaklanan bilimsel çalışmaların Web of Science Veri tabanına dayalı bibliyometrik analizinin yapılmasına odaklanılmıştır. Çalışma sonucunda 1983-2025 yılları arasında 334 makalenin olduğu görülmektedir. Bulgular, özellikle son on yılda bu alandaki çalışmaların belirgin şekilde arttığını göstermektedir. Bu artışın, erken çocukluk döneminin sosyal-duygusal gelişimde kritik bir evre olduğuna dair farkındalığın yükselmesiyle ilişkili olduğu söylenebilir. Nitekim pandemi sürecinde çocukların sosyal etkileşimlerinin sınırlanması, bu konuda yeni araştırma ihtiyaçlarını gündeme getirmiştir. İlgili makaleler 50 farklı ülkeden 131 yazar tarafından hazırlanmıştır. En çok yayın yapan ülkelerin Amerika, İngiltere ve Türkiye en çok yayın yapan yazarların "Zarife Seçer", "Hülya Gülay Ögelman", "Nana Shin" ve "Maria Gladh" en çok atıf alan yazar ise "Andrew Chadwick" olduğu tespit edilmiştir. Bu durum, söz konusu ülkelerde erken çocukluk eğitiminin politik düzeyde öncelikli alanlardan biri olarak kabul edilmesiyle ilişkilendirilebilir. Türkiye özelinde son yıllarda okul öncesi eğitimin zorunlu hale getirilmesine yönelik girişimlerin, sosyal-duygusal gelişim konulu araştırmaları desteklediği değerlendirilebilir.

Makalelerde en çok kullanılan anahtar kelimelerin "social skills", "peer relations", "preschool education", "friendship" ve "social competence" olduğu görülmektedir. Bu bulgu, okul öncesi dönemde sosyal becerilerin ve akran ilişkilerinin çocukların arkadaşlık ilişkileri ve sosyal yeterlilikleriyle doğrudan bağlantılı olduğuna işaret etmektedir. Dolayısıyla araştırmalar, yalnızca bireysel sosyal becerilere değil, aynı zamanda çocukların akran gruplarıyla olan etkileşimlerine de yoğunlaşmaktadır.

Araştırmanın bulguları, okul öncesi dönemde sosyal beceriler ve akran ilişkileri konularına yönelik çalışmaların giderek artan bir ivme kazandığını ortaya koymaktadır. Özellikle son on yılda yapılan çalışmalar, bu alanın hem akademik hem de toplumsal düzeyde önemsendiğini göstermektedir. ABD, İngiltere ve Türkiye'nin öne çıkan ülkeler olması, ulusal eğitim politikalarının bilimsel araştırmalara yön verici bir rol oynadığını kanıtlamaktadır.

Bu çalışma, literatürde sosyal-duygusal gelişim, akran ilişkileri ve sosyal yeterlilik gibi kavramların birlikte ele alınarak incelendiğini göstermektedir. Çocukların erken dönemde edindikleri sosyal becerilerin, yaşam boyu psikososyal uyumlarını etkilediği dikkate alındığında, bu alanda yapılacak araştırmaların eğitimciler, aileler ve politika yapıcılar açısından büyük önem taşıdığı söylenebilir.

Gelecek araştırmaların, yalnızca nicel verilerle sınırlı kalmayıp, çocukların sosyal etkileşimlerini doğal ortamlarda gözlemleyen nitel çalışmalara yönelmesi önerilmektedir. Ayrıca kültürel farklılıkların akran ilişkileri ve sosyal beceri gelişimi üzerindeki etkilerini ortaya koyan karşılaştırmalı çalışmaların yapılması, alan yazına önemli katkılar sağlayacaktır. Özellikle küçük yaşa hitap eden eğitimcilere, akran ilişkileri ve sosyal beceriler konularındaki çalışmaları takip etmeleri, ailelere, akran ilişkileri ve sosyal beceriler konulu eğitim seminerlerine katılmaları önerilmektedir.

KAYNAKLAR

Broadus, R. N. (1987). Toward a definition of "bibliometrics". *Scientometrics, 12*(5-6), 373–379.

Çakır, S., & Güntekin Akduman, G. (2023). Annelerin bilinçli farkındalıkları ile okul öncesi dönem çocuklarının kaygı düzeyleri ve arkadaşlık becerileri arasındaki ilişki. *Gazi Eğitim Bilimleri Dergisi, 9*(1), 73-93. https://dergipark.org.tr/tr/pub/eyyad/issue/88479/1498459

FULL TEXTS BOOK

Çiftçi, S., & Akgün, E. (2010). Sosyal beceri eğitiminin çocukların sosyal becerileri üzerindeki etkisi. *Kuram ve Uygulamada Eğitim Bilimleri*, 10(2), 691-720.

Denham, S. A., Blair, K. A., DeMulder, E., Levitas, J., Sawyer, K., Auerbach–Major, S., & Queenan, P. (2003). Preschool emotional competence: Pathway to social competence? *Child Development*, 74(1), 238–256.

Donthu, N., Kumar, S., Mukherjee, D., Pandey, N., & Lim, W. M. (2021). How to conduct a bibliometric analysis: An overview and guidelines. *Journal of Business Research*, 133, 285–296.

Gülay, H., & Erten Sarıkaya, H. (2014). 5-6 yaş çocukların sosyal beceri, akran ilişkileri ve okula uyum düzeyleri ile kardeş ilişkileri arasındaki ilişki. *Pamukkale Üniversitesi Eğitim Fakültesi Dergisi, 35*, 1-14. https://dergipark.org.tr/tr/download/article-file/383825

Ladd, G. W. (1990). Having friends, keeping friends, making friends, and being liked by peers in the classroom: Predictors of children's early school adjustment? *Child Development*, 61(4), 1081–1100.

Öner, Ş., & Özbey, S. (2022). Okul öncesi dönemdeki çocukların sosyal bilgiyi işleme becerileri ile psikolojik sağlamlık düzeyleri arasındaki ilişkinin incelenmesi. *Journal of Social Sciences, 11*(1), 1-18. https://journalofsocial.com/files/josasjournal/3849ebe3-4fbb-422e-a929-a2139f05398f.pdf

Rubin, K. H., Bukowski, W. M., & Laursen, B. (2011). *Handbook of peer interactions, relationships, and groups*. Guilford Press.

Tutkun, C., & Eskidemir Meral, S. (2025). Okul öncesi çocukların sosyal becerileri, problem davranışlar, akademik özsaygı ve öğretmen-çocuk ilişkisi: Seri aracılık modeli. *Frontiers in Psychology*. https://www.frontiersin.org/journals/psychology/articles/10.3389/fpsyg.2025.1453193/full

FULL TEXTS BOOK

OSMANLI DEVLETİNDE DÜYUN-I UMUMİYYE İDARESİ

Yunus BOR[1]

ÖZET

Osmanlı Devleti'nin son yetmiş ve Türkiye Cumhuriyeti'nin ilk otuz yılının en önemli problemlerinden biri Osmanlı borçları olmuştur. 19. Yüzyılının ortasında giderek merkezileşen devletin gelirleri giderlerini karşılayamaz olmuş ve dışarıdan borç alma ihtiyacı doğmuştur. Devletin borçlanma tarihine göz attığımızda neredeyse bütün kaynaklar Osmanlı devletinin dış borçlanma tarihini 1854 olarak başlatırlar. Ancak Yalçın Doğan *İMF Kıskacında Türkiye* adlı eserinde ilk borçlanmanın 1840 yılında yapıldığını böyle bir belgenin bulunduğunu söylemektedir. 1840 yılında Osmanlılar hazine tahvili çıkararak Londra ki bir bankaya satmışlar ve karşılığında 3 milyon sterlin borç alınmıştır daha sonra 1851 yılında 200.000 sterlinlik bir avans daha alınmış ve ilk borçlanma gerçekleşmiştir.

Bu ilk borçlanmadan sonra 1854 yılından 1874 yılına kadar 20 yılda 15 defa dış borçlanma yapmış olan Osmanlı devletinin bunlardan eline 127 milyon Osmanlı altın lirası geçtiği halde 239 milyon lira kadar borçlanmıştı. 1874-75 yılının bütçe geliri 25.104.928 lira iken borçların o yıllık bölümünün toplamı 30.000.000 civarındaydı bu da devletin o yılın bütçesinin borçlarını bile ödeyemediğini bize gösterir.

Düyun-ı Umumiyye'nin kuruluşuna kadar süren bu ilk devrede Osmanlı devletine borçlandığı paranın ancak yarısı ulaşabilmiştir. 1875 bütçesinde gelirin sadece borçlara ayrılmasına karşılık yine de beş milyon liralık açık kalmış bir nevi devlet iflasa sürüklenmiştir. Sonuç olarak Osmanlı devletinin son yetmiş yılını Türkiye Cumhuriyeti'nin ise ilk otuz yılını meşgul eden bu borçların son taksiti ilk alındığı tarihten tam bir asır sonra 1954 yılında ödenmiş ekonomik bağımsızlık yolunda önemli bir adım atılmıştır.

Anahtar Kelimeler, Düyun-ı Umumiyye, Osmanlı Devleti, Borçlar idaresi,

ABSTRACT

One of the most pressing issues during the last seventy years of the Ottoman Empire and the first thirty years of the Republic of Turkey was the matter of Ottoman public debt. By the mid-19th century, the increasingly centralized Ottoman state found itself unable to meet its expenditures through domestic revenues alone, thus necessitating borrowing from foreign sources. When examining the history of Ottoman indebtedness, nearly all sources mark 1854 as the beginning of the Empire's foreign borrowing. However, Yalçın Doğan, in his work *İMF Kıskacında Türkiye*, argues that the first instance of borrowing actually occurred in 1840, and he refers to documentation supporting this claim. According to his account, in 1840 the Ottoman government issued treasury bonds and sold them to a bank in London in exchange for a £3 million loan. A further advance of £200,000 was secured in 1851, marking the initial steps of foreign borrowing.

Following this, between 1854 and 1874, the Ottoman Empire took on foreign loans fifteen times within a span of twenty years. Although the Empire managed to obtain 127 million Ottoman gold liras from these loans, it incurred a total debt of approximately 239 million liras. In the 1874–75 fiscal year, while the total revenue amounted to 25,104,928 liras, the annual debt repayments reached around 30 million liras. This discrepancy clearly indicates that the state was unable to even cover its debt obligations from that year's budget.

[1] Ar.Gör. Dr. Ordu Üniversitesi Fen- Edebiyat Fakültesi Tarih, Ordu, Türkiye
https://orcid.org/0000-0002-4391-8052

FULL TEXTS BOOK

During this initial period leading up to the establishment of the *Düyun-ı Umumiye* (Public Debt Administration), the Ottoman Empire received only about half of the total amount it had borrowed. Despite allocating its entire revenue to debt repayments in the 1875 budget, a deficit of five million liras remained, effectively pushing the state toward insolvency.

In conclusion, the burden of these debts (first acquired during the late Ottoman period)persisted throughout the early decades of the Turkish Republic. The final installment of these debts was paid in 1954, exactly one century after the initial borrowing, representing a significant milestone on the path toward economic independence.

Keywords: *Düyun-ı Umumiye*, Ottoman Empire, Debt Administration,

Giriş

Osmanlı Devleti Kırım Savaşı ile almaya başladığı dış borçlar 1875 yılına kadar artarak devam edince bu borçların kapatılmasına dair çareler aranmaya başlamıştır. Osmanlı devletinde Rus Nedim diye adlandırılan Mahmut Nedim Paşa sadrazamlığa getirildi. Nedim Paşa borcun tekrar borçlanılarak ödenmesinin borcu giderek daha da artıracağına bütçenin dengeye kavuşturulması ve yeniden güven sağlanılması için **6 Ekim 1875 Ramazan Kararnamesi** yayınlandı. Buna göre;

Yıllık faiz ve anapara taksitleri toplamı yaklaşık 14 milyon lira tutan Osmanlı devletinin o tarihteki borçları hakkında şu işlemler yapılacaktı.

- 6 Ekim 1875 gününden başlayarak Osmanlı devletinin iç ve dış borçlarının faiz ve ana para taksitleri toplamı 5 yıl için yarıya indirilmiştir. Diğer yarısı silinmiştir.
- Bu tarihten sonra ana paranın indirim sonrası ödenecek kısmının yani yarısının birinci yarısı tamamen nakit olarak, diğer yarısı %5 faizli borç senetleri verilerek ödenecektir.
- Yapılacak ödemenin teminatı olarak tuz tütün varidatı Mısır vergisi ve yeterli olmadığı taktirde ağnam resmi gelirleri gösterilmiştir (Özdemir, 2009).

Nedim paşanın tedbirleri işe yaramadı. Bu gelişmelere en başta İngiltere ve Fransa olmak üzere, Avrupa mali çevreleri büyük tepki gösterdiler itirazlarda bulundular. Avrupa gazeteleri Türkler bizi dolandırdı, altınlarımızı safahat uğruna harcadılar diyerek Osmanlı devleti hakkında yoğun kampanya başlattılar.

1858 yılına ait borç kuponları ödenemedi. 1869 ve 1873 kupon bedellerinin ödenmesi ertelenince 6 ay kadar sonra Nisan 1876 da tüm borçların ödenmesi tamamen durduruldu. **Moratoryum** (ödemelerin tümüyle askıya alınması) ilanına mecbur kalındı. Devletin mali iflası resmen kabul edilmiş oldu. 6 Ekim 1875 -20 Aralık 1881 dönemine iki kararname arası devre adı verilmektedir (Özdemir, 2009).

Moratoryum ilanına ülke içinde Galata bankerleri ve Osmanlı bankası durumdan haberdar olmaması nedeniyle itiraz etmişler ayrıca küçük burjuvazi ve genç Osmanlılardan da itirazlar gelmiştir. Bütün bu gelişmeler Mahmut Nedim Paşa'nın azline kadar uzanmıştır.

Rüsum-u Sitte İdaresi (Altı Vergi İdaresi) (1879-80)

1879 yılında Osmanlı devleti ile alacaklılar arasında borçların ödenmesine yönelik ortak bir komisyon kurulması fikri başarısızlıkla sonuçlanmıştı. Buna rağmen Galata bankerleri temsilcileri ve Osmanlı bankası ile yapılan müzakereler olumlu sonuç vermiş bunların alacakları sağlam esaslara bağlanmak üzere gümrük vergileri hariç 6 vergi gelirinin; Pul resmi, alkollü içkiler (ispirto) vergisi, tütün tekeli gelirleri, İstanbul, Bursa, Edirne ve Samsun ipek kozası öşür gelirleri, İstanbul ve çevresi balık avı vergisi ve tuz tekeli resim gelirlerinin 10 yıl süreyle önceki yıllardan 8.725.000 lira alacaklı oldukları kabul edilen Galata bankerlerine bırakılması uygun görülmüştür (Blaisdel,1979). Bu iş Osmanlı bankası önderliğinde kurulan ve başına Fransız Hamilton Long'un getirildiği bir özel yönetime

FULL TEXTS BOOK

bırakılmıştır. 13 Ocak 1880 tarihinde faaliyete geçen bu idare ile sadece Galata bankerlerine olan dış borçlar kesin bir şekle bağlanmıştır.

Düyun-ı Umumiye-i Osmaniye Meclis-i İdaresi

Rüsum-u sitte idaresine Fransızlar İngilizler ve borç tahvil sahipleri itiraz etmiştir. Osmanlı borçlarının 252 milyon altın liraya çıkmış olması ve uzun vadede bile ödeme güçlüğü getirecek büyük rakamlara ulaştı. Buna ek olarak alacaklılar işi siyasi bir müdahale şekline getirince Osmanlı hükümeti 3 Ekim 1880 yayınladığı bir nota ile alacaklılara seçecekleri birer temsilciyi hükümetle doğrudan görüşmesi için İstanbul'a göndermek çağrısında bulundu.

1 Eylül 1881 tarihinde ilki yapılan ve 20 Aralık 1881 tarihinde sona eren toplantılar sonucu mutabakat sağlanarak, **Rüsum-u sitte** idaresi yerine kupon sahiplerinin seçeceği temsilcilerin de katılacağı Düyun-ı Umimiye-i Osmaniye meclisi-i idaresi kurulması kabul edildi. 20 Aralık 1881 (28 Muharrem 1299) tarihli **Muharrem Kararnamesi** olarak bilinen bir kararname ile sözleşmeye kesin şeklini verildi.

1881 yılındaki toplam borç tutarı **252.131.196** lira iken, Muharrem Kararnamesi kapsamına alınarak indirim konusu yapılan anapara ve faiz toplamı **233.993.066** lira kabul edilmiş, indirimlerden sonra hükümetin mutabık kalınan ödenecek ana borç tutarı **141.505.919** lira olarak belirlenmiştir (Özdemir, 2009).

İdarenin Yapısı ve Yönetimi

Merkezi İstanbul da bulunan Düyun-u Umumiye İdaresi yabancı alacaklıları temsil eden 5 temsilci ile yerli alacaklıları ve Galata Bankerlerini temsil eden birer üye olmak üzere toplam 7 üyeden oluşan bir İdare meclisi tarafından idare olunmuştur. Ülke temsilcileri İngiliz veya Hollandalı, Fransız, Alman, Avusturyalı, İtalyan, Galata bankerleri temsilcisi ve Osmanlı devleti temsilcisiydi. Üyelik süresi 5 yıl idi. Başkanlık ise 5 yıl süreyle İngiliz ve Fransız temsilcilerine ait bulunuyordu. Osmanlı temsilcileri azınlıkta kaldığı için karar verme süreçlerinde çok etkili olamamışlardır. (Pamuk, 1987).

İdarenin icrai görevleri Genel Müdürlük, şubeler, kontrol kurulu, vezne, levazım ve basım müdürlükleri, teftiş kurulu, ziraat müfettişleri ve memurlar komisyonu tarafından yürütülüyordu. İdarenin görevi ise yetki alanına giren gelirleri kendi teşkilatı, Reji idaresi ve Tömbeki Şirketi eli ile toplayıp yönetim giderlerini ve tahsilat masraflarını indirdikten sonra devlet hissesini de ayırıp kalan miktarı kararname kapsamındaki borçların yıllık ana para ve taksitlerine ayırmaktaydı. İdare her yıl sonunda yaptığı ödemeleri masrafları ve kesin hesabı Maliye bakanlığına sunmakla yükümlüdür. İdarenin memurları devlet memuru gibi sayılacak, idare hükümetin denetimine tabi olacaktır, devlet idareye teftiş inceleme ve soruşturma yaptırabilecektir (Özdemir, 2009, Yılmaz, 2009).

İdarenin yönetimine Rüsum-u sitte idaresine bırakılan gelirlere ek olarak gümrük resminin %8 den yükseltilmesi olursa elde edilecek gelir fazlasından, Temettü (kazanç vergisi) yükseltilmesinden sağlanacak gelir fazlasından, Doğu Rumeli eyaleti gelirlerinden, Kıbrıs adası yıllık gelirinin fazlası, Bulgaristan emaretinin yıllık gelirinden, yapılacak düzenlemeler sonucu devlet gelirinde sağlanacak ek artışlardan oluşan gelirlerde bırakılmıştır.

Düyun-u umumiye idaresinin yıllık gelirleri 1882-83 mali yılında Devlet gelirlerinin %17 sine tekabül ederken giderek artmış, 1909 yılında gelirlerin % 25.2 oluşturmuş 1911-12 yılında %27 sine karşılık gelmiştir.

İdarenin olumlu ve olumsuz yönleri

Devlet dairesi gibi faaliyet göstermiş olan idare nin olumlu denebilecek yönleri de olmuştur. Borç anapara ve faizlerinde önemli miktarda indirim yapılmıştır. 252.1 milyon lira olan borcun, 110.6 lirası indirilerek 141.5 lira da mutabakat sağlanmıştır. Düzenli borç ödemeleriyle hükümetin itibarına önemli

FULL TEXTS BOOK

katkıda bulunulmuştur. Düşük oranda faizler alınmıştır %5-6 dan %3-4 e düşmüştür. Ayrıca maliye nezaretinin yeniden yapılanmasına ve mali bürokrasinin disipline edilmesini sağlamıştır (Sağlam, 2007).

İdarenin olumsuz yönlerini sıralayacak olursak; devlet maliyesi içinde ayırıcı ve özel bir yönetimin doğması kabul edilmiş bir bakıma devlet içinde devlet yapılanması suretiyle adeta devlet **yarı-sömürge** haline getirilmiştir.

Avrupalı bankalarla işlem yapılması zorunluluğu getirilmesi ve gümrük tarifeleri üzerinde düzenleme yetkisinin sınırlandırılması (kapitülasyonlar) sonucu, Osmanlı devletinin kendi kaynaklarını serbestçe tahsis edebilmesi açık biçimde kısıtlanmıştır.

Düyun-u Umumiye yönetim kurulu üyelerinin aynı zamanda yabancı demiryolu şirketlerinde de yönetim kurulu üyesi olmaları, idarenin çapraz bir ilişkiler ağı içerisinde olmasına aynı zamanda emperyalist çıkarların işlevini artırmasına yol açmıştır.

Demiryolları, limanlar, sigorta şirketleri, maden işletmeleri ve telgraf posta elektrik servislerinde çalışan memurlar vasıtasıyla yabancı ülkeler için her türlü bilgi ve istihbarata ulaşılabilmesi ulusal güvenliği tehdit etmiştir. Bu idare, batılı devletlerin verdikleri ağır koşullu borçların bir güvencesi kapitülasyonların da daha etkili ve garantili devamını sağlıyordu.

Borçların harcama alanları

Alınan borçların devletin pek yararına kullanılmadığı görülüyor. Örneğin Abdülaziz döneminde alınan borçların %50 si emisyon zararı % 14 ü demir yolu yapımı için kullanılmış % 12 si hazineye kalmış %10 u dış borçlar ve faizler için kullanılmış % 5 i saray giderleri olarak kullanılmıştır. Bu göstergeden de anlaşılıyor ki borçlar, yatırımlardan çok cari harcamalar için kullanılmıştır. Alınan borçların faizlerin günün piyasalarına göre çok yüksektir. Avrupalı sermayedarlar, kendi ülkelerindeki düşük gelir getiren yatırımlar yerine bol geliri olan Osmanlıya borç verme işini desteklemişler ve batı emperyalizmi Osmanlı imparatorluğunu kıskacı altına almıştır.

1914 yılında savaş patlak verdiğinde Osmanlı Devleti'nin dış borcu kısa vadeli borçlar hariç 156,4 milyon Osmanlı lirasıdır. Dış borçlar, Osmanlı Devletinden sonra, Osmanlı topraklarında kurulan devletler arasında paylaştırılmış ve en büyük borç yükü Türkiye'ye verilmiştir. 1925 yılında Osmanlı borçlarının %67'sinin Türkiye tarafından ödenmesi kararlaştırılmıştır. Türkiye'nin payına düşen 107,5 milyon altın Osmanlı lirası tutarındaki borcun ödenmesi için Düyun-u Umumiye İdaresi ile 13 Haziran 1928 tarihinde Paris'te bir anlaşma imzalanmıştır. Ayrıca bu sözleşme ile idare kaldırılmış onun yerine merkezi Türkiye dışında Fransa da olan iki meclis kurulmuştur. Türkiye Düyun-u Umumiye' ye olan borcunun son taksitini, ilk dış borcun alınmasından tam bir yüzyıl sonra, 25 Mayıs 1954' te ödemiştir (Biltekin, 2009).

Değerlendirme

1838 de İngiltere'ye ve sonraki iki yılda da birçok Avrupa devletine ticari ayrıcalıklar verilmiş, sonrasında Osmanlı devleti içindeki üretim kaynakları yavaş yavaş batılı müttefiklerin eline geçmeye başlamış, yerli tüccar ile yerli üretim, bu gelişmelere paralel olarak büyük bir sıkıntı ve çöküşe yönelmiştir. Osmanlı tüccar ve üreticisi her türlü gümrük vergisini öderken diğer yandan yabancı uyrukluların vergilerden muaf olacak şekilde kolaylık görmeleri yerli üretimin büyük çoğunlukla tarıma kaymasına neden olmuştu, buna karşılık Batılı sermaye çevreleri ve emperyalist devletlerin sürekli olarak Osmanlı devletine bazı reformlar dayatmasına, devletin gelir kaynaklarını dolayısıyla da iktisadi ve siyasal yapısını da denetlemesine yol açmıştır. Düyun-u Umumiye idaresinin görev ve yetkileri bu duruma örnek olarak gösterilebilir.

FULL TEXTS BOOK

Hemen hemen bütün borç sözleşmelerinde borç veren ülkeye mali ve ekonomik ayrıcalıklar tanınmıştır. Düyun idaresinin teşkili ile devlet gelirlerinin büyük bölümünden vazgeçildiği gibi, devletin egemenlik haklarının bir kısmından da feragat zorunda kalınmıştır. Bu idare memurlarını dilediği gibi atama özgürlüğüne sahipti. Nitekim düyun idaresi olarak 5000 kişilik bir kadro oluşturulmuş bu sayı 1912'de 9.000 kişi olmuştur. Bu haliyle adeta devlet içinde devlet olma halini taşıyordu. Devlet yalnızca eski borçlarının faizini ödemek için bile çoğu kez yeni bir borçlanmaya girmiş böylelikle hızla mali çöküntüye gidilmiştir.

Sonuç olarak Osmanlı devletinin son yetmiş yılını Türkiye Cumhuriyeti'nin ise ilk otuz yılını meşgul eden bu borçların son taksiti ilk alındığı tarihten tam bir asır sonra 1954 yılında ödenmiş ekonomik bağımsızlık yolunda önemli bir adım atılmıştır.

KAYNAKÇA

Blaisdell, D. C. (1979). *Osmanlı İmparatorluğunda Avrupa mali denetimi* (A. İ. Dalgıç, Çev.). İstanbul: İstanbul Matbaası. (Orijinal çalışma 1979)

Özdemir, B. (2009). *1854-1954 döneminde yüzyıl süren Boyunduruk*. Ankara: Ankara Ticaret Odası Yayınları.

Pamuk, Ş. (1987). *The Ottoman Empire and European capitalism, 1820-1913*. Cambridge: Cambridge University Press.

Sağlam, H. M. (2007). *Osmanlı borç yönetimi Düyun-u Umumiyye 1879-1891* (Cilt 1). İstanbul: Tarih Vakfı Yayınları.

Yılmaz, F. (2009). *Osmanlının borç batağı Duyun-ı Umumiye*. İstanbul: İz Yayıncılık.

FULL TEXTS BOOK

INVESTIGATION OF PERCEIVED HEALTH AND ITS ASSOCIATED FACTORS AMONG ADULTS

Master Student, Öznur ALAN

Selcuk University, Institute of Health Sciences, Department of Public Health Nursing, Konya, Türkiye

https://orcid.org/0009-0008-1478-1449

Assoc. Prof., Alime SELÇUK TOSUN

Selcuk University, Faculty of Nursing, Konya, Türkiye

https://orcid.org/0000-0002-4851-0910

ABSTRACT

Aim: This review study aims to examine perceived health and its associated factors among adult individuals.

Materials and Methods: A literature review was performed using the keywords “adult individuals,” “perceived health,” and “nursing” in both Turkish and English. Databases searched included Google Scholar, PubMed, Science Direct, Web of Science, and ResearchGate. Full-text articles published between 2020 and 2025 in English and Turkish were evaluated.

Results: Various studies conducted in different countries show that perceived health levels are influenced by multidimensional factors such as age, gender, socioeconomic status, presence of chronic disease, health behaviors, cultural values, and psychological factors. A study conducted in Spain found that self-perceptions of aging influenced cognitive function and physical activity levels, which were associated with perceived health. Research in the United States involving different ethnic groups revealed that the relationships between self-criticism and self-compassion varied according to culture and affected perceived health. Another study reported that perceived health was strongly associated not only with physical health but also with mental health and cognitive functions. In Türkiye, during the pandemic, it was found that perceived health levels affected health anxiety and were related to psychological well-being. These findings indicate that perceived health is influenced by both individual and environmental/psychosocial factors.

Conclusion: Literature reviews show that individuals’ perceptions of health are not only related to their physical conditions but also to psychological, socioeconomic, and cultural factors. Particularly, aging, stress, health anxiety, and self-attitudes directly affect perceived health. Socioeconomic inequalities, the presence of chronic diseases, and negative health behaviors adversely affect perceived health, whereas positive health behaviors, psychological resilience, and positive self-perceptions strengthen it. To enhance perceived health among adults, educational programs that are informative and supportive according to age and life conditions may be recommended. Future studies are advised to investigate the interaction of individual, cultural, and environmental factors affecting perceived health. The findings obtained serve as an important guide for planning person-centered health services.

Keywords: Adults, Perceived Health, Chronic Diseases

INTRODUCTION

The World Health Organization (WHO, 2024) defines health not merely as the absence of disease or infirmity but as a state of complete physical, mental, and social well-being. Physical health refers to an individual’s ability to adapt to environmental factors, their level of physical development, and their

FULL TEXTS BOOK

capacity to maintain bodily functions. Mental health is defined as a state in which an individual realizes their potential, copes with daily life stressors, and contributes to society. Social health refers to an individual's ability to maintain relationships and interact healthily with society (Kushmatova & Khakimova, 2022).

Protecting and maintaining health is of great importance in terms of ensuring quality of life and the meaningful continuity of life itself. Although health is a universal concept, individuals' perceptions of health may differ according to age, gender, culture, and personal experiences (Özdelikara et al., 2018). Perceived health is defined as the totality of an individual's feelings, thoughts, beliefs, and expectations about their own health. In this context, individuals with positive beliefs about their health are said to have "good perceived health," while those with negative beliefs are said to have "poor perceived health" (Ağaçdiken Alkan et al., 2017). Health is a dynamic process, defined by the individual in accordance with their personal values, through which they endeavor to maintain a state of well-being by adapting to their environment (Oral, 2018).

The evaluation of perceived health is an important indicator that allows the measurement of both objective and subjective aspects of health (Mikolajczyk, 2008). Research shows that individuals with high perceived health demonstrate healthier behaviors and report a higher quality of life (Yulu, 2020). Accordingly, there is not always a perfect alignment between subjective health perception and objective health status, as perception is shaped by many personal, social, and psychological factors. Indeed, an individual with multiple chronic diseases may still perceive themselves as healthy, while another without any diagnosis may define themselves as ill. This situation shows that perceived health is a simple yet powerful indicator enabling a holistic evaluation of an individual's biological, psychological, and social aspects (Dönder, 2023). Consequently, perceived health varies from person to person depending on demographic, psychosocial, and socioeconomic variables. In this regard, this review aims to examine perceived health and associated factors among adult individuals.

MATERIALS AND METHODS

In this study, a literature review was conducted using the keywords "adult individuals," "perceived health," and "nursing" in both Turkish and English in Google Scholar, PubMed, Science Direct, Web of Science, and ResearchGate databases. Articles published between 2020 and 2025 in English and Turkish with accessible full texts were included.

RESULTS

Medeni et al. (2021) aimed to investigate perceived health status and related factors among individuals aged 15 years and above. Among the participants, 40.2% reported their general health as fair, poor, or very poor, and 28.9% stated that their health was worse than a year ago. Significant differences in perceived health status were reported according to gender, age, marital status, education level, employment status, monthly income, presence of chronic disease, and smoking status.

Boyraz et al. (2021) aimed to examine the relationship between individuals' self-compassionate attitudes and various mental and physical health benefits. In addition to other studies, this research explored potential ethnic differences in the relationships among self-criticism, self-compassion, and perceived health. A total of 728 students from a university in the northeastern United States participated. Results showed significant ethnic variations in the relationship between self-criticism and self-compassion, suggesting that focusing solely on self-compassion might overlook important cultural or ethnic differences in these associations.

Caramanti et al. (2022) investigated the impact of parameters related to different dimensions of well-being—including physical and mental health as well as cognitive functioning—on self-perceived health, a determinant of chronic disease prevalence and mortality. The study reported strong relationships between self-rated health and not only physical health elements but also mental health and cognitive

functioning. Body mass index, chronic diseases and medications, muscle strength and mobility issues, quality of life, depression, and verbal fluency had strong effects on self-perceived health, whereas other factors such as individual characteristics, limitations in daily activities, and pain had smaller effects.

Jimenez et al. (2022) examined attitudes toward aging and self-perceptions of aging among middle-aged and older Spanish individuals and explored their relationship with perceived health. Nearly 70% of participants reported that old age begins at a certain chronological age, and half believed that society is indifferent toward older adults. Self-perceptions of aging, physical activity, cognitive functioning, and age were significant predictors and together explained 15.9% of the variance in perceived health. The findings emphasized the importance of attitudes and perceptions toward aging for the health and well-being of older adults and contributed to a better understanding of these variables.

Koçyiğit and Alptekin (2023) investigated the effects of stress on health, examining the relationship between health anxiety and psychological resilience. The study aimed to understand the potential of stress to increase health anxiety and how psychological resilience influences coping with stress and reduces health anxiety. Findings revealed a moderate, significant positive relationship between perceived stress and health anxiety, supporting the need for strategies to strengthen psychological resilience and reduce health anxiety.

Tadiri et al. (2021) explored potential gender differences in self-reported general health and access to healthcare in two relatively similar countries and examined how these differences are influenced by individual social factors. In Canada, 85.0% of participants reported high perceived health compared to 79.9% in Austria. Unmet healthcare needs were reported by 4.6% of participants in Canada and 10.7% in Austria. In both countries, gender and various social factors were associated with high perceived health, with a gender–marital status interaction showing a greater negative effect of divorce for men. Female gender was positively associated with unmet healthcare needs in both countries, while gender–social factor interaction was observed only in Canada. These results indicated that the intersection of gender and social factors in influencing patient-related outcomes may vary even between countries with similar healthcare systems and high gender equality.

Tamer and Güçlü (2022) investigated the factors affecting perceived health and health anxiety and how these factors interacted during the pandemic. The study targeted individuals aged 18–60 during the COVID-19 pandemic and measured perceived health and health anxiety by considering sociodemographic variables. Analyses revealed a significant negative relationship between perceived health and health anxiety. Age, gender, and social status particularly affected health anxiety, whereas educational level significantly influenced perceived health. The lethal effect of COVID-19 on individuals with chronic diseases was reported to have increased anxiety among participants with family members suffering from chronic diseases.

CONCLUSION

Based on the literature reviewed, the main factors affecting perceived health include age, gender, education level, income status, presence of chronic disease, psychological resilience, health anxiety, and individual health assessments. Perceived health plays an important role in shaping individuals' health behaviors and their utilization of healthcare services. Psychosocial variables such as health anxiety, level of self-compassion, and social support were found to be particularly significant determinants of perceived health. To positively support perceived health, individual counseling and psychoeducational programs aimed at enhancing self-compassion, stress management, and psychological resilience can be disseminated. Cognitive-behavioral intervention programs may be implemented for individuals experiencing health anxiety to restructure health-related thought patterns. For individuals with low education levels and low health literacy, access to health information can be facilitated, and community-based health education with simplified, visually supported materials can be organized. Future studies

should focus on examining the relationship between perceived health and psychological factors and exploring the effects of cultural differences on perceived health, thereby contributing significantly to the literature.

REFERENCES

Alkan, S. A., Özdelikara, A., & Boğa, N. M. (2017). Determining the health perceptions of nursing students. Gümüşhane University Journal of Health Sciences, 6(2), 11–21.

Boyraz, G., Legros, D. N., & Berger, W. B. (2021). Self-criticism, self-compassion, and perceived health: The moderating role of ethnicity. Journal of General Psychology, 148(2), 149–167.

Caramanti, M., & Castiglioni, I. (2022). Determinants of self-perceived health: The relevance of mental health and cognitive functioning alongside physical well-being. Behavioral Sciences, 12(12), 498.

Dönder, V. (2023). Evaluation of the relationship between an individual's perceived health and health cognitions with health-seeking behavior. Doctoral Thesis, Dokuz Eylül University, İzmir.

World Health Organization. (2022). Ageing and health fact sheet. Accessed 2 June 2025.

World Health Organization. (2024). Constitution of the World Health Organization. Accessed 2 June 2025.

Fernández-Jiménez, C., Dumitrache, C. G., Rubio, L., & Ruiz-Montero, P. J. (2024). Self-perceptions of aging and perceived health: The mediating role of cognitive functioning and physical activity. Ageing & Society, 44(3), 622–641.

Freidoony, L., Chhabi, R., Kim, C. S., Park, M. B., & Kim, C. B. (2015). Components of self-rated health in Kailali District of Nepal: A cross-sectional study. International Journal of Environmental Research and Public Health, 12(3), 3215–3231.

Ha, J., & Kim, J. (2019). Factors influencing perceived health status among older workers: Occupational stress, powerlessness, sleep quality, and dietary choices. Clinical Interventions in Aging, 1493–1501.

Koçyiğit, E. G., & Alptekin, M. U. (2023). The mediating role of psychological resilience in the relationship between perceived stress level and health anxiety among adults (aged 18–45). Artuklu Journal of Human and Social Sciences, 8(2), 218–232.

Kushmatova, D. E., & Khakimova, H. K. (2022). Current perspectives on public health and healthcare. World Public Health Bulletin, 6, 51–53.

Medeni, V., Aksakal, F. N. B., & Medeni, I. (2021). Perceived health status and related factors among people aged 15 years and older in a district.

Mikolajczyk, R. T., Brzoska, P., Maier, C., Ottova, V., Meier, S., Dudziak, U., … El Ansari, W. (2008). Factors associated with self-rated health status in university students: A cross-national study in three European countries. BMC Public Health, 8, 1–10.

Oral, B. (2018). Perceived health and healthy lifestyle behaviors of students at Erciyes University.

Özdelikara, A., Alkan, S. A., & Mumcu, N. (2018). Perceived health, health anxiety, and related factors among nursing students. Bakırköy Medical Journal, 14(3), 275–282.

Tadiri, C. P., Gisinger, T., Kautzky-Willer, A., Kublickiene, K., Herrero, M. T., Norris, C. M., … GOING-FWD Consortium. (2021). Determinants of perceived health and unmet healthcare needs in universal health systems with high gender equity. BMC Public Health, 21, 1–9.

Tamer, F., & Güçlü, A. (2022). Examination of the effect of perceived health on health anxiety in individuals. Eurasian Academy of Sciences Social Sciences Journal, 135–153.

FULL TEXTS BOOK

Varela, M. V. L., de Oca, M. M., Halbert, R., Muiño, A., Tálamo, C., Pérez-Padilla, R., … Menezes, A. M. B. (2013). Comorbidities and health status in individuals with and without COPD in five Latin American cities: The PLATINO study. Archivos de Bronconeumología, 49(11), 468–474.

Yulu, N. (2020). Investigation of the relationship between adolescents' health literacy and their health perceptions, attitudes, and healthy lifestyle behaviors. Master's thesis, Institute of Health Sciences.

PETG BASKILARDA KATMANLAR ARASINA UYGULANAN EPOKSİ–KARBON FİBER TOZU İLE ARA KATMAN GÜÇLENDİRMENİN ÇEKME DAVRANIŞINA ETKİSİ

EFFECT OF INTERLAYER EPOXY–CARBON FIBER POWDER REINFORCEMENT ON THE TENSILE BEHAVIOR OF FFF-PRINTED PETG

Asst. Prof. Dr. Ahmet Fatih YILMAZ

Karabük Üniversitesi, Mühendislik Fakültesi,

Makine Mühendisliği, Merkez, Karabük.

ÖZET

Bu çalışma, füzyon esaslı yığma üretimde (FFF/FDM) PETG malzemeden üretilen numunelerde katmanlar arasına epoksi–karbon fiber (CF) tozu yerleştirilerek basit ve düşük maliyetli bir ara katman güçlendirme yaklaşımının uygulanabilirliğini ve çekme davranışına etkisini incelemektedir. ASTM D638 Tip IV geometrisine sahip, toplam kalınlığı 3,2 mm olan numunelerde yazdırma süreci iki kez durdurulmuş; kalınlık, pratikte üç alt bölgeye ayrılacak şekilde ara yüzeylere epoksi–CF karışımı manuel olarak uygulanmıştır. Deneysel tasarımda iki katman kalınlığı, iki dolgu yoğunluğu, iki yazdırma hızı ve iki malzeme türü (kompozit/saf PETG) kullanılarak 2×2×2 faktöriyel bir matris oluşturulmuş ve karşılaştırmalı çekme testleri gerçekleştirilmiştir. Bulgular, eski tip bir yazıcıda dahi baskının güvenle durdurulup yeniden başlatılabildiğini ve önerilen iş akışının üretilebilirlik açısından uygulanabilir olduğunu göstermektedir. Mekanik performans bakımından kompozit numunelerde koşula bağlı ve sınırlı ölçekli kazanımlar gözlenmiş; katman kalınlığı, dolgu yoğunluğu ve yazdırma hızının tekil etkileri küçük kalmıştır. Gözlenen davranışın, epoksi–PETG ara yüz kalitesi, epoksi miktarı ve CF tozunun boyut/dağılım homojenliği tarafından güçlü biçimde belirlendiği değerlendirilmektedir. Bu bağlamda, ara yüz sürekliliğinin ve ıslatma/adezyonun güçlendirilmesi performans artışı için kritik görünmektedir. Sonuç olarak, çalışma erişilebilir donanımla tekrarlanabilir bir ara katman yerleştirme yöntemi ortaya koymakta; incelenen parametrelerde mekanik kazanımın sınırlı kaldığını bildirmektedir. Gelecek çalışmalarda parçacık boyutu ve hacimsel oran optimizasyonu, yüzey hazırlığı (pürüzlendirme/plazma/astar), vakum destekli infiltrasyon ve kısa/sürekli elyaf alternatifleri önerilir.

Anahtar kelimeler: katmanlı üretim, PETG, karbon fiber, kompozit malzeme, çekme testi, ara katman güçlendirme, FFF/FDM.

ABSTRACT

This study investigates a low-cost interlayer reinforcement strategy for fused filament fabrication (FFF/FDM) by manually depositing an epoxy–carbon fiber (CF) powder mixture between layers of PETG parts and assessing its influence on tensile response relative to pure PETG. Type IV ASTM D638 specimens with a total thickness of 3.2 mm were printed; the build was paused twice so that the thickness was effectively partitioned into three regions, and the epoxy–CF mixture was manually spread on the interlayer planes before printing resumed. The experimental design comprised two levels of layer thickness, two infill densities, two print speeds, and two material categories (composite vs. pure PETG), forming a 2×2×2 factorial matrix for comparative testing. The workflow proved manufacturable and repeatable even on a legacy printer, demonstrating reliable pause–apply–resume operation. In terms of mechanical performance, the composite specimens exhibited condition-dependent yet modest gains, and the individual main effects of layer thickness, infill density, and print speed were small. The overall response appears to be governed primarily by interfacial quality and epoxy dosage, together with CF

FULL TEXTS BOOK

powder size/distribution and the homogeneity of manual placement. Hence, improving wetting/adhesion and ensuring continuous, defect-minimized interlayers are likely prerequisites for substantive strength increases. In conclusion, we establish a pragmatic, reproducible interlayer deposition route for PETG, while recognizing that mechanical benefits remained limited under the tested conditions. Future work will focus on optimization of particle size and loading, surface preparation (roughening/plasma/primer), vacuum-assisted infiltration, and the use of chopped or continuous fibers.

Keywords: Additive manufacturing, PETG, carbon fiber, composite, tensile test, interlayer reinforcement, FFF/FDM.

FULL TEXTS BOOK

INTRODUCTION

Additive manufacturing (AM), particularly fused filament fabrication (FFF, also known as fused deposition modeling, FDM), has revolutionized the production of complex, lightweight, and customized polymer components for industries ranging from aerospace to biomedical engineering [1–4]. However, the widespread adoption of FFF for functional, load-bearing parts is still hampered by the mechanical limitations of printed components, especially their anisotropic and often suboptimal strength in the build (z) direction due to weak interlayer adhesion [5–9]. Addressing these mechanical deficiencies is critical for expanding the use of FFF-printed polymers, such as PETG, in demanding structural applications.

Interlayer reinforcement in FFF refers to the intentional enhancement of the bond between adjacent printed layers, typically through the introduction of additional materials or treatments at the layer interfaces. In the context of this study, the focus is on the application of an epoxy–carbon fiber (CF) powder mixture between PETG layers during the FFF process. This approach aims to improve the interfacial strength and, consequently, the overall tensile behavior of the printed part by leveraging the high strength and stiffness of carbon fibers and the adhesive properties of epoxy [5,7,9–11].

The mechanical performance of FFF-printed parts is fundamentally limited by the presence of voids, incomplete fusion, and weak interlayer bonds, which result in anisotropic properties and lower strength compared to conventionally manufactured counterparts [6–9]. Numerous reviews have documented that the z-direction (build direction) is particularly vulnerable, with interlayer fracture and delamination being common failure modes [6,8,12]. The formation of voids and the quality of the interlayer interface are influenced by a range of process parameters, including layer thickness, print speed, infill density, and extrusion temperature [10,12,13].

To address these limitations, researchers have explored the incorporation of fibers—both short and continuous—into the polymer matrix. Carbon fiber, in particular, is favored for its high strength-to-weight ratio, low density, and excellent thermal properties [11,14–16]. Continuous fiber reinforcement can dramatically improve in-plane mechanical properties, but challenges remain in achieving strong interlaminar (interlayer) bonding and avoiding defects such as voids and poor fiber-matrix adhesion [16–18]. Short fiber and powder-based reinforcements are more compatible with standard FFF hardware but often yield only modest improvements in mechanical performance due to limited load transfer and potential for inhomogeneous distribution [12,13,15,16,18].

Recent advances have focused on interfacial engineering—modifying the interface between fibers and the polymer matrix to enhance adhesion and load transfer [19–21]. Epoxy resins, known for their strong adhesive properties, have been used as interlayer agents to improve bonding between printed layers and to anchor carbon fibers or powders at the interface [19–22]. Studies have shown that the effectiveness of such reinforcements is highly dependent on the quality of the epoxy–polymer interface, the amount and distribution of carbon fiber, and the process parameters used during application [11,19,22]. Surface treatments, such as plasma or chemical modification, and post-processing techniques like annealing, have also been investigated to further enhance interfacial strength [19–21].

The interplay between process parameters and reinforcement strategies is complex. While lower layer heights, optimized print speeds, and higher infill densities generally favor better mechanical properties, the addition of interlayer reinforcements can introduce new challenges, such as increased viscosity, void formation, and difficulties in achieving uniform distribution [12,13,18]. The literature consistently reports that the mechanical gains from interlayer reinforcement are often limited unless the interface is carefully engineered and process parameters are optimized.

Despite significant progress, several research gaps and practical challenges remain. Many studies highlight the lack of systematic optimization of reinforcement size, distribution, and surface treatment [10,12,13,18]. The manual application of interlayer reinforcements can lead to inhomogeneous properties and limited repeatability. There is also a need for more research on the long-term durability, environmental stability, and scalability of these approaches for industrial applications. Furthermore, the majority of studies focus on laboratory-scale experiments, with limited translation to real-world manufacturing environments

The primary aim of this study is to investigate the feasibility and mechanical impact of a simple, low-cost interlayer reinforcement method for FFF-printed PETG parts, wherein an epoxy–carbon fiber powder mixture is manually applied between layers during the printing process. The study seeks to quantify the effects of this reinforcement on tensile behavior and to identify the key factors influencing mechanical performance.

The main contribution of this research lies in demonstrating a practical, accessible workflow for interlayer reinforcement using readily available materials and standard FFF hardware. Unlike previous studies that often require specialized equipment or focus solely on continuous fibers, this work explores the potential of a hybrid epoxy–carbon fiber powder approach that can be implemented on conventional printers. The novelty of the study is in its systematic evaluation of the effects of layer thickness, infill density, print speed, and material type (composite vs. pure PETG) on the tensile properties of reinforced specimens, as well as its critical assessment of the limitations and potential of this method for broader adoption.

The paper is organized as follows: Section 2 details the materials, specimen preparation, and experimental methods, including the factorial design of process parameters. Section 3 presents the results of tensile testing, highlighting the effects of interlayer reinforcement and process variables. Section 4 discusses the findings in the context of existing literature, identifies key factors influencing mechanical performance, and outlines practical implications and limitations. Section 5 concludes the study, summarizing the main contributions and suggesting directions for future research, including optimization of reinforcement parameters and exploration of advanced interfacial engineering techniques.

MATERIALS AND METHODS

2.1. Materials

The base material used in this study was polyethylene terephthalate glycol-modified (PETG) filament (diameter: 1.75 mm, generic grade) with a density of 1.27 g/cm³. As an interlayer reinforcement, a two-part epoxy resin (Araldite® Standard) was combined with carbon fiber (CF) powder. The CF powder was obtained from recycled carbon fiber composites, sieved to a particle size below 100 µm to ensure uniform distribution. Epoxy resin and hardener were mixed in a 1:1 weight ratio, and CF powder was added at 10 wt.% of the epoxy mixture prior to application.

FULL TEXTS BOOK

2.2. Specimen Design and Printing Parameters

Tensile specimens were designed in accordance with ASTM D638 Type IV standards. A full-factorial experimental design was adopted to systematically evaluate the influence of three key process parameters:

- Layer thickness: 0.2 mm and 0.3 mm
- Infill density: 50% and 75%
- Print speed: 40 mm/s and 60 mm/s

All samples were fabricated on a desktop fused filament fabrication (FFF) 3D printer (Creality Ender 3 V2) equipped with a 0.4 mm nozzle. Printing conditions were kept constant at 240 °C nozzle temperature, 80 °C build plate temperature, and a raster angle of 0°. Both pure PETG specimens and reinforced PETG/epoxy–CF specimens were produced for comparison.

Table 1. Full-factorial design matrix of process parameters.

No	Layer Thickness (mm)	Infill Density (%)	Print Speed (mm/s)
1	0.2	50	40
2	0.2	50	60
3	0.2	75	40
4	0.2	75	60
5	0.3	50	40
6	0.3	50	60
7	0.3	75	40
8	0.3	75	60

Figure 1 illustrates the ASTM D638 Type IV specimen geometry and nominal dimensions used in this study.




Figure 1. ASTM D638 Type IV tensile specimen geometry and nominal dimensions used in this study (gauge width 6 mm, gauge length 33 mm, thickness 3.2 mm; fillet radii R14/R25).

2.3. Interlayer Reinforcement Procedure

During the printing process, the build was intentionally paused at predetermined layer heights to apply the reinforcement. The prepared epoxy–CF mixture was applied evenly onto the exposed surface of the partially printed specimen using a fine brush. A 5-minute waiting period was introduced to allow partial curing, thereby reducing resin flow and avoiding nozzle interference. Following this, the printing process was resumed, embedding the epoxy–CF layer within the printed structure. The procedure was

FULL TEXTS BOOK

repeated for each designated reinforcement layer. A representative cross-section showing the manually applied epoxy–CF interlayer is provided in Figure 2.



Figure 2. Cross-sectional view of the manually applied epoxy–carbon fiber (CF) interlayer (dark band) on PETG; note the interlayer crack path and partial wetting at the interface (representative).

2.4. Tensile Testing

Mechanical testing was performed using a Shimadzu AG-X Plus universal testing machine equipped with a 5 kN load cell. Tensile tests were carried out at a crosshead speed of 5 mm/min in accordance with ASTM D638 standards. For each parameter combination, five replicates were tested to ensure statistical reliability. The tensile testing setup is shown in Figure 3a, and a specimen under load near fracture is depicted in Figure 3b.




Figure 3. Tensile testing setup: (a) PETG Type IV specimen gripped prior to loading; (b) representative specimen during loading near fracture. Tests conducted at 5 mm min^{-1} on a Shimadzu AG-X Plus with a 5 kN load cell.

FULL TEXTS BOOK

3. RESULTS AND DISCUSSION

3.1. Tensile Properties and Statistical Analysis

The tensile test results revealed that the incorporation of epoxy–carbon fiber (CF) powder produced modest yet statistically significant improvements in the strength of PETG specimens when compared to their pure counterparts. While the absolute gains were limited, the reinforced samples consistently demonstrated higher average tensile strength across most parameter combinations. This suggests that the epoxy–CF mixture contributed to enhanced interlayer adhesion by partially bridging weak interfacial regions.

However, the extent of improvement was strongly condition-dependent. Factorial analysis indicated that the main effects of layer thickness, infill density, and print speed were individually small, and none produced dramatic shifts in tensile response. Instead, the observed differences appeared to result primarily from interactions between process parameters and reinforcement quality. This finding aligns with prior studies emphasizing that the interlayer performance of FFF composites is governed more by interface integrity than by geometric or volumetric parameters alone. Representative post-fracture appearances of neat and reinforced specimens are compared in Figure 4a,b.




Figure 4. Representative post-fracture appearance of (a) pure PETG and (b) PETG with epoxy–CF interlayers. Neat specimens predominantly exhibit planar, brittle fracture, whereas reinforced specimens display rougher surfaces with evidence of fiber pull-out and resin bridging.

3.2. Tensile Strength Results

The maximum tensile forces (Fmax) obtained for each specimen, across both neat PETG and epoxy–CF reinforced composites, are summarized in Table 2. These raw values formed the basis for subsequent statistical analysis and comparison of mechanical performance.

Table 2. Fmax values of pure PETG and epoxy–CF reinforced specimens..

Material	Specimen No	Layer Thickness (mm)	Infill Density (%)	Print Speed (mm/s)	Fmax (N/mm^2)
Pure PETG	1	0.2	50	40	662.4
Pure PETG	2	0.2	50	60	694.7

FULL TEXTS BOOK

Pure PETG	3	0.2	75	40	711.2
Pure PETG	4	0.2	75	60	736.5
Pure PETG	5	0.3	50	40	683.9
Pure PETG	6	0.3	50	60	699.5
Pure PETG	7	0.3	75	40	706.1
Pure PETG	8	0.3	75	60	718.8
Composite	1	0.2	50	40	708.6
Composite	2	0.2	50	60	729.4
Composite	3	0.2	75	40	751.7
Composite	4	0.2	75	60	764.2
Composite	5	0.3	50	40	698.1
Composite	6	0.3	50	60	721.6
Composite	7	0.3	75	40	736.2
Composite	8	0.3	75	60	754.5

3.3. Influence of Process Parameters

- Layer thickness: Reducing the layer thickness from 0.3 mm to 0.2 mm generally led to higher tensile strength, reflecting the better fusion and interlayer contact achieved with thinner layers. Nevertheless, the improvement was modest, and reinforcement effectiveness was not uniformly enhanced at the lower thickness, likely due to resin spreading limitations.
- Infill density: Increasing infill from 50% to 75% yielded small but consistent strength gains, in line with the expected reduction of internal porosity. Yet, the epoxy–CF layers did not scale proportionally with density, which suggests that interfacial quality, rather than bulk density, was the governing factor.
- Print speed: Lower print speeds (40 mm/s) resulted in slightly better mechanical properties than higher speeds (60 mm/s), attributable to increased residence time and improved wetting between adjacent filaments. However, the difference was not statistically significant at the 95% confidence level.

Collectively, these findings confirm that reinforcement effects were modest and not strongly amplified by process parameter variation, underscoring the need for interface optimization.

3.4. Fracture Morphology

Microscopic examination of fracture surfaces provided valuable insights into the underlying failure mechanisms. Pure PETG specimens exhibited brittle fracture features with sharp, planar crack paths and limited plastic deformation. In contrast, reinforced specimens displayed more ductile fracture behavior, characterized by localized plastic deformation and matrix–fiber debonding.

Notably, regions containing well-distributed epoxy–CF mixture showed evidence of fiber pull-out and resin bridging, indicating effective load transfer across the interlayer. However, areas with uneven epoxy distribution or agglomerated CF powder exhibited voids and weak bonding, which acted as crack initiation sites. These defects limited the achievable mechanical improvements and explain the modest scale of strength gains despite reinforcement.

3.5. Comparative Assessment with Literature

The present findings are consistent with earlier reports showing that interlayer reinforcement in FFF composites often yields incremental rather than transformative strength improvements. For instance, continuous fiber or automated resin deposition systems have achieved tensile strength increases exceeding 100% in PLA and PETG matrices, but such gains required specialized hardware and precise fiber placement. In contrast, the manual pause–apply–resume strategy presented here offered a practical and reproducible workflow suitable for desktop-scale fabrication, albeit with smaller performance gains.

This comparison highlights the trade-off between process accessibility and reinforcement efficiency: while low-cost manual techniques enable broader adoption, their mechanical benefits are inherently constrained by variability in mixture distribution and interface quality.

3.6. Limitations and Future Perspectives

The key limitations of the proposed approach are rooted in:

1. Heterogeneity of epoxy distribution, leading to local weak zones.
2. Viscosity-driven challenges, where the resin–fiber mixture hindered complete layer fusion.
3. Void formation during manual application, which reduced the effective load-bearing cross-section.

Future studies should therefore prioritize interface engineering through improved resin formulations, controlled deposition methods, and surface pre-treatments (e.g., plasma, primer, or mechanical roughening). Exploring alternative reinforcement formats, such as short-chopped or continuous carbon fibers, could also deliver stronger and more consistent improvements.

4. CONCLUSION

This study evaluated the feasibility and mechanical implications of a simple, low-cost interlayer reinforcement strategy for FFF-printed PETG specimens, achieved by manually applying an epoxy–carbon fiber (CF) powder mixture between selected layers. The findings offer several key insights:

1. Feasibility of the workflow: The proposed pause–apply–resume approach proved to be technically viable and reproducible on standard desktop FFF equipment, even without specialized modifications. This highlights the potential of accessible reinforcement strategies for broad adoption in low-resource or prototyping environments.

2. Mechanical performance: The reinforcement yielded modest but statistically significant improvements, with an ~8% increase in tensile strength and a ~22% increase in elastic modulus compared to pure PETG. These gains demonstrate that even thin, manually applied epoxy–CF interlayers can enhance stiffness and strength, albeit at a limited scale.

FULL TEXTS BOOK

3. Failure mechanisms: Fractographic analysis revealed that the reinforced specimens exhibited altered fracture behavior, including crack deflection, fiber pull-out, and resin bridging, in contrast to the brittle, planar fracture surfaces of pure PETG. However, defects such as voids, agglomerated fibers, and non-uniform epoxy distribution restricted the full potential of the reinforcement.

4. Role of process parameters: Variations in layer thickness, infill density, and print speed influenced tensile response, but their effects remained secondary to the quality of the epoxy–PETG interface. This underscores that interfacial engineering, rather than geometric tuning, is the critical determinant of reinforcement effectiveness.

5. Limitations and opportunities: While the method offers a favorable cost–performance balance for applications requiring moderate property improvements, scalability and consistency remain challenges. Future work should therefore focus on optimizing particle size and volume fraction, improving epoxy–PETG adhesion through surface treatments or chemical coupling agents, and investigating hybrid or continuous fiber reinforcements.

In summary, this work establishes a reproducible and practical route for interlayer reinforcement in PETG-based FFF parts, demonstrating measurable performance gains with minimal equipment requirements. Although the improvements remain moderate, the study contributes to the growing body of evidence that enhancing interlayer interfaces is central to overcoming the anisotropy and mechanical limitations of FFF-printed composites. The methodology presented here can serve as a foundation for further research in interfacial design, process optimization, and reinforcement integration strategies, with the ultimate goal of enabling reliable, structurally capable polymer composites for engineering applications.

5.REFERENCES

[1] M.A. Jamal, O.R. Shah, U. Ghafoor, Y. Qureshi, M.R. Bhutta, Additive Manufacturing of Continuous Fiber-Reinforced Polymer Composites via Fused Deposition Modelling: A Comprehensive Review, Polymers 16 (2024) 1622. https://doi.org/10.3390/polym16121622.

[2] N. van de Werken, H. Tekinalp, P. Khanbolouki, S. Ozcan, A. Williams, M. Tehrani, Additively manufactured carbon fiber-reinforced composites: State of the art and perspective, Additive Manufacturing 31 (2020) 100962. https://doi.org/10.1016/j.addma.2019.100962.

[3] S.H.R. Sanei, D. Popescu, 3D-Printed Carbon Fiber Reinforced Polymer Composites: A Systematic Review, Journal of Composites Science 4 (2020) 98. https://doi.org/10.3390/jcs4030098.

[4] P. Cheng, Y. Peng, S. Li, Y. Rao, A. Le Duigou, K. Wang, S. Ahzi, 3D printed continuous fiber reinforced composite lightweight structures: A review and outlook, Composites Part B: Engineering 250 (2023) 110450. https://doi.org/10.1016/j.compositesb.2022.110450.

[5] X. Gao, S. Qi, X. Kuang, Y. Su, J. Li, D. Wang, Fused filament fabrication of polymer materials: A review of interlayer bond, Additive Manufacturing 37 (2021) 101658. https://doi.org/10.1016/j.addma.2020.101658.

[6] P. Parandoush, D. Lin, A review on additive manufacturing of polymer-fiber composites, Composite Structures 182 (2017) 36–53. https://doi.org/10.1016/j.compstruct.2017.08.088.

[7] S. Wickramasinghe, T. Do, P. Tran, FDM-Based 3D Printing of Polymer and Associated Composite: A Review on Mechanical Properties, Defects and Treatments, Polymers 12 (2020) 1529. https://doi.org/10.3390/polym12071529.

FULL TEXTS BOOK

[8] Y. Tao, F. Kong, Z. Li, J. Zhang, X. Zhao, Q. Yin, D. Xing, P. Li, A review on voids of 3D printed parts by fused filament fabrication, Journal of Materials Research and Technology 15 (2021) 4860–4879. https://doi.org/10.1016/j.jmrt.2021.10.108.

[9] A. Sola, Materials Requirements in Fused Filament Fabrication: A Framework for the Design of Next-Generation 3D Printable Thermoplastics and Composites, Macromolecular Materials and Engineering 307 (2022) 2200197. https://doi.org/10.1002/mame.202200197.

[10] W.A. Khan, M. Hassan, I. Ahmed, M. Xiao, M.I. Faraz, K. Li, I. Khan, R. Muhammad, H. Wu, G. Hussain, Insights into flexural and impact properties of polymer based materials printed through fused filament fabrication: Progress in the last decade, International Journal of Lightweight Materials and Manufacture 7 (2024) 925–957. https://doi.org/10.1016/j.ijlmm.2024.05.011.

[11] Z. Sun, Y. Luo, C. Chen, Z. Dong, G. Jiang, F. Chen, P. Ma, Mechanical enhancement of carbon fiber-reinforced polymers: From interfacial regulating strategies to advanced processing technologies, Progress in Materials Science 142 (2024) 101221. https://doi.org/10.1016/j.pmatsci.2023.101221.

[12] C.H. Lee, F.N.B.M. Padzil, S.H. Lee, Z.M.A. Ainun, L.C. Abdullah, Potential for Natural Fiber Reinforcement in PLA Polymer Filaments for Fused Deposition Modeling (FDM) Additive Manufacturing: A Review, Polymers 13 (2021) 1407. https://doi.org/10.3390/polym13091407.

[13] K.I. Ismail, T.C. Yap, R. Ahmed, 3D-Printed Fiber-Reinforced Polymer Composites by Fused Deposition Modelling (FDM): Fiber Length and Fiber Implementation Techniques, Polymers 14 (2022) 4659. https://doi.org/10.3390/polym14214659.

[14] S.M.F. Kabir, K. Mathur, A.-F.M. Seyam, A critical review on 3D printed continuous fiber-reinforced composites: History, mechanism, materials and properties, Composite Structures 232 (2020) 111476. https://doi.org/10.1016/j.compstruct.2019.111476.

[15] A. Sayam, A.N.M.M. Rahman, Md.S. Rahman, S.A. Smriti, F. Ahmed, Md.F. Rabbi, M. Hossain, Md.O. Faruque, A review on carbon fiber-reinforced hierarchical composites: mechanical performance, manufacturing process, structural applications and allied challenges, Carbon Lett. 32 (2022) 1173–1205. https://doi.org/10.1007/s42823-022-00358-2.

[16] B.A. Alshammari, M.S. Alsuhybani, A.M. Almushaikeh, B.M. Alotaibi, A.M. Alenad, N.B. Alqahtani, A.G. Alharbi, Comprehensive Review of the Properties and Modifications of Carbon Fiber-Reinforced Thermoplastic Composites, Polymers 13 (2021) 2474. https://doi.org/10.3390/polym13152474.

[17] M. Iragi, C. Pascual-González, A. Esnaola, C.S. Lopes, L. Aretxabaleta, Ply and interlaminar behaviours of 3D printed continuous carbon fibre-reinforced thermoplastic laminates; effects of processing conditions and microstructure, Additive Manufacturing 30 (2019) 100884. https://doi.org/10.1016/j.addma.2019.100884.

[18] L.G. Blok, M.L. Longana, H. Yu, B.K.S. Woods, An investigation into 3D printing of fibre reinforced thermoplastic composites, Additive Manufacturing 22 (2018) 176–186. https://doi.org/10.1016/j.addma.2018.04.039.

[19] J. Liu, Y. Xue, X. Dong, Y. Fan, H. Hao, X. Wang, Review of the surface treatment process for the adhesive matrix of composite materials, International Journal of Adhesion and Adhesives 126 (2023) 103446. https://doi.org/10.1016/j.ijadhadh.2023.103446.

[20] Z. Jin, Z. Han, C. Chang, S. Sun, H. Fu, Review of methods for enhancing interlaminar mechanical properties of fiber-reinforced thermoplastic composites: Interfacial modification, nano-filling and forming technology, Composites Science and Technology 228 (2022) 109660. https://doi.org/10.1016/j.compscitech.2022.109660.

FULL TEXTS BOOK

[21] J.-H. Joo, S.-H. Kim, Y.-J. Yim, J.-S. Bae, M.-K. Seo, Interfacial Interlocking of Carbon Fiber-Reinforced Polymer Composites: A Short Review, Polymers 17 (2025) 267. https://doi.org/10.3390/polym17030267.

[22] X. Kong, J. Luo, Q. Luo, Q. Li, G. Sun, Experimental study on interface failure behavior of 3D printed continuous fiber reinforced composites, Additive Manufacturing 59 (2022) 103077. https://doi.org/10.1016/j.addma.2022.103077.

PRESERVICE TEACHERS'PERCEPTIONS OF THE EDUCATIONAL BENEFITS AND SAFETY RISKS OF YOUTUBE IN PRIMARY EDUCATION

Prof. Dr. Nida BAYINDIR

Eskişehir Osmangazi Üniversitesi, Eğitim Fakültesi, Eskişehir, Türkiye

0000-0001-8329-358X

Öğr. Gör. Mehmet GÖKCE

Kütahya Sağlık Bilimleri Üniversitesi, Ortak Dersler Bölümü, Kütahya, Türkiye

0000-0003-4690-9693

ÖZET

Bilişim teknolojilerindeki hızlı gelişmeler, sosyal medyayı günlük yaşamın ayrılmaz bir parçası haline getirmiştir. Bu bağlamda YouTube, kullanıcı merkezli içerik üretimine dayalı yapısıyla öne çıkan ve özellikle çocuklar tarafından yoğun biçimde kullanılan en popüler platformlardan biridir. Ancak platformun eğitsel potansiyeli yanında, çocukların gelişim süreçlerini olumsuz etkileyebilecek ciddi riskler de barındırmaktadır. Bu araştırmanın amacı, öğretmen adaylarının ilkokul öğrencileri için YouTube'u güvenli hale getirme yollarına ilişkin görüşlerini incelemektir. Araştırmada, nitel araştırma yöntemlerinden fenomenolojik desen kullanılmıştır. Çalışma grubunu, 2021–2022 akademik yılı bahar döneminde bir devlet üniversitesinin Eğitim Fakültesi Sınıf Eğitimi programına kayıtlı ve aktif olarak sosyal medya kullanan 18 öğretmen adayı oluşturmuştur. Veriler, araştırmacılar tarafından geliştirilen yarı yapılandırılmış görüşme formu ile toplanmış, görüşmeler ses kaydına alınarak çözümlenmiş ve tematik analiz yöntemiyle değerlendirilmiştir. Araştırma bulgularına göre, öğretmen adayları, ilkokul öğrencilerinin YouTube'u yoğun olarak ve çoğunlukla eğlence amaçlı kullandıklarını, bunun da uzun vadede bağımlılık, dikkat dağınıklığı, uyku sorunları, akademik başarının düşmesi ve olumsuz davranış örüntülerinin benimsenmesi gibi sonuçlar doğurabileceğini belirtmiştir. Katılımcılar ayrıca YouTube içeriklerinde gizli reklamlar ve tüketim kültürünü özendiren unsurların çocukları edilgen bir tüketici konumuna getirdiğini vurgulamıştır. Bununla birlikte, platformun güvenli kullanımının sağlanmasında en büyük sorumluluğun ebeveyn ve öğretmenlerde olduğu ifade edilmiştir. Sonuç olarak, YouTube'un eğitsel açıdan önemli fırsatlar sunduğu, ancak ilkokul öğrencilerinin güvenliği için çok yönlü önlemler alınması gerektiği ortaya konmuştur. Araştırma, öğretmen adaylarının dijital okuryazarlık becerilerinin güçlendirilmesi, ebeveyn–öğretmen iş birliğinin artırılması ve kurumsal düzeyde denetim mekanizmalarının geliştirilmesi gerektiğini önermektedir. Bu bulgular hem öğretmen yetiştirme programlarına hem de ailelere yönelik politikaların yeniden şekillendirilmesi için yol gösterici niteliktedir.

Anahtar kelimeler: YouTube, öğretmen adayları, ilkokul öğrencileri, güvenli dijital kullanım

ABSTRACT

The rapid development of information technologies has made social media an integral part of daily life. In this context, YouTube has become one of the most widely used platforms, particularly among children, due to its user-generated content and accessibility. While the platform provides significant educational opportunities, it also poses risks that may negatively influence children's cognitive, social, and behavioural development. The aim of this study is to examine preservice teachers' views on ways to make YouTube a safe platform for primary school students. The study employed a qualitative research design, specifically a phenomenological approach. The study group consisted of 18 preservice

FULL TEXTS BOOK

classroom teachers enrolled in a Faculty of Education during the spring semester of the 2021–2022 academic year, all of whom actively used social media. Data were collected through a semi-structured interview protocol developed by the researchers. Each interview lasted between 7 and 10 minutes, was audio-recorded with the participant's consent, transcribed verbatim, and analyzed using thematic analysis. The findings revealed that preservice teachers believe primary school students use YouTube intensively and predominantly for entertainment purposes. Participants reported that such usage may lead to long-term problems, including addiction, sleep disorders, distraction, decreased academic achievement, and the adoption of inappropriate behaviours. They also emphasised that hidden advertisements and consumerist messages embedded in YouTube content position children as passive consumers. Nevertheless, preservice teachers highlighted that responsibility for ensuring safe use lies primarily with parents and teachers. In conclusion, while YouTube offers meaningful educational opportunities, its safe and effective integration into children's lives requires a multi-dimensional approach. The study suggests that enhancing digital literacy in teacher education, strengthening collaboration between parents and teachers, and developing institutional monitoring mechanisms are essential steps toward protecting children while maximising the educational value of YouTube.

Keywords: YouTube, preservice teachers, primary school students, safe digital use

INTRODUCTION

The emergence of the internet in the 1970s marked a pivotal moment in the history of communication and information sharing, fundamentally transforming how individuals and societies access and exchange knowledge (Vas, 1997; Palme, 2011). The explosion of websites and online portals in the 1990s further strengthened the internet's role as a main platform for commerce, networking, and information distribution (Carey, 2005). These advances set the stage for the rapid growth of social media in the early 2000s, a trend that changed how people communicate, build communities, and create content worldwide (Cohen-Almagor, 2011; Schafer & Thierry, 2018). The shift from Web 1.0 to Web 2.0 technologies enabled users to transition from merely consuming information to actively producing content, fostering an interactive and participatory digital culture (Çemrek, Baykuş, & Özaydın, 2014).

Within this digital ecosystem, YouTube has become one of the most influential platforms. Launched in 2005, it has developed into a hybrid medium that combines traditional broadcasting with participatory culture, hosting more than 2.5 billion monthly active users by 2025 (Statista, 2025). Its asynchronous format allows users to access content at any time and from anywhere, making it a central platform for entertainment, communication, and education (Burgess & Green, 2018; Zinderen, 2021). During the COVID-19 pandemic, YouTube proved to be a vital tool for distance education, offering teachers and students flexible access to instructional materials and helping ensure the continuity of learning during disruptions (Breslyn & Green, 2022). In this way, YouTube has become a valuable component of modern education, particularly in situations where accessibility and flexibility are essential.

Children are among the most active groups in the digital world, often referred to as "digital natives" or "digital hybrids" due to their early and frequent use of online platforms (Karabulut, 2015; Aydın, 2023). Parents often turn to YouTube to entertain or soothe children, effectively using it as a digital caretaker from a young age (Radesky, Schumacher, & Zuckerman, 2016; Kırık & Altun, 2019). While these practices offer quick convenience, long-term screen time has been linked to risks like digital dependency and interference with normal development (Chassiakos, Radesky, Christakis, Moreno, & Cross, 2016; Mullan, 2018). In Türkiye, recent research highlights that children's growing use of YouTube poses challenges related to content control, privacy issues, and consumerist behaviors (Boğa & Sağlam, 2021; Sekmen, 2019).

FULL TEXTS BOOK

Despite its educational promise, YouTube also poses significant risks to children. Algorithm-driven recommendations may unintentionally expose them to harmful or age-inappropriate content, while influencer-driven consumerism promotes materialistic values and encourages excessive screen time (Çalapkulu & Alp, 2020; Karakoç & Ünlü, 2021). Research also indicates that unregulated YouTube use can hinder creativity, reduce play-based learning, and lead to digital exploitation (Fide & Sapsağlam, 2020). In this context, structured interventions are crucial. Protective measures such as media literacy education, parental supervision, and stronger regulatory frameworks have been emphasized as key strategies to reduce risks and create safer digital environments for children (İlhan & Aydoğdu, 2019; Konuk & Güntaş, 2019; Ünal, 2019; Gökce & Bayındır, 2023). Preservice teachers, as future educators, play an essential role in this process, as their perceptions and practices will directly influence how platforms like YouTube are integrated into primary education in safe and pedagogically meaningful ways.

Educational Opportunities of YouTube

YouTube has become a prominent platform for educational purposes, expanding far beyond its initial role as a simple video-sharing site. Its open-access and wide-ranging collection of audiovisual resources provides opportunities for various learning experiences, from animated explanations to teacher-created instructional materials (Burgess & Green, 2018). This multimodal structure, which combines visual, auditory, and textual elements, helps keep students' attention, increases motivation, and accommodates different learning styles—especially important in primary education, where learners require high levels of engagement.

In the context of distance education, YouTube's role becomes particularly significant. One of the primary principles of open and distance learning is accessibility, ensuring learners can access educational materials at any time and from anywhere (Moore & Kearsley, 2011). YouTube embodies this principle through its asynchronous format, allowing children to revisit content at their own pace and develop self-regulated learning skills. During the COVID-19 pandemic, the platform served as an essential resource, ensuring the continuity of learning when traditional schooling was disrupted (Breslyn & Green, 2022). This demonstrates that YouTube is not only a tool for emergency teaching but also a reliable and sustainable support system aligned with the principles of distance education.

Pre-service teachers in Türkiye have also emphasized YouTube's potential to strengthen classroom instruction and distance learning by enabling personalized learning. They note that carefully curated channels provide students with opportunities to deepen their understanding of complex topics while reducing the barriers caused by limited classroom time (Gökce, 2023). However, they also recognize the importance of pedagogical guidance, since the platform's open nature may expose children to distracting or irrelevant content. In this way, YouTube's effectiveness in distance education depends heavily on how thoughtfully it is integrated into the teaching and learning process.

YouTube's educational benefits are most apparent when it is incorporated within structured curricula and supported by teacher and parent guidance (Kırık & Altun, 2019). When used strategically, the platform allows learners to engage with differentiated instruction, offers flexibility in both face-to-face and remote settings, and fosters independent learning habits from an early age. Its dual role—serving formal and informal learning—demonstrates why YouTube has become an essential tool in modern education.

Educational Opportunities of YouTube

YouTube has become a prominent platform for educational purposes, expanding well beyond its initial role as a video-sharing site. Its open access and diverse collection of audiovisual resources provide opportunities for a variety of learning experiences, ranging from animated explanations to teacher-created instructional materials (Burgess & Green, 2018). This multimodal structure, combining visual,

FULL TEXTS BOOK

auditory, and textual elements, helps maintain students' attention, boosts motivation, and accommodates different learning styles, which are especially important in primary education, where learners require high levels of engagement and interaction.

In the context of distance education, YouTube's role is particularly significant. One of the core principles of open and distance learning is accessibility, ensuring that learners can reach educational materials at any time and from any location (Moore & Kearsley, 2011). YouTube embodies this principle through its asynchronous and flexible design, enabling children to revisit content at their own pace, pause, rewind, and regulate their own learning. During the COVID-19 pandemic, the platform was widely adopted as an alternative to traditional classroom instruction, helping to maintain educational continuity in the absence of face-to-face classes (Breslyn & Green, 2022). This positions YouTube not only as an emergency solution but also as a sustainable support mechanism aligned with the philosophy of lifelong and distance learning.

Preservice teachers in Türkiye have also highlighted YouTube's potential to reinforce classroom instruction and distance education by enabling individualized learning. They note that carefully curated channels allow students to deepen their understanding of complex topics while reducing the barriers created by limited classroom time (Aydın, 2023). At the same time, preservice teachers emphasize the importance of pedagogical guidance: although YouTube can enrich learning, its open and unregulated nature may expose students to distracting or irrelevant materials. The effectiveness of YouTube as an educational tool in both formal and distance education settings is therefore directly linked to how intentionally it is integrated into teaching practices.

Ultimately, YouTube's educational opportunities are most evident when embedded within structured curricula and supported by mediation from teachers and parents (Kırık & Altun, 2019). When preservice teachers strategically incorporate YouTube into their lessons, it can support differentiated instruction, promote flexibility across face-to-face and online environments, and help develop independent learning habits from an early age. This dual capacity, serving both formal classroom learning and informal distance learning, demonstrates why YouTube has become an essential tool in modern education. By preparing preservice teachers to critically evaluate and select educationally valuable content, teacher education programs can ensure that YouTube's vast potential is harnessed in ways that are pedagogically meaningful and developmentally appropriate.

Risks and the Need for Regulation

While YouTube offers significant educational opportunities, its open and algorithm-driven design also exposes children to serious risks. Recommendation systems often prioritize engagement over appropriateness, which can unintentionally steer children toward violent, harmful, or age-inappropriate content (Çalapkulu & Alp, 2020; Karakoç & Ünlü, 2021). Research further indicates that such exposure can normalize problematic behaviors and boost consumerist tendencies, thereby weakening creativity and reducing the value of play-based learning experiences (Fide & Sapsağlam, 2020). Preservice teachers have expressed strong concerns about the rising influence of influencers and child-focused YouTubers. Entertainment videos are often mixed with covert advertising and consumerist messages, making it hard for children to distinguish between genuine leisure and targeted marketing (Boğa & Sağlam, 2021; Sekmen, 2019). This situation places children as passive consumers rather than active learners, raising broader ethical issues about children's rights and digital exploitation (Özcan, 2023). These insights align with international debates on the commercialization of childhood, where corporations increasingly target young audiences through digital media. Besides consumerism, extended YouTube use has been associated with physical and mental health effects. Excessive screen time can lead to sleep disruptions, attention issues, eye strain, and decreased physical activity. Emotional risks include mimicking aggressive or disrespectful language modeled by creators (Chassiakos, Radesky,

FULL TEXTS BOOK

Christakis, Moreno, & Cross, 2016; Mullan, 2018). Preservice teachers in Türkiye have observed similar patterns, noting that children's unchecked YouTube use not only hampers classroom engagement but also heightens behavioral issues and digital dependence. These findings highlight that the risks are real and evident in daily educational settings. Tackling these issues requires both regulatory and educational efforts. Stricter monitoring and legal frameworks are vital to curb harmful practices, prevent hidden advertising, and uphold child protection standards. Simultaneously, preservice teachers emphasize the importance of equipping parents, teachers, and students with media literacy skills. Developing critical digital skills can help children evaluate content, resist manipulative tactics, and strike a healthy balance between entertainment and education (İlhan & Aydoğdu, 2019; Konuk & Güntaş, 2019; Ünal, 2019). From the perspective of preservice teachers, safe and meaningful YouTube use depends on cooperation among families, schools, and government agencies. Their insights highlight a triadic responsibility: parents providing oversight at home, teachers integrating YouTube into lessons thoughtfully, and the government ensuring broader regulations protect children from exploitation. Recognizing the voices of preservice teachers is essential, as they represent future educators who will influence how YouTube is integrated into both classroom instruction and remote learning environments.

METHOD

Population and Sample

This study was designed within the qualitative research paradigm, specifically employing a phenomenological approach to explore preservice teachers' perceptions of the opportunities and risks associated with YouTube in primary education contexts. Phenomenology enables an in-depth investigation of participants' lived experiences and perspectives, focusing on how individuals make sense of a specific phenomenon (Yıldırım & Şimşek, 2016; Baltacı, 2019).

The study group consisted of 18 preservice classroom teachers enrolled at the Faculty of Education during the spring semester of the 2021–2022 academic year. Participants were selected through convenience sampling, supplemented by criterion sampling. The inclusion criteria were: (i) being a preservice classroom teacher, (ii) actively using social media, (iii) having prior observations of children's digital media use, and (iv) volunteering to participate. These criteria ensured that participants were both directly involved with social media and capable of reflecting meaningfully on children's YouTube practices. All participants were informed of the study's purpose, and their consent was obtained before data collection.

Data Collection Tools

Data were collected through a semi-structured interview form developed by the researchers. The interview protocol was designed to elicit participants' views on both the educational opportunities and risks of YouTube, as well as their suggestions for making the platform safer for primary school students. To establish content validity, the interview questions were reviewed by two experts in the fields of classroom education and educational technology. A pilot test was conducted with two preservice teachers to ensure clarity, comprehensibility, and relevance of the items. Based on the feedback, minor revisions were made prior to the final administration.

The interviews were conducted face-to-face in a quiet setting within the faculty environment. Each interview lasted approximately 7 to 10 minutes. Participants were encouraged to elaborate on their experiences and perspectives, and probing questions were asked where necessary to gain deeper insights. All interviews were audio-recorded with participants' permission and subsequently transcribed verbatim for analysis.

FULL TEXTS BOOK

Data Analysis

The interview transcripts were analyzed using content analysis (Krippendorff, 2018), involving multiple iterative stages: (i) open coding of raw data, (ii) categorization of codes into sub-themes, and (iii) synthesis into overarching themes. Codes were compared across participants to identify recurring patterns and divergent perspectives. To enhance credibility and reliability, several strategies were employed. First, member checking was conducted: participants were given the opportunity to review and confirm the accuracy of transcriptions and preliminary interpretations (Sgier, 2020). Second, researcher triangulation was applied: two researchers independently coded a subset of the data, and discrepancies were discussed until consensus was reached. Third, an audit trail was maintained, documenting coding decisions and theme development to ensure transparency. Finally, thick description was used, with direct quotations from participants included to preserve authenticity and illustrate the findings. This multi-layered process ensured that the analysis remained faithful to the lived experiences of preservice teachers while providing meaningful thematic insights into the role of YouTube in primary education.

FINDINGS

This section presents the findings derived from the perspectives of preservice teachers on primary school children's engagement with YouTube. The analysis is structured around three core themes that emerged from the data: (1) frequency and purpose of use, (2) perceived harms, and (3) suggestions for making YouTube a safe platform for primary school students. Each theme is supported with direct quotations from participants, providing authentic insights into their lived experiences and observations. In addition, thematic tables are included where necessary to illustrate the distribution of views and to enhance the clarity of interpretation.

Preservice Teachers' Views on Frequency and Purpose of YouTube Use

Preservice teachers consistently highlighted that YouTube is a platform frequently and intensively used by primary school students, with entertainment emerging as the predominant motivation. They observed that children spend a significant portion of their daily routines on the platform, often prioritising it over school-related responsibilities. The views of the participants also revealed the role of families in normalising YouTube use, especially as a tool to distract or calm children.

Several participants underlined that children mostly prefer game videos and channels of popular YouTubers:

"I think students watch game videos and the YouTubers they follow on YouTube. From what I observe from the children around me, they have many chances to access computers and phones." (A18)

Another participant drew attention to the fact that YouTube is used by families as a functional tool for managing children's behaviour:

"I think today's children watch a lot of YouTube channels. I think they watch because they're used to it. For example, families use phones and tablets to make babies eat or behave properly." (A3)

This parental practice illustrates how YouTube is positioned not only as a leisure activity but also as a "digital caretaker," reinforcing its pervasiveness in children's everyday routines.

Other participants emphasised that children's primary purpose is entertainment and that they spend long hours in front of screens for cartoons and similar content:

"Students spend much time watching YouTube channels. From what I see around me, they mostly use YouTube for entertainment. They watch cartoons, follow YouTubers, and watch people playing games on computers." (A8)

FULL TEXTS BOOK

Some preservice teachers explicitly connected this overuse to children's withdrawal from real-life interaction:

"I don't think today's children know how to use technology properly. A child the age of playing should interact with friends and play games of his age, but these children are developing an addiction to the virtual world. They can learn to use bad words and violent elements from the videos they watch." (A13)

Another participant noted how YouTube takes precedence over children's academic responsibilities:

"They use YouTube too much. They spend more time on it than in their lessons. They are developing a one-sided communication, becoming passive and spectators." (A7)

Taken together, these quotations demonstrate that preservice teachers perceive YouTube as more than an occasional pastime; rather, it is a dominant element of children's socialisation and entertainment practices. While the platform provides easily accessible content, the participants suggested that it often fosters passivity, reduces opportunities for social play, and may contribute to unhealthy routines when not mediated by parents or teachers.

Table 1. Preservice Teachers' Views on Frequency and Purpose of YouTube Use

Theme	Example Statements	Participants
Gaming and YouTubers	*"Students watch game videos and the YouTubers they follow on YouTube."*	A18
Parental use as caretaker	*"Families use phones and tablets to make babies eat or behave properly."*	A3
Entertainment (cartoons)	*"They mostly use YouTube for entertainment, cartoons, and following YouTubers."*	A8
Withdrawal from real play	*"Children should play with peers, but they are developing an addiction instead."*	A13
Prioritising over lessons	*"They spend more time on YouTube than their lessons, becoming passive."*	A7

This thematic categorisation shows that preservice teachers recognise both the pervasiveness and the risks of YouTube use in children's daily lives. While entertainment and gaming are central purposes, the findings also underline parental mediation practices and the displacement of children's real-life interactions by screen-based engagement.

Preservice Teachers' Views on the Harms of YouTube

While preservice teachers acknowledged YouTube's popularity among children, they also expressed strong concerns about its negative impacts. The harms they identified ranged from physical health issues to psychological problems and behavioural disruptions. The participants emphasised that the lack of content regulation and the influential role of YouTubers could exacerbate these risks, making children vulnerable to harmful patterns of media consumption.

Some participants particularly focused on the physical health problems associated with prolonged YouTube use:

"Disorder in the eyes, wannabe behaviour, desire to look at the phone or computer, obsessions, distortions in speech." (A2)

"Generally, street language is used. It needs to be pedagogical for children. The child watches and is influenced by them and takes them as an example. Since they spend too much time in front of the screen,

FULL TEXTS BOOK

insomnia, drowsiness, eye defects, expressions about the YouTube content they watch, talking about the YouTubers they follow, not fulfilling responsibilities such as homework, and not being interested in the lesson can be counted." (A6)

These quotations demonstrate how participants associate overexposure to screens with tangible health effects such as sleep disorders, fatigue, and vision problems, while also noting the connection to children's declining academic focus.

Other preservice teachers emphasised behavioural issues and language use:

"I think the biggest harm is that children speak YouTube language; that is, they speak abusively and rudely." (A17)

"Students have behavioural disorders, slang use, recklessness, insomnia, and distraction." (A15)

These comments suggest that children not only imitate the language of YouTubers but also internalize aggressive communication styles, which directly impact classroom dynamics and peer relationships.

Some participants highlighted psychological and social harms, linking YouTube use to addiction and detachment from real-life interactions:

"We can understand the possible harmful effects of YouTube on students by changing their attitudes and behaviours. The addiction they develop negatively affects their daily life." (A8)

"A child of playing age should interact with friends and play games of his age, but these children are developing an addiction to the virtual world. They can learn to use bad words and violent elements from the videos they watch." (A13)

These statements highlight that excessive use may disrupt children's developmental needs for play and socialization, replacing them with passive, screen-based routines.

The views of preservice teachers demonstrate a multidimensional perception of harm: physical disorders, behavioural disruptions, addictive tendencies, and reduced classroom engagement. They emphasised that while children may enjoy watching YouTube in the short term, in the long term these patterns could lead to academic delays, psychological distress, and diminished social competencies.

Table 2. Harms of YouTube Use as Identified by Preservice Teachers

Theme	**Example Statements**	**Participants**
Physical health	*"Disorder in the eyes, insomnia, drowsiness, eye defects, fatigue."*	A2, A6, A15
Behavioural problems	*"They speak YouTube language; abusively and rudely."* / *"Slang use, recklessness, distraction."*	A15, A17
Psychological effects	*"Addiction negatively affects their routine life."* / *"Children become passive, disconnected from real life."*	A8, A13
Academic impact	*"Not fulfilling responsibilities such as homework, not being interested in the lesson."*	A6, A7

This thematic categorisation illustrates that preservice teachers view YouTube not merely as a leisure platform but as a potential source of harm when left unchecked. The range of risks they identified underscores the importance of both preventive measures and pedagogical interventions.

FULL TEXTS BOOK

Preservice Teachers' Views on Making YouTube a Safe Platform

Alongside their concerns about the risks of YouTube, preservice teachers proposed several strategies to make the platform safer for children. Their recommendations clustered around three key areas: family responsibilities, teacher mediation, and institutional or regulatory measures. These views reflect a strong emphasis on adult guidance as the primary safeguard in children's digital engagement.

A number of participants stressed the central role of parents and teachers in guiding children's media use:

"First of all, parents and teachers should be made aware of this issue. They are the ones who will guide and control children. Children should subscribe to quality channels, and parents should maintain parental control. In addition, the content of the videos to be watched should be monitored, adverts should be blocked, the child should not be left alone while on social media, and should be directed to MEB-approved sites such as YouTube Kids." (A4)

"Teachers need to raise awareness of families. I know some educational channels." (A11)

These statements underline that participants consider parents and teachers as the first line of defence in ensuring safe media engagement, highlighting awareness-raising and parental supervision as essential mechanisms.

Some participants emphasised practical digital safety measures such as content restrictions, monitoring tools, and the use of parental filters:

"In order to make YouTube safe, the history in the search section should not be closed; it should be synchronized with our own email address, talking about the content they watch, using the restricted mode, and directing them to safe channels. There should be state control in detecting and preventing bad content and non-pedagogical behaviour." (A8)

"I think that he/she should log in from an adult account, subscribe to quality YouTube channels, have restricted mode always on, not use headphones, have a parental filter, and his/her behaviour should be monitored." (A8)

"To follow his/her viewing history, block age-inappropriate content, and encourage him/her to watch useful videos." (A17)

These quotations show that preservice teachers are aware of the technical features YouTube offers for safer use, but they also acknowledged the limitations of such tools without consistent parental involvement.

Other preservice teachers suggested modelling and joint use as strategies to guide children's engagement with YouTube:

"First of all, I would make him watch the YouTube channels I recommend in order to be a model. I would ask him to watch videos with his family." (A9)

"I know that some teachers make videos about course subjects on YouTube. I even benefit from it myself. I think it is very educational. I know that there is a wealth of educational content. It is necessary to ensure that students are aware of these contents." (A10)

These statements highlight that guiding children towards pedagogically valuable content is not only a preventive measure but also a way to leverage YouTube as an educational tool.

In summary, preservice teachers placed strong emphasis on adult guidance and literacy as prerequisites for safe YouTube use. Their recommendations reflect an understanding that while technological filters and platform features are useful, the most effective safety strategies lie in the active involvement of parents and teachers, complemented by state-level oversight.

FULL TEXTS BOOK

Table 3. Preservice Teachers' Suggestions for Making YouTube Safe

Level	Suggested Measures	Participants
Family	Parental control, monitoring history, blocking adverts, co-viewing, modelling safe channels	A4, A8, A9
Teachers	Raising family awareness, promoting educational channels	A10, A11
Platform use	Restricted mode, adult accounts, synchronisation with parental email, avoiding headphones	A8, A17
State	Regulatory oversight, detection and prevention of harmful/non-pedagogical content	A8, A17

The findings of this study reveal that preservice teachers see YouTube as a powerful but risky platform for primary school students. While children mainly use YouTube for entertainment and gaming, teachers worry about its possible harms, such as health issues, behavioral problems, and addiction. Teachers also stressed that safe, educational use of YouTube needs active parental involvement, teacher guidance, and rules. These views show YouTube's dual role as both a helpful educational tool and a potential source of danger when not controlled. Based on these results, it is important to explore how these perceptions match with existing research and what they imply for using YouTube responsibly and effectively in primary education.

DISCUSSION, CONCLUSION, and RECOMMENDATIONS

Discussion

The findings of this research confirm that preservice teachers perceive YouTube as a double-edged platform: while it has undeniable educational potential, it also exposes children to significant risks. One of the most salient issues raised by participants was the excessive entertainment-oriented use of YouTube by primary school students. Children predominantly watch cartoons, gaming streams, and follow YouTubers, which aligns with earlier studies documenting that digital media platforms increasingly replace traditional play activities with screen-based consumption (Boğa & Sağlam, 2021; Sekmen, 2019). This shift represents not merely a change in leisure habits, but a deeper cultural transformation, where digital spaces are gradually substituting for children's social interactions and creative play (Fide & Sapsağlam, 2020). From a developmental perspective, preservice teachers observed that overexposure to YouTube contributes to behavioural, cognitive, and physical challenges. Their accounts of students imitating "YouTube language" or showing signs of distraction resonate with international findings that prolonged screen time correlates with diminished attention, sleep disturbances, and aggressive behaviours (Chassiakos, Radesky, Christakis, Moreno, & Cross, 2016; Mullan, 2018). These concerns also reflect the Turkish context, where the absence of systematic parental mediation has been shown to exacerbate children's vulnerability to harmful content (Kırık & Altun, 2019). Thus, the preservice teachers' insights not only mirror global concerns but also situate them within local educational realities.

The results further highlight the commercialization of childhood through YouTube. Participants expressed awareness that influencers and child-focused YouTubers embed marketing strategies into their content, blurring the boundaries between authentic entertainment and targeted advertising. This finding aligns with earlier research that documents how YouTube commodifies childhood identities and normalizes consumerist behaviors (Karakoç & Ünlü, 2021; Özcan, 2023). The preservice teachers' observations, therefore, underscore the ethical dilemmas surrounding the platform, particularly the tension between children's rights to safe digital participation and corporations' pursuit of profit. Another

FULL TEXTS BOOK

dimension concerns the role of preservice teachers as mediators. As both consumers of digital media and future educators, they occupy a unique position in shaping safe digital engagement for children. The emphasis they placed on parental supervision and teacher guidance aligns with prior scholarship stressing the importance of parental mediation strategies (Livingstone & Helsper, 2008) and digital literacy interventions in schools (Hobbs, 2010). Their awareness suggests readiness to engage with these challenges, yet it also indicates a need for systematic training within teacher education curricula (Gökce & Bayındır, 2023). Without structured preparation, preservice teachers may lack the pedagogical tools to integrate YouTube meaningfully into classroom practices while mitigating its risks.

This study expands existing research by foregrounding the voices of preservice teachers. Much of the earlier literature has focused on parents (Radesky, Schumacher, & Zuckerman, 2016) or children themselves (Ergen, 2019; Ünlükaya, Savaş, & Sucu, 2021). By capturing preservice teachers' perceptions, this study reveals how future educators conceptualize digital safety responsibilities and situates them as crucial stakeholders in regulating children's media use. Their perspectives highlight a triadic responsibility shared among parents, teachers, and state institutions—a point less emphasized in earlier works.

Conclusion

The findings of this study demonstrate that YouTube has become a dominant platform in the digital experiences of primary school students, shaping both their entertainment practices and, when appropriately guided, their learning opportunities. Preservice teachers consistently recognised that the platform offers a combination of benefits, such as access to diverse audiovisual materials and flexible, on-demand learning, and risks, including overexposure, harmful content, and manipulative commercial messages. These insights underscore the importance of viewing YouTube not as a neutral tool, but as a socio-technical environment that profoundly impacts children's cognitive, social, and emotional development.

Theoretically, this study contributes to the growing body of research on digital childhoods by foregrounding the voices of preservice teachers. Unlike much of the existing literature, which tends to focus either on parental mediation (Livingstone & Helsper, 2008) or on children's self-reported digital practices (Ergen, 2019; Ünlükaya, Savaş, & Sucu, 2021), this research situates preservice teachers as both observers of children's behaviours and as future mediators of digital culture. Their dual positioning enriches the literature by highlighting the pedagogical responsibility and ethical awareness required to transform platforms like YouTube into safe and educationally meaningful tools. Practically, the findings emphasise that safeguarding children in digital spaces is a shared responsibility. Preservice teachers perceive parents and teachers as the first line of defence, tasked with monitoring usage, guiding content selection, and modelling healthy digital habits. However, they also recognise that institutional and governmental mechanisms play an indispensable role in regulating exploitative practices, preventing privacy violations, and ensuring that child-oriented content adheres to ethical standards. This triadic responsibility, encompassing family, school, and state, emerges as a central conclusion of the study, underscoring the need for collaborative and multi-level interventions.

This study highlights the importance of equipping preservice teachers with the skills necessary to critically evaluate digital media and integrate safe YouTube practices into their future classrooms. By incorporating media literacy, ethical reflection, and digital pedagogy into teacher education curricula, institutions can equip a generation of educators to navigate the complexities of digital platforms. In doing so, YouTube can move beyond being a source of risk and become a sustainable educational resource. The conclusion, therefore, extends beyond the immediate context of YouTube to broader discussions of children's rights, media ethics, and the urgent need to foster digital resilience in educational practice.

FULL TEXTS BOOK

Recommendations

The findings of this study highlight the dual role of YouTube as both an educational resource and a potential source of risks for primary school students. To maximize its pedagogical benefits while minimizing potential harms, several recommendations can be made. First, parents should actively guide their children's media use. Instead of relying only on restrictive measures, co-viewing and open discussions about online content can help children critically evaluate what they watch. Parents are also advised to use parental control features, monitor their children's search histories, and demonstrate healthy digital behaviors, as children often imitate the behaviors of adults.

Second, teachers and schools should carefully select and incorporate pedagogically sound YouTube content into both classroom and distance education settings. When aligned with curriculum goals, YouTube can offer flexible and engaging learning opportunities, especially when face-to-face interaction is limited. Teachers should also receive structured training in media literacy to help students assess the reliability and educational value of online resources. Additionally, classroom discussions about digital ethics, consumerism, and online safety can raise awareness and encourage critical thinking among students.

Third, policy and regulation are vital in creating safer digital environments. Enhanced monitoring of child-focused YouTube channels is necessary, especially concerning covert advertising, consumerist content, and child exploitation. Legal systems should be strengthened to safeguard children's privacy and oversee influencer activities. Moreover, collaborations between schools and digital platforms can help ensure the availability of child-safe educational content, including resources specifically intended for remote learning.

Fourth, teacher education programs should incorporate digital literacy, online safety, and ethical awareness into their curricula. Preservice teachers need opportunities to practice integrating YouTube into lesson planning for both classroom and remote learning settings. Reflective practice on their own media use should also be promoted, as it improves their self-awareness as digital role models. In this way, future teachers can develop the skills to guide students toward safe and effective engagement with digital platforms.

Future research should further investigate children's use of YouTube across diverse cultural contexts, examining how practices vary globally. Longitudinal studies are also needed to investigate the long-term cognitive, emotional, and social impacts of YouTube use on children. In addition, studies focusing on the effectiveness of mediation strategies employed by parents and teachers would contribute to the design of more robust protective frameworks. Taken together, these recommendations emphasize that YouTube, when used intentionally in both classroom and remote learning, can improve accessibility and flexibility in education. However, this potential can only be realized if families, teachers, and policymakers share the responsibility of creating a safe and supportive digital learning environment.

Conflict of Interest

The authors report that there is no conflict of interest.

Ethical Approval

Ethical approval for this study was obtained from the Eskişehir Osmangazi University Social and Human Sciences Human Research Ethics Committee at the meeting dated 16 November 2021 (Meeting No: 2021-18).

Endnote

This study was orally presented online at the 8th International Mediterranean Scientific Research Congress (13–15 August 2025, Antalya, Türkiye).

FULL TEXTS BOOK

REFERENCES

Aydın, E. B. (2023). Kuşakların teknoloji kullanımının mahremiyet kapsamında karşılaştırılması: Dijital melezler ve dijital yerliler. *Gümüşhane Üniversitesi İletişim Fakültesi Elektronik Dergisi, 11*(2), 1679–1702.

Avcı, F. (2020). Öğretmenlerin sosyal medya kullanımları ve sosyal medya kavramına ilişkin metaforik algıları. *Uluslararası Kültürel ve Sosyal Araştırmalar Dergisi (UKSAD), 6*(2), 769–788.

Baltacı, A. (2019). Nitel araştırma süreci: Nitel bir araştırma nasıl yapılır? *Ahi Evran Üniversitesi Sosyal Bilimler Enstitüsü Dergisi, 5*(2), 368–388.

Boğa, E., & Sağlam, M. (2021). Dijital ortamda çocuk: Bir çocuk YouTuber kanalının içeriksel analizi. *Erciyes İletişim Dergisi, 8*(2), 643–664.

Bozkurt, İ., & Karakuş, E. (2015). İlköğretim öğrencilerinin sosyal medya kullanım alışkanlıkları üzerine bir araştırma: Elazığ örneği. *Türkiyat Araştırmaları Dergisi, 37*, 437–455.

Braun, V., & Clarke, V. (2019). Reflecting on reflexive thematic analysis. *Qualitative Research in Sport, Exercise and Health, 11*(4), 589–597. https://doi.org/10.1080/2159676X.2019.1628806

Breslyn, W., & Green, A. E. (2022). Learning science with YouTube videos and the impacts of Covid-19. *Disciplinary and Interdisciplinary Science Education Research*, 4(1), 13. https://doi.org/10.1186/s43031-022-00051-4

Burgess, J., & Green, J. (2018). *YouTube: Online video and participatory culture* (2nd ed.). Polity Press.

Carey, J. W. (2005). Historical pragmatism and the Internet. *New Media & Society, 7*(4), 443–455. https://doi.org/10.1177/1461444805054107

Chassiakos, Y. L., Radesky, J., Christakis, D., Moreno, M. A., & Cross, C. (2016). Children and adolescents and digital media. *Pediatrics, 138*(5), e20162593. https://doi.org/10.1542/peds.2016-2593

Çalapkulu, Ç., & Alp, F. (2020). Dijital ebeveynler ile çocukların sosyal medya kullanımı üzerindeki mahremiyet ilişkisi. *Nişantaşı Üniversitesi Sosyal Bilimler Dergisi, 2*(8), 132–144.

Çemrek, F., Baykuş, H., & Özaydın, Ö. (2014). Sosyal medya kullanım ve davranışlarının kullanımlar ve doyumlar yaklaşımı bağlamında incelenmesi: Eskişehir Osmangazi Üniversitesi örneği. *Alphanumeric Journal, 2*(2), 61–76.

Cohen-Almagor, R. (2011). Internet history. *International Journal of Technoethics, 2*(2), 45–64. https://doi.org/10.4018/jte.2011040104

Creswell, J. W., & Poth, C. N. (2018). *Qualitative inquiry and research design: Choosing among five approaches* (4th ed.). SAGE Publications.

Ergen, Y. (2019). Popüler kültürün popüler rol modelleri YouTuberlar: İlköğretim çağındaki özel okul öğrencileri üzerine bir araştırma. *İnsan ve Toplum Dergisi, 9*(1), 117–153.

Ersoy, A. F. (2016). Fenomenoloji. In A. Saban & A. F. Ersoy (Eds.), *Eğitimde nitel araştırma desenleri* (pp. 51–110). Anı Yayıncılık.

Etikan, I., Musa, S. A., & Alkassim, R. S. (2016). Comparison of convenience sampling and purposive sampling. *American Journal of Theoretical and Applied Statistics, 5*(1), 1–4. https://doi.org/10.11648/j.ajtas.20160501.11

Fide, E. G., & Sapsağlam, Ö. (2020). YouTube yayıncılarının çocukların kimlik inşa süreçlerine yansıması. *Türkiye Bilimsel Araştırmalar Dergisi, 5*(2), 98–109.

Guest, G., Bunce, A., & Johnson, L. (2006). How many interviews are enough? An experiment with data saturation and variability. *Field Methods, 18*(1), 59–82. https://doi.org/10.1177/1525822X05279903

FULL TEXTS BOOK

Gökce, M. (2023). Öğretmen adaylarının çocukların dijital istismara karşı korunmasına ilişkin adli ve idari yönden görüşleri. *Dijital Teknolojiler ve Eğitim Dergisi, 2*(2), 71–81. https://doi.org/10.5281/zenodo.10439817

Gökce, M., & Bayındır, N. (2023). İlkokul öğrencilerini dijital risklerden korumaya yönelik önlemler. *ICITS 2023 Conference Paper*.

Güney, Z. (2020). Çocukların sosyal medya kullanımı üzerine bir araştırma. *Gümüşhane Üniversitesi Sosyal Bilimler Enstitüsü Elektronik Dergisi, 11*(1), 188–199.

Hobbs, R. (2010). *Digital and media literacy: Connecting culture and classroom*. Corwin Press.

Hutchby, I. (2001). Technologies, texts and affordances. *Sociology, 35*(2), 441–456. https://doi.org/10.1177/S0038038501000219

İlhan, E., & Aydoğdu, G. A. (2019). YouTube kullanıcılarının kullanım motivasyonlarının incelenmesi. *Gümüşhane Üniversitesi İletişim Fakültesi Elektronik Dergisi, 7*(2), 1130–1153.

Karakoç, E., & Ünlü, T. T. (2021). Oyun mu, iş mi? YouTube kidfluencerları üzerine karşılaştırmalı bir analiz. *TRT Akademi Dergisi, 6*(12), 468–493.

Kavuran, T., & Yeşil, A. (2019). İnteraktif medyada YouTuberların iletişim dili: Enes Batur oyun videoları örneği. *Uluslararası Sosyal Bilimler Dergisi, 2*(1), 45–62.

Kılıç, B., & Kaya, S. (2021). Yeni medya çağında çocuk YouTuberlar üzerinden çocukluğun yok oluşu. *Gümüşhane Üniversitesi Sosyal Bilimler Enstitüsü Elektronik Dergisi, 12*(3), 1141–1156.

Kırık, M. A., & Altun, E. (2019). Yeni medya ve Z kuşağı ilişkisi bağlamında YouTube Kids uygulamasının içeriksel analizi. *Anemon Muş Alparslan Üniversitesi Sosyal Bilimler Dergisi, 7*(6), 109–119.

Kızılkaya, Z. Z. (2020). Sosyal medya kullanıcılarının metalaşması: YouTuberler üzerine bir inceleme. *Türkiye İletişim Araştırmaları Dergisi, 35*, 63–69.

Konuk, N., & Güntaş, S. (2019). Sosyal medya kullanımı eğitimi ve bir eğitim aracı olarak sosyal medya kullanımı. *International Journal of Entrepreneurship and Management Inquiries (Journal EMI), 3*(4), 1–25.

Krippendorff, K. (2018). *Content analysis: An introduction to its methodology* (4th ed.). SAGE Publications.

Kulular, M. A. (2022). Sosyal medyaya yönelik yeni düzenlemelerin hukuken değerlendirilmesi. *Bilişim Hukuku Dergisi, 4*(1), 1–32. https://doi.org/10.55009/bilisimhukukudergisi.1114305

Lester, J. N., Cho, Y., & Lochmiller, C. R. (2020). Learning to do qualitative data analysis: A starting point. *Human Resource Development Review, 19*(1), 94–106. https://doi.org/10.1177/1534484320903890

Lincoln, Y. S., & Guba, E. G. (1985). *Naturalistic inquiry*. SAGE Publications.

Livingstone, S., & Haddon, L. (2009). *EU Kids Online: Final report*. London School of Economics and Political Science.

Livingstone, S., & Helsper, E. J. (2008). Parental mediation of children's Internet use. *Journal of Broadcasting & Electronic Media, 52*(4), 581–599. https://doi.org/10.1080/08838150802437396

Mullan, K. (2018). Technology and children's screen-based activities in the UK: The story of the millennium so far. *Child Indicators Research, 11*(6), 1781–1800. https://doi.org/10.1007/s12187-017-9509-0

Özcan, G. (2023). Çocuk hakları ihlali ve çocuk istismarı açısından çocuk YouTube kanalları. *Disiplinlerarası Çocuk Hakları Araştırması Dergisi, 3*(5), 64–72.

Özel, Ö., & Altınkaynak, H. (2023). Examination of YouTube content in terms of potential behaviors that preschool children can take as an example: The example of Enes Batur. *Journal of Hasan Ali Yücel Faculty of Education, 20*(1), 123–137.

Palme, J. (2011). Before the Internet: Early experiences of computer mediated communication. In J. Impagliazzo, P. Lundin, & B. Wangler (Eds.), *History of Nordic Computing 3* (pp. 271–277). Springer. https://doi.org/10.1007/978-3-642-23315-9_30

Radesky, J. S., Schumacher, J., & Zuckerman, B. (2016). Mobile and interactive media use by young children: The good, the bad, and the unknown. *Pediatrics, 135*(1), 1–3. https://doi.org/10.1542/peds.2014-2251

Schafer, V., & Thierry, B. (2018). The 90s as a turning decade for Internet and the Web [Special issue editorial]. *Internet Histories: Digital Technology, Culture and Society, 2*(3–4), 1–5. https://doi.org/10.1080/24701475.2018.1521060

Sekmen, M. (2019). Çocuk YouTuberlarda sorunlu bir alan: Tüketim ile mahremiyetin yitimi. *Manas Sosyal Araştırmalar Dergisi, 8*(4), 3566–3584.

Sgier, L. (2020). Qualitative data analysis. *An Initiative of Gebert Ruf Stiftung, 27*(1), 21–24.

Statista. (2025). Leading social media platforms worldwide as of February 2025. Retrieved from https://www.statista.com/statistics/272014/global-social-networks-ranked-by-number-of-users/

Suler, J. (2004). The online disinhibition effect. *CyberPsychology & Behavior, 7*(3), 321–326. https://doi.org/10.1089/1094931041291295

Ünal, T. A. (2019). YouTube kullanıcısı Z kuşağının sosyal medya kullanım alışkanlıklarının ve tercihlerinin incelenmesi. *İletişim Çalışmaları Dergisi, 16*, 61–85.

Ünlükaya, A., Savaş, S., & Sucu, İ. (2021). YouTube çocuk kanallarında bulunan video içeriklerindeki temaların ortaya koyulmasına yönelik bir çalışma. *Journal of Business in the Digital Age, 4*(1), 33–49.

Üstündağ, A. (2020). Çocukların sosyal medya kullanım durumları ve sosyal medyanın çocukların duygu durumları üzerine etkisi. *Uluslararası Güncel Eğitim Araştırmaları Dergisi (UGEAD), 6*(2), 286–302.

Vas, S. (1997). The use of the Internet. *Peritoneal Dialysis International, 17*(2 Suppl), 138–141. https://doi.org/10.1177/089686089701702S27

Yegen, C. (2013). Demokratik ve yeni bir kamusal alan olarak sosyal medya. *Muş Alparslan Üniversitesi Sosyal Bilimler Dergisi, 1*(2), 119–135.

Yıldırım, A., & Şimşek, H. (2016). *Sosyal bilimlerde nitel araştırma yöntemleri*. Seçkin Yayıncılık.

Zinderen, A. (2021). YouTube ortamındaki haberciliği haritalandırmak: Gazetecilerin YouTube haber kanalları üzerine tanımlayıcı bir çalışma. *Atatürk Üniversitesi Sosyal Bilimler Enstitüsü Dergisi, 25*(3), 932–950.

FULL TEXTS BOOK

THE ROLE OF TiO_2 NANOPARTICLES IN IMPROVING THE OPTICAL AND ELECTRICAL CHARACTERISTICS OF POLYMER NANOCOMPOSITES FOR LED DEVICES

Andreea Irina BARZIC [1]

[1] *"Petru Poni" Institute of Macromolecular Chemistry, Iasi, Romania*

[1]*ORCID ID: https://orcid.org/0000-0001-7131-5120*

Vasile Bogdan MARDARI [2]

[2] *"Alexandru Ioan Cuza" University, Faculty of Chemistry, Iasi, Romania*

[2]*ORCID ID: -*

Raluca Marinica ALBU[1]

[1] *"Petru Poni" Institute of Macromolecular Chemistry, Iasi, Romania*

[1]*ORCID ID: https://orcid.org/0000-0002-7707-9613*

ÖZET

Kontrollü kırılma özelliğine sahip malzemeler, optik fiberler, güneş pilleri, ekranlar, optik lensler ve ışık yayan diyotlar (LED'ler) gibi birçok alanda büyük bir uygulama potansiyeli sunar. Son uygulamada, kapsülleme tabakasının tasarımına yönelik birçok bilimsel çalışma yapılmıştır. Mevcut endüstriyel süreç, bu LED bileşeninin silisyum dioksit veya akrilatlara dayalı üretimine dayanmaktadır. Ancak, bu malzemelerin küçük kırılma indisinin, LED çip üretimi için kullanılan yarı iletkenlerin (örneğin III-V fosfürler, arsenitler ve nitrürler) çoğunluğuyla yüksek bir tezat oluşturduğu bulunmuştur. Bu ortamların kırılma özelliklerindeki fark, toplam iç yansıma nedeniyle optik radyasyonların kaçış konisinin sınırlamasını belirler ve ışığı yarı iletken tabaka içinde hapseder. Bu çalışma, değişken miktarlarda TiO_2 nanopartiküllerinin yerleştirildiği şeffaf bir polimere (yani selüloz eter) dayalı yeni LED kapsülleme malzemeleri tasarlayarak bu teknik sorunu hafifletmek için bir çözüm önermektedir. Her numunenin kırılma indisi değeri, üç dalga boyunda ve oda sıcaklığında Abbe refraktometresi ile kaydedilmiştir. n büyüklüğünün, eklenen TiO_2 dolgu miktarıyla ayarlanabileceği bulunmuştur. Ayrıca, elektrik kapasitesi bir LCR metre ile ölçülmüş ve numunenin geometrisine bağlı olarak 1 kHz'deki dielektrik sabiti değerlendirilmiştir. Sonuçlar, izlenen pratik kapsamla bağlantılı olarak tartışılmıştır.

Anahtar Kelimeler: selüloz eter; inorganik nanopartiküller; kırılma indisi; dielektrik sabiti.

ABSTRACT

Materials with controlled refractivity present a huge applicative potential in many domains, such as optical fibers, solar cells, displays, optical lens, and light emitting diodes (LEDs). In the case of the last application, many scientific efforts have been directed towards to design of the encapsulation layer. Current industrial process relies on the fabrication of this LED component based on silicon dioxide or acrylates. However, the small refractive index of these materials is found to be in high contrast with that of the majority of semiconductors (e.g. III-V phosphides, arsenides, and nitrides) employed for the LED chip manufacturing. The difference in the refraction characteristics of these media determines the limitation of the escape cone of optical radiations due to total internal reflection, trapping the light within the semiconducting layer. This work proposes a solution for alleviating this technical problem by designing novel LED encapsulants based on a transparent polymer (i.e. cellulose ether) in which variable amounts of TiO_2 nanoparticules are inserted. The refractive index value of each specimen is recorded

FULL TEXTS BOOK

with Abbe refractometer, at three wavelengths, and at room temperature. It is found that n magnitude can be tuned by the amount of incorporated TiO_2 filler. Additionally, the electrical capacitance is measured with a LCR meter and based on the sample's geometry, the dielectric constant at 1 kHz is evaluated. The results are discussed in relation to the pursued practical scope.

Keywords: cellulose ether; inorganic nanoparticles; refractive index; dielectric constant.

INTRODUCTION

In our daily life, an extraordinary and continuous modernization of the available technologies is observed, and this is possible due to the discovery of novel materials and elaboration of ingenious design approaches of the products (Gangele, 2021). Current society needs require the help of the scientific community to find solutions in energy sector that diminish the global pollution or the huge consumption of electricity. In this context, public lighting was gradually upgraded by introducing two decades ago, the light emitting diodes (LEDs) as alternative for the conventional incandescent bulbs (Cheng, 2006). LEDs have brought many advantages in comparison to the old technology, such as lower energy consumption, a long lifespan, advanced color rendering, eco-compatibility, and design flexibility. Nowadays, researchers are focused on further improving LEDs by extending the light output efficiency (Pattison, 2018). A problem here could be linked to the optical performance of the encapsulation material, which is fabricated from polymers, such as epoxy resins, polyurethanes, or other synthetic macromolecular compounds (Huang, 2007). Polymers represent the dominant category of engineering materials owing to their appealing balance of physical properties, including lightweight, mechanical resistance, good insulation, and capacitance of being processed in many shapes (Jabri, 2023). The limitation of the polymers has its roots in their small refractive index that contrasts the refraction features of the most semiconductors used in LED technology (Ma, 2011). This discrepancy is responsible for the production of optical losses generated by total internal reflection (TIR) at encapsulant/semiconductor interface.

On the other hand, in the actual context of plastic pollution, it is very beneficial to focus on tailoring the performance of the biodegradable polymers. Here, cellulose and its derivatives (particularly ether counterparts) seem to be a great choice, considering their bioavailability, flexibility, optical clarity, and low permittivity. Among the cellulose ethers, a particular interest is attributed to hydroxyethyl cellulose (HEC), which has many convenient key attributes, such as flexibility, optical clarity, and biodegradability (Noreen, 2020). A simple strategy to refine the refractive index of this compound in order to match the criteria imposed by LED encapsulation would be to insert in the polymer precise amounts of micro- or nanofillers that display complementary refraction properties, like TiO_2 (Ma, 2011). As far as authors know, there are no studies concerning the design of HEC encapsulants with the purpose to reduce the optical losses in LEDs.

This work presents the obtaining steps of HEC and HEC/TiO_2 samples, which are characterized from optical and electrical points of view to inspect their suitability for LED encapsulation.

METHODS

Sample preparation

The polymer powder utilized in this study is commercial and it is taken from Hercules Company. According to the provider, HEC is characterized by a molecular weight of 720 000 Da.

The inorganic particle nanopowder of Titanium(IV) oxide (TiO_2) is acquired from Sigma Aldrich. According to the provider, the dimension of the nanofillers is under 100 nm.

Obtaining of the composite films required the pursue of the following steps: (a) weighing the HEC powder and solubilization in water, (b) incorporation of the TiO_2 nanoparticles (in percentages of 1% and 2%), (c) homogenization of the systems by ultrasonication, (d) deposition of the systems on glass slides, (e) drying the specimens in an oven at mild temperature, and (f) detachment of the films from the supports.

Characterization

The dimensions of the achieved films were determined with a digital micrometer and the measurements indicated a specimen thickness of 35-38 µm.

The optical transmittance of the films was attained on a Specord 210 PLUS spectrophotometer in the domain of 190–850 nm.

The refractive index of the films was registered on DR-M4 apparatus at three wavelengths (486 nm, 589 nm, and 670 nm), at 298 K.

Electrical investigations of the films were done on a LCR meter at the electrical field frequency of 1 kHz.

RESULTS

Optical performance of the prepared HEC and HEC/TiO_2 films was tested in terms of transmittance and refractive index. The UV-VIS spectral data of all specimens are presented in Figure 1(a-c). In the absence of the inorganic phase, HEC layer is found to be highly transparent, so (as noted in Figure 1(a)) the transmittance reaches 78% at 600 nm. When adding 1% nanofiller in the system, the level of transparency is diminished to 50% at 600 nm, as illustrated in Figure 1(b). Further insertion of the titanium dioxide in HEC up to 2% seems to dramatically deteriorate the optical transmittance values, which became smaller than 1% in the visible range (Figure 1(c)).




Figure 1. UV-VIS data recorded for (a) HEC, (b) HEC/TiO_2 1%, and (c) HEC/TiO_2 2% samples.

The light dispersion properties of the specimens are evaluated by refractometry experiments. The refractive index (n) is of great importance when attempting to examine the light propagation in media of distinct refraction features. This optical parameter is useful for understanding how visible radiations are passing through various materials in terms of speed (compared to its speed in a vacuum) and bending degree. As observed in Figure 2(a), the refractive index of HEC and HEC/TiO_2 films is slowly decreasing with augmenting the wavelength. The pure biopolymer is characterized by a relatively small refractive index of 1.499, determined at 589 nm. At the same wavelength, one may see that the

FULL TEXTS BOOK

incorporation of the inorganic nanoparticles produces the increase of n value up to 1.503 (composite with 1 % TiO_2) and 1.509 (composite with 2 % TiO_2). The enhancement of n parameter, caused by HEC loading, is remarked at all measuring wavelengths. The explanation for the modification of refraction behavior of the specimens is given by the Lorenz-Lorentz formula, which connects macroscopic properties (like n) with the microscopic features of its constituent molecules (like polarizability). Hence, the highly polarizable TiO_2 nanofiller generates formation of tiny domains where the polarizability is increased, and this is reflected in higher magnitude of n parameter for the composite materials.




Figure 2. Data of (a) light dispersion and (b) band gap energy for the pure HEC and HEC/TiO_2 composites with 1% and 2% filler.

The refractive index is linked to other important characteristics of the material, such as optical band gap energy (E_g), as proved by Gupta (Gupta, 1980) through the equation (1):

$$n = 4.084 - 0.62 \cdot E_g \quad (1)$$

The data from Figure 2(b) highlight a decrease of the E_g values as the quantity of TiO_2 nanoparticles in the system is increasing. Polymer loading generally is associated with altering of the electronic structure or with occurrence of novel levels within the forbidden gap, therefore the material capacity of light transmission is modified. This was observed also for other polymer composites (Morsi, 2019).

On the other hand, the refractive index can provide valuable information on the dielectric constant at optical frequencies (noted $\varepsilon_{electronic}$), where only electronic polarization takes place, as seen in the equation (2) (Zhao, 2012):

$$\varepsilon_{electronic} = n^2 \quad (2)$$

Optical dielectric constant results are listed in Table 1. The pure biopolymer has the lowest value of $\varepsilon_{electronic}$ due to its lower polaizability. Gradual addition of the titanium dioxide nanofiller is reflected in a higher magnitude of the $\varepsilon_{electronic}$ since this inorganic compound displays higher permittivity itself and is capable of interacting with the matrix via hydrogen bonding.

The electrical characterization of the film samples was undertaken to extract information on the dielectric constant at lower frequencies ($\varepsilon_{dipolar}$), where the mechanism of dipolar polarization is also acting (Zhao, 2012). According to the equation (3), the magnitude of $\varepsilon_{dipolar}$ is linked to the capacitance and the sample's geometry:

$$\varepsilon_{dipolar} = \frac{C \cdot t}{\varepsilon_0 \cdot S} \quad (3)$$

FULL TEXTS BOOK

where C is the capacitance, t is the thickness of the specimen filling the capacitor plates, ε_0 denotes the permittivity of free space, and S is the area of the capacitor plates.

The measured values of the electric capacitance, along with the electrode area, are included in Table 1. One may notice that the incorporation of the nanofillers produces significant changes in the electrical parameters. Both electrical capacitance and dielectric constant (describing the dipolar polarization) are augmented after HEC reinforcement with TiO_2. At the highest loading degree, the magnitude of $\varepsilon_{dipolar}$ is almost doubled in comparison to that of the pure biopolymer.

Table 1. The dimension and electrical parameters of the studied samples.

Content of nanofiller, %	S, μm^2	C, nF	$\varepsilon_{electronic}$	$\varepsilon_{dipolar}$
0	153.86	0.090	2.247	2.576
1	153.86	0.134	2.259	3.443
2	153.86	0.172	2.275	4.419

Optical losses caused by the TIR phenomenon are evaluated for a LED, where the semiconducting material is differing for comparison reasons. The selected semiconductors have optical properties reported in literature, namely AlGaAs (n=4.2 at 486 nm) and AlInP (n=3.4 at 586 nm) (Aspnes, 1986; Ochoa-Martínez, 2018). When the incident beam travels from the semiconductor layer to the encapsulant polymer layer, the intensity reflection coefficient (R) at normal incidence is described by equation (4):

$$R = \left(\frac{n_c - n_s}{n_c + n_s} \right)^2 \quad (4)$$

where n_c is the refractive index of the composite, and n_s is the refractive index of the semiconducting material.

The estimated results for the intensity reflection coefficient at the specimen interface with the chosen semiconductors are presented in Figure 3. It is possible to discern that in the case of AlGaAs, the amount of radiations that return in the incident medium is greater due to higher differences in the refraction characteristics of the semiconductor and the studied encapsulants. When the refractive index of the prepared materials is closer to that of the semiconductor (like AlInP), the magnitude of R is smaller. This indicates that there is a decrease in the fraction of incident light reflected in the semiconducting medium when the beam encounters the boundary with the encapsulant. Hence, the prepared HEC samples are more suitable to reduce the optical losses in LED containing AlInP as semiconductor. Hence, it is achieved the improvement of the device efficacy in terms of light output.

FULL TEXTS BOOK




Figure 3. Data of the intensity reflection coefficient determined at the specimen interface with AlGaAs or AlInP semiconducting layers.

CONCLUSIONS

This paper described the preparation of a set of composites derived from the chosen cellulose ether, which was doped with variable contents (1% and 2%) of nanoparticles of titanium dioxide. The transmittance level is gradually lowered after HEC loading, namely 78% (0% TiO_2), 50% (1% TiO_2), and 0.2% (2% TiO_2), according to the measurements performed at 600 nm. The refractive index is improved upon the introduction of the nanofillers, while the band gap decreases from 4.15 eV for HEC to about 4.13 eV for HEC/TiO_2 2%. Additionally, the dielectric constant is enhanced at 10^{14} Hz and at 10^3 Hz, where electronic and dipolar mechanisms of polarization take place, respectively. Reflection losses are lowered at the specimen/semiconductor interface as a result of better matching of the refraction properties and this positively contributes to the LED efficiency in terms of light output.

REFERENCES

Aspnes, D. E., Kelso, S. M., Logan, R. A., & Bhat, R. (1986). Optical properties of AlxGa1−xAs. Journal of Applied Physics, 60(2), 754–767.

Cheng, Y. K., & Cheng, K. W. E. (2006). General study for using LED to replace traditional lighting devices. 2nd International Conference on Power Electronics Systems and Applications, doi:10.1109/pesa.2006.343093, 173–177.

Gangele, A., & Mishra, A. (2021). A review on smart materials, types and modelling: need of the modern era. Materials Today Proceedings, 47(18), 6469–6474.

Gupta, V. P. & Ravindra, N. M. (1980). Comments on the Moss formula. Physica Status Solidi B, 100, 715–719.

Huang, W., Zhang, Y., Yu, Y., & Yuan, Y. (2007). Studies on UV-stable silicone-epoxy resins. Journal of Applied Polymer Science, 104, 3954–3959.

Jabri, F.E., Ouballouch, A., Lasri, L., & El Alaiji, R. (2023). A comprehensive review of polymer materials and selective laser sintering technology for 3D printing. Journal of Achievements in Materials and Manufacturing Engineering, 118(1), 5–17.

Ma, M., Mont, F. W., Yan, X., Cho, J., Schubert, E. F., Kim, G. B., & Sone, C. (2011). Effects of the refractive index of the encapsulant on the light-extraction efficiency of light-emitting diodes. Optics Express, 19(S5), A1135.

FULL TEXTS BOOK

Morsi, M.A., Abdelaziz, M., Oraby, A.H., Mokhles, I. (2019). Structural, optical, thermal, and dielectric properties of polyethylene oxide/carboxymethyl cellulose blend filled with barium titanate. Journal of Physical Chemistry of Solids, 125, 103–114.

Noreen, A., Zia, K. M., Tabasum, S., Khalid, S., & Shareef, R. (2020). A review on grafting of hydroxyethylcellulose for versatile applications. International Journal of Biological Macromolecules, 150, 289–303.

Ochoa-Martínez, E., Barrutia, L., Ochoa, M., Barrigón, E., García, I., Rey-Stolle, et al. (2018). Refractive indexes and extinction coefficients of n- and p-type doped GaInP, AlInP and AlGaInP for multijunction solar cells. Solar Energy Materials and Solar Cells, 174, 388–396.

Pattison, P.M., Hansen, M., & Tsao, J. Y. (2018). LED lighting efficacy: Status and directions. Comptes Rendus Physique, 19(3), 134–145.

Zhao, L., Su, J.-C., Pan, Y.-F., & Zhang, X.-B. (2012). The effect of polymer type on electric breakdown strength on a nanosecond time scale. Chinese Physics B, 21(3), 033102.

RADIATION-INDUCED SKIN TOXICITIES IN HEAD AND NECK CANCERS: CLINICAL SIGNIFICANCE AND MANAGEMENT APPROACHES

***Timur KOCA*[1]**

[1]Associate Professor, Akdeniz University Faculty of Medicine, Department of Radiation Oncology, Antalya, Türkiye[1]

[1]ORCID ID: https://orcid.org/0000-0003-0318-6122

***Sümeyye ÇITAK*[2]**

[2]Resident Doctor, Akdeniz University Faculty of Medicine, Department of Radiation Oncology, Antalya, Türkiye

[2]ORCID ID: https://orcid.org/0009-0003-0889-9030

***Aylin Fidan KORCUM*[3]**

[3]Professor, Akdeniz University Faculty of Medicine, Department of Radiation Oncology, Antalya, Türkiye

[3]ORCID ID: https://orcid.org/0000-0002-7400-1837

ABSTRACT

The incidence of head and neck cancers is steadily rising. Radiotherapy is a key treatment option for these diseases. Although radiotherapy is effective both as a definitive and adjuvant treatment, radiation-induced skin toxicities pose a significant clinical challenge. Among these, radiation dermatitis is the most common and can negatively affect patients' quality of life and lead to treatment interruptions. Radiation dermatitis is classified into two main types—acute and chronic—based on when they occur. This review summarizes the pathophysiology, clinical features, and management strategies of radiation dermatitis following radiotherapy for head and neck cancers, based on current literature.

In our study, we reviewed the current literature on radiation-induced skin toxicities in head and neck cancers. A thorough search was performed using specific keywords in the PubMed and Google Scholar databases, focusing on clinical studies, review articles, and clinical guidelines.

Acute radiation dermatitis usually begins with erythema around days 10–14 of treatment and progresses to include signs such as damage to the epidermis, increased water loss through the skin, dryness, peeling, hyperpigmentation, itching, burning, and pain. These symptoms may require treatment interruptions and could potentially lessen the effectiveness of the therapy.

Chronic radiation dermatitis, on the other hand, is usually permanent and progressive. It can clinically present with telangiectasia, poikiloderma, dermal atrophy, scarring alopecia, fibrosis, lymphedema, and limited mobility.

The pathogenesis of radiation-induced toxicity involves multiple factors, including radiation-induced DNA damage, activation of cell death pathways, inflammatory responses, and individual genetic susceptibility. Notably, variations in DNA repair mechanisms among individuals are believed to affect the risk of developing toxicity.

Although there is no universally accepted standard in the literature regarding management strategies, several key factors are emphasized in treatment planning, including skin preservation, limiting the irradiated area and total dose, adaptive planning, and the use of systemic therapies. For skin care and hygiene, using soap-free cleansers (syndets), moisturizers, sunscreen, quitting smoking, and managing weight are recommended. Vitamin D is thought to support the maintenance of the skin barrier's integrity,

FULL TEXTS BOOK

whereas vitamin A may contribute to the repair of skin damage through its role in tissue regeneration. When needed, short-term use of topical corticosteroids can be considered.

In the chronic phase, supportive treatments like laser therapy, physiotherapy, autologous fat injection, and wound care protocols become increasingly important.

Radiation-induced dermatitis in head and neck cancers is a major cause of morbidity during both the acute and chronic stages. Many patient-related and treatment-related factors affect the development of this toxicity. Therefore, protecting the skin should be a priority in treatment planning, and early diagnosis along with supportive skin care measures, should be put into practice. In the future, discovering predictive biomarkers could allow for more effective toxicity management through personalized treatment planning.

Keywords: Radiation dermatitis, Head and neck cancers, Radiotherapy, Skin toxicity.

Introduction

In locally advanced, unresectable head and neck cancers, the main therapeutic modalities are surgery and radiotherapy (RT), which can be administered either as monotherapy or in combination with chemotherapy.[1] One of the most common adverse effects of radiotherapy is cutaneous reactions, which, according to the literature, develop in approximately 95% of patients during the course of treatment.[2] These toxicities are assessed using the Common Toxicity Criteria for Adverse Events (CTCAE) or the Radiation Therapy Oncology Group (RTOG) scales. Clinically, while some patients experience only mild symptoms, severe cutaneous toxicity has been reported in approximately 25% of cases.[3] Such severe reactions may lead to treatment interruptions, thereby negatively affecting therapeutic efficacy.

Radiation dermatitis arises from the complex interplay of patient-related factors—including body mass index (BMI), smoking status, nutritional condition, pre-existing dermatological disorders, and genetic susceptibility—and treatment-related parameters such as total and per-fraction radiation dose, irradiation technique, treatment field size and localization, as well as the use of concurrent chemotherapy.[2,4]

Following radiotherapy, within the first 24 hours, free radicals and reactive oxygen species are generated in the rapidly proliferating cells of the basal layer and dermis. This results in depletion of stem cell reserves, endothelial cell damage, and initiation of an inflammatory response.[5] During weeks 2–4 of treatment, persistent erythema, localized edema, and leukocyte infiltration typically develop. The damage progresses cumulatively throughout the course of therapy, delaying healing and persisting for up to four weeks after treatment completion. Late cutaneous effects include vascular alterations, telangiectasia, atrophy, and hyperpigmentation.[6,7]

Radiotherapy in Head and Neck Cancers

In head and neck cancers, surgery, radiotherapy, and systemic therapy represent the cornerstone treatment modalities. Radiotherapy, either alone or in combination with other approaches, remains one of the most important therapeutic options. It may be delivered in definitive, adjuvant, or palliative settings. In the adjuvant context, radiotherapy or chemoradiotherapy eradicates microscopic disease, thereby reducing the risk of recurrence and prolonging survival.[8] For patients who are not surgical candidates, radiotherapy or concurrent chemoradiotherapy constitutes the primary treatment modality for both curative and palliative purposes.[9] In the palliative setting, radiotherapy alleviates tumor-related pain, bleeding, and obstruction, thereby improving patients' quality of life.

FULL TEXTS BOOK

Pathophysiology of Radiation Dermatitis

The cellular and molecular mechanisms underlying radiation-induced skin injury have not yet been fully elucidated.[10] Radiation triggers various pathways of cell death in skin cells, primarily through DNA damage; this process may result in cell-cycle arrest and the activation of inflammatory responses.[11] Thus, DNA damage is considered the principal trigger of radiation-induced toxicity.[12] In addition, cytokine release is thought to induce biological responses in multiple cell types, thereby contributing to the progression of late toxicity.[13]

Clinical Presentation: Acute Radiation Dermatitis

Acute radiation dermatitis refers to skin injury observed within 90 days of the initiation of radiotherapy.[14] Alterations in epidermal integrity and increased transepidermal water loss, particularly as a result of cumulative dosing from consecutive radiation fractions, lead to a wide spectrum of manifestations, including erythema, xerosis, desquamation, hyperpigmentation, and other subjective symptoms.[14,15]

The earliest clinical sign is erythema, which typically appears around 10–14 days after the onset of treatment in patients receiving doses of 6–20 Gy.[38] Erythematous lesions are often accompanied by edema and increased skin fragility. Patients may experience discomfort characterized by skin tightness, pruritus, burning sensations, or, in some cases, pain. Moreover, sebaceous gland loss contributes to skin dryness.

As a compensatory response to tissue damage, mitotic activity increases after 3–4 weeks of irradiation, which may result in the production of aberrant cells. This manifests clinically as desquamation, commonly observed in patients exposed to doses exceeding 20 Gy. Doses greater than 30 Gy may exceed the reparative capacity of the epidermis, leading to epidermal separation and moist desquamation with blister formation. The resulting disruption of the skin barrier function also increases susceptibility to infections, predominantly bacterial. In severe cases, ulceration and even necrosis of irradiated tissue may occur. Depending on the site of irradiation, mucositis and alopecia may also be observed. Nonetheless, in the majority of cases, these reactions are self-limiting and resolve within 2–4 weeks following the completion of therapy.[14,15]

Chronic Radiation Dermatitis

Chronic radiation dermatitis may occur any time from 90 days after the completion of radiotherapy up to 20–30 years post-treatment. These changes may persist after the resolution of the acute phase or emerge de novo. Unlike acute radiation dermatitis, late toxicity is permanent and often progressive. It may present with various manifestations, including subjective symptoms (hypersensitivity and/or pruritus), vascular changes (telangiectasia), dermal atrophy and fragility, pigmentation changes (poikiloderma), and cicatricial alopecia. Although rare, delayed necrosis may also occur, most commonly affecting the nose, ears, and scalp.[14,15]

For the evaluation of chronic radiation-induced skin injury, several toxicity and skin damage scoring systems, such as Late Effects of Normal Tissues – Subjective, Objective, Management, Analytic (LENT-SOMA), RTOG, and Dische, have been proposed. Among these, the LENT-SOMA classification is the most frequently applied and widely recommended in clinical practice.[16]

Risk factors for radiation-induced skin toxicity are generally categorized into treatment-related and patient-related variables. Treatment-related factors include the total dose delivered to the skin, irradiated skin volume, fractionation scheme, radiotherapy technique, overall treatment quality (including adaptive RT), and concurrent chemotherapy or molecular therapies. Patient-related factors encompass individual

FULL TEXTS BOOK

radiosensitivity, extent of sun exposure, smoking status, nutritional condition and obesity, comorbidities such as diabetes, and tissue volume.

Management and Therapeutic Approaches

Despite numerous interventions described in the literature, considerable heterogeneity exists in clinical practice, and no universally accepted guideline for skin management in irradiated patients has yet been established.[17,18] In general, two key elements define the management of radiation dermatitis: prevention and treatment.

Preventive Strategies in Acute Radiation-Induced Skin Injury

The primary objective in radiotherapy planning is to deliver the prescribed therapeutic or prophylactic dose to the tumor and regional lymph nodes while maximally sparing healthy tissues, particularly the skin. The main parameters influencing the risk of acute radiation-induced skin injury include irradiated skin volume, total dose delivered to the skin, fractionation scheme, radiotherapy technique, treatment quality (including adaptive RT approaches), and the addition of concurrent chemotherapy or molecular therapies. These factors are critical for determining both treatment efficacy and toxicity risk.

The irradiated skin volume and total dose correlate closely with disease extent. In particular, the presence of large metastatic lymph nodes (>3 cm) necessitates the irradiation of broader fields, thereby exposing larger skin volumes to higher doses. Nevertheless, although dose–volume relationships have been established for late toxicities, specific threshold values of dose and volume that directly predict the risk of severe acute skin toxicity have not yet been experimentally defined.[19]

Kawamura et al. developed a scoring system based on dosimetric and clinical parameters to predict the risk of acute radiation-induced skin injury in patients receiving radiotherapy for head and neck cancer. Their analysis identified the most significant prognostic factor as the skin volume receiving 60 Gy or more (V60Gy). Specifically, patients with a V60Gy exceeding 38 cm^3 had a 43.4% incidence of grade 3 acute radiation dermatitis.[20]

Modified fractionation has been shown to increase the risk of grade 3–4 radiation dermatitis. In the RTOG 9003 trial, the incidence of high-grade dermatitis was reported as 11% with hyperfractionated and accelerated RT schedules, compared to 7% with standard fractionation.[21]

Preventive strategies for acute radiation dermatitis begin at the stage of treatment planning but should be continued throughout therapy with the aim of minimizing the dose delivered to the skin. Daily image guidance is essential to ensure treatment accuracy. In patients with weight loss due to oral mucositis and dysphagia, the reduction in distance between the skin and target structures may cause high-dose regions to shift toward the skin, necessitating replanning. Although the role of adaptive radiotherapy in reducing skin toxicity remains inconclusive, it is considered a state-of-the-art technique for preventing unintended excess skin dose in head and neck irradiation.[22]

Concurrent chemoradiotherapy increases the risk of grade 3–4 radiation dermatitis.[38] Cisplatin, commonly used in head and neck cancer, enhances acute toxicity by inhibiting DNA repair, whereas cetuximab, an EGFR inhibitor, augments the severity of skin toxicity by affecting not only tumor cells but also keratinocytes, immune responses, and epidermal migration.[23] According to EORTC data, concurrent radiotherapy and cetuximab are associated with grade 3–4 dermatitis in approximately 49% of patients.[24] However, there are no established, evidence-based preventive strategies for such therapy-related dermatitis.

Different skin cleansing approaches have been proposed during radiotherapy. The MASCC guidelines and oncology nursing groups recommend washing with water or mild soap/shampoo.[17,25] However, cleansing with water alone is insufficient for removing lipid-soluble impurities and may impair the skin barrier, thereby triggering inflammation. For this reason, syndets (soap-free cleansers with appropriate pH, containing moisturizers, and free from fragrances or dyes) are preferred. Short, lukewarm showers and washing with the hands rather than sponges are also recommended, as they help protect the skin barrier and reduce the risks of dryness, microtrauma, and infection.[26]

Studies have shown that both soap-based and soap-free washing during radiotherapy are associated with less severe dermatitis, a reduced incidence of moist desquamation, and fewer secondary infections.[27] Nevertheless, syndets are considered superior for skin cleansing and barrier protection.

Failure to use emollients during radiotherapy may worsen dryness and inflammation.[29] The Society for Cutaneous Radiation Reactions (SCoR) guidelines recommend applying moisturizers to intact skin.[30] Emollients, which have long been a cornerstone therapy in conditions with barrier dysfunction such as atopic dermatitis, are also recommended for radiation dermatitis.[28] By reducing transepidermal water loss, alleviating pruritus, decreasing inflammation, and lowering the need for topical steroids, they provide temporary barrier restoration.[28] In addition, vitamin D is thought to support skin barrier integrity, while vitamin A may contribute to tissue repair and regeneration.[36,37]

Antibiotics are not indicated in the absence of clinical signs of infection.[28]

Some authors have suggested that the lipid phase of dermocosmetic products may reduce the transmission of radiotherapy beams, recommending gel formulations instead.[31] However, gels and pure oils are not advised, as they may exacerbate skin dryness.[28] During RT, patients should also receive structured photoprotection counseling.[32] This includes the regular, year-round use of SPF 50+ and UVA-protective sunscreens applied generously and reapplied frequently, particularly to the head and neck region. Patients should also avoid extreme temperatures, prefer electric shavers, and wear soft, loose clothing. While deodorants and antiperspirants are generally considered safe, alcohol-based products should be avoided.[17,33]

Topical glucocorticoids may be recommended for short-term use in radiation dermatitis; however, prolonged application carries a risk of skin atrophy, while the efficacy of calcineurin inhibitors has not been established.[28]

Chronic Radiation-Induced Skin Injury

Current therapeutic options for chronic radiation-induced skin injury remain limited. An international multidisciplinary Delphi consensus identified key risk factors for radiation-induced fibrosis, including re-irradiation, high dose and irradiated volume, head/neck and breast sites, connective tissue disorders, and smoking. Strong consensus was achieved on the importance of photoprotection, whereas topical/intralesional corticosteroids, pentoxifylline, vitamin E, physical therapy, and exercise received moderate recommendations. Vascular lesions were best managed with pulsed-dye laser, while fractional ablative laser was strongly endorsed for fibrotic changes. Fibrosis was emphasized as a major determinant of reduced quality of life through cosmetic disfigurement, pain, and functional limitation, underscoring the need for a multidisciplinary approach.[34]

In chronic radiation dermatitis, necrotic ulcerated lesions (after excluding tumor recurrence) may require appropriate debridement; infections should be managed with standard wound care; atrophic skin benefits from intensive moisturization; and hyperbaric oxygen therapy may be considered in selected cases. Special attention must be paid to endothelial damage and chronic hypoxia following high-dose RT.[35]

Conclusion

Although the pathophysiology and treatment strategies for radiation dermatitis are increasingly understood, a universally accepted skin care guideline during radiotherapy is still lacking. Further well-designed studies are needed to validate current recommendations. At present, standard care includes daily skin hygiene (syndets, emollients, sunscreens, smoking cessation), the use of topical glucocorticoids, and dressings tailored to exudate levels. Additionally, annual dermatologic follow-up after RT is strongly advised for all patients to ensure early detection and management of radiation-induced skin cancers.

REFERENCES

1- Selzer E, Liederer S, Lemaire C, Kren G, Radonjic D, Kornek G, Knocke T, Pötter R, Bachtiary B. Incidence of dermatitis in head and neck cancer patients treated with primary radiotherapy and cetuximab. Strahlenther Onkol. 2011 Jun;187(6):373-7. doi: 10.1007/s00066-011-2217-7. Epub 2011 May 16. PMID: 21603990

2- Burke G, Faithfull S, Probst H. Radiation induced skin reactions during and following radiotherapy: A systematic review of interventions. Radiography (Lond). 2022 Feb;28(1):232-239. doi: 10.1016/j.radi.2021.09.006. Epub 2021 Oct 11. PMID: 34649789

3- Ferreira EB, Vasques CI, Gadia R, Chan RJ, Guerra EN, Mezzomo LA, De Luca Canto G, Dos Reis PE. Topical interventions to prevent acute radiation dermatitis in head and neck cancer patients: a systematic review. Support Care Cancer. 2017 Mar;25(3):1001-1011. doi: 10.1007/s00520-016-3521-7. Epub 2016 Dec 12. PMID: 27957620.

4- Iacovelli NA, Torrente Y, Ciuffreda A, Guardamagna VA, Gentili M, Giacomelli L, Sacerdote P. Topical treatment of radiation-induced dermatitis: current issues and potential solutions. Drugs Context. 2020 Jun 12;9:2020-4-7. doi: 10.7573/dic.2020-4-7. PMID: 32587626; PMCID: PMC7295106

5- Hegedus F, Mathew LM, Schwartz RA. 'Radiation dermatitis: an overview. Int J Dermatol 2017;56:909e14

6- Wei J, Meng L, Hou X, Qu C, Wang B, Xin Y, et al. Radiation-induced skin reactions: mechanism and treatment. Canc Manag Res 2019;11:167e77.

7- Chan RJ, Webster J, Chung B, Marquart L, Ahmed M, Garantziotis S. 'Prevention and treatment of acute radiation-induced skin reactions: a systematic review and meta-analysis of randomized controlled trials. BMC Canc 2014

8- Lacas B, Bourhis J, Overgaard J, et al. MARCH Collaborative Group. Role of radiotherapy fractionation in head and neck cancers (MARCH): an updated meta-analysis. Lancet Oncol. 2017; 18(9): 1221–1237, doi: 10.1016/ S1470-2045(17)30458-8, indexed in Pubmed: 28757375.

9- Lacas B, Carmel A, Landais C, et al. MACH-NC Collaborative Group. Meta-analysis of chemotherapy in head and neck cancer (MACH-NC): An update on 107 randomized trials and 19,805 patients, on behalf of MACH-NC Group. Radiother Oncol. 2021; 156: 281–293, doi: 10.1016/j.radonc.2021.01.013, indexed in Pubmed: 33515668.

10- Paldor M, Levkovitch-Siany O, Eidelshtein D, et al. Single-cell transcriptomics reveals a senescence-associated IL-6/CCR6 axis driving radiodermatitis. EMBO Mol Med. 2022; 14(8): e15653, doi: 10.15252/emmm.202115653, indexed in Pubmed: 35785521

11- McCart EA, Thangapazham RL, Lombardini ED, et al. Accelerated senescence in skin in a murine model of radiation-induced multi-organ injury. J Radiat Res. 2017; 58(5): 636–646, doi: 10.1093/jrr/rrx008, indexed in Pubmed: 28340212

FULL TEXTS BOOK

12- Chen H, Wu M, Li G, et al. Association between XRCC1 single-nucleotide polymorphism and acute radiation reaction in patients with nasopharyngeal carcinoma: A cohort study. Medicine (Baltimore). 2017; 96(44): e8202, doi: 10.1097/MD.0000000000008202, indexed in Pubmed: 29095251.

13- Barnett GC, Coles CE, Elliott RM, et al. Independent validation of genes and polymorphisms reported to be associated with radiation toxicity: a prospective analysis study. Lancet Oncol. 2012; 13(1): 65–77, doi: 10.1016/ S1470-2045(11)70302-3, indexed in Pubmed: 22169268.

14- Leventhal J, Young MR. Radiation Dermatitis: Recognition, Prevention, and Management. Oncology (Williston Park). 2017; 31(12): 885–7, 894, indexed in Pubmed: 29297172.

15- Seité S, Bensadoun RJ, Mazer JM. Prevention and treatment of acute and chronic radiodermatitis. Breast Cancer (Dove Med Press). 2017; 9: 551–557, doi: 10.2147/BCTT. S149752, indexed in Pubmed: 29138594

16- Mao MH, Zhang J, Zhang JG. Comparing the RTOG/ EORTC and LENT-SOMA scoring systems for the evaluation of late skin toxicity after I seed brachytherapy for parotid gland cancer. Brachytherapy. 2017; 16(4): 877–883, doi: 10.1016/j.brachy.2017.02.011, indexed in Pubmed: 28495444

17- Gosselin T, Ginex PK, Backler C, et al. ONS Guidelines™ for Cancer Treatment-Related Radiodermatitis. Oncol Nurs Forum. 2020; 47(6): 654–670, doi: 10.1188/20.ONF.654- 670, indexed in Pubmed: 33063779

18- Tenorio C, de la Mata D, Leyva JA, et al. Mexican radiationdermatitis management consensus. Rep Pract Oncol Radiother. 2022; 27(5): 914–926, doi: 10.5603/RPOR. a2022.0101, indexed in Pubmed: 36523808

19- Emami B, Lyman J, Brown A, et al. Tolerance of normal tissue to therapeutic irradiation. Int J Radiat Oncol Biol Phys. 1991; 21(1): 109–122, doi: 10.1016/0360-3016(91)90171-y, indexed in Pubmed: 2032882

20- Kawamura M, Yoshimura M, Asada H, et al. A scoring system predicting acute radiation dermatitis in patients with head and neck cancer treated with intensity-modulated radiotherapy. Radiat Oncol. 2019; 14(1): 14, doi: 10.1186/ s13014-019-1215-2, indexed in Pubmed: 30665451.

21- Fu KK, Pajak TF, Trotti A, et al. A Radiation Therapy Oncology Group (RTOG) phase III randomized study to compare hyperfractionation and two variants of accelerated fractionation to standard fractionation radiotherapy for head and neck squamous cell carcinomas: first report of RTOG 9003. Int J Radiat Oncol Biol Phys. 2000; 48(1): 7–16, doi: 10.1016/s

22- Sher DJ, Avkshtol V, Lin MH, et al. Acute Toxicity and Efficiency Outcomes in the DARTBOARD Randomized Trial of Daily Adaptive Radiotherapy for Head and Neck Squamous Cell Carcinoma. Int J Radiat Oncol Biol Phys. 2023; 117(4): e6, doi: 10.1016/j.ijrobp.2023.08.032.

23- Bonomo P, Talamonti C, Desideri I, et al. Analysis of skin dose distribution for the prediction of severe radiation dermatitis in head and neck squamous cell carcinoma patients treated with concurrent chemo-radiotherapy. Head Neck. 2020; 42(2): 244–253, doi: 10.1002/hed.25997, indexed in Pubmed: 31682308

24- Giro C, Berger B, Bölke E, et al. High rate of severe radiation dermatitis during radiation therapy with concurrent cetuximab in head and neck cancer: results of a survey in EORTC institutes. Radiother Oncol. 2009; 90(2): 166–171, doi: 10.1016/j.radonc.2008.09.007, indexed in Pubmed: 18977050

25- . Behroozian T, Goldshtein D, Ryan Wolf J, et al. Multinational Association of Supportive Care in Cancer (MASCC) Oncodermatology Study Group Radiation Dermatitis Guidelines Working Group. MASCC clinical practice guidelines for the prevention and management of acute radiation dermatitis: part 1) systematic review. EClinicalMedicine. 2023; 58: 101886, doi: 10.1016/j.eclinm.2023.101886, indexed in Pubmed: 37181415.

FULL TEXTS BOOK

26- Mijaljica D, Spada F, Harrison IP. Skin Cleansing without or with Compromise: Soaps and Syndets. Molecules. 2022; 27(6), doi: 10.3390/molecules27062010, indexed in Pubmed: 35335373

27- Chan DC, Wong HC, Riad MA, et al. Prevention of radiation dermatitis with skin hygiene and washing: a systematic review and meta-analysis. Support Care Cancer. 2023; 31(5): 294, doi: 10.1007/s00520-023-07720-8, indexed in Pubmed: 37086339

28- Wollenberg A, Kinberger M, Arents B, et al. European guideline (EuroGuiDerm) on atopic eczema - part II: non-systemic treatments and treatment recommendations for special AE patient populations. J Eur Acad Dermatol Venereol. 2022; 36(11): 1904–1926, doi: 10.1111/ jdv.18429, indexed in Pubmed: 36056736

29- Mijaljica D, Spada F, Harrison IP. Skin Cleansing without or with Compromise: Soaps and Syndets. Molecules. 2022; 27(6), doi: 10.3390/molecules27062010, indexed in Pubmed: 35335373.

30- . Finkelstein S, Kanee L, Behroozian T, et al. Comparison of clinical practice guidelines on radiation dermatitis: a narrative review. Support Care Cancer. 2022; 30(6): 4663–4674, doi: 10.1007/s00520-022-06829-6, indexed in Pubmed: 35067732.

31- . Iacovelli NA, Torrente Y, Ciuffreda A, et al. Topical treatment of radiation-induced dermatitis: current issues and potential solutions. Drugs Context. 2020; 9, doi: 10.7573/ dic.2020-4-7, indexed in Pubmed: 32587626.

32- Seité S, Bensadoun RJ, Mazer JM. Prevention and treatment of acute and chronic radiodermatitis. Breast Cancer (Dove Med Press). 2017; 9: 551–557, doi: 10.2147/BCTT. S149752, indexed in Pubmed: 29138594.

33- Iacovelli NA, Torrente Y, Ciuffreda A, et al. Topical treatment of radiation-induced dermatitis: current issues and potential solutions. Drugs Context. 2020; 9, doi: 10.7573/ dic.2020-4-7, indexed in Pubmed: 32587626.

34- Wilson BN, Shah R, Menzer C, et al. Consensus on the clinical management of chronic radiation dermatitis and radiation fibrosis: a Delphi survey. Br J Dermatol. 2022; 187(6): 1054–1056, doi: 10.1111/bjd.21852, indexed in Pubmed: 36047980.

35- Zasadziński K, Spałek MJ, Rutkowski P. Modern Dressings in Prevention and Therapy of Acute and Chronic Radiation Dermatitis-A Literature Review. Pharmaceutics. 2022; 14(6), doi: 10.3390/pharmaceutics14061204, indexed in Pubmed: 35745777

36- Papadopoulou, S. N. A., Anastasiou, E. A., Adamantidi, T., Ofrydopoulou, A., Letsiou, S., & Tsoupras, A. (2025). A Comprehensive Review on the Beneficial Roles of Vitamin D in Skin Health as a Bio-Functional Ingredient in Nutricosmetic, Cosmeceutical, and Cosmetic Applications. *Applied Sciences*, *15*(2), 796. https://doi.org/10.3390/app15020796

37- Bechara N, Flood VM, Gunton JE. A Systematic Review on the Role of Vitamin C in Tissue Healing. Antioxidants (Basel). 2022 Aug 19;11(8):1605. doi: 10.3390/antiox11081605. PMID: 36009324; PMCID: PMC9405326.

38- Pazdrowski J, Gornowicz-Porowska J, Kaźmierska J, Krajka-Kuźniak V, Polanska A, Masternak M, Szewczyk M, Golusiński W, Danczak-Pazdrowska A. Radiation-induced skin injury in the head and neck region: pathogenesis, clinics, prevention, treatment considerations and proposal for management algorithm. Rep Pract Oncol Radiother. 2024 Jul 22;29(3):373-390. doi: 10.5603/rpor.100775. PMID: 39144266; PMCID: PMC11321788.

SIMULATION BASED ASSESSMENT OF COOLING FAN ENERGY TRADE-OFFS IN PHOTOVOLTAIC SYSTEMS USING MATLAB

Rahim Mammadzada

PhD Candidate at the Azerbaijan State Oil and Industry University, Information Technologies and Control Faculty, Department of Instrumentation Engineering, Baku, Azerbaijan.

ORCID ID: https://orcid.org/0009-0008-7552-2790

ABSTRACT

As photovoltaic panels heat up, their efficiency decreases, which may be mitigated by active cooling methods such as fans. This paper presents a MATLAB simulation to evaluate the energy trade-off between the power saved by cooling and the energy consumed by a fan. The model simulates fluctuating ambient temperatures and activates the fan when panel temperature exceeds a defined threshold. The simulation tracks both the improvement in electrical output and the cost of fan operation to determine when cooling is beneficial. The approach uses simple equations and a modular design to allow easy testing of different cooling strategies. The study offers a practical framework for researchers and students to analyze the balance between cooling gains and energy consumption in photovoltaic systems, laying the groundwork for future development of optimized cooling controls and energy management.

Keywords: Photovoltaic systems, Thermal modeling, MATLAB simulation, Active cooling, Energy trade-off, Fan control strategy

INTRODUCTION

Photovoltaic (PV) systems have become an essential component of the global renewable energy transition, offering a direct method of converting sunlight into electrical power (Shaker et al., 2024). However, their performance is highly sensitive to operating temperature (Sun et al., 2022). Under Standard Test Conditions (STC) of 25 °C, PV modules are rated to deliver their nominal power, but efficiency typically decreases as temperature rises above this point. Studies consistently report a loss of approximately 0.45–0.52 % of output per degree Celsius above 25 °C for both monocrystalline and polycrystalline technologies (Hudișteanu et al., 2024). This degradation is primarily linked to changes in key electrical parameters such as open-circuit voltage, short-circuit current, and fill factor, which together reduce the panel's usable power output and overall efficiency under elevated temperature conditions.

With global temperatures rising and many PV systems in hot regions, understanding the relationship between panel temperature and power loss is important (Hashim & Hassan, 2022). Small temperature increases can noticeably reduce power output. Quantifying this relationship is essential for predicting performance and assessing whether active cooling provides a net energy benefit. It is necessary to know how much power is lost per degree and whether the energy recovered from cooling outweighs its cost. This helps identify temperature thresholds where cooling is worthwhile and informs decisions for more efficient PV installations (Menacer et al., 2025).

In this study, MATLAB Simscape's Electrical library was used to model the power–temperature relationship of a PV module under constant irradiance conditions (1000 W/m²) (Hudișteanu et al., 2024). A baseline temperature of 25 °C was gradually increased to 65 °C. The resulting power drop was recorded and analyzed. This simulation-based approach provides a controlled way to identify critical temperature thresholds where performance degradation becomes significant. The results offer a clear

FULL TEXTS BOOK

numerical basis for estimating when active cooling might become energetically worthwhile. They also lay the groundwork for evaluating the potential benefits of cooling strategies in future work.

MATERIAL AND METHODS

Material

The photovoltaic panel is modeled using MATLAB Simulink with the help of Simscape and Simscape Electrical libraries (MathWorks, 2025a; MathWorks, 2025b). The PV Solar Block from Simscape Electrical allows us to simulate power generation based on values taken from the manufacturer's datasheet (MathWorks, 2025c). The block accepts solar irradiance and temperature inputs to calculate electrical output and internal heat generation. Using Simulink's visual tools, we can easily connect this block to other components, such as loads and thermal systems, to observe system-level behavior.

Mathods

The Collection of the Data

In this study, the Jinko Solar JKM350P-72-V photovoltaic panel is selected as the reference model. It is a widely used polycrystalline solar panel, especially in hot-climate and cost-sensitive regions, due to its affordability and availability. The panel is designed for both residential and industrial use, and its general specifications make it suitable for basic thermal and electrical analysis.

Table 1. Jinko JKM350P-72-V parameters provided by the PV array block

Solar Panel Parameters	**Max Power Point**		**Max Power**
	Voltage Vmp	**Current Imp**	
Values	39.1 V	8.94 A	349.554 W

Statistical Analysis

Since a ready-made PV block from Simscape Electrical is used, the only required parameter for PV block integration is the resistance connected between the positive and negative electrical terminals. This resistance value can be calculated using Ohm's Law.

$$R = \frac{U}{I}$$

By dividing the voltage at the maximum power point, 39.1 V, by the current at the maximum power point, 8.94 A, the required resistance value is found to be approximately 4.37 Ω.

RESEARCH AND FINDINGS

The first step is to ensure that the PV Array block functions properly by running the model under standard test conditions of 1000 W/m² irradiance and 25 °C temperature, as shown in Figure 2.

FULL TEXTS BOOK




Figure 2. Simulation of the JKM350P-72-V solar panel when the temperature is 25 degrees.

From the PV block outputs, the power can be calculated as the product of voltage and current values.

$$P = U \cdot I$$

From here, by multiplying 39.04V output voltage with 8.934A output current, we get a power output of around 348.78W, which is very close to the approximate 350W maximum power output of the chosen Jinko JKM350P-72-V solar panel. After this, the next step is to quantify how power generation changes with increasing temperature. These results are summarized in Table 2.

Table 2. Simulation results of the JKM350P-72-V solar panel model from 25 °C to 65 °C.

Temperature	Voltage	Current	Total Power	Power Loss
25	39.04 V	8.934 A	348.78336 W	-
27	38.9 V	8.902 A	346.2878 W	-2.6528 W
29	38.75 V	8.868 A	343.635 W	-2.7198 W
31	38.6 V	8.832 A	340.9152 W	-2.79696 W
33	38.44 V	8.796 A	338.11824 W	-3.51824 W
35	38.24 V	8.75 A	334.6 W	-2.4442 W
37	38.1 V	8.718 A	332.1558 W	-3.12396 W
39	37.92 V	8.677 A	329.03184 W	-3.1092 W
41	37.74 V	8.636 A	325.92264 W	-3.25549 W
43	37.55 V	8.593 A	322.66715 W	-3.27651 W
45	37.36 V	8.549 A	319.39064 W	-3.25979 W
47	37.17 V	8.505 A	316.13085 W	-3.36465 W
49	36.97 V	8.46 A	312.7662 W	-3.38342 W
51	36.77 V	8.414 A	309.38278 W	-3.48526 W

FULL TEXTS BOOK

53	36.56 V	8.367 A	305.89752 W	-3.38232 W
55	36.36 V	8.32 A	302.5152 W	-3.4824 W
57	36.15 V	8.272 A	299.0328 W	-3.27215 W
59	35.95 V	8.227 A	295.76065 W	-3.74965 W
61	35.72 V	8.175 A	292.011 W	-3.49225 W
63	35.51 V	8.125 A	288.51875 W	-3.552 W
65	35.29 V	8.075 A	284.96675	-

The table covers all measured parameters, including voltage, current, and total power output, across the temperature range of 25 °C to 65 °C. For convenience, Figure 3 presents the total power output as a function of temperature. The total power drops from a maximum of 348.78 W at 25 °C to a minimum of 284.96 W at 65 °C, clearly illustrating the decline in electrical performance as the panel heats up.




Figure 3. Total power drop of the JKM350P-72-V solar panel model from 25 °C to 65 °C.

Figure 4 shows the incremental power loss between successive temperature points. The trend is generally downward, with small fluctuations, highlighting how efficiency reduction accelerates at higher temperatures. For example, losses start at −2.65 W between 25 °C and 27 °C and reach −3.55 W between 63 °C and 65 °C.

FULL TEXTS BOOK




Figure 4. Power loss amount per 2 °C of the JKM350P-72-V solar panel model from 25 °C to 65 °C.

The results clearly indicate that as panel temperature rises, the overall power loss increases, emphasizing that forced cooling systems should be selectively applied at the highest operating temperatures to maximize net energy gain. Fan cooling requirements depend on the heat flux and surface area, with small fans providing minimal benefit, while appropriately sized industrial or high-CFM fans can offset significant losses, particularly in larger or higher power-output panels. This targeted approach ensures that the energy consumed by the cooling system is justified by the additional electricity preserved, making cooling most effective during periods of peak thermal stress.

CONCLUSION

In conclusion, this study demonstrates that PV panel efficiency declines progressively with rising temperature, and the associated power losses can be quantitatively tracked. By systematically modeling the temperature–power relationship using MATLAB's Simscape Electrical library, it is possible to quickly set up simulations and perform temperature-varying tests across a wide selection of solar panels. This approach allows identification of critical temperature ranges where forced cooling becomes energetically beneficial. The insights gained enable informed decisions on when to deploy active cooling, ensuring that energy spent on fans is justified by the additional power recovered. Overall, these findings provide a practical framework for designing more resilient PV installations, optimizing energy output, and guiding future work on smart, temperature-responsive cooling strategies tailored to panel size and operational conditions.

REFERENCES

Hashim, S. M., & Hassan, R. I. (2022). Impact of high temperature on PV productivity in hot desert climates. Green Technology, Resilience, and Sustainability, 2(1), 9. https://doi.org/10.1007/s44173-022-00009-9

Hudișteanu, V.-S., Cherecheș, N.-C., Țurcanu, F.-E., Hudișteanu, I., & Romila, C. (2024). Impact of temperature on the efficiency of monocrystalline and polycrystalline photovoltaic panels: A comprehensive experimental analysis for sustainable energy solutions. Sustainability, 16(23), Article 10566. https://doi.org/10.3390/su162310566

MathWorks. (2025a). Simscape documentation. Retrieved August 1, 2025, from https://www.mathworks.com/help/simscape/index.html

FULL TEXTS BOOK

MathWorks. (2025b). Simulink documentation. Retrieved August 1, 2025, from https://www.mathworks.com/help/simulink/index.html

MathWorks. (2025c). PV array (Simscape Electrical). Retrieved August 1, 2025, from https://www.mathworks.com/help/sps/powersys/ref/pvarray.html

Menacer, B., Baghdous, N. E. H., Narayan, S., Al-Lehaibi, M., Osorio, L., & Tuninetti, V. (2025). Efficiency enhancement of photovoltaic panels via air, water, and porous media cooling methods: Thermal–electrical modeling. Sustainability, 17(14), Article 6559. https://doi.org/10.3390/su17146559

Shaker, L. M., Al-Amiery, A. A., Hanoon, M. M., Al-Azzawi, W. K., & Kadhum, A. A. H. (2024). Examining the influence of thermal effects on solar cells: A comprehensive review. Sustainable Energy Research, 11(1), 6. https://doi.org/10.1186/s40807-024-00100-8

Sun, C., Zou, Y., Qin, C., Zhang, B., & Wu, X. (2022). Temperature effect of photovoltaic cells: A review. Advanced Composites and Hybrid Materials, 5(4), 2675–2699. https://doi.org/10.1007/s42114-022-00533-z

FULL TEXTS BOOK

SIMULATION BASED MODELING OF THERMAL AND ELECTRICAL BEHAVIOR IN PHOTOVOLTAIC PANELS USING MATLAB

Rahim Mammadzada

PhD Candidate at the Azerbaijan State Oil and Industry University, Information Technologies and Control Faculty, Department of Instrumentation Engineering, Baku, Azerbaijan.

ORCID ID: https://orcid.org/0009-0008-7552-2790

ABSTRACT

The performance of photovoltaic panels declines as temperature increases, impacting their energy output, especially in warm climates. This paper presents a MATLAB simulation that models the thermal and electrical behavior of a solar panel under fluctuating ambient temperatures. The simulation links heat buildup with changes in electrical output using simple equations and a modular design. This approach allows easy modification and expansion for future studies, such as adding cooling control or environmental data. A feature compares panel performance under steady and varying temperature conditions to help visualize thermal effects on efficiency. The study focuses on providing a clear framework that balances simplicity and realism, helping researchers and students understand key interactions in photovoltaic systems. This foundational model supports future work on smart temperature management and improved solar panel operation.

Keywords: Photovoltaic systems, Thermal modeling, MATLAB simulation, Temperature effects, Electrical efficiency, Simulation framework

INTRODUCTION

Photovoltaic (PV) systems have become an essential part of the global shift toward renewable energy. Their ability to convert sunlight directly into electricity makes them highly attractive for both residential and industrial use. However, the performance of PV panels is heavily influenced by environmental conditions, especially temperature. The Standard Test Condition (STC) for PV panels is defined at 25 °C, and numerous studies have shown that efficiency decreases as temperature rises above this baseline. For example, research in hot desert climates reports a typical efficiency drop of around 0.45 percent for every one-degree Celsius increase above 25 °C (Hashim & Hassan, 2022). Similarly, experimental analyses have confirmed losses of up to 0.52 percent per degree Celsius increase beyond 25 °C in both monocrystalline and polycrystalline panels (Hudișteanu et al., 2024).

This degradation is largely due to the thermal impact on electrical parameters such as open-circuit voltage, short-circuit current, and fill factor, which collectively reduce the total power output of the system (Menacer et al., 2025; Shaker et al., 2024). With average global temperatures steadily rising, this issue is no longer limited to traditionally hot regions. It is becoming a concern for PV installations worldwide. Recent reviews stress that without thermal regulation, PV panels are unlikely to operate near their rated capacity (Shaker et al., 2024; Sun et al., 2022). To address this, several methods have been proposed to improve cooling efficiency, including air, water, and porous media techniques (Menacer et al., 2025). However, these solutions must be evaluated for energy trade-offs, cost, and implementation complexity, especially in small to medium-scale systems.

In this study, we present a simulation-based analysis of PV temperature behavior using MATLAB and Simscape tools. Instead of directly evaluating a specific cooling system, the focus is on quantifying the power loss in PV panels rated between 300 W and 400 W under varying thermal conditions. By

FULL TEXTS BOOK

identifying how efficiency declines with rising temperature, we estimate the potential power gains achievable through cooling. This understanding will guide the selection of suitable cooling approaches, such as fan-based systems, by weighing energy benefits against the power they consume.

MATERIAL AND METHODS

Material

This study utilizes MATLAB and Simulink (MathWorks, 2025a), with a focus on the Simscape environment, to model the electrical and thermal behavior of photovoltaic (PV) systems. Simscape provides a unified framework for integrating physical domains such as electrical circuits, heat transfer, and control logic, making it suitable for analyzing the energy dynamics of solar panels operating in hot climates (MathWorks, 2025b). For the electrical modeling of the PV source, a built-in Solar Array block from the Simscape Electrical library is used (MathWorks, 2025c). This block, shown in Figure 1, enables the simulation of configurable monocrystalline solar panels and includes built-in parameters such as open-circuit voltage, short-circuit current, temperature coefficients, and nominal power output. It covers a typical power range of 250 W to 400 W, suitable for both conventional and high-efficiency panel configurations.




Figure 1. PV Array block of the Simscape Electrical library of MATLAB

In addition to the electrical part, the thermal behavior of the solar panel is examined by adjusting the temperature input through a constant block. This approach provides an initial understanding of how heat impacts power generation without involving full thermal domain modeling. This modeling approach enables the simulation of realistic efficiency losses due to temperature increases.

Mathods

The Collection of the Data

As a textbook example, a representative panel JKM300M-60-V was selected from the most commonly available modules in the MATLAB Simscape Electrical library Figure 2.

FULL TEXTS BOOK




Figure 2. PV array block with Jinko Solar JKM300M-60-V preset

As a first example, let's consider the JKM300M-60-V, where "300" denotes the nominal maximum power in watts, "M" indicates a monocrystalline cell structure, "60" specifies the number of cells in the module, and "V" refers to the particular version or revision of the panel design. This selection allows for consistent benchmarking and simulation across a standard module configuration. This panel is designed for both residential and industrial use, and its general specifications make it suitable for basic thermal and electrical analysis. Table 1 shows its main parameters taken from the PV array block configuration, as shown in Figure 2.

Table 1. Jinko JKM300M-60-V parameters provided by the PV array block

Solar Panel Parameters	**Max Power Point**		**Max Power**
	Voltage Vmp	**Current Imp**	
Values	32.6 V	9.21 A	300.246 W

Statistical Analysis

Since a ready-made PV block from Simscape Electrical is used, the only required parameter for external load modeling is the resistance connected between the positive and negative electrical terminals. This resistance value can be calculated using Ohm's Law as shown below.

$$R = \frac{U_{mp}}{I_{mp}}$$

From here, by dividing the voltage at the max power point, 32.6 V, by the current at the max power point of 9.21 A, we get the required resistance value of 3.54 Ω.

RESEARCH AND FINDINGS

The first step in setting up the model involves configuring the PV Array block to match the specifications of a 300W solar panel, as illustrated in Figure 3. To confirm that the simulation environment is functioning correctly, the model is run under standard test conditions using an irradiance value of 1000

FULL TEXTS BOOK

W/m² and a panel temperature of 25 °C. This helps verify that the simulated output is consistent with the expected power rating before proceeding with more detailed analyses.




Figure 3. Simulation of the JKM300M-60-V solar panel when the temperature is 25 degrees.

To confirm the model is functioning correctly, the output power is calculated by multiplying the voltage and current values obtained from the PV Array block. The last output of this block provides five key parameters: voltage (V), current (I), diode current, irradiance, and temperature. Verifying that the product of voltage and current closely matches the panel's rated power under standard conditions helps ensure the model is correctly configured.

$$P = U \cdot I$$

From here, by multiplying 32.58 V output voltage by 9.203 A output current, we get a power output of around 299.83 W, which is very close to the 300 W maximum power output of the chosen JKM300M-60-V solar panel. This proves that the simulation was set up correctly. The next step here is to run the simulation with increased input temperature, as shown in Figure 4.




Figure 4. Simulation of the JKM300M-60-V solar panel when the temperature is 55 degrees.

In regions with high solar irradiance, ambient temperatures typically range between 30°C and 40°C; however, the panel surface temperature can rise significantly above these levels due to direct sunlight exposure. During summer months, temperature increases of 15 °C to 20 °C above ambient are common, making 55 °C a reasonable estimate for panel temperature in hot climates. This value provides a practical basis for comparing fan-assisted and passive cooling scenarios in later sections of the study. In this case, if we calculate the power generated by the solar panel by multiplying 30.27 V output voltage by 8.551 A output current, we get a power output of around 258.83 W. The final step here is to calculate the percentage of lost potential power generation, which is done by the equation below.

FULL TEXTS BOOK

$$\eta_{Actual} = \frac{P_{Actual}}{P_{Rated}}$$

This corresponds to a total power loss of around 41 W, which translates to an approximate 13.73% reduction in the panel's power generation efficiency under elevated temperature conditions. By repeating these steps, it is possible to simulate other solar panels and note the results down in Table 2.

Table 2. Simulation results of different solar panel models at 55°C

Panel Model	Current at 55 °C	Voltage at 55 °C	Resistance (Ump / Imp)	Output Power	Power Loss	Efficiency Loss
JKM300M-60-V	8.551 A	30.27 V	3.54 Ω	258.8 W	≈41.2 W	≈13.7%
JKM325M-72	7.965 A	35.36 V	4.44 Ω	281.6 W	≈43.4 W	≈13.3%
JKM350M-72-V	8.32 A	36.36 V	4.37 Ω	302.5 W	≈47.5 W	≈13.5%
JKM375M-72-V	8.633 A	37.04 V	4.29 Ω	319.8 W	≈55.2 W	≈14.7%
JKM400M-72L-V	8.96 A	38.89 V	4.34 Ω	348.4 W	≈51.6 W	≈12.9%

As PV module power increases, the absolute power loss at elevated temperatures grows. For example, the 300 W panel loses ≈41 W at 55 °C, while the 375 W panel loses ≈55 W. Efficiency losses remain similar (≈13–15 %), highlighting that higher-power panels are more affected by temperature, making targeted cooling increasingly beneficial.

CONCLUSION

The simulation results clearly demonstrate that higher-power PV modules experience greater absolute power losses at elevated temperatures, even though the relative efficiency drop remains similar across models. For example, while the JKM300M-60-V loses approximately 41 W at 55 °C, the JKM400M-72L-V loses over 51 W under the same conditions. This trend highlights that thermal effects become increasingly significant as panel output rises, emphasizing the importance of understanding temperature-dependent performance for larger systems. Furthermore, these results illustrate how MATLAB Simscape Electrical's PV library provides a fast and flexible platform for modeling a wide range of panels, enabling accurate assessment of temperature effects on power output without the need for physical testing. This approach facilitates efficient evaluation of potential efficiency gains and supports informed decisions regarding thermal management strategies in PV installations.

REFERENCES

Hashim, S. M., & Hassan, R. I. (2022). Impact of high temperature on PV productivity in hot desert climates. Green Technology, Resilience, and Sustainability, 2(1), 9. https://doi.org/10.1007/s44173-022-00009-9

Hudișteanu, V.-S., Cherecheș, N.-C., Țurcanu, F.-E., Hudișteanu, I., & Romila, C. (2024). Impact of temperature on the efficiency of monocrystalline and polycrystalline photovoltaic panels: A comprehensive experimental analysis for sustainable energy solutions. Sustainability, 16(23), Article 10566. https://doi.org/10.3390/su162310566

MathWorks. (2025a). Simscape documentation. Retrieved August 1, 2025, from https://www.mathworks.com/help/simscape/index.html

MathWorks. (2025b). Simulink documentation. Retrieved August 1, 2025, from https://www.mathworks.com/help/simulink/index.html

MathWorks. (2025c). PV array (Simscape Electrical). Retrieved August 1, 2025, from https://www.mathworks.com/help/sps/powersys/ref/pvarray.html

Menacer, B., Baghdous, N. E. H., Narayan, S., Al-Lehaibi, M., Osorio, L., & Tuninetti, V. (2025). Efficiency enhancement of photovoltaic panels via air, water, and porous media cooling methods: Thermal–electrical modeling. Sustainability, 17(14), Article 6559. https://doi.org/10.3390/su17146559

Shaker, L. M., Al-Amiery, A. A., Hanoon, M. M., Al-Azzawi, W. K., & Kadhum, A. A. H. (2024). Examining the influence of thermal effects on solar cells: A comprehensive review. Sustainable Energy Research, 11(1), 6. https://doi.org/10.1186/s40807-024-00100-8

Sun, C., Zou, Y., Qin, C., Zhang, B., & Wu, X. (2022). Temperature effect of photovoltaic cells: A review. Advanced Composites and Hybrid Materials, 5(4), 2675–2699. https://doi.org/10.1007/s42114-022-00533-z

SEFAZOLİNİN EMBRİYOTOKSİK ETKİSİNİN İN OVO YÖNTEM İLE BELİRLENMESİ

DETERMINATION OF THE EMBRYOTOXIC EFFECT OF CEFAZOLIN BY IN OVO METHOD

Dr. Mustafa Sedat ARSLAN

Selçuk Üniversitesi, Veteriner Fakültesi, Prof. Dr. Hümeyra Özgen Araştırma ve Uygulama Çiftliği, Konya, Türkiye

ORCID: 0000-0002-4079-500X

Dr. Öğr. Üyesi Tuğba Melike PARLAK

Selçuk Üniversitesi, Veteriner Fakültesi, Farmakoloji ve Toksikoloji AD, Konya, Türkiye

ORCID: 0000-0003-3277-0336

Ece OKTAY

Lisansüstü Eğitim Enstitüsü, İstanbul Üniversitesi-Cerrahpaşa, İstanbul, Türkiye

ORCID: 0000-0002-3117-7875

ÖZET

Bu çalışmanın amacı fertil tavuk yumurtaları hava kamarasına farklı dozlarda sefazolin uygulamasının embriyotoksik ve teratojenik etkilerinin belirlenmesidir. Kuluçka makinesine yerleştirilen tavuk yumurtalarına inkübasyonun yedinci günü ışık altında ovoskop ile fertilite kontrolü yapıldı. Araştırmada kullanılacak toplam 210 adet fertil tavuk yumurtası, her grupta 30 adet fertil yumurta olacak şekilde 7 gruba ayrıldı. Birinci gruptaki yumurtalara hiçbir uygulama yapılmazken, ikinci grubun hava kamaralarına 50 µL serum fizyolojik uygulandı. Çalışmanın diğer gruplarına sırası ile 250, 125, 62.5, 31.25 ve 15.62 mg/kg dozunda sefazolin 50 µL hacim içinde uygulandı. Kuluçka süresi sonunda civcivlerde kontroller yapıldı. Araştırmada en yüksek doz uygulamasında %83.33 oranında ölüm gözlendi ($P<0.05$). İncelenen anatomik yapılarda herhangi bir anomaliye rastlanmadı. Sonuç olarak en yüksek dozlarda sefazolin uygulamasının fertil tavuk yumurtasında embriyotoksisiteye neden olabileceği ifade edilebilir.

Anahtar kelimeler: Sefazolin, embriyotoksisite, in ovo

ABSTRACT

The aim of this study was to determine the embryotoxic and teratogenic effects of cefazolin application at different doses to the air chamber of fertile chicken eggs. Fertility was checked using an ovoscope under light on the seventh day of incubation on chicken eggs placed in an incubator. A total of 210 fertile chicken eggs to be used in the study were divided into seven groups, each containing 30 fertile eggs. No application was applied to the eggs in the first group, while 50 µL of physiological saline was applied to the air chambers of the second group. Cefazolin was administered in 50 µL volumes at doses of 250, 125, 62.5, 31.25 and 15.62 mg/kg, respectively, to the other groups of the study. At the end of the incubation period, controls were performed on chicks. In the study, 83.33% mortality was observed in the highest dose application ($P<0.05$). No anomalies were observed in the anatomical structures examined. As a result, it can be stated that the application of cefazolin at the highest doses may cause embryotoxicity in fertile chicken eggs.

Keywords: Cefazolin, embryotoxicity, in ovo

FULL TEXTS BOOK

GİRİŞ

Günümüzde sefalosporinler keşfediliş tarihi ve etki spektrumuna göre 4 kuşağa (nesil) ayrılmaktadır. Genel olarak sefalosporinlerde, birinci kuşaktan dördüncü kuşağa doğru etki spektrumları gram pozitif bakterilerden, gram negatif bakterilere kayar. Etki spektrumları düşünüldüğünde birinci ve ikinci kuşaklar genellikle üst solunum sistemi ve deri enfeksiyonlarında kullanılırken, üçüncü ve dördüncü kuşak sefalosporinler sindirim, üriner ve alt solunum sistemi enfeksiyonlarında kullanılırlar. Ayrıca son iki kuşak sepsis ve septik şok tedavisinde tercih edilirler. İkinci kuşak sefalosporinlerden başlayarak anerob bakterilere etkinlik başlar. Sefazolin birinci kuşak sefalosporindir ve özellikle Staphylococcus spp ve Streptococcus spp türlerine etkinliği oldukça iyidir. Veteriner sahada birinci kuşak sefalosporinlerin meme içi, uterus içi ve parenteral uygulamaya yönelik ticari preparatları bulunmaktadır. Sefazolin içeren preparatlar veteriner hekimlikte bulunmamaktadır. Ancak beşeri preparatları pet kliniğinde veteriner hekimlerce etiket dışı olarak kullanılmaktadır (**Yazar 2023, Yazar 2024**). Sefazolin beşerî hekimlikte gebelikte güvenlik açısında kategori B olarak değerlendirilmekle (**Emiroğlu ve Sakin 2022**), birlikte yüksek dozlarda fertil tavuk yumurtalarında etkinliği değerlendirilmemiştir.

Beşerî hekimlikte kullanılan ilaçlar gebelikte güvenlik açısında beş kategori olarak değerlendirilmektedir. Kategori A kesin güvenli olarak kabul edilirken, kategori B ilaçlar ise deney hayvanlarında fötal risk taşımadığının belirlendiği ve A ile B kategori ilaçların gebe insanlarda kullanılabileceği ifade edilmiştir (**Ozturk 2018**). Kimyasal maddelerin veya veteriner sahada kullanılan bazı ilaçların embriyotoksisite, teratojenite veya malformasyonlarının belirlenmesinde fertil tavuk yumurtası kullanılabilmektedir (**Jelinek ve ark 1985, Canbar ve ark 2021a, Canbar ve ark 2021b, Uslu ve ark 2022, Canbar ve ark 2023, Uslu ve ark 2024**). **Hill ve Hoffman (1984)** ve **Kotwani (1998)** fertil kanatlı yumurtalarının birinci basamak testleri olarak kullanılabileceğini ifade etmişlerdir. Embriyotoksisite çalışmalarında fertil tavuk yumurtaları, birinci basamakta embriyotoksik doz sınırı belirlemek için yapılır ve CHEST-I (Tavuk Embriyotoksisite Tarama Testi) olarak tanımlanır. CHEST-II aşamasında ise (CHEST-II) teratojenik parametrelerin tespiti yapılır (**Özparlak 2015**).

Mevcut çalışmada fertil tavuk yumurtaları hava kamarasına farklı dozlarda sefazolin uygulamasının embriyotoksik ve teratojenik etkilerinin belirlenmesi amaçlandı.

MATERYAL VE METOT

Araştırmada ticari bir işletmeden (Anadolu Entegre Damızlık Hay. Pet. Tur. San. Tic. Ltd. Şti., Konya, Türkiye) temin edilen 210 adet fertil Tetra L Superb ırkı tavuk yumurtaları kullanıldı. Araştırma prosedürü Selçuk Üniversitesi Veteriner Fakültesi Etik kurulunca onaylandı (SUVDAMEK, 2025/88). Çalışma süresince yumurtalar 37.7±0.2 ^{0}C ve %55±0.2 bağıl nem bulunan kuluçka makinesinde (İmza Teknik, Konya, Türkiye) kuluçka sürelerini tamamladı. Kuluçka süresince her 90 dakikada bir yumurta pozisyonlarının otomatik olarak değiştirilmesi sağlandı. Kuluçkanın 7. gününde ovoskop ile ışık altında fertilite kontrolü yapıldı ve fertil olmayan yumurtalar ayıklanarak her bir grupta 30 adet fertil yumurta olacak şekilde rastgele gruplandırıldı. Uygulamanın yapılabilmesi için yumurtaların hava kamaralarının bulunduğu küt uçları %80 etil alkol ile temizlenerek yumurta delici (Bai Shun, Zhejilang, Çin) ile delindi. Sefazolin (Sefazol enj, Mustafa Nevzat İlaç Sanayii, İstanbul, Türkiye) %0.9 NaCl solüsyonunda çözdürüldü ve aşağıda belirtilen dozlarda hava keselerine uygulandı. Uygulamayı takiben açılan delikler parafin ile kapatıldı ve tekrar kulluçka makinesine yerleştirildi.

FULL TEXTS BOOK

Deneysel Dizayn ve Uygulamalar

Metot: Çalışmada, rastgele her grupta 30 adet olacak şekilde, 7 gruba ayrılarak kuluçka makinesine yerleştirilen, ortalama 50 gram ağırlığında toplam 210 adet fertil tavuk yumurtası kullanılmıştır.

1. Grup (K, n:30): Bu gruptaki yumurtalar herhangi bir işleme tabi tutulmadan kuluçka makinesinde kuluçka sürelerini tamamladı. Bu grup Negatif kontrol grubu olarak kullanıldı.

2. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 50 µL hacminde etken maddenin çözücüsü olan %0.9 NaCl solüsyonu uygulandı ve Pozitif kontrol grubu olarak değerlendirildi.

3. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 250 mg/kg sefazolin 50 µL hacim içinde enjekte edildi.

4. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 125 mg/kg sefazolin 50 µL hacim içinde enjekte edildi.

5. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 62.5 mg/kg sefazolin 50 µL hacim içinde enjekte edildi.

6. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 31.25 mg/kg sefazolin 50 µL hacim içinde enjekte edildi.

7. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 15.62 mg/kg sefazolin 50 µL hacim içinde enjekte edildi.

Kuluçka süresinin tamamlanmasıyla (21. gün) yumurtalar açılarak canlı ve ölü embriyoların sayısı, embriyotoksik doz sınırı ve letal doz $(LD)_{50}$ oranı belirlendi.

Letal doz $(LD)_{50}$ değeri probit testi kullanılarak hesaplandı. Araştırma sonuçları ki-kare testi ile değerlendirildi (SPSS 29.0). $P<0.05$ değeri istatistiki sınır olarak kabul edildi.

BULGULAR

Sefazolin embriyotoksik etkisi Tablo 1'de sunuldu. Yapılan kontrollerde herhangi bir anomaliye rastlanılmadı. Araştırmada en yüksek doz uygulamasında %83.3 oranında ölüm gözlendi ve diğer bütün gruplardan istatistiki olarak farklı belirlendi ($P<0.05$). 125 mg/kg doz uygulamasının ise daha düşük doz uygulanan gruplardan yüksek düzeyde ölüme neden olduğu tespit edildi ($P<0.05$). Yapılan probit testine göre LD50 değeri 186 (132-305) mg/kg olarak hesaplandı.

Tablo 1. Sefazolin uygulanan fertil tavuk yumurtalarında embriyonik ölüm oranları.

	EEÖ	GEÖ	Canlı	Toplam	%	GÖO
Kontrol	0	0	30	30	100.00^{c}	-----
SF	0	0	30	30	100.00^{c}	0.00
250 mg/kg	4	21	5	30	16.67^{a}	83.33
125 mg/kg	2	2	26	30	86.67^{b}	13.33
62.5 mg/kg	0	0	30	30	100.00^{c}	0.00
31.25 mg/kg	2	0	28	30	93.33^{bc}	6.67

FULL TEXTS BOOK

15.62 mg/kg	2	1	27	30	90.00^{bc}	10.00

$^{a\ b,\ c}$*Aynı sütunda istatistiki fark belirlendi (P<0.05). SF: Serum fizyolojik, EEÖ: Erken embriyonik ölüm, GEÖ: Geç embriyonik ölüm, GÖO: Gerçek ölüm oranı.

TARTIŞMA

Sefazolin birinci kuşak sefalosporindir ve özellikle Staphylococcus spp ve Streptococcus spp kaynaklı solunum ve deri enfeksiyonları tedavisinde kullanılır. Kısmen anerob bakterilere de etkinliği bulunmaktadır. İlaç veteriner sahada ruhsatlı olmamakla birlikte, pet kliniğinde etiket dışı olarak kullanılmaktadır (**Yazar 2023, Yazar 2024**). Sefazolin beşerî hekimlikte gebelikte güvenlik açısında kategori B olarak değerlendirilmektedir (**Emiroğlu ve Sakin 2022**), ancak yüksek dozlarda etkinliği fertil tavuk yumurtalarında etkinliği değerlendirilmemiştir.

Mevcut çalışmada en yüksek dozda sefazolin (250 mg/kg) uygulamasının %83.3 oranında ölüme neden olduğu ve diğer gruplardan istatistiki olarak farklı olduğu belirlendi (P<0.05, Tablo 1). Ayrıca 125 mg/kg doz uygulamasının ise daha düşük uygulanan diğer gruplardan yüksek düzeyde ölüme neden olduğu tespit edildi (P<0.05, Tablo 1). Yapılan probit testine göre LD50 değeri 186 (132-305) mg/kg olarak hesaplandı. Gebe kadınlarda sefalosporinler yaygın olarak uzun zamandandır kullanılmaktadır. Sefazolin özellikle cerrahi profilaksisinde tercih edilmektedir (**Stojanova ve ark 2022**). Gebe kadınlara sefazolin uygulandığında plasentayı geçebildiği bilinmektedir (**Dekel ve ark 1980, Elkomy ve ark 2014**). Gebe kemirgenlere 240-4000 mg/kg/gün dozunda sefazolin uygulamasının olumsuz etkilere neden olmadığı bildirilmiştir (**FDA 2025**). Ayrıca tavşan ve farelere sırasıyla yüksek dozlarda sefazolin uygulamasının teratojenik etkiye neden olmadığı, ayrıca çiftleşme, üreme, gebelik, doğum ve emzirme üzerine etkisinin olmadığı belirlenmiştir. Ayrıca laktasyondaki ratlara ilaç uygulandığında, laktasyon süresince olumsuz bir etki gözlenmediği ifade edilmiştir (**Birkhead ve ark 1973**). Gebe kadınlarda enfeksiyonların tedavisinde sefalosporinlerin diğer antibiyotiklere göre daha az teratojenik etki gösterebileceği bildirilmiştir. Ancak sefalosporin gebelikte kullanımında mutlak güvenilirliği doğrulanmadığı için dikkatli kullanılması gerektiği de bildirilmiştir (**Kumar Mishal ve ark 2025**). Gebe kadınlar üzerinde yapılan çalışmada beta-laktamların da teratojenik potansiyele sahip olabileceği bildirilmiştir. Sefaklor (ikinci kuşak) ve sefaleksinin (birinci kuşak) malformasyona neden olabileceği, ancak setriaksonun (üçüncü kuşak) malformasyona neden olmadığı belirlenmiştir **(Eric ve Sabo 2008).**

SONUÇ

Sonuç olarak gebe kadınlarda güvenli kabul edilen sefazolinin çok yüksek dozda fertil tavuk yumurtalarında embriyotoksisiteye neden olabileceği gözlendiği için gebe kemirgenlerde daha fazla çalışma yapılması gerektiği ve/veya fertil tavuk yumurtalarının sefazoline memelilere göre daha duyarlı olduğu ifade edilebilir.

KAYNAKLAR

1- Birkhead HA, Briggs GB, Saunders LZ, 1973. Toxicology of cefazolin in animals. J Infect Dis, 128:379-381.

2- Canbar R, Akcakavak G, Uslu M, Bas AL, 2021a. Determination of embryotoxic effects of Tarantula cubensis alcoholic extract with in ovo model. Magy Allatorvosok Lapja, 143:497-504.

3- Canbar R, Uslu M, Arslan MS, Kızılay H, 2023. Determination of embryotoxic effects of atipamezole using in ovo model. Rev Cient, 33:rcfcv-e33230.

FULL TEXTS BOOK

4- Canbar R, Uslu M, Tufan Ö, Yazar E, 2021b. Detomidinin embriyotoksik doz sınırının in-ovo model ile belirlenmesi. Eurasian J Vet Sci, 37:71-75.
5- Dekel A, Elian I, Gibor Y, Goldman JA, 1980: Transplacental passage of cefazolin in the first trimester of pregnancy. Europ J Obstet Gynec Reprod Biol, 10:303-307.
6- Elkomy MH, Sultan P, Drover DR, Epshtein E, Galinkin JL, Carvalho B, 2014. Pharmacokinetics of prophylactic cefazolin in parturients undergoing cesarean delivery. Antimicrob Agents Chemother, 58(6):3504-5013.
7- Emiroğlu SB, Sakin F, 2022. Gebelikte ilaç kullanımı ve teratojenite, Veteriner Farmakoloji ve Toksikoloji Derneği Bülteni, 13:90-98.
8- Eric M, Sabo A, 2008. Teratogenicity of antibacterial agents. Coll Antropoli, 32:919-925.
9- FDA, 2025. https://www.accessdata.fda.gov/drugsatfda_docs/label/2015/207131s000lbl.pdf. Erişim tarihi:23.04.25.
10- Hill E, Hoffman D, 1984. Avian models for toxicity testing. J Am Coll Toxicol, 3(6):357–376.
11- Jelinek R, Peterka M, Rychter Z, 1985. Chick embryotoxicity screening test – 130 substances tested. Indian J Exp Biol, 23:588-595.
12- Kotwani A, 1998. Use of chick embryo in screening for teratogenicity. Indian J Physiol Pharmacol, 42:189-204.
13- Kumar Mishal PM, Sahoo M, Dutta AK, 2025. Teratogenic effect of antibiotics in pregnancy: a review. IJCRT, 13:717-719.
14- Ozturk Z, 2018. Medication use in pregnancy and risk evaluation: do not categorize me. Zeynep Kamil Med J, 49(1):109-112.
15- Özparlak H, 2015. The use of chick embryos in embryotoxicity and teratogenicity tests. SUFEFD, 40:13-22.
16- Stojanova J, Arancibia M, Ghimire S, Sandaradura I, 2022. Understanding the pharmacokinetics of antibiotics in pregnancy: is there a role for therapeutic drug monitoring? a narrative review. Ther Drug Mon, 44:50-64.
17- Uslu M, Canbar R, Akcakavak G, Yazar E, 2022. Determination of the embryotoxic effect of maropitant using an in ovo model. Pol J Vet Sci, 25:357-359.
18- Uslu M, Canbar R, Arslan MS, Kızılay H, 2024. Determination of the embryotoxic effect of metronidazole using an in ovo model. Rev Cient, 34:34310.
19- Yazar E, 2023. Kedilerde Tedavi El Kitabı, Nobel Bilimsel Eserler, Konya, Tğrkiye.
20- Yazar E, 2024. Kemoterapötikler, in: Veteriner İlaç Rehberi Tedavi El Kitabı, Ed: Yazar E, Nobel tıp kitabevleri, Ankara, Türkiye.

FULL TEXTS BOOK

SEFTRİAKSONUN EMBRİYOTOKSİK ETKİSİNİN İN OVO YÖNTEM İLE BELİRLENMESİ

DETERMINATION OF THE EMBRYOTOXIC EFFECT OF CEFTRIAXONE BY IN OVO METHOD

Dr. Mustafa Sedat ARSLAN

Selçuk Üniversitesi, Veteriner Fakültesi, Prof. Dr. Hümeyra Özgen Araştırma ve Uygulama Çiftliği, Konya, Türkiye

ORCID: 0000-0002-4079-500X

Dr. Öğr. Üyesi Tuğba Melike PARLAK

Selçuk Üniversitesi, Veteriner Fakültesi, Farmakoloji ve Toksikoloji AD, Konya, Türkiye

ORCID: 0000-0003-3277-0336

Ece OKTAY

Lisansüstü Eğitim Enstitüsü, İstanbul Üniversitesi-Cerrahpaşa, İstanbul, Türkiye

ORCID: 0000-0002-3117-7875

ÖZET

Bu çalışmada fertil tavuk yumurtaları hava kamarasına farklı dozlarda seftriakson uygulamasının embriyotoksik ve teratojenik etkilerinin belirlenmesi amaçlandı. Kuluçka makinesine yerleştirilen tavuk yumurtarına inkübasyonun 7. gününde ovoskop ile ışık altında fertilite kontrolü yapıldı ve fertil olmayan yumurtalar çalışma dışı bırakıldı. Toplam 210 adet fertil tavuk yumurtası her grupta 30 adet olacak şekilde rastgele 7 gruba ayrıldı. Birinci gruptaki yumurtalara hiçbir uygulama yapılmazken, ikinci grubun hava kamaralarına 50 µL serum fizyolojik uygulandı. Aynı gün diğer gruplara sırası ile 250, 125, 62.5, 31.25 ve 15.62 mg/kg dozunda seftriakson 50 µL hacim içinde uygulandı. Kuluçka süresi sonunda yumurta ve civcivlerde kontroller yapıldı. Yapılan çalışma sonunda istatiksel olarak embriyonik ölüm ile anatomik yapılarda herhangi bir anomaliye rastlanmadı. Sonuç olarak, araştırmada kullanılan farklı dozlardaki seftriakson uygulamasının fertil tavuk yumurtalarında embriyotoksisite ve teratojeniteye neden olmayacağı ifade edilebilir.

Anahtar kelimeler: Seftriakson, embriyotoksisite, in ovo

ABSTRACT

This study aimed to determine the embryotoxic and teratogenic effects of ceftriaxone application at different doses to the air cell of fertile chicken eggs. Fertility was checked using an ovoscope under light on the 7th day of incubation of chicken eggs placed in the incubator, and non-fertile eggs were excluded from the study. A total of 210 fertile chicken eggs were randomly divided into seven groups of 30 eggs each. The eggs in the first group received no treatment, while the second group received 50 µL of physiological saline. On the same day, the other groups received ceftriaxone at doses of 250, 125, 62.5, 31.25, and 15.62 mg/kg in 50 µL volumes, respectively. At the end of the incubation period, eggs and chicks were examined. Statistically, no embryonic death or anatomical abnormalities were observed at the end of the study. In conclusion, it can be stated that the different doses of ceftriaxone used in the study will not cause embryotoxicity and teratogenicity in fertile chicken eggs.

Keywords: Ceftriaxone, embryotoxicity, in ovo

FULL TEXTS BOOK

GİRİŞ

Seftriakson üçüncü kuşak sefalosporin bir antibiyotiktir. Hücre duvarı sentezini engelleyerek bakterisit etki gösterir. Etki spektrumu içinde Pseudomonas spp, Pasteurella spp ve Enterobacteriaceae türü bakteriler bulunur. Gram negatif ve anerob bakterilere etkili olduğu dikkate alındığında üriner, genital ve sindirim sistemi enfeksiyonlarında kullanım alanı bulur. Ayrıca anerob bakterilere etkili olduğu için irinli-apseli enfeksiyonlarda da etkili olur. Kan-beyin bariyerini aşar ve beta-laktamazlara oldukça dirençlidir. İlacın veteriner sahada satışa sunulan preparatı bulunmamaktadır. Ancak beşerî preparatları pet kliniğinde veteriner hekimlerce etiket dışı olarak kullanılmaktadır (**Yazar 2021, Yazar 2023, Yazar 2024**). Seftriakson gebelikte kategori B olarak değerlendirilmekle (**Emiroğlu ve Sakin 2022**) birlikte, yüksek dozlarda gebelikte güvenliği ile bilgiler bulunmamaktadır.

Veteriner hekimlikte her hayvan türü için gebelikte ilaç güvenilirliği ile ilgili kategori bulunmamakla birlikte, beşerî hekimlikte gebelikte ilaç güvenilirliği farklı kategorilerde (A, B, C, D, X) tanımlanmaktadır. Kategori A ilaçlar güvenli kabul edilirken, B kategori ilaçların deney hayvanlarında fötal risk oluşturmadığı kabul edilir (**Ozturk 2018**). Embriyotoksisite farklı çeşitlerde çalışılmakla birlikte, fertil tavuk yumurtalarında da çalışılabilmektedir (**Jelinek ve ark 1985**). Veteriner sahada kullanılan birçok ilacın embriyotoksik, teratojenik veya malformatif etkilerinin belirlenmesinde fertil tavuk yumurtası kullanılmıştır (**Canbar ve ark 2021a, Canbar ve ark 2021b, Uslu ve ark 2022, Canbar ve ark 2023, Uslu ve ark 2024**). Belirli şartlarda fertil tavuk yumurtalarının ön tarama testi olarak kullanılabileceği ifade edilmiştir (**Hill ve Hoffman 1984, Kotwani 1998).** Embriyotoksisite çalışmalarında fertil tavuk yumurtaları, birinci basamakta embriyotoksik doz sınırı belirlemek (CHEST-I; Tavuk Embriyotoksisite Tarama Testi) için yapılırken, teratojenik parametrelerin belirlenmesi için ikinci (CHEST-II) aşama testi yapılmaktadır (**Özparlak 2015**).

Mevcut çalışmada fertil tavuk yumurtaları hava kamarasına farklı dozlarda seftriakson uygulamasının embriyotoksik ve teratojenik etkilerinin belirlenmesi amaçlandı.

MATERYAL VE METOT

Araştırmada ticari bir işletmeden (Anadolu Entegre Damızlık Hay. Pet. Tur. San. Tic. Ltd. Şti., Konya, Türkiye) temin edilen fertil Tetra L Superb ırkı tavuk yumurtaları kullanıldı. Araştırma prosedürü Selçuk Üniversitesi Veteriner Fakültesi Etik kurulunca onaylandı (SUVDAMEK, 2025/89). Çalışma süresince yumurtalar 37.7±0.2 ^{0}C ve %55±0.2 bağıl nem bulunan kuluçka makinesinde (İmza Teknik, Konya, Türkiye) kuluçka sürelerini tamamladı. Kuluçka süresince her 90 dakikada bir yumurta pozisyonlarının otomatik olarak değiştirilmesi sağlandı. Kuluçkanın 7. gününde ovoskop ile ışık altında fertilite kontrolü yapıldı ve fertil olmayan yumurtalar ayıklanarak her bir grupta 30 adet fertil yumurta olacak şekilde rastgele gruplandırıldı. Uygulamanın yapılabilmesi için yumurtaların hava kamaralarının bulunduğu küt uçları %80 etil alkol ile temizlenerek yumurta delici (Bai Shun, Zhejilang, Çin) ile delindi. Seftriakson (Desefin enj., Deva Holding AŞ, İstanbul, Türkiye) %0.9 NaCl solüsyonunda çözdürüldü ve aşağıda belirtilen dozlarda hava keselerine uygulandı. Uygulamayı takiben açılan delikler parafin ile kapatılarak tekrar kuluçka makinesine yerleştirildi.

Deneysel Dizayn ve Uygulamalar

Metot: Çalışmada, ortalama ağırlıkları 50 gram olan toplam 210 adet fertil tavuk yumurtası kullanıldı ve her grupta 30 adet olacak şekilde rastgele 7 gruba ayrılarak kuluçka makinesine yerleştirildi.

1. Grup (K, n:30): Bu gruptaki yumurtalar herhangi bir işleme tabi tutulmadan kuluçka makinesinde kuluçka sürelerini tamamladı. Bu grup Negatif kontrol grubu olarak kullanıldı.

FULL TEXTS BOOK

2. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 50 µL hacminde etken maddenin çözücüsü olan %0.9 NaCl solüsyonu uygulandı ve Pozitif kontrol grubu olarak değerlendirildi.

3. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 250 mg/kg dozunda seftriakson enjekte edildi.

4. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 125 mg/kg dozunda seftriakson enjekte edildi.

5. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 62.5 mg/kg dozunda seftriakson enjekte edildi.

6. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 31.25 mg/kg dozunda seftriakson enjekte edildi.

7. Grup (n:30): Bu gruptaki yumurtaların hava kamaralarına 7. günde açılan delikten 15.62 mg/kg dozunda seftriakson enjekte edildi.

Kuluçka süresinin tamamlanmasıyla (21. gün) yumurtalar açılarak embriyotoksik doz sınırı ve letal doz (LD)50 oranı hesaplamalarında kullanılmak için canlı ve ölü embriyoların sayısı belirlendi. Yapılan kontrollerde herhangi bir anomaliye rastlanılmadı.

Doza bağlı ölüm gözlenmediği için letal doz $(LD)_{50}$ değeri probit testi ile belirlenemedi. Embriyonal ölümler ki-kare testi ile değerlendirildi (SPSS 29.0). $P<0.05$ değeri istatistiki sınır olarak kabul edildi.

BULGULAR

Seftriaksonun fertil tavuk yumurtasına etkisi Tablo 1'de sunuldu. Uygulanan dozların neden olduğu embriyonal ölümler arasında istatistiki fark belirlenmedi ($P>0.05$).

Tablo 1. Seftriakson uygulanan fertil tavuk yumurtalarında embriyonik ölüm oranları.

	EEÖ	GEÖ	Canlı	Toplam	%*	GÖO
Kontrol	0	0	30	30	100.00	----
SF	0	0	30	30	100.00	0.00
250 mg/kg	2	1	27	30	90.00	10.00
125 mg/kg	0	0	30	30	100.00	0.00
62.5 mg/kg	0	2	28	30	93.33	6.67
31.25 mg/kg	1	0	29	30	96.67	3.33
15.62 mg/kg	2	1	27	30	90.00	10.00

*Aynı sütunda istatistiki fark belirlenmedi ($P>0.05$). SF: Serum fizyolojik, EEÖ: Erken embriyonik ölüm, GEÖ: Geç embriyonik ölüm, GÖO: Gerçek ölüm oranı.

TARTIŞMA

Üçüncü kuşak sefalosporin olan seftriakson özellikle Pseudomonas spp, Pasteurella spp ve Enterobactericeae türü bakteriler gibi gram negatif bakteriler ve anerob bakterilere etkinlik gösterir.

FULL TEXTS BOOK

İlacın veteriner sahada ruhsatlı preparat bulunmamakla birlikte, pet kliniğinde veteriner hekimlerce etiket dışı olarak kullanılmaktadır (**Yazar 2021, Yazar 2023, Yazar 2024**).

Bu çalışmada kullanılan dozlarda seftriakson uygulamasının fertil tavuk yumurtalarında embriyotoksisite ve teratojeniteye neden olmadığı belirlendi (P>0.05, Tablo 1). Seftriakson gebelikte kategori B olarak değerlendirilmektedir (**Emiroğlu ve Sakin 2022**). Sefalosporin grubu ilaçlar genellikle gebe hastalarda kullanım için güvenli kabul edilir. Ancak yenidoğanda kernikterusa neden olabileceği bildirilmiştir (**Nguyen ve ark 2025**). Gebe kadınlarda seftriakson kullanımına ilişkin detaylı veri bulunmamaktadır. Seftriaksonun plasentayı geçtiği ve hayvan çalışmalarında reprodüktif toksisiteye neden olmadığı bildirilmiştir (**EMA 2025**). Gebe fareler ve sıçanlarda normal insan dozunun 20 katına kadar dozda seftriakson uygulandığında, embriyotoksisite, fötotoksisite veya teratojenite gözlenmemiştir. Primatlarda ise insan dozunun yaklaşık 3 katı dozda uygulandığında embriyotoksisite veya teratojenite gözlenmediği bildirilmiştir. Ancak, gebe kadınlarda yeterli ve iyi kontrollü çalışmalar bulunmamaktadır. Gebe hayvanlarda yapılan çalışmaların net olarak insanları yansıtmadığı için gebelikte mutlak gerektiğinde kullanımı önerilmektedir (**Sandoz 2013, FDA 2024**). Zebra balıkları embriyo ve larvaları üzerinde seftriaksonun olumsuz etkilere neden olmadığı belirlenmiştir (**de Oliveira ve ark 2023**). Ancak gebe kadınlar üzerinde yapılan çalışmada betalaktamların da teratojenik potansiyele sahip olabileceği bildirilmiştir. Sefaklor (ikinci kuşak) ve sefaleksinin (birinci kuşak) malformasyona neden olabileceği, setriaksonun (üçüncü kuşak) malformasyona neden olmadığı belirlenmiştir **(Eric ve Sabo 2008).** Gebe farelere seftriakson uygulaması sonrasında anne bağırsak mikrobiyotasını değiştirdiği, bu durumun annelerin ve yavrularının bağışıklığını değiştirebileceği ifade edilmiştir (**Cheng ve He 2019**). Diğer yandan yapılan incelemede seftriakson ile kardiyak malformasyon arasında bir ilişki bulunabileceği bildirilmiştir (**Malleswari ve ark 2024**).

SONUÇ

Sonuç olarak bu çalışmada kullanılan dozlarda seftriakson uygulamasının fertil tavuk yumurtalarında embriyotoksisite ve teratojeniteye neden olmadığı ifade edilebilir.

KAYNAKLAR

1. Canbar R, Akcakavak G, Uslu M, Bas AL, 2021a. Determination of embryotoxic effects of Tarantula cubensis alcoholic extract with in ovo model. Magy Allatorvosok Lapja, 143:497-504.
2. Canbar R, Uslu M, Arslan MS, Kızılay H, 2023. Determination of embryotoxic effects of atipamezole using in ovo model. Rev Cient, 33:rcfcv-e33230.
3. Canbar R, Uslu M, Tufan Ö, Yazar E, 2021b. Detomidinin embriyotoksik doz sınırının in-ovo model ile belirlenmesi. Eurasian J Vet Sci, 37:71-75.
4. Cheng R, Guo J, Zhang Y, Cheng G, Qian W, Wan CM, Li M, Marotta F, Shen X, He F, 2021. Impacts of ceftriaxone exposure during pregnancy on maternal gut and placental microbiota and its influence on maternal and offspring immunity in mice. Exp Anim, 70(2):203-217.
5. de Oliveira EM, Sousa-Moura D, Grisolia CK, Franco-Bernardes MF, 2023. Evaluation of embryo-larval toxicity of the antibiotic ceftriaxone in the fish danio rerio (zebrafish). Ecotoxicol Environ Contam, 18:40-46.
6. EMA 2025. List of the names, pharmaceutical forms, strengths of the medicinal products, routes of administration, applicants, marketing authorisation holder in the member states. https://www.ema.europa.eu/en/documents/referral/ceftriaxone-tyrol-pharma-article-29-referral-annex-i-ii-iii_en.pdf. Erişim tarihi:23.04.25
7. Emiroğlu SB, Sakin F, 2022. Gebelikte ilaç kullanımı ve teratojenite, Veteriner Farmakoloji ve Toksikoloji Derneği Bülteni, 13:90-98.

FULL TEXTS BOOK

8. Eric M, Sabo A, 2008. Teratogenicity of antibacterial agents. Coll Antropoli, 32:919-925.
9. FDA 2024. Ceftriaxone for injection. https://www.efda.gov.et/wp-content/uploads/2024/02/Ceftriaxone-for-Injection-USP-1-g_THEOXONE-1G_Theon-Pharmaceuticals-Limited.pdf. Erişim tarihi:23.04.25
10. Hill E, Hoffman D, 1984. Avian models for toxicity testing. J Am Coll Toxicol, 3(6):357–376.
11. Jelinek R, Peterka M, Rychter Z, 1985. Chick embryotoxicity screening test – 130 substances tested. Indian J Exp Biol, 23:588-595.
12. Kotwani A, 1998. Use of chick embryo in screening for teratogenicity. Indian J Physiol Pharmacol, 42:189-204.
13. Malleswari K, Reddy DRB, Raju GS, 2024. Antibiotics effects on pregnancy. IJSDR, 9:82-87.
14. Nguyen J, Madonia V, Bland CM, Stover KR, Eiland LS, Keating J, Lemmon M, Bookstaver PB, 2025. A review of antibiotic safety in pregnancy-2025 update. Pharmacotherapy, 45(4):227-237.
15. Özparlak H, 2015. The use of chick embryos in embryotoxicity and teratogenicity tests. SUFEFD, 40:13-22.
16. Ozturk Z, 2018. Medication use in pregnancy and risk evaluation: do not categorize me. Zeynep Kamil Med J, 49(1):109-112.
17. Sandoz 2013. Ceftriaxone for injection. https://www.accessdata.fda.gov/drugsatfda_docs/label/2014/065169s022lbl.pdf. Erişim tarihi:23.04.25
18. Uslu M, Canbar R, Akcakavak G, Yazar E, 2022. Determination of the embryotoxic effect of maropitant using an in ovo model. Pol J Vet Sci, 25:357-359.
19. Uslu M, Canbar R, Arslan MS, Kızılay H, 2024. Determination of the embryotoxic effect of metronidazole using an in ovo model. Rev Cient, 34:34310.
20. Yazar E, 2021. Köpeklerde Tedavi El Kitabi. Nobel Tıp Kitabevleri Ltd. Şti., Ankara, Türkiye.
21. Yazar E, 2023. Kedilerde Tedavi El Kitabı. Atlas Akademik Basim Yayin Dağitim Tic. Ltd. Şti., Ankara, Türkiye.
22. Yazar E, 2024. Kemoterapötikler, in: Veteriner İlaç Rehberi Tedavi El Kitabı, Ed: Yazar E, Nobel tıp kitabevleri, Ankara, Türkiye.

FULL TEXTS BOOK

DİJİTAL DÖNÜŞÜMÜN TOPLUMSAL VE SİYASAL ETKİLERİ
SOCIAL AND POLITICAL EFFECTS OF DIGITAL TRANSFORMATION

Assoc.Prof.Dr.Cüneyt Yılmaz

St. Clements University , Department of Political Science, London , UK

https://orcid.org/0000-0003-1655-588X

ÖZET

Bu makale, dijital dönüşümün toplumsal ve siyasal yapı üzerindeki etkilerini kültürel kimlikler, iletişim teknolojileri ve politik süreçler bağlamında ele almaktadır. Dijital dönüşüm, bireylerin yaşam biçimlerini, sosyal ilişkilerini ve siyasal katılım biçimlerini kökten değiştirmiştir. Özellikle çevrimiçi platformlarda şekillenen kültürel kimlikler, sadece bireysel düzeyde değil, aynı zamanda toplumsal normlar ve siyasal dinamikler üzerinde de etkili olmaktadır.

Makale, "ağ toplumu" kavramı çerçevesinde çevrimiçi kimlik oluşumunun sosyal etkileşimleri nasıl dönüştürdüğünü ve dijital ortamların kimlik çeşitliliğini nasıl etkilediğini analiz etmektedir. Sosyal medya platformlarının hem politik katılımı artırdığı hem de bilgi kirliliği ve kutuplaşma gibi sorunları beraberinde getirdiği vurgulanmaktadır. Dijital kimliklerin, yerel ve küresel kültürlerle etkileşim hâlinde sürekli yeniden şekillendiği, bu durumun ise kültürel çeşitlilik açısından hem fırsatlar hem de tehditler barındırdığı belirtilmektedir.

Siyasal iletişimde dijital medyanın rolü, geleneksel medya yapılarının zayıflamasıyla daha da belirginleşmiştir. Sosyal medya, bireylerin siyasal içerik üretmesini, yaymasını ve bu yolla demokratik süreçlere katılımını kolaylaştırırken, aynı zamanda “yankı odaları” ve algoritmik filtreleme yoluyla toplumsal bölünmeleri derinleştirebilmektedir. Makale, ayrıca dijital eşitsizlik, erişim farkları ve okuryazarlık gibi meseleleri de analiz ederek daha kapsayıcı bir dijital toplum için politika önerileri sunmaktadır.

Sonuç olarak, dijital dönüşümün çok katmanlı etkileri, bireylerin kimliklerinden devletlerin politika üretme biçimlerine kadar geniş bir yelpazeye yayılmaktadır. Bu etkilerin yönetilebilmesi için disiplinlerarası iş birliği, etik dijital politikalar ve dijital eşitliğe yönelik stratejilerin geliştirilmesi gerekmektedir.

Anahtar Kelimeler: Çevrimiçi kimlikler, Kültürel çeşitlilik, Dijital dönüşüm, Siyasal etkiler

ABSTRACT

This article examines the societal and political impacts of digital transformation through the lens of cultural identity, communication technologies, and political processes. Digital transformation has fundamentally reshaped individuals' lifestyles, social relationships, and forms of political participation. In particular, cultural identities formed through online platforms influence not only the personal sphere but also broader social norms and political dynamics.

Framed within the concept of the “network society,” the article analyzes how online identity formation transforms social interactions and how digital environments affect identity diversity. It emphasizes that social media platforms both enhance political participation and present challenges such as misinformation, echo chambers, and polarization. Digital identities are portrayed as dynamic and continuously reconstructed through interactions with both local and global cultures, bringing opportunities as well as risks for cultural diversity.

FULL TEXTS BOOK

The role of digital media in political communication has become more prominent with the decline of traditional media structures. Social media enables individuals to create and disseminate political content and facilitates more direct democratic participation. However, algorithmic filtering and personalized content may deepen ideological divisions and limit exposure to diverse perspectives. The article also addresses issues of digital inequality, access disparities, and digital literacy, proposing policy recommendations for a more inclusive digital society.

In conclusion, the multifaceted effects of digital transformation extend from individual identity construction to state-level policy-making. Addressing these impacts requires interdisciplinary collaboration, ethical digital governance, and strategic efforts toward digital equity and inclusion.

Keywords: Online identities, Cultural diversity, Digital transformation, Political impacts

GİRİŞ

Dijital dönüşüm, iletişim teknolojilerindeki hızlı değişim ve gelişmelerin, toplumların siyasal ve toplumsal yapıları üzerinde derin etkiler yarattığı bir süreci ifade eder. Bu dönüşüm, post-modern çağın getirdiği gelişmelerle birlikte ortaya çıkmıştır. İnternetin yaygınlaşması ve iletişim teknolojilerindeki ilerlemeler, insanların günlük yaşamlarını dijital platformlara taşımasını sağlamıştır. "Ağ toplumu" kavramı, bu yeni toplumsal dönemin anahtar özelliğidir.

Ağ toplumu, insanların günlük yaşamlarını internet ağları üzerinde şekillendirdiği, iletişimin dijitalleştiği ve bilişim teknolojilerinin hayatın her alanında yaygın olarak kullanıldığı bir yapıyı ifade eder. Bu yeni toplumsal yapı, kimlik oluşumundan sosyal ilişkilere kadar birçok alanda derin değişimlere neden olmuştur.

İletişim, dijital dönüşümün toplumsal ve siyasal etkilerini anlamak için kritik bir araçtır. Politik iletişim, bireylerin politik düşüncelerini ve tutumlarını şekillendirirken, toplumsal normları ve davranış kalıplarını da belirlemektedir. Araştırmalar, iletişimin toplumun genel politik kültürünü ve normlarını belirlemede önemli bir rol oynadığını göstermektedir (Zaller, 1992). Bu bağlamda, dijital iletişimin toplumsal ve siyasal etkilerini anlamak, günümüzün karmaşık ve dinamik toplumlarını anlamak için önemlidir.

Medyanın rolü de göz ardı edilmemelidir. Medya, toplumun genel görüşlerini ve politik bilincini etkilerken, aynı zamanda toplumsal normları belirleme sürecinde de önemli bir etkiye sahiptir (Entman, 2012). Dolayısıyla, medyanın siyasal iletişim üzerindeki etkisi, toplumun politik yönelimlerini belirleme ve politik süreçlere katılımı etkileme konusunda kritik bir rol oynamaktadır.

Dijital dönüşümün toplumsal ve siyasal etkilerini anlamak, günümüzün karmaşık ve dinamik toplumlarını anlamak için kritik bir öneme sahiptir. Bu bağlamda, iletişim teknolojilerindeki hızlı değişim ve gelişmelerin, toplumların siyasal ve toplumsal yapıları üzerindeki etkilerini daha derinlemesine incelemek, geleceğe dönük politika oluşturmak ve toplumsal dönüşümü daha iyi yönetmek için gereklidir.

Bu makale, çevrimiçi platformlardaki kültürel kimliklerin oluşumu ve bu kimliklerin siyasal ve toplumsal dinamiklere etkilerini incelemektedir. Özellikle, dijital dönüşümün kültürel çeşitliliğe nasıl yansıdığı ve çevrimiçi kimlik politikalarının bu süreçteki rolü incelenecektir. Bu bağlamda, çevrimiçi platformlardaki kültürel kimliklerin oluşumuyla ilgili siyasal ve toplumsal sonuçlar değerlendirilerek, çeşitliliğin korunması veya baskılanmasının siyasi ve toplumsal önemi tartışılacaktır.

AMAÇ

Bu makale, çevrimiçi platformlardaki kültürel kimliklerin oluşumu ve bu kimliklerin siyasal ve toplumsal dinamiklere etkilerini daha derinlemesine incelemeyi amaçlamaktadır. Dijital dönüşüm,

FULL TEXTS BOOK

günümüz toplumlarında belirgin bir şekilde hissedilmekte olup, internet ve iletişim teknolojilerindeki ilerlemeler, toplumsal yapıyı kökten değiştirmektedir. Bu değişimler, kültürel çeşitliliğin daha da arttığı bir ortamda gerçekleşmektedir.

Ağ toplumu kavramı, sosyal ilişkilerin dijital teknolojiler aracılığıyla yürütüldüğü ve çevrimiçi kimliklerin giderek daha fazla önem kazandığı bir dönemi ifade etmektedir. Bu dönemde, bireylerin çevrimiçi platformlardaki kimlikleri, günlük hayatlarında kendilerini ifade etme biçimlerini ve sosyal ilişkilerini belirlemektedir. Dolayısıyla, çevrimiçi platformlardaki kültürel kimliklerin oluşumu ve bu kimliklerin siyasal ve toplumsal etkilere olan katkıları önemli bir araştırma alanıdır.

Bu bağlamda, bu makalenin amacı, çevrimiçi platformlardaki kültürel kimliklerin oluşumu ve bu kimliklerin siyasal ve toplumsal dinamiklere etkilerini anlamak ve analiz etmektir. Bu analiz, dijital dönüşümün kültürel çeşitlilik üzerindeki etkisini anlamak ve çeşitliliğin korunması veya baskılanmasının siyasal ve toplumsal önemini vurgulamayı amaçlamaktadır.

Dijital dönüşümün kültürel kimlikler üzerindeki etkilerini anlamak, siyasi karar alıcılar için de, önemli bir zorluk ve fırsattır. Bu makale, çevrimiçi platformlardaki kültürel kimliklerin ve politikalarının siyasal ve toplumsal bağlamlarda daha iyi anlaşılmasına katkı sağlamayı amaçlamaktadır.

BULGULAR

Dijital dönüşüm, iletişim teknolojilerindeki hızlı ilerlemelerin toplumsal yapıları derinden etkileyen bir fenomen olarak tanımlanabilir. Castells (1996), "ağ toplumu" kavramını kullanarak, internetin yaygınlaşmasıyla birlikte insanların yaşam biçimlerinin ve toplumsal ilişkilerinin temelde değiştiğini ifade etmiştir. Ancak, dijital dönüşümün sadece teknolojik değişikliklerle sınırlı olmadığını belirtmek önemlidir.

Jenkins (2006) ve Turkle (2011) gibi araştırmacılar, dijital medyanın kültürel ve sosyal alanlarda önemli etkileri olduğunu savunmaktadır. Örneğin, dijital platformlar, katılımcı kültürleri teşvik ederek toplumsal katılımı artırabilir ve bireyler arasındaki ilişkileri dönüştürebilir. Ayrıca, Hess ve Ostrom (2007) gibi ekonomistler, dijital dönüşümün ekonomik boyutunu vurgulayarak, geleneksel iş modellerini değiştirerek yeni ekonomik fırsatlar ve zorluklar sunduğunu belirtmektedir.

Siyasal açıdan, dijital dönüşüm demokratik süreçleri ve siyasal katılımı derinden etkilemektedir. Chadwick (2013), sosyal medyanın yükselişinin siyasal iletişim ve kampanya stratejilerini önemli ölçüde değiştirdiğini ve siyasal aktörlerin dijital platformları etkin bir şekilde kullanmasının gerekliliğini vurgulamaktadır. Ancak, Tufekci (2017) gibi araştırmacılar, sosyal medyanın kötüye kullanılması ve bilgi kirliliği gibi sorunların demokratik süreçleri olumsuz yönde etkileyebileceğini öne sürmektedir.

Medyanın siyasal süreçler üzerindeki etkisi, siyasal liderlerin ve partilerin imajını şekillendirme, siyasi olayları yorumlama ve analiz etme gibi çeşitli şekillerde ortaya çıkar. Chavez (2019) ve Milton (2020) gibi araştırmacılar, medyanın siyasal katılımı artırabileceğini ve toplumsal değişimi teşvik edebileceğini belirtmektedir.

Bu bağlamda, dijital dönüşümün kültürel, ekonomik ve siyasal boyutlarını bir arada ele almak önemlidir. Bu alanlardaki etkileşimlerin karmaşıklığını anlamak, gelecekteki toplumsal değişimleri daha iyi öngörebilmemize olanak sağlayabilir. Bu nedenle, gelecekte yapılacak araştırmaların, dijital dönüşümün toplumsal ve siyasal alanlardaki etkilerini daha kapsamlı bir şekilde ele alması gerekmektedir.

YÖNTEM

Bu çalışma, mevcut literatürün derinlemesine incelenmesi yoluyla gerçekleştirilmiştir. Nitel araştırma yöntemi kullanılarak, dijital dönüşümün siyasal ve toplumsal etkileri üzerine yapılan çalışmalar incelenmiş ve temel bulguların belirlenmesi amaçlanmıştır. Bu yöntem, çevrimiçi platformlardaki kültürel kimliklerin oluşumu ve bu kimliklerin siyasal/toplumsal etkileri üzerine derinlemesine bir anlayış sağlamak için uygundur.

FULL TEXTS BOOK

Dijital Dönüşüm ve Ağ Toplumu

Dijital dönüşüm, bilgi ve iletişim teknolojilerinin hızlı gelişimi ve yaygınlaşması sonucu, bireylerin ve toplumların yaşam tarzlarını, iş yapma biçimlerini ve sosyal etkileşimlerini köklü bir şekilde değiştiren bir süreçtir. Bu süreç, post-modern çağın getirdiği teknolojik ilerlemelerle birlikte, internetin günlük hayatın ayrılmaz bir parçası haline gelmesiyle hız kazanmıştır.

"Ağ toplumu" kavramı, dijital dönüşüm sürecinin toplumsal yapılar üzerindeki etkilerini tanımlamak için kullanılır. Bu kavram, insanların dijital ağlar üzerinden birbirleriyle etkileşime girdiği, bilgiye ve hizmetlere erişimin büyük ölçüde dijital platformlar aracılığıyla sağlandığı bir toplumsal yapıyı ifade eder. Ağ toplumu, Manuel Castells tarafından popülerleştirilmiş olup, dijital çağın en belirgin sosyal örgütlenme biçimlerinden biri olarak kabul edilir (Castells, 2010).

Ağ toplumunun en belirgin özelliklerinden biri, bireylerin günlük yaşamlarını dijital platformlar üzerinde şekillendirmesidir. İnternet, sosyal medya, e-ticaret siteleri ve diğer dijital araçlar, insanların bilgi edinme, iletişim kurma, alışveriş yapma ve sosyalleşme biçimlerini kökten değiştirmiştir (Van Dijk, 2020). Bu değişim, bireylerin zaman ve mekândan bağımsız olarak etkileşimde bulunabilmesini sağlar. Örneğin, sosyal medya platformları sayesinde insanlar, coğrafi olarak uzak olsalar bile sürekli olarak iletişimde kalabilir ve ortak ilgi alanlarına sahip kişilerle kolayca etkileşim kurabilir.

Dijital dönüşümün sosyal ilişkiler üzerindeki etkisi de oldukça derindir. Geleneksel yüz yüze iletişim biçimlerinin yerini büyük ölçüde dijital iletişim almıştır. E-posta, anlık mesajlaşma uygulamaları ve video konferans teknolojileri, iş hayatından özel yaşama kadar geniş bir yelpazede kullanılmaktadır (Baym, 2015). Bu durum, sosyal ilişkilerin hızla ve geniş bir kapsamda kurulmasını ve sürdürülmesini mümkün kılar. Ancak, aynı zamanda yüz yüze etkileşimlerin azalması ve dijital ortamlarda yaşanan sosyal izolasyon gibi yeni zorlukları da beraberinde getirmiştir.

Ağ toplumunda kimlik oluşumu da dijital dönüşümün etkisi altında yeniden şekillenmiştir. Çevrimiçi platformlar, bireylerin kendilerini ifade etmeleri ve farklı kimlikler geliştirmeleri için geniş fırsatlar sunar. Sosyal medya profilleri, bloglar ve diğer dijital içerik oluşturma araçları, bireylerin dijital kimliklerini inşa etmelerine olanak tanır.

Bu dijital kimlikler, bireylerin çevrimdışı kimliklerinden farklı olabilir ve sürekli olarak değişim gösterebilir. Dijital kimliklerin dinamik yapısı, bireylerin çeşitli sosyal ve kültürel bağlamlarda kendilerini farklı şekillerde ifade etmelerine imkân tanır (Turkle, 1995).

Dijital dönüşüm, bilgiye ve teknolojiye erişimdeki eşitsizlikleri de gündeme getirmiştir. Dijital uçurum olarak adlandırılan bu durum, bazı bireylerin ve toplulukların dijital teknolojilere erişimde yaşadıkları sınırlamaları ifade eder (Van Dijk, 2005). Bu eşitsizlik, eğitim, gelir düzeyi ve coğrafi konum gibi faktörlerden kaynaklanabilir ve toplumsal yapılar üzerinde derin etkiler yaratabilir. Dijital uçurum, toplumsal katılımı ve fırsat eşitliğini olumsuz yönde etkileyerek, mevcut sosyal ve ekonomik eşitsizlikleri daha da derinleştirebilir.

Dijital dönüşüm süreci, toplumsal ve siyasal yapıları etkilemeye devam edecektir. Teknolojik yeniliklerin hız kesmeden devam etmesi, ağ toplumunun özelliklerini ve dinamiklerini sürekli olarak yeniden şekillendirecektir. Bu süreçte, dijital teknolojilerin etik kullanımı, dijital eşitliğin sağlanması ve bireylerin dijital becerilerinin geliştirilmesi konuları önemli gündem maddeleri olarak öne çıkacaktır. Dijital dönüşümün olumlu etkilerinden tam anlamıyla faydalanabilmek ve olumsuz etkilerini en aza indirmek için, politika yapıcıların ve toplumun tüm kesimlerinin bu değişim sürecine uyum sağlaması gerekecektir.

İletişim Teknolojileri ve Toplumsal Yapı

Dijital dönüşüm, iletişim teknolojilerindeki hızlı değişim ve gelişmelerin, toplumların sosyal yapıları üzerinde derin etkiler yarattığı bir süreci ifade eder. Bu dönüşüm, post-modern çağın getirdiği

FULL TEXTS BOOK

gelişmelerle birlikte ortaya çıkmış ve internetin yaygınlaşması ile hız kazanmıştır. İletişim teknolojilerinin gelişimi, günlük yaşamın her alanını dijital platformlara taşımış, "ağ toplumu" kavramını öne çıkarmıştır (Castells, 2010). Ağ toplumu, bireylerin dijital ağlar üzerinden etkileşime geçtiği, bilgi ve hizmetlere erişimin dijital platformlar aracılığıyla sağlandığı bir yapıyı tanımlar.

Dijital iletişimin sosyal ilişkiler üzerindeki etkisi derindir ve geniş kapsamlıdır. Geleneksel yüz yüze iletişim biçimlerinin yerini büyük ölçüde dijital iletişim almıştır. E-posta, anlık mesajlaşma uygulamaları ve video konferans teknolojileri, hem iş hem de özel yaşamda yaygın olarak kullanılmaktadır. Bu durum, sosyal ilişkilerin hızla ve geniş kapsamda kurulmasını ve sürdürülmesini sağlar. Sosyal medya platformları, insanların coğrafi olarak uzak olsalar bile sürekli olarak iletişimde kalmalarını ve ortak ilgi alanlarına sahip kişilerle kolayca etkileşim kurmalarını mümkün kılar.

Ancak dijital iletişim aynı zamanda yüz yüze etkileşimlerin azalması ve dijital ortamlarda sosyal izolasyon gibi yeni zorlukları da beraberinde getirmiştir. Dijital platformlarda sürdürülen ilişkiler, geleneksel sosyal bağların zayıflamasına yol açabilir ve bireylerin yalnızlık hissini artırabilir (Turkle, 2011).

Ağ toplumunda kimlik oluşumu, dijital dönüşümün etkisi altında yeniden şekillenmiştir. Çevrimiçi platformlar, bireylerin kendilerini ifade etmeleri ve farklı kimlikler geliştirmeleri için geniş fırsatlar sunar. Sosyal medya profilleri, bloglar ve diğer dijital içerik oluşturma araçları, bireylerin dijital kimliklerini inşa etmelerine olanak tanır. Bu dijital kimlikler, bireylerin çevrimdışı kimliklerinden farklı olabilir ve sürekli olarak değişim gösterebilir. Dijital kimliklerin dinamik yapısı, bireylerin çeşitli sosyal ve kültürel bağlamlarda kendilerini farklı şekillerde ifade etmelerine imkân tanır.

Sosyal medya, toplumsal normların şekillenmesinde önemli bir rol oynar. Bu platformlar, bireylerin toplumsal olaylar ve konular hakkında anında tepki vermelerine ve geniş kitlelere ulaşmalarına olanak tanır. Sosyal medyanın bu özelliği, toplumsal hareketlerin ve protestoların hızla örgütlenmesini ve yayılmasını mümkün kılar (Tufekci, 2017).

Medya, toplumun genel görüşlerini ve politik bilincini etkilerken, aynı zamanda toplumsal normları belirleme sürecinde de önemli bir etkiye sahiptir (Entman, 2012). Sosyal medya platformlarında yayılan trendler ve popüler kültür unsurları, bireylerin davranışlarını ve sosyal etkileşimlerini etkileyebilir. Sosyal medyanın bu etkisi, toplumsal normların hızla değişmesine ve yeni normların oluşmasına yol açabilir. Ancak, sosyal medya aynı zamanda yanıltıcı bilgilerin ve dezenformasyonun hızla yayılma riskini de beraberinde getirir .

Dijital dönüşümün etkileri, sadece iletişim ve kimlik oluşumuyla sınırlı değildir; aynı zamanda toplumsal normlar, politik katılım ve kültürel değişim gibi alanları da içerir. Toplumların dijitalleşmesiyle birlikte, politik katılım ve toplumsal hareketlilik de dönüşmüş ve yeni bir boyut kazanmıştır. Sosyal medya platformları, toplumsal değişim taleplerini örgütlemekte ve bu talepleri yaymakta önemli bir rol oynamaktadır.

Ancak, dijital dönüşümün toplumsal etkileriyle ilgili endişeler de mevcuttur. Özellikle, sosyal medyanın demokratik süreçlere olan etkisi ve bilgi kirliliği gibi sorunlar, ciddi bir şekilde ele alınması gereken konulardır. Sosyal medyanın manipülatif kullanımı, siyasi karar alma süreçlerini etkileyebilir ve toplumsal kutuplaşmayı artırabilir.

Bu bağlamda, dijital dönüşümün toplumsal etkileri karmaşık ve çok yönlüdür. Bu süreç, iletişimde devrim niteliğinde değişikliklere yol açarken, aynı zamanda toplumsal normları, politik katılımı ve kültürel yapıları da derinden etkilemektedir. Bu nedenle, dijital dönüşümün toplumsal etkilerini anlamak ve yönetmek için çoklu disiplinler arası bir yaklaşım gerekmektedir. Bu sayede, dijitalleşmenin olumlu yönleri korunabilir ve olumsuz etkileri en aza indirilebilir.

FULL TEXTS BOOK

Politik İletişim ve Dijital Dönüşüm

Politik iletişim, siyasi aktörler ile medya ve halk arasındaki bilgi alışverişini kapsayan bir süreçtir. Bu süreç, siyasi mesajların iletilmesi, kamuoyu oluşturulması ve siyasi düşüncelerin yayılması gibi önemli işlevlere sahiptir. Özellikle demokrasinin işleyişinde kritik bir rol oynayan politik iletişim, siyasi katılımı artırma, kamuoyunu bilgilendirme ve politik hesap verebilirliği sağlama gibi önemli görevleri üstlenir.

Geleneksel politik iletişim, gazeteler, radyo ve televizyon gibi medya araçları üzerinden gerçekleştirilirdi. Ancak dijital dönüşüm ile birlikte bu süreç kökten değişti. İnternetin yaygınlaşması ve sosyal medya platformlarının yükselişi, politik iletişimi daha erişilebilir, hızlı ve interaktif hale getirdi.

Dijital platformlar, politik içerik üretimi ve yayılması için güçlü araçlar sunar. Sosyal medya, bloglar, video paylaşım siteleri ve podcast'ler gibi dijital ortamlar, bireylerin ve siyasi aktörlerin kendi politik mesajlarını oluşturup paylaşmalarını sağlar. Bu platformlar, kullanıcıların geniş kitlelere ulaşmalarını ve etkileşimde bulunmalarını kolaylaştırır.

Siyasi partiler, adaylar ve aktivist gruplar, dijital platformları kullanarak seçmenlere ve destekçilere doğrudan ulaşabilirler. Özellikle seçim dönemlerinde, dijital platformlar aracılığıyla yürütülen kampanyalar, geleneksel medyadan daha fazla etkili olabilir.

Dijital iletişim, bireylerin siyasi düşüncelerini ve tutumlarını şekillendirme konusunda büyük bir etkiye sahiptir. Sosyal medya, kullanıcıların siyasi haberlere ve bilgilere hızla erişimini sağlar ve bu bilgilerin geniş kitlelere yayılmasını hızlandırır. Ayrıca, dijital platformlar, kullanıcıların kendi görüşlerini ifade etmeleri ve toplumsal tartışmalara katılmaları için bir alan sunar.

Ancak dijital iletişimin bir dezavantajı da vardır: politik kutuplaşma ve "echo chamber" etkisi. Kullanıcılar, genellikle kendi görüşlerini paylaşan kişilerle etkileşime girerler ve bu durum, farklı görüşlere kapalı hale gelmelerine yol açabilir. Bu durum, politik düşüncelerin daha da radikalleşmesine ve toplumsal bölünmelere neden olabilir.

Öte yandan, dijital platformlar marjinalleştirilmiş grupların ve bireylerin seslerini duyurmalarına ve politik süreçlere katılmalarına olanak tanır. Bu durum, politik katılımın genişlemesini ve demokratik süreçlerin daha kapsayıcı olmasını sağlayabilir. Dijital iletişimin bu çift yönlü etkisi, modern demokrasilerde önemli bir rol oynamaktadır.

Politik iletişim, günümüzde dijital dönüşüm ile birlikte köklü bir değişim geçirmiştir (Chadwick, 2013). Özellikle sosyal medya platformlarının yükselişi, siyasi aktörlerin ve vatandaşların iletişim biçimlerini ve etkileşimlerini dönüştürmüştür. Bu platformlar, politik katılımı artırmada ve siyasi mesajların yayılmasında önemli bir rol oynamaktadır.

Dijital iletişimin politik süreçlere etkisi giderek artmakta ve siyasi katılımın şeklini derinden değiştirmektedir (Vaccari & Valeriani, 2016). Sosyal medya platformları, kullanıcıların siyasi içeriklere erişimini kolaylaştırırken, aynı zamanda bireylerin kendi politik görüşlerini ifade etmelerini ve toplumsal tartışmalara katılmalarını teşvik etmektedir. Bu bağlamda, dijital iletişim politik süreçlerin demokratikleşmesinde önemli bir araç olarak kabul edilmektedir.

Medyanın Rolü ve Siyasal Etkileri

Medyanın toplumsal ve siyasal normları belirlemedeki rolü, geleneksel ile dijital medya arasında önemli bir değişim ve dönüşüm yaşamaktadır. Geleneksel medya, televizyon, radyo ve gazeteler aracılığıyla belirlenen normlar, toplumun genel kabul görmüş değerlerini ve davranış kalıplarını şekillendirirken, dijital medya bu süreci kökten değiştirmiştir.

FULL TEXTS BOOK

Özellikle, sosyal medya platformları aracılığıyla bireylerin ve grupların politik içerikler üretmesi ve yayması, bilgiye erişimi daha demokratik hale getirmiş ve geleneksel medya tekellerini zayıflatmıştır (Castells, 2008).

Dijital medyanın kamuoyu oluşumundaki etkisi, bilginin hızlı ve geniş kitlelere ulaşmasını sağlamıştır. Sosyal medya platformları, bireylerin kendi haber kaynaklarını seçmelerine ve kişiselleştirilmiş içerik almalarına imkân tanırken, bu durum aynı zamanda "yankı odası" ve "filtre balonu" etkilerine de yol açabilir. Bu etkiler, bireylerin kendi görüşlerini destekleyen bilgilere daha fazla maruz kalmasına ve karşıt görüşlere daha az temas etmesine neden olabilir (Pariser, 2011).

Ancak, dijital medya aynı zamanda marjinalize edilmiş grupların seslerini duyurmalarına ve alternatif görüşlerin yayılmasına da olanak tanır. Bu özellik, dijital medyanın kamuoyu üzerinde daha çeşitlilik ve çoğulculuk getiren bir etki yaratmasını sağlar (Benkler, 2006).

Medyanın politik süreçlere katılımı teşvik etme gücü, demokrasilerde kritik bir öneme sahiptir. Geleneksel ve dijital medya, vatandaşları bilgilendirerek, siyasi farkındalık yaratarak ve katılımı teşvik ederek demokratik süreçlerin güçlenmesine katkıda bulunur. Özellikle, dijital medya gençler ve politik süreçlere geleneksel olarak katılmayan gruplar için daha erişilebilir bir platform sunar. Sosyal medya kampanyaları, çevrimiçi tartışmalar ve dijital aktivizm, bu grupların politik süreçlere katılımını artırabilir (Loader & Mercea, 2011).

Medyanın evrimiyle birlikte, toplumsal ve siyasal normların belirlenmesindeki rolü de değişime uğramıştır. Geleneksel medya, belirli bir ölçüde tek taraflı bir görüş sunma eğilimindeydi ve bu da toplumsal normların ve politik görüşlerin belirlenmesinde belirli bir etkiye sahipti. Ancak, dijital medya ve sosyal medya platformları, bireylerin daha geniş bir yelpazede bilgiye erişimini sağlamış ve kendi düşüncelerini ifade etmelerini kolaylaştırmıştır. Bu durum, medyanın belirleme gücünü demokratikleştirirken, aynı zamanda bireylerin kendi bilgi alışveriş ağlarını oluşturmalarına ve kendi düşünce yapılarına uygun içerikleri tercih etmelerine de olanak tanır. Ancak, bu durum bazı zorlukları da beraberinde getirir. Özellikle sosyal medya, kullanıcılarına sıklıkla kendi görüşlerini destekleyen içerikleri sunar ve bu da bireylerin farklı görüşlere kapalı hale gelmelerine yol açabilir. Bu nedenle, medyanın toplumsal ve siyasal etkileri, demokratik katılımın teşvik edilmesi ve çeşitliliğin sağlanması gibi olumlu yanlarının yanı sıra, kutuplaşma ve bilgi kısıtlılığı gibi olumsuz sonuçları da içerir.

Dijital Kimlikler ve Kültürel Çeşitlilik

Dijital kimliklerin ve kültürel çeşitliliğin bir araya geldiği çevrimiçi platformlar, bireylerin ve grupların kimliklerini ifade etme ve şekillendirme süreçlerinde önemli bir rol oynar. Sosyal medya, bloglar, forumlar gibi dijital mecralar, kullanıcıların çeşitli yollarla kendi kültürel kimliklerini sergilemelerine olanak tanır. Bu platformlar, kimlik oluşturma süreçlerini daha etkileşimli ve dinamik hale getirir; çünkü kullanıcılar farklı kültürel ifadeleri keşfetme ve kendi kimliklerini bu çerçevede yeniden tanımlama fırsatı bulurlar.

Dijital ortamlar, kullanıcıların hem global hem de yerel kültürlerle etkileşim kurmalarına olanak sağlar. Bu da, kimliklerin sadece yerel veya ulusal bağlamlarla sınırlı kalmayıp, küresel bir perspektifte yeniden şekillendiği anlamına gelir. Özellikle gençler arasında, dijital platformlar aracılığıyla kültürel kimliklerin ifade edilmesi ve paylaşılması yaygın hale gelmiştir.

Ancak, dijital dönüşümün kültürel çeşitliliğe etkisi hem olumlu hem de olumsuz olabilir. Dijital platformlar, azınlık kültürlerinin ve marjinal grupların seslerini duyurmalarına yardımcı olur ve kültürel çeşitliliği destekler. Ancak, bu platformların büyük ölçüde dominant kültürel güçlerin kontrolünde olması, kültürel homojenleşmeye ve bazı kültürel ifadelerin marjinalleştirilmesine yol açabilir.

Çevrimiçi kimlik politikaları da, dijital platformlarda oluşturulan ve paylaşılan kimliklerin siyasal ve toplumsal dinamikler üzerinde önemli bir etkiye sahiptir. Bu politikalar, bireylerin çevrimiçi

FULL TEXTS BOOK

kimliklerini nasıl oluşturduklarını, nasıl algılandıklarını ve bu kimliklerin toplumsal ilişkilerde nasıl bir rol oynadığını kapsar.

Siyasi aktivizm ve toplumsal hareketler açısından, dijital platformlar önemli bir rol oynar. Bu platformlar, bireylerin ve grupların politik görüşlerini ifade etmeleri ve örgütlenmeleri için kritik araçlar sunar. Ancak, çevrimiçi kimlik politikaları aynı zamanda toplumsal kutuplaşmayı artırabilir ve dijital ortamda yayılan yanlış bilgiler ve nefret söylemleri gibi sorunlara yol açabilir.

Dijital Dönüşümün Siyasal ve Toplumsal Sonuçları

Dijital dönüşüm, siyasal ve toplumsal katılımın şeklini derinden değiştirmektedir. Çevrimiçi kimlikler, bireylerin siyasi görüşlerini ifade etme, politik tartışmalara katılma ve toplumsal hareketlere destek verme gibi yeni olanaklar sunmaktadır (Bennett & Segerberg, 2012). Özellikle gençler, politik katılım için çevrimiçi platformları yoğun bir şekilde kullanmaktadırlar. Ancak, bu eğilimin gölgesinde, internet erişimi ve dijital okuryazarlık eksikliği gibi faktörler bazı grupların siyasi süreçlere eşit katılımını engellemektedir (Norris, 2001).

Dijital platformlar, kültürel ve politik çeşitliliğin korunması veya baskılanması açısından çifte bir potansiyel taşımaktadır. Bu platformlar, azınlık gruplarının ve marjinal toplulukların sesini duyurmasına yardımcı olarak toplumsal ve politik çeşitliliği destekleyebilirler (Castells, 2009). Ancak, içerik moderasyon politikaları ve algoritmaların ticari ve politik etkilere maruz kalarak bazı görüşlerin ve kültürel ifadelerin marjinalleştirilmesine yol açabileceği hususunda dikkat gerekmektedir (Gillespie, 2018).

Sosyal medya platformları, kullanıcıların etkileşimlerini izleyerek ve analiz ederek kişiselleştirilmiş içerik sunmaktadırlar. Bu durum, kullanıcıların sadece kendi görüşlerini destekleyen içeriklerle karşılaşmasına ve yankı odaları içinde kalmasına neden olabilir. Sonuç olarak, toplumsal kutuplaşma artabilir ve farklı görüşler arasındaki diyalog zorlaşabilir (Sunstein, 2001).

Dijital dönüşüm, teknolojik gelişmelerin hızına ve yönüne bağlı olarak gelecekte siyasal ve toplumsal yapıları derinden etkilemeye devam edecektir. Yapay zekâ, büyük veri ve blockchain gibi teknolojiler, siyasal süreçleri ve toplumsal dinamikleri yeniden şekillendirebilecektir. Ancak, bu süreçte toplumsal eşitsizliklerin derinleşme riski de bulunmaktadır. Teknolojik yeniliklere erişimdeki eşitsizlikler, toplumsal katmanlar arasındaki dijital uçurumun genişlemesine yol açabilir (Van Dijk, 2020).

Bu bağlamda, geleceğe dönük politikalar oluştururken, dijital dönüşümün bu çift yönlü etkilerini dikkate almak hayati önem taşımaktadır. Demokratik katılımı artırma ve toplumsal çeşitliliği desteklemenin yanı sıra, dijital eşitsizlikler ve kutuplaşma gibi sorunları da ele almak gerekmektedir.

SONUÇ

Dijital dönüşümün toplumsal ve siyasal etkileri, literatürde yer alan geçmiş araştırmalar yolu ile birçok açıdan incelenmiştir. Öncelikle, çevrimiçi kimliklerin siyasal katılım üzerindeki rolü ele alınmıştır. Çevrimiçi platformlar, bireylerin siyasi görüşlerini ifade etmeleri ve politik süreçlere katılmaları için önemli bir alan sağlamaktadır. Ancak, internet erişimi ve dijital okuryazarlık eksikliği gibi faktörler, siyasal katılımın eşitlikçi bir şekilde gerçekleşmesini engelleyebilir. Bu nedenle, dijital katılımı teşvik etmek ve dijital eşitsizlikleri azaltmak için politika önlemleri alınmalıdır.

İkinci olarak, dijital platformlarda çeşitliliğin korunması veya baskılanması ele alınmıştır. Dijital platformlar, kültürel ve politik çeşitliliğin desteklenmesinde önemli bir rol oynayabilirler. Ancak, içerik moderasyon politikaları ve algoritmaların etkisi altında, bazı görüşlerin ve kültürel ifadelerin marjinalleştirilme riski bulunmaktadır. Bu nedenle, içerik moderasyonu ve algoritma tasarımı konusunda şeffaflık ve hesap verebilirlik sağlayacak politikaların geliştirilmesi gerekmektedir.

Son olarak, dijital dönüşümün geleceğe dönük toplumsal ve siyasal etkileri incelenmiştir. Teknolojik gelişmelerin hızına ve yönüne bağlı olarak, siyasal süreçler ve toplumsal dinamikler önemli ölçüde

FULL TEXTS BOOK

değişebilir. Ancak, bu süreçte toplumsal eşitsizliklerin derinleşme riski bulunmaktadır. Bu nedenle, teknolojiye erişimdeki eşitsizlikleri azaltmak ve dijital demokrasiyi güçlendirmek için politika önlemleri alınmalıdır.

Dijital dönüşümün toplumsal ve siyasal etkileri, dikkatlice incelenmesi gereken karmaşık bir konudur. Gelecekte, daha adil ve kapsayıcı bir dijital ortamın oluşturulması için politika yapıcıların ve sivil toplumun birlikte çalışması gerekmektedir. Bu çerçevede, teknoloji şirketleri, akademisyenler, hükümetler ve sivil toplum kuruluşları arasında iş birliğini teşvik eden politikaların benimsenmesi önemlidir. Bu şekilde, dijital dönüşümün toplumsal ve siyasal alanlarda olumlu etkileri arttırılabilir ve olumsuz etkileri azaltılabilir.

Öte yandan, gelecekteki araştırmaların, dijital dönüşümün toplumsal ve siyasal etkilerini daha derinlemesine anlamak için belirli alanlara odaklanması önemlidir. Bu bağlamda, aşağıdaki öneriler göz önünde bulundurulabilir:

-Dijital Katılımın Değerlendirilmesi: Dijital dönüşümün siyasal katılım üzerindeki etkilerini daha iyi anlamak için çeşitli demografik gruplar arasındaki dijital katılımı değerlendiren çalışmalara ihtiyaç vardır.

-Dijital Kutuplaşmanın İncelenmesi: Dijital dönüşümün toplumsal kutuplaşma üzerindeki etkilerini anlamak için daha fazla araştırma yapılmalıdır. Bu, algoritmik sınıflandırma, filtre balonları ve bilgi yankı odaları gibi konuları içerebilir.

-Dijital Eşitsizliklerin Belirlenmesi: Dijital dönüşümün toplumsal eşitsizliklere etkilerini daha iyi anlamak için daha fazla araştırmaya ihtiyaç vardır. Özellikle dijital okuryazarlık, internet erişimi ve teknolojiye erişimdeki eşitsizliklerin incelenmesi önemlidir.

Dijital dönüşümün toplumsal ve siyasal etkileri göz önüne alındığında, politika yapıcıların ve toplumsal yöneticilerin aşağıdaki önerilere dikkat etmeleri önemlidir:

-Dijital Okuryazarlık Eğitimi: Toplumun dijital okuryazarlık düzeyini artırmak için eğitim ve farkındalık kampanyaları düzenlenmelidir. Bu, bireylerin dijital ortamlarda daha bilinçli ve eleştirel olmalarını sağlayabilir.

-Dijital Erişim ve Altyapının Geliştirilmesi: Dijital eşitsizlikleri azaltmak için internet erişimi ve teknoloji altyapısının geliştirilmesi gerekmektedir. Özellikle kırsal bölgelerde ve düşük gelirli topluluklarda dijital erişim sağlanması önemlidir.

-Veri Mahremiyeti ve Güvenliği Politikalarının Güçlendirilmesi: Dijital ortamlarda veri mahremiyeti ve güvenliğini korumak için daha etkili politika ve düzenlemeler oluşturulmalıdır. Bu, bireylerin kişisel verilerinin korunmasını ve dijital ortamlarda güvenli bir deneyim yaşamalarını sağlayabilir.

KISITLILIK

Araştırma daha önce yapılmış ve literatüre geçmiş bilimsel araştırmalar ile sınırlıdır

BİLDİRİMLER

Değerlendirme: İç ve dış danışmanlarca değerlendirilmiştir.

Çıkar Çatışması: Yazar bu makale ile ilgili herhangi bir çıkar çatışması bildirmemiştir.

Finansal Destek: Yazar bu makale ile ilgili herhangi bir malî destek kullanımı bildirmemiştur.

ETİK BEYAN

Current Science dergisinin yayın etiği; bilimsel araştırma ve yayınların dürüstlük, açıklık, nesnellik, başkalarının bulgu ve yaratılarına saygı gibi temel ilkelere uygun biçimde yapılmasını sağlamayı amaçlayan ve bunun gerçekleşmesi için çalışmalar yapan sosyal bilimleri alanında bu ilkeleri amaç edinmiş ulusal tabanlı bilimsel bir dergidir. Helsinki bildirgesi kriterleri göz önünde bulundurulmuştur.

FULL TEXTS BOOK

KAYNAKÇA

Alvarez, R. M., & Hall, T. E. (2004). Point, Click, and Vote: The Future of Internet Voting. Washington D.C.: Brookings Institution Press.

Altun, A. (2017). Dijital Dönüşüm ve Ağ Toplumu. İstanbul: Beta Yayıncılık.

Aktay, H. (2022). Dijital Demokrasi ve Etik Sorunlar. Bursa: Etik Yayınları.

Baym, N. K. (2015). Personal connections in the digital age. Polity.

Benkler, Y. (2006). The Wealth of Networks: How Social Production Transforms Markets and Freedom. Yale University Press.

Bimber, B. (2014). Digital Media in the Obama Campaigns of 2008 and 2012: Adaptation to the Personalized Political Communication Environment. Journal of Information Technology & Politics, 11(2), 130-150.

Brown, E. (2021). The Political Implications of Online Identity Formation. Journal of Political Science, 45(2), 213-230.

Castells, M. (2008). The New Public Sphere: Global Civil Society, Communication Networks, and Global Governance. The Annals of the American Academy of Political and Social Science, 616(1), 78-93.

Castells, M. (2009). Communication Power. Oxford University Press.

Castells, M. (2010). The rise of the network society: The information age: Economy, society, and culture (Vol. 1). John Wiley & Sons.

Chadwick, A. (2006). Internet Politics: States, Citizens, and New Communication Technologies. Oxford: Oxford University Press.

Chadwick, A. (2013). The Hybrid Media System: Politics and Power. Oxford University Press.

Chen, P. (2021). Digital Media and the Future of Elections. Sydney: University of Sydney Press.

Çelik, A. (2023). Dijital Medya ve Seçmen Davranışları. Ankara: Politika Yayınları.

Çoban, B. (2013). Gezi Parkı ve Sosyal Medya. İstanbul: Toplumsal Hareket Yayınları.

Ellison, N. B., & Boyd, D. (2013). Sociality through social network sites. In W. H. Dutton (Ed.), The Oxford handbook of internet studies (pp. 151-172). Oxford University Press.

Entman, R. M. (2012). Scandal and Silence: Media Responses to Presidential Misconduct. Polity.

Floridi, L. (2014). The Fourth Revolution: How the infosphere is reshaping human reality. Oxford University Press.

Garcia, M. (2018). Online Identity and Social Movements: A Comparative Analysis. Social Movement Studies, 20(3), 345-362.

Gillespie, T. (2018). Custodians of the Internet: Platforms, Content Moderation, and the Hidden Decisions that Shape Social Media. Yale University Press.

Gökalp, S. (2023). Siyasal İletişimde Dijitalleşme. İstanbul: Siyasal Yayınları.

Gürsel, Z. (2016). The Politics of Technology and Its Influence on Public Opinion. Ankara: Politika Yayınları.

Howard, P. N. (2006). New Media Campaigns and the Managed Citizen. Cambridge University Press.

Howard, P. N., & Hussain, M. M. (2011). The Role of Digital Media in the Arab Spring. Journal of Democracy, 22(3), 35-48.

FULL TEXTS BOOK

Jones, K. (2020). Social Interaction Theory and Online Identity Formation. Communication Studies, 30(4), 501-518.

Karahasanoğlu, S. (2019). Siyasal İletişim ve Teknolojik Değişimler. İstanbul: Siyasal Yayınları.

Kreiss, D. (2016). Prototype Politics: Technology-Intensive Campaigning and the Data of Democracy. Oxford: Oxford University Press.

Marwick, A., & Lewis, R. (2017). Media Manipulation and Disinformation Online. Data & Society Research Institute.

Mercuri, R. T. (2002). Electronic Vote Tabulation Checks & Balances. Washington D.C.: Brookings Institution Press.

Nakamura, L. (2002). Cybertypes: Race, Ethnicity, and Identity on the Internet. Routledge.

Norris, P. (2001). Digital divide: Civic engagement, information poverty, and the internet worldwide. Cambridge University Press.

Papacharissi, Z. (2010). A private sphere: Democracy in a digital age. Polity.

Pariser, E. (2011). The Filter Bubble: How the New Personalized Web Is Changing What We Read and How We Think. Penguin Press.

Smith, J. (2019). The Role of Culture in Online Identity Formation. Cultural Studies, 25(1), 89-104.

Strömbäck, J. (2008). Four Phases of Mediatization: An Analysis of the Mediatization of Politics. The International Journal of Press/Politics, 13(3), 228-246.

Sunstein, C. R. (2001). Republic.com. Princeton University Press.

Tufekci, Z. (2017). Twitter and Tear Gas: The Power and Fragility of Networked Protest. New Haven: Yale University Press.

Turkle, S. (1995). Life on the screen: Identity in the age of the Internet. Simon and Schuster.

Turkle, S. (2011). Alone together: Why we expect more from technology and less from each other. Basic Books.

Vaccari, C., & Valeriani, A. (2016). Follow the Leader! Direct and Indirect Flows of Political Communication during the 2013 Italian General Election Campaign. New Media & Society, 18(4), 516-534.

Van Dijk, J. A. G. M. (2020). The Digital Divide. Polity.

Van Dijk, J. (2020). The network society. Sage Publications.

West, D. M. (2019). The future of work: Robots, AI, and automation. Brookings Institution Press.

STRENGTHENING FEED SECURITY IN WEST AFRICA: UNLEASHING THE STRATEGIC POTENTIAL OF FINANCIAL INSTITUTIONS FOR SUSTAINABLE LIVESTOCK SYSTEMS

Dr. Emmanuel Ugochukwu ANASO

Department of Animal Science, Federal University of Agriculture Mubi, Adamawa, Nigeria

ABSTRACT

Feed security is a critical component of sustainable livestock production and food systems in West Africa. With the growing challenges of climate change, rising input costs, and fluctuating grain availability, the role of financial institutions in safeguarding feed security has become increasingly vital. This paper explores how financial institutions contribute to enhancing feed security across the subregion through strategic financing, risk mitigation, and investment in feed value chains. Key roles include providing access to credit for smallholder livestock farmers and feed producers, supporting agribusiness development through agricultural financing schemes, and enabling insurance products that cushion the effects of feed-related supply shocks. Moreover, financial institutions facilitate public–private partnerships and donor-funded programs aimed at improving infrastructure, feed processing, and storage. Case studies from Nigeria, Ghana, and Senegal illustrate how tailored financial products—such as warehouse receipt systems, input financing, and microcredit—have improved access to quality feed and reduced volatility in feed markets. Despite these efforts, challenges such as limited financial literacy, high interest rates, and inadequate rural banking infrastructure persist. The study recommends the development of inclusive financial models, digital finance tools, and policy support to strengthen the role of financial institutions in achieving feed security in West Africa.

Keywords; Feed security, financial institutions, Livestock production, Agricultural financing, Access to credit, Feed value chains, Agribusiness development, Risk management

Introduction

The livestock industry serves as a cornerstone of rural economies, nutritional well-being, and economic growth across West Africa (Anaso et al., 2024b). It plays an essential role in enhancing household earnings, generating employment, and supplying a substantial proportion of dietary proteins to communities (FAO, 2020; Anaso et al., 2021a,b). Livestock production in the region is closely integrated into mixed crop–livestock farming systems, where animals not only provide manure to maintain soil fertility and traction for cultivation but also function as a form of financial security and capital reserve for rural households.


Despite its importance, the sector is constrained primarily by the persistent challenge of feed scarcity and rising costs (Olafadehan et al., 2023). Feed expenditures represent the dominant share of production costs, often comprising between 60% and 70% of the total operational budget (FAO, 2018; Akinmoladun et al., 2019; Anaso et al., 2025a). Hence, achieving feed security—characterized by stable and affordable access to high-quality feed resources—is indispensable for improving productivity and maintaining competitiveness within the livestock sector (Anaso et al., 2025b,c).

Multiple factors contribute to feed insecurity in West Africa. Climatic variability, frequent drought episodes, and unpredictable rainfall negatively affect both forage and grain output; global market volatility increases the price of imported feed inputs such as soybean meal and maize; and systemic

FULL TEXTS BOOK

inefficiencies in supply chains delay feed distribution and exacerbate cost pressures (Njie et al., 2021; Anaso & Olafadehan, 2025).

Efforts to enhance feed security have traditionally focused on technological solutions such as the development of drought-resistant forage cultivars, the promotion of unconventional feed resources like crop residues and agro-industrial by-products, and improved feed conservation practices including silage production and haymaking (Tarawali et al., 2011; Ayantunde et al., 2018). While these interventions are valuable, they are insufficient on their own. Without complementary financial mechanisms—such as accessible credit facilities, risk-mitigation tools like insurance, and investment in feed-related infrastructure—farmers and feed processors struggle to adopt and scale these innovations sustainably. As a result, financial institutions play a pivotal role in ensuring resilient feed systems and safeguarding the region's livestock sector.



2. The Role of Financial Institutions in Feed Security

Financial institutions (FIs)—ranging from commercial banks to microfinance providers and development finance institutions—shape feed security through three main avenues:

A. Access to Credit and Working Capital

Access to affordable financing empowers stakeholders across the feed value chain—including producers, processors, and marketers—to acquire raw materials in large quantities, upgrade processing technologies, and sustain continuous production flows. In Nigeria, for instance, the Central Bank's Anchor Borrowers' Programme (ABP) has enhanced grain availability by facilitating input credit and creating linkages between smallholder farmers and agro-processors (CBN, 2023). Likewise, in Ghana, the Agricultural Development Bank provides specialized loan schemes for the poultry and aquaculture

FULL TEXTS BOOK

industries, allowing significant cost reductions through bulk procurement of feed resources (ADB, 2022).

B. Risk Mitigation

Risk-mitigation tools like index-linked weather insurance and price volatility coverage help shield producers from the adverse impacts of climatic extremes and market instability. In Nigeria, the Agricultural Insurance Corporation (NAIC) offers subsidized policies that offset financial losses caused by drought, floods, and pest infestations (NAIC, 2022). These initiatives provide farmers with greater confidence to sustain or scale up livestock operations even under unpredictable conditions.

C. Value Chain Development and Infrastructure Financing

Through financing essential infrastructure—including feed processing plants, storage units, and transportation networks—financial institutions (FIs) play a key role in minimizing post-harvest losses, stabilizing seasonal price fluctuations, and enhancing producers' access to markets. Regional bodies such as the ECOWAS Bank for Investment and Development (EBID) have backed agribusiness initiatives aimed at strengthening rural infrastructure, thereby increasing the efficiency and reliability of feed distribution systems (EBID, 2021).

3. Case Studies

- **Nigeria:** The CBN's ABP has indirectly strengthened feed security by financing grain production, thereby ensuring a steady supply to feed manufacturers.
- **Ghana:** Agricultural Development Bank input financing for poultry and aquaculture producers has improved feed affordability and reduced unit costs.
- **Senegal:** The introduction of warehouse receipt systems for maize and groundnuts has allowed farmers to store grain securely and access credit against stored commodities, stabilizing seasonal feed prices.

4. Challenges Limiting Financial Institutions' Impact

Despite promising initiatives, systemic barriers limit FI effectiveness:

- **High Interest Rates** discourage long-term investment in feed production and processing (World Bank, 2021).
- **Collateral Requirements** often exclude smallholders without formal land titles.
- **Limited Rural Banking Penetration** leaves many livestock producers beyond the reach of formal credit systems.
- **Low Financial Literacy** hampers effective utilization of financial products (Aidoo & Fromm, 2015).

5. Recommendations for Strengthening FI Contributions

- **Inclusive Financial Models** tailored to smallholder realities, cooperatives, and commercial producers.
- **Digital Financial Tools** leveraging mobile platforms for microloans, savings, and insurance access (GSMA, 2020).
- **Policy and Regulatory Reforms** to lower agricultural lending rates and incentivize agribusiness financing.
- **Public–Private Partnerships** to co-invest in feed production, processing, and distribution infrastructure.

FULL TEXTS BOOK

6. Conclusion

Ensuring feed security in West Africa requires addressing both technical and financial constraints. Although advancements in feed production technologies are vital, their widespread adoption largely depends on access to affordable financing, effective risk mitigation strategies, and targeted investments across the feed value chain. Financial institutions play a pivotal role in this process by offering tailored credit facilities, implementing innovative insurance solutions, and supporting infrastructure development. Achieving sustainable feed systems will require an integrated strategy that combines financial innovation with supportive policy frameworks and capacity-building initiatives, thereby positioning the livestock sector as a resilient and reliable contributor to regional food and nutrition security.

REFERENCES

- ADB (2022). *Annual Report 2021/2022*. Agricultural Development Bank of Ghana.
- Aidoo, R., & Fromm, I. (2015). Willingness to pay for weather insurance in northern Ghana. *Journal of Social and Development Sciences*, 6(1), 1–15.
- Akinmoladun, F. O., et al. (2019). Feed resources and feeding practices for sustainable livestock production in Nigeria. *International Journal of Veterinary Science and Animal Husbandry*, 4(6), 13–20.
- Anaso, E.U., Olafadehan, O.A. (2025). Nutritive value, feed intake and blood profile of buckling goats fed diets containing Pleurotus ostreatus biodegraded sugarcane scrapings. Archiva Zootechnica, 28:1, 61-76, 2025 DOI: 10.2478/azibna-2025-0004.
- Anaso, E.U., Olafadehan, O.A., Chibuogwu, I.C. (2025a). Nutritive effects and physiologic responses of rabbits supplemented with Camel's Foot (Piliostigma thonningii) essential oil-based diet. Veterinarski Archiv, 2025. DOI: 10.24099/vet.arhiv.2669
- Anaso, E.U., Olafadehan, O.A., Chibuogwu, I.C. (2025b). Nutrient digestibility, growth performance, cecal microbial profile, and fermentation characteristics of rabbit bucks fed Piliostigma thonningii essential oil supplemented diet. Biotechnology in Animal Husbandry, 41(1), 37-55, 2025 https://doi.org/10.2298/BAH2302205K
- Anaso, E.U., Olafadehan, O.A., Oluwafemi, R.A. (2021b). Biodegradation of agro industrial by-products as a panacea to feed scarcity in Nigeria amidst farmer insecurity and global pandemic. Proceedings of the 46th Annual Conference of the Nigerian Society for Animal Production, pp. 616-620.
- Anaso, E.U., Olafadehan, O.A., Shoyombo, A.J. (2021a). Semen characteristics of Kano Brown bucks fed white rot fungi (Pleurotus ostreatus) biodegraded sugarcane scrapings based diets. Proceedings of the 46th Annual Conference of the Nigerian Society for Animal Production, March 14-18, 2021, pp. 625-628. 18.
- Anaso, E.U., Olafadehan, O.A., Shoyombo, A.J., Emeka, F.S. (2024). Body weight, scrotal parameters, and semen characteristics of Kano Brown bucks fed Pleurotus ostreatus solid state fermented sugarcane scrapings. Turkish Journal of Food and Agricultural Science, DOI: https://doi.org/10.53663/turjfas.1486383.
- Ayantunde, A. A., et al. (2018). Improving livestock feeding in West Africa: Strategies and recommendations. *ILRI Policy Brief*.
- CBN (2023). *Anchor Borrowers' Programme Guidelines*. Central Bank of Nigeria.

FULL TEXTS BOOK

- EBID (2021). *Annual Report 2021*. ECOWAS Bank for Investment and Development.
- FAO (2018). *Africa Sustainable Livestock 2050: Feed resources in West Africa*. Food and Agriculture Organization of the United Nations.
- FAO (2020). *Livestock Sector Brief – West Africa*. FAO Regional Office for Africa.
- GSMA (2020). *The Mobile Economy: Sub-Saharan Africa 2020*. GSM Association.
- NAIC (2022). *Corporate Profile*. Nigerian Agricultural Insurance Corporation.
- Njie, M., et al. (2021). Climate change and livestock feed systems in West Africa. *Pastoralism*, 11(12), 1–15.
- Olafadehan, O.A., Anaso, E.U., Shoyombo, A.J., Okunade, S.A. (2023). Body thermoregulation and serum metabolic profile of Kano Brown bucks fed Pleurotus ostreatus biodegraded sugarcane scrapings. Biotechnology in Animal Husbandry, 39(1), 61-72. 37 https://doi.org/10.2298/BAH2202081R
- Tarawali, S., et al. (2011). Pathways for sustainable development of mixed crop–livestock systems: Taking a livestock value chain approach. *Livestock Science*, 139(1-2), 11–21.
- World Bank (2021). *Nigeria Agriculture Public Expenditure Review*. Washington, DC: World Bank.

FULL TEXTS BOOK

SÜRTÜNME KARIŞTIRMA PROSESİNİN METAL MATRİS KOMPOZİTLERİN MİKROYAPISI ve MEKANİK ÖZELLİKLERİ ÜZERİNDEKİ ETKİSİ

EFFECT of FRICTION STIR PROCESSING on MICROSTRUCTURE and MECHANICAL PROPERTIES of METAL MATRIX COMPOSITES

İsmail VAROL[1]

Yüksek Lisans Öğrencisi,Dokuz Eylül Üniversitesi,Mühendislik Fakültesi, Makina Mühendisliği,İzmir,Türkiye

Fatih KAHRAMAN[2]

Profesor Doktor, Dokuz Eylül Üniversitesi,Mühendislik Fakültesi, Makina Mühendisliği,İzmir,Türkiye

[2]*ORCID ID: https://orcid.org/0000-0002-1075-9171*

ÖZET

Sürtünme Karıştırma Prosesi (SKP), sürtünme karıştırma kaynağı (SKK) temeline bağlı olarak geliştirilen, malzemelerin yüzeye yakın bölgelerinde mikroyapı iyileştirmesi sağlayan bir katı hal yöntemidir. Diğer yöntemlere kıyasla sarf malzeme tüketimine uğramayan döner bir takım aracılığıyla malzemede sürtünmeye bağlı olarak gelişen yoğun plastik deformasyon ve ısıl etkileşim oluşturularak tane incelmesi, faz homojenliği ve yoğunluk artışı sağlanır. Bu süreç, özellikle alüminyum alaşımları ve kompozit malzemelerde çökelti dağılımını düzenlerken, işlenen bölgede sertlik ve aşınma direnci gibi mekanik özellikleri de artırır. SKP ile farklı elementlerin karıştırma bölgesine dahil edilmesiyle yüzey veya hacimsel alaşım modifikasyonu mümkün hale gelir. Bu yönüyle SKP, havacılık, otomotiv ve taşımacılık gibi sektörlerde yüksek performanslı hafif malzemelerin mikroyapısal olarak geliştirilmesinde yaygın olarak kullanılmaktadır.Sürtünme karıştırma prosesiyle üretimi sağlanan kompozit malzemelerin mikroyapısal ve mekanik özellikleri , SKP parametrelerine (takım dönme devri,takım ilerleme hızı,paso sayısı,dalma derinliği,karıştırıcı uç tipi) ve matris – takviye partikülü eklentisine bağlı olarak güçlü yada zayıf olarak değişim gösterebilmektedir.

Bu araştırmada, sürtünme karıştırma prosesi ile üretilen metal matrisli kompozitlerde meydana gelen mikroyapısal değişimlerin, kompozitlerin mekanik dayanımı üzerindeki etkileri değerlendirilecektir. Ayrıca, SKP işlemine bağlı olarak oluşan ve malzeme performansını olumsuz yönde etkileyen iç yapı kusurlarının oluşum mekanizmaları da çalışmanın kapsamı içerisinde incelenecektir.

Anahtar Kelimeler: Sürtünme karıştırma prosesi, metal matrisli kompozit, iç yapı kusurları, mekanik özellikler.

ABSTRACT

Friction Stir Processing (FSP) is a solid-state technique developed based on the principles of Friction Stir Welding (FSW), enabling microstructural refinement in the near-surface regions of materials. Unlike other methods, it utilizes a non-consumable rotating tool that induces intense plastic deformation and thermal exposure through friction, resulting in grain refinement, phase homogeneity, and densification. This process is particularly effective in aluminum alloys and composite materials, enhancing precipitate distribution and improving mechanical properties such as hardness and wear resistance in the processed zones. Through FSP, surface or bulk alloy modification can also be achieved by incorporating various elements into the stirred region. Owing to these advantages, FSP is widely used in the aerospace, automotive, and transportation industries for microstructural enhancement of high-performance lightweight materials.

FULL TEXTS BOOK

The microstructural and mechanical properties of composite materials fabricated through FSP can vary significantly depending on processing parameters such as tool rotational speed, traverse speed, number of passes, plunge depth, and pin geometry, as well as the type and distribution of matrix and reinforcement particles.

In this study, the effects of microstructural changes induced by friction stir processing on the mechanical strength of metal matrix composites will be investigated. Additionally, the formation mechanisms of internal defects that may negatively affect material performance due to the FSP process will also be examined as part of the research scope.

Keywords: Friction stir processing, metal matrix composite, internal defects, mechanical properties.


TEŞEKKÜR

Bu çalışmanın her aşamasında bilgi ve tecrübesiyle hem akademik hem de mesleki gelişimime büyük katkı sağlayan danışman hocam Sayın Prof. Dr. Fatih KAHRAMAN'a en içten teşekkürlerimi sunarım.

FYL-2024-3456 numaralı bu çalışmada Bilimsel Araştırma Projesi desteğinden dolayı Dokuz Eylül Üniversitesi BAP koordinatörlüğüne ve deneysel çalışmalarım sırasında analizlerimi gerçekleştirmem de yardımcı olan Sayın Arş. Gör. Dr. Coşkun YOLCU ve Dokuz Eylül Üniversitesi Bilim Teknoloji Uygulama ve Araştırma Merkezine teşekkür ederim.

GİRİŞ

Metal matrisli kompozitler, mühendislik malzemelerinden kullanım aşamasında beklenen özellikleri karşılamak amacıyla biri metal (saf ya da alaşım durumda) olmak üzere en az iki farklı malzemenin (metal, ametal, seramik vb.) belirli yöntemler dahilinde birleşimi ile elde edilen malzemelerdir. Metal matrisli kompozitleri oluşturan malzemeler fiziksel ve kimyasal olarak birbirlerinden farklı özellikler göstermektedirler (Chawla & Chawla, 2006). Bileşen malzemelerin herhangi birinden tek başına elde edilemeyen özelliği sağlamak amacıyla üretilen MMK'de ana malzeme metal matris iken diğer malzemeler ise takviye görevi gören mikro ya da nano boyutlarda partiküller, kısa fiberler, sürekli fiberler ya da sürekli tabakalar olabilmektedir (Şekil 1). Matris malzemesi takviye elemanlarını bir arada tutar ve malzemenin şekilsel bütünlüğünü korur. Takviyeler ise matrise belirli oranda ve boyutlarda eklenerek kompozit yapıdan beklenen özellikleri sağlamaktadırlar (Kalemtaş, 2014) (Gürbüz & Mutuk, 2019).



Şekil 1. Farklı Türdeki Metal Matrisli Kompozitlerin Şematik Gösterimi (Chawla & Chawla, 2006)

Metal matrisli kompozitlerde otomotiv (partikül takviyeli fren diskleri) ve havacılık endüstrisi başta olmak üzere ağırlık azaltmak için yapılan çalışmalar, düşük ağırlığa sahip malzemelerin kullanıldığı tasarımların optimizasyonu için çok büyük itici güç olmaktadır. Bu kapsamda, alüminyum alaşımları; yüksek mukavemet/ağırlık oranı, iyi yorulma dayanımı ve aşınma direnci, yüksek sünekliğe sahip matris

FULL TEXTS BOOK

yapıları, kolay işlenebilirlik özellikleri ve uygun maliyetleri sayesinde MMK üretiminde matris malzemesi olarak en yaygın tercih edilen metal haline gelmiştir (Sharma, Bhandari, Aherwar, Rimašauskienė, & Bretotean, 2020). Metal matrisli kompozit elemanları birçok farklı yöntemle üretilebilmekte olup, kullanılacak yöntemin seçiminde parçaların geometrisi, parçadan beklenen özellikler, tercih edilecek matris malzemesi ile takviye elemanının türü, şekli ve boyutu dikkate alınmaktadır. MMK'lerde takviye malzemelerinin yapı içinde homojen dağılımı ise önemli fakat güç bir durumdur (Kraiklang, Onwong, & Santhaweesuk, 2020) (Sharifitabar, Sarani, Khorshahian, & Afarani, 2011).

Piston üretiminde kullanılan seramik takviyeli metal matrisli kompozitler, matris malzemelerinin tek başına sunduğu özelliklere kıyasla aşınmaya karşı belirgin derecede daha yüksek direnç sergilemektedir. Ayrıca, düşük termal genleşme katsayıları sayesinde dar toleransların uygulanmasına imkân tanımakta; bu durum, izin verilebilir maksimum basınç değerlerinin artmasına ve termal iletkenlik özelliklerinin iyileşmesine katkı sağlamaktadır (Blaza & Lozica, 2015).



Şekil 2. Metal matris esaslı kompozit motor bloğu, pistonu ve fren diski (Blaza & Lozica, 2015)

Metalik malzemelerin nanopartiküller ile takviye edilmesi, son yıllarda benzersiz mekanik ve fiziksel özelliklere sahip yeni kompozitlerin geliştirilme potansiyeli nedeniyle önemli bir araştırma ilgisi uyandırmıştır. Metal matrisli kompozitlerin üretim yöntemleri genel olarak katı hal ve sıvı hal olmak üzere iki kategoriye ayrılmaktadır. Sıvı hal işlemler, malzemelerin ergime sıcaklığının üzerinde gelişirken, katı hal prosesler malzemelerin ergime sıcaklığı altında gelişmektedir.


Bu proseslere örnek olarak;

- Sıkıştırmalı döküm,
- İnfiltrasyon tekniği,
- Toz metalurjisi,
- Sürtünme karıştırma prosesi,

teknikleri metal matrisli kompozit üretimi yöntemlerinde sık kullanılan proseslerdir. Kompozit malzemelerde üstün mekanik özelliklerin elde edilebilmesi için, takviye edici partiküllerin kompozitlerin metal matrisi içerisinde homojen bir şekilde dağılması gerekmektedir. Ancak, seramik nanopartiküllerin metaller içinde homojen dağılımını sağlamak zordur. Sıvı hal işlemleri sırasında nanopartiküller, geniş yüzey alanına sahip olmaları ve ergimiş metallerde düşük ıslanabilirlikleri nedeniyle, iri kümeler halinde toplanma eğilimindedir. Bu bağlamda kompozit üretiminde ki bu tür problemlerin üstesinden gelebilecek, daha efektif ayarlanabilir işlem parametrelerine sahip olan sürtünme karıştırma kaynağı esaslı sürtünme karıştırma prosesi (SKP) karşımıza çıkmaktadır.

FULL TEXTS BOOK

SKP, 1991 yılında The Welding Institute (TWI) tarafından geliştirilen bir katı hal işleme tekniğidir ve malzeme boyunca hareket eden aşınmayan döner bir aletten oluşur. Geleneksel alet, yapılacak modifikasyonun derecesine bağlı olarak, numune kalınlığı ile uyumlu uzunlukta tasarlanmış özel bir omuz ve pime sahiptir. İşlem sırasında dönen pim katı iş parçasına daldırılırken, dönen omuz malzemenin üst yüzeyiyle temas eder ve alet, belirlenen hat boyunca hareket eder (Şekil 3). Aletin hareketi sırasında oluşan sürtünme ısısı ve mekanik karıştırma, geçtiği bölgede mikro yapısal değişikliklere neden olur. Bu deformasyon süreci, sürekli dinamik yeniden kristalleşme ile birlikte gerçekleşir ve tane incelmesine yol açar. Ayrıca, aletin hareketi, SKP ile işlenmiş bölgede ince, eş eksenli tanelerin homojen bir şekilde dağılmasını sağlar (Patel, Vairis, Badheka, & Li, 2019).




Şekil 3. a) SKP prensibinin şematik gösterimi b) SKP bölgelerinin kesit görünümü (Patel, Vairis, Badheka, & Li, 2019)

Metal matrisli kompozit üretiminde karşılaşılan problemlerden biri, kompozit mukavemetini önemli ölçüde etkileyen, parçacıkların matris içindeki dağılımıdır. Parçacıkların matris içinde kümelenmesi , mikro boşluk oluşumuna ve yük altında çatlak oluşumu başlangıcı görevi görmesine dolayısıyla, kompozitlerin mekanik özelliklerinin düşmesine sebep olabilmektedir. Kompozit yapısında mikroyapı kusurlarının oluşmasına ve malzemenin dayanımının düşmesinde sürtünme karıştırma prosesi parametreleri ve matris-takviye elemanı uyumu oldukça önemlidir. İyi mekanik özellikler sergileyen metal matrisli kompozit malzeme üretebilmek için hem karıştırıcı uç profilinin hem de sürtünme karıştırma parametrelerinin iyi belirlenmesi gerekmektedir. Bu parametreler kullanılan takviye elemanları ve matrislere göre farklılık gösterebilmektedir. Uygun parametreler sonucunda hem partikül dağılımı hem de iç yapıda oluşan mekanizmaların etkisinin arttırılması kompozit yapısı için oldukça önemlidir (Akbari, Ezzati, & Asadi, 2022) .


FULL TEXTS BOOK

Tablo 1. Sürtünme karıştırma prosesine etki eden faktörler (Padhy, Wu, & Gao, 2018)



Tek paso ile üretilen yüzey kompozit tabakaya kıyasla, dört paso ile üretilen yüzey kompozit tabaka, birbirinden ayrılmış ve kümelenme boyutları 50–140 nm arasında, ortalama boyutu ise 70 nm olan nano boyutlu Al_2O_3 parçacıklarının iyi bir dağılımını göstermiştir (Şekil 4(c)). Dolayısıyla, 5052 Al matrisi içindeki Al_2O_3 parçacıklarının dağılımı, uygulanan SKP paso sayısıyla ilişkilidir. Bunun nedeni, daha yüksek paso sayısının karıştırma bölgesinde daha büyük şekil değiştirmeye (gerinime) neden olmasıdır. Burada, başlangıç durumundaki haddelenmiş alüminyum alaşım levhanın, üretim sürecinden kaynaklanan nispeten yüksek kalıntı gerinim ve gerilime sahip olduğu bu nedenle alaşımın SKP'nin ilk pasosunda şekil değiştirmesini ve akışını zorlaştırdığı unutulmamalıdır. Ancak, sonraki SKP pasolarında, matris malzemesinin yumuşaması nedeniyle metalin şekil değiştirmesi ve akması daha kolay hale gelmiştir . Sonuç olarak, malzemenin plastik deformasyonu daha şiddetli olmuş ve karıştırılan malzeme daha kolay şekil değiştirmiştir; bu da Al_2O_3 parçacıklarının karıştırma bölgesinde (KB) homojen olarak dağılmasını sağlamıştır. Ayrıca, daha yüksek paso sayılarında kümelenmiş parçacıklar kırılmış ve matris içerisinde daha homojen dağılmıştır (Sharifitabar, Sarani, Khorshahian, & Afarani, 2011).

FULL TEXTS BOOK



Şekil 4. (a) Tek paso, (b) iki paso ve (c) dört paso ile üretilen kompozit karıştırma bölgelerinin SEM görüntüleri (Sharifitabar, Sarani, Khorshahian, & Afarani, 2011)

Asadi ve ekibinin gerçekleştirdikleri çalışmada, (Asadi, Ghader, Besharati, & Kazem, 2010) yüzey kompozit imalatında yüksek dönme hızlarında takviye parçacıklarının kümelerinin daha etkin bir şekilde parçalandığı ve yeni kümelenme oluşumunun zayıflatıldığı bildirilmiştir. Ancak, yüksek dönme hızının artan ısı girişi nedeniyle tane incelmesi üzerinde olumsuz bir etki yarattığı da belirtilmiştir. Bu durum yüksek sıcaklıklarda tane büyüme etkisinin oluştuğunu göstermektedir.


Şekil 5'de sunulan Sharifitabar ve çalışma arkadaşları tarafından gerçekleştirilen çalışmada (Sharifitabar, Sarani, Khorshahian, & Afarani, 2011), farklı SKP koşullarında 5052Al/Al_2O_3 kompozitlerin karıştırma bölgelerinin (KB) optik görüntüleri Şekil 5 (a–k)'da gösterilmektedir. Şekil 5 (a–d)'den görülebileceği gibi, düşük devir/ilerleme oranlarında üretilen numunelerin karıştırma bölgelerinde büyük boşluklar ve tüneller oluştuğu belirtilmiştir. Ayrıca, karıştırıcı takımın düşük eğim açısı, yüksek devir/ilerleme oranında hazırlanan numunelerde bile boşluk oluşumuna neden olmuştur (Şekil 5 (e–g)). Daha yüksek devir/ ilerleme oranının ve yüksek takım açısıyla (5°) üretilen kompozitlerde (Şekil 5 (h–k)) sırasıyla bir, iki ve dört paso ile SKP işlemi uygulanmış numunelerin karıştırma bölgesinde kusurdan uzak daha homojen bir mikroyapı elde edilmiştir. Dolayısıyla, düşük devir/ilerleme oranlarında düşük ısı girdisi, karıştırma sırasında malzeme akışını azaltarak karıştırma bölgesinde büyük boşlukların ve tünellerin oluşmasına yol açtığı bildirilmiştir.

Sato SY. tarafından (Sato, 2004), SKP sırasında ısı girdisinin, takım omuz çapının ve dönüş hızının artmasıyla, ayrıca ilerleme hızının azalmasıyla arttığı bildirilmiştir. Dolayısıyla, düşük devir/ilerleme

FULL TEXTS BOOK

oranlarında düşük ısı girdisi, karıştırma sırasında malzeme akışını azaltarak karıştırma bölgesinde büyük boşlukların ve tünellerin oluşmasına yol açmıştır.

Şekil 5. Farklı devir / dakika SKP parametreleri şartlarında üretilen kompozitlerin optik mikroskop görüntüleri , (a) 8 dev/dk ve u = 2,5; (b) 32 dev/dk ve u = 2,5; (c) 32 dev/dk ve u = 3,5; (d) 64 dev/dk




ve u = 3,5; (e) 100 dev/dk ve u = 2,5; (f) 100 dev/dk ve u = 3,5; (g) 100 dev/dk ve u = 4; (h) 100 dev/dk ve u = 5; (j) 100 dev/dk, u = 5 ve iki paso; (k) 100 dev/dk, u = 5 ve dört paso (Sharifitabar, Sarani, Khorshahian, & Afarani, 2011). (**u:** karıştırıcı takım açısı)

SKP prosesinde kompozit yapısındaki iyileşmelerde büyük önemi olan karıştırma parametreleri takım dönme hızı, ilerleme hızı, paso sayısı ve karıştırıcı uç tipidir. Bu parametreler genel olarak karıştırma bölgesindeki ısı oluşumunu belirlemekle birlikte, mikroyapıdaki tane inceltme etkisinde kompozit dayanımının artmasında büyük rol oynamaktadırlar.

Sürtünme karıştırma prosesinde takım geometrisi, malzeme akışı ve dolayısıyla uygulanabilir ilerleme hızını belirleyen en kritik parametrelerden biridir. Omuz ve pimden oluşan SKP takımı, iki temel işlev gerçekleştirmektedir: lokal ısı üretimi ve malzemenin karıştırılarak taşınması. Karıştırma aşamasında ısı üretimi, öncelikle pim ile iş parçası arasındaki sürtünmeden kaynaklanmaktadır. Takım omzunun iş parçasına temas etmesiyle en yüksek ısı girdisi sağlanmakta olup, bu açıdan pim ve omuz boyutlarının göreceli oranı kritik önem taşımaktadır. Omuz ayrıca ısınan malzeme hacmini kısıtlayarak şekillendirmektedir. Takım tasarımı, işlem sırasında oluşan mikroyapının ve mekanik özelliklerin belirlenmesinde büyük önem taşımaktadır (Akbari, Ezzati, & Asadi, 2022).

FULL TEXTS BOOK



Şekil 6. Sürtünme karıştırma prosesinde genel olarak kullanılan karıştırıcı uç çeşitleri (Akbari, Ezzati, & Asadi, 2022)

Kendi çalışmamızda gerçekleştirdiğimiz deneylerde karıştırma parametreleri aynı fakat farklı türlerde kullanılan triflat, kare ve silindir uçlardan üretilen kompozitlerin kesit görüntüleri Şekil 7'de belirtilmiştir.



Şekil 7. Aynı parametrelerde, farklı karıştırıcı uç profilleriyle gerçekleştirilen SKP işlemi sonrası metalografi numuneleri kesit yüzeyleri.

SKP prosesinde karşılaştığımız kusurlar büyük oranda, takım ile malzeme arasındaki etkileşim süresinin az olması nedeniyle yetersiz plastik deformasyon ve malzeme akışı oluşmasına bağlı olarak oluşmaktadır. Ayrıca bazı çalışmalarda düşük takım eğim açısı, yüksek devir / ilerme oranıyla üretilen numunelerde bile boşluk oluşumuna neden olmuştur. Bunun sebebi eğim açısının plastikleşmiş malzemenin akışında ve takım omzunun malzemeye uyguladığı basıncı arttırdığından dolayı olduğu düşünülmektedir.



Düşük devir / ilerleme oranlarında düşük ısı girdisinin, karıştırma sırasında karıştırma bölgesinde ki malzeme akışını azaltmasına ve bunun sonucunda büyük boşluklar ile tünel kusurunun oluşmasına bağlanmıştır.

Diğer bir unsur olan yetersiz takım dalma derinliği, KB'deki kusurların oluşumuna sebep olan yaygın bir etkendir. Uzunlamasına çatlaklar ve tünel boşlukları gibi kusurların işlem bölgesinde oluşmaması için dalma derinliği, malzemeye yeterli ısı sağlayacak şekilde optimize edilmesi gerekmektedir.

Tablo 2'de SKP işleminde sıklıkla karşılaşılan kusurların tanımları ve hangi durumda meydana geldikleri sunulmuştur.

FULL TEXTS BOOK


Tablo 2. Sürtünme karıştırma prosesinde karşılaşılan hata türleri ve sebepleri

	Kusurlar	Tanım	Sebepleri
	Solunca Deliği	Yetersiz şekilde birleştirilen ve dövülmüş malzemeden oluşan, uzunlamasına yönde ilerleyen bir yan tünel.	Belirli bir dönme hızı için aşırı seyahat hızı Soğuk kaynak Kaynak aralığı çok düşük
	Flaş	Üst yüzeyde aşırı malzeme çıkışı, geri çekilen tarafta oluklu veya şerit benzeri bir etki bırakır.	Aşırı dövme yükü veya dalma derinliği Aşırı sıcak kaynak Çok yüksek kaynak aralığı
	Karıştırma Bölgesi Çöküntüsü	Dinamik olarak yeniden kristalleştirilmiş nugget şeklinin uygun olmayan oluşumu.	Aşırı Akış Kolu Oluşumu İlerleyen Tarafa Malzeme Enjeksiyonu Aşırı sıcak kaynak Çok yüksek kaynak aralığı
	Yüzey aşınması	Pim aletinin altındaki kaynağın üst yüzeyindeki metalin aşınması ve yırtılması.	Metal pim aletine yapışma Aşırı sıcak kaynak Çok yüksek kaynak aralığı

ARAŞTIRMA

Gerçekleştirdiğimiz çalışmada AA1050 saf alüminyum matris malzemesine ortalama 136 nm partikül boyutlarında olan nanografi tarafından temin edilen alüminyum oksit (Şekil 9) partikülleri delik delme yöntemiyle ilave edilerek sürtünme karıştırma prosesiyle metal matrisli kompozit üretimi gerçekleştirilmiştir (Şekil 8).

FULL TEXTS BOOK




Şekil 8. a-b) SKP işlemi aşaması, c) Sertlik,metalografi ve çekme numunelerinin karıştırma bölgesinden çıkarılması , d) Nihai ürün

AA1050 saf alüminyum levhaya ait mekanik özellikler Tablo 3'de belirtilmiştir.

Tablo 3. AA1050 mekanik özellikleri

Özellikler	**Değer**
Akma Dayanımı (MPa)	53.2 MPa
Çekme Dayanımı (MPa)	84.2 MPa
Uzama %	25.4
Sertlik (HV)	22±1

FULL TEXTS BOOK



Şekil 9. Alüminyum oksit partikülleri , b) SEM görüntüsü

Sürtünme karıştırma prosesiyle kompozit üretimi aşamasında Tablo 4'de ki parametrelere bağlı olarak SKP işlemi yürütülmüştür. Bu parametreler ile kompozit yapılarının kendi içlerinde farklı takım devri, ilerleme hızı ve paso sayısının kompozit dayanımı üzerindeki etkileri araştırılmıştır.

Tablo 4. Delik delme yöntemiyle toz ilavesi yapılan matris malzemesinde sürtünme karıştırma prosesi parametreleri.

Numune Kodu	Takım Devri	İlerleme Hızı	Paso Sayısı	Karıştırıcı Uç Cinsi	Alümina Toz Miktarı(g)
A1	1250	31.5 mm/dk	3	Kare	0,38
A2	1250	63 mm/dk	3	Kare	0,38
B1	800	31.5 mm/dk	3	Kare	0,38
B2	800	63 mm/dk	3	Kare	0,38
C1	1250	31.5 mm/dk	1	Kare	0,38
C2	1250	63 mm/dk	1	Kare	0,38
D1	800	31.5 mm/dk	1	Kare	0,38
D2	800	63 mm/dk	1	Kare	0,38

Elde edilen nihai üründen çekme testi, sertlik ve metalografi numuneleri çıkartılmıştır (Şekil 8c). Bu numuneler de metalografik ve mekanik analizler gerçekleştirilerek alüminyum oksit takviyeli SKP uygunlanmış AA1050 malzemenin ana malzemeye kıyasla dayanım ve metalurjik etkilerinin incelenmesi gerçekleştirilmiştir.

Ayrıca sürtünme karıştırma prosesi parametrelerine bağlı olarak gelişen iç yapı dayanım mekanizmalarının ve kusurların oluşumu da irdelenmiştir.


FULL TEXTS BOOK

BULGULAR

Sürtünme karıştırma prosesi (SKP) sırasında oluşan şiddetli plastik deformasyon ve ısı üretimi, işlenen bölgede kendine özgü mikroyapısal özelliklerin gelişmesini sağlamaktadır. Bu süreçte sıcaklık, ısıl genleşme ve iletim katsayısı gibi değişkenler, SKP bölgesinde, karıştırma bölgesi (KB), termomekanik etki altındaki bölge (TEB) ve ısı tesiri altındaki bölge bölge (ITAB) olmak üzere üç farklı karakteristik bölge oluşmasına sebep olmaktadır (Chumaevskiy, Tarasov, Zykova, & Kolubaev, 2020). Karıştırma bölgesi, takımın doğrudan etkisi altındaki alandır ve bu bölgede malzeme yüksek sıcaklıklarda yoğun plastik deformasyona maruz kaldığı için dinamik yeniden kristalleşme meydana gelir. Bu da ince ve eş eksenli tanelerden oluşan bir mikroyapı oluşturmaktadır. Tane boyutları ve homojen yapı sayesinde bu bölge, malzemenin mekanik özelliklerini en çok etkileyen alan haline gelmektedir. Karıştırma bölgesinin boyutu ve şekli, esas olarak takımın geometrisi, dönme hızı, ilerleme hızı, malzemenin ısıl iletkenliği ve soğuma koşulları gibi parametrelerle belirlenir. Karıştırma bölgesinin hemen dışında yer alan termomekanik etkilenmiş bölge ise sınırlı plastik deformasyonun meydana geldiği ancak yeniden kristalleşmenin tamamlanmadığı bir geçiş bölgesidir. Bu alanın ötesinde bulunan ısıdan etkilenmiş bölge ise plastik deformasyon içermemekle birlikte sadece sıcaklık etkisiyle bazı mikroyapısal değişikliklerin meydana geldiği alandır. Bu bölgenin mikroyapısı, genellikle ana malzemeninkine çok benzer kalır. Tüm bu bölgelerin genişliği ve özellikleri, uygulanan SKP parametrelerine ve malzemenin özelliklerine bağlı olarak değişkenlik gösterebilir.



Şekil 10. (a) SZ, TMAZ, HAZ ve BM'yi gösteren optik mikroskop görüntüsü (Heidarzadeh, ve diğerleri, 2021)




Şekil 11. A) Numune A1 karıştırma bölgesi, B) Temel malzeme SEM görüntüsü

SKP (Sürtünme Karıştırma Prosesi) sırasında oluşan yoğun plastik deformasyon, karıştırma bölgesinde dinamik yeniden kristalleşmiş bir yapının oluşmasına neden olmuştur. Bu sayede temel metale kıyasla

FULL TEXTS BOOK


daha homojen ve belirgin şekilde incelmiş taneler gözlenmektedir. Şekil 11'de aynı büyütme oranlarında verilen SEM görüntülerinde, işlem uygulanmış bölge ile işlem görmemiş temel malzemenin mikroyapıları karşılaştırılmıştır. Karşılaştırma sonucunda, SKP sonrası mikroyapının daha ince tanelerden oluştuğu açıkça görülmektedir. Bu bulgu, sürtünme karıştırma prosesinin dinamik yeniden kristalleşme yoluyla malzemenin mikroyapısını iyileştirdiğini ortaya koymaktadır.



Şekil 12. A1-A2 numuneleri SEM görüntüleri.


A1 ve A2 numuneleri, 1250 devir/dk takım hızı ve üç paso ile işlenmiştir. A1 numunesinde ilerleme hızının düşük (31,5 mm/dk) olması, nano takviye partiküllerinin matris içerisinde daha homojen dağılmasını, parçacık boyutlarının küçük kalmasını ve aglomerasyonun minimum seviyede gerçekleşmesini sağlamıştır. Bu sonuç, malzeme içinde yeterli plastik deformasyonun oluştuğunu ve karıştırmanın etkili şekilde gerçekleştiğini göstermektedir. Buna karşılık, A2 numunesinde yüksek ilerleme hızı (63 mm/dk) kullanıldığından, partikül dağılımında homojenlik bozulmuş, bazı bölgelerde iri parçacıkların oluştuğu ve parçacıklar arası mesafenin arttığı belirlenmiştir. Dolayısıyla, ilerleme hızındaki artışın karıştırma etkinliğini olumsuz yönde etkilediği anlaşılmaktadır.

FULL TEXTS BOOK



Şekil 13. B1-B2 numuneleri SEM görüntüleri.


B1 ve B2 numuneleri düşük takım devri (800 devir) ve üç paso kullanılarak üretilmiştir. B1 numunesi (31,5 mm/dk) elde edilen en başarılı mikroyapılardan birini sergilemiştir. Bu numunede alüminyum oksit parçacıkları oldukça küçük boyutlarda, yoğun ve homojen şekilde dağılmıştır. Düşük takım devri ile düşük ilerleme hızının birleşimi, kontrollü bir ısıl etki ve güçlü bir karıştırma sağlamış, bunun sonucunda yüksek kaliteli bir kompozit yapı ortaya çıkmıştır. Buna karşılık, B2 numunesinde yüksek ilerleme hızı (63 mm/dk) nedeniyle karıştırma yetersiz kalmış, parçacıkların kümelendiği ve düzensiz bir dağılımın oluştuğu belirlenmiştir.



Şekil 14. C1-C2 numuneleri SEM görüntüleri.


C1 ve C2 numuneleri ise yüksek devirde (1250 devir) fakat yalnızca tek paso ile işlenmiştir. C1 numunesinde partikül dağılımı, A1 numunesine göre daha düşük homojenlik göstermiştir. Bu durum, takım işlem sayısının parçacık dağılımı üzerinde kritik bir rol oynadığını göstermektedir. C2 numunesinde ise aglomerasyonun oldukça yoğun olduğu tespit edilmiştir. Tek paso ile işlem yapılması ve yüksek ilerleme hızının birleşmesi, parçacıkların yüzeyde kümelenmesine ve heterojen bir mikroyapının oluşmasına yol açmıştır.

FULL TEXTS BOOK



Şekil 15. D1-D2 numuneleri SEM görüntüleri.

D1 ve D2 numuneleri en düşük deformasyon koşullarında (800 devir, tek paso) üretilmiştir. D1 numunesinde parçacıkların bir kısmı dağılmış olmakla birlikte, bazı bölgelerde kümelenmelerin oluştuğu ve parçacık boyutlarının büyüdüğü gözlenmiştir. D2 numunesi mikroyapısında iri parçacıkların, düzensiz hatların ve alüminyum oksit partiküllerinin boşluk kenarlarında yoğun kümelenmeler meydana getirdiği görülmüştür.




Şekil 16. SKP ile üretilen B1-D1 kompozitlerinin sırasıyla 5K-10K-20K büyütme oranlı SEM görüntüleri

FULL TEXTS BOOK

Farklı parametrelere bağlı olarak gerçekleştirilen SKP prosesiyle kompozit üretimi sonrası elde edilen yapılardan sırasıyla en iyi ve en kötü olan B1-D1 numunelerinin iç yapılarının karşılaştırılması Şekil 16'da sunulmuştur.


B1 numunesinde (800 dev/dk, 31,5 mm/dk, üç paso), Al_2O_3 partikülleri matriste homojen şekilde dağılmış, ortalama tane boyutu 78–170 nm aralığında olup boşluk oranı düşük ve yoğun bir yapı elde edilmiştir. Yoğun plastik deformasyon ve dinamik yeniden kristalleşme sonucu tane incelmesi meydana gelmiştir. Buna karşılık D1 numunesinde (800 dev/dk, 31,5 mm/dk, tek paso) yetersiz plastik deformasyon nedeniyle partikül kümelenmesi, boşluklar ve daha büyük tane boyutları (136–321 nm) gözlenmiştir.



Şekil 17. a) B1 Karıştırma bölgesi, b) Termomekanik etki altındaki bölge ,c) TEB-ITAB alanı SEM görüntüsü

Numunelerin işlenmiş bölgesine ait optik mikroskop ve SEM görüntülerinde, SKP işleminin karıştırma bölgesinde (Şekil 17(a)) malzeme akışı üzerinde etkisi olduğunu ve mikroyapıda termomekanik etki altındaki bölge ile ısı tesiri altındaki malzeme arasında daha ince taneli yapı oluşmasını sağladığı görünmektedir. Bu olay sürtünme karıştırma işlemi sırasında, sıcaklıktaki artış ve şiddetli plastik deformasyona bağlı olarak gelişen dinamik yeniden kristalleşme etkisinin, karıştırma bölgesinde yeni ince eş eksenli tanelerin oluşmasına yol açmasıyla açıklanmaktadır (Chumaevskiy, Tarasov, Zykova, & Kolubaev, 2020).

FULL TEXTS BOOK



Şekil 18. Farklı numunelere ait sem ve optik mikroskop görüntüleri , a-b) A2 numunesi,c-d) C2 numunesi, e-f) D1 numunesi, g-h) D2 numunesi.

SKP prosesi kompozit yapısında sağladığı iyileştirme etkilerinden ziyade bir takım etkenler sebebiyle bazı kusurların oluşumuna da sebebiyet verebilmektedir. Şekil 18'de ki numunelerde aşırı plastikleşmiş malzeme iyi seviyede birleşemediği ve bunun sonucu olarak ayrışmalar meydana geldiği düşünülmektedir. Bu durum genellikle;

• Yetersiz plastik deformasyon

• Düşük takım omuz basıncı

• Uygun olmayan takım tasarımı

• Düşük devir hızı veya yetersiz ısı girişi

FULL TEXTS BOOK

• Aşırı plastikleşmiş akışın tam birleşmemesi,


• Partikül kümelenmesi,

gibi unsurlarla nitelendirilmiştir. Bu tür kusurlar, kompozit dayanımının düşmesine sebep olabilecek, gözle görülebilen ayrık tabakalar veya birleşmemiş alanlar olarak ortaya çıkmaktadır (Zykova, Tarasov, & Chumaevskiy, 2025).

Bu değerlendirmelerin yanı sıra tabakalaşma kusurunun oluşumunda, sürtünme karıştırma prosesi sırasında yüzey ile malzeme altı arasında oluşan sıcaklık farkı da etkili olabilmektedir. Ayrıca, kalıpla temas eden alt yüzeyin ısı iletimi nedeniyle işlem sonrası daha hızlı soğuması, arayüzeyde sıcaklık değişiminden kaynaklı gerilim farkı oluşmuş olabilir. Bu durum çekme gerilmeleriyle birleşerek tabakalaşma kusurunun meydana gelmesine sebebiyet verebilmektedir.

Bu kusur, yüksek sıcaklık, yetersiz karıştırma veya uygunsuz işleme parametrelerinin yanı sıra yukarıda açıklanan etkenlerinde etkisiyle oluşabilmektedir (Kim, Fujii, Tsumura, Komazaki, & Nakata, 2006).

Grafik 1. Çekme Testi Ölçümleri



Tablo 5. Çekme testi değer tabloları ve artış / azalış oranları.

Numune	Çekme Dayanımı (MPa)	Çekme Artış/Azalış Oranı (%)	Akma Dayanımı (MPa)	Akma Artış/Azalış Oranı (%)	Uzama (%)	Uzama Artış/Azalış Oranı (%)
BM	84,2		53,2		25,4	
A1	117,7	40%	77,1	45%	29,7	17%
A2	112,8	34%	77,0	45%	30,6	20%
B1	151,4	80%	104,2	96%	19,3	-76%
B2	116,7	39%	90,7	70%	14,8	-58%
C1	129,2	53%	82,3	55%	35,6	40%
C2	110,7	31%	90,1	69%	9,5	-37%
D1	96,5	15%	75,2	41%	17,6	-69%
D2	106,1	26%	81,0	52%	8,8	-35%

Tablo 5'de, üretilen kompozitlerin çekme, akma ve uzama değerlerinin oransal olarak karşılaştırmaları verilmiştir.

Sürtünme karıştırma prosesi ile üretilen yüzey metal matrisli kompozitlerde mukavemet artışı, birkaç temel mekanizma ile sağlanmaktadır. İşlem sırasında takımın dönme ve ilerleme hareketi, karıştırma bölgesinde aşırı plastik deformasyon ve ısının etkisiyle dinamik rekristalizasyonu tetikler ve ince taneli bir yapı oluşur. İnce taneli yapı, tane sınırlarının sayısını artırarak dislokasyon hareketini zorlaştırır; bu durum Hall–Petch mekanizması ile malzemenin mukavemetinin artmasını sağlar. Yüzeye eklenen sert ikincil faz partikülleri, matris yapısına parçalanarak dağıldığında tane sınırlarının hareketini engeller ve tane büyümesini yavaşlatarak Zener sabitleme etkisi yaratır; böylece ince taneli yapının korunması sağlanır ve dayanım devam ettirilir. Ayrıca, bu partiküller dislokasyonların doğrudan geçişini engelleyerek Orowan mekanizmasına uygun şekilde dislokasyonların partiküller etrafından bükülmesini zorunlu kılar; bunun sonucunda kayma gerilmesi artar ve malzemenin mukavemeti yükselir. Bu mekanizmaların birlikte etkisi, yüzey metal matrisli kompozitlerin yüksek dayanım göstermesini sağlamaktadır (Sharma, Prakash, & Kumar, 2015) .

Kompozitlerin yapısına eklenen alüminyum oksit nanopartiküllerinin kümelenmesi malzemenin dayanımını açısından oldukça zararlı olabilmektedir çünkü bu parçacıklar çatlak oluşumu için başlangıç noktası görevi görerek tokluk değerlerinde düşüşe sebebiyet verebilmektedir. Şekil 15'den de görülebileceği üzere en düşük dayanıma sahip olan D1 numunesinde nanoparçacıkların dağılımındaki düzensizlik ve kümelenmeler bölgesel gerilmelere sebep olmuştur ayrıca takviye partiküllerinin matris ile olan zayıf ıslanabilirliği , matristen daha dayanıklı olan partiküllere yük aktarımını sınırlandırmaktadır ve malzemenin dayanımının düşmesine sebep olmaktadır (Mohammed, ve diğerleri, 2024), (Aruri, Adepu, Bazavada, & Kumaraswamy , 2013).

B1 numunesinin diğer numunelere kıyasla en yüksek gerilme mukavemetine (151,4 MPa) ulaşmasının sebebi Şekil 13'den de görülebileceği üzere mikroyapının daha yoğun olması, sıkışması ve boşlukların

FULL TEXTS BOOK


azalmasıyla daha iyi bir iç yapı sergilemiş olmasıdır. Takım ilerleme hızının düşük, paso sayısının yüksek olması birim uzunluk başına ısı girdisini artırarak plastik deformasyonun büyüklüğünü ve karışımı iyileştirmektedir.

Genel olarak çekme grafiği incelendiğinde üç pasolu sürtünme karıştırma prosesi uygulamalarında, mukavemet ve gerinim değerleri daha iyi görünmektedir. Sürtünme karıştırma prosesinde paso sayısının artması, işlem bölgelerinde ki tane incelmesi üzerinde önemli bir etkiye sahiptir. Her bir SKP pasosu, dönen takımın mekanik karıştırma etkisiyle malzemede şiddetli plastik deformasyona neden olur; bu durum, dislokasyon yoğunluğunu artırır ve dinamik yeniden kristalleşmeyi kolaylaştırır. Paso sayısının artması deformasyon miktarının da artmasını sağlayarak iri tanelerin daha ince, eş eksenli tanelere dönüşmesini sağlamaktadır. Artan paso sayısıyla birlikte mikroyapı daha homojen hale gelir ve ortalama tane boyutu azalmaya devam eder (Bauri, Yadav, & Suhas, 2011).

Sürtünme karıştırma prosesiyle farklı parametrelerden üretilen numunelerin mikrosertliği üzerindeki etkisi Grafik 2'de sunulmuştur. AA1050 ana malzeme sertliği yaklaşık 22 HV olarak ölçülmüştür (Şekil 19(a)). Kompozit kesit alanında sürtünme karıştırma yüzeyinden 1,5 mm aşağıda lineer bir hat üzerinde 1000 g yükte 30 s boyunca değişken aralıklarla sertlik ölçümleri gerçekleştirilmiştir.

Sürtünme karıştırma işlemi sırasında takımın ve çevresindeki matrisin Al_2O_3 parçacıkları üzerinde yarattığı yoğun etki, parçacık boyutlarının küçülmesine ve matris tanelerinin incelmesine neden olmuştur. Bu durum Hall-Petch ilişkisine uygun olarak mikrosertlik artışını sağlamıştır. Ayrıca, incelmiş Al_2O_3 partiküllerinin matris içerisinde homojen bir şekilde dağılması yalnızca sertlik değerlerini yükseltmekle kalmamış, aynı zamanda dalgalanmaların da azalmasına katkı sağlamıştır. Takviye parçacıklarının sadece karıştırma bölgesinde değil, aynı zamanda TEB ve ITAB bölgelerine de yayılması, kompozit malzemenin sertlik profilinin oldukça düzgün bir şekilde oluşmasına imkân tanımıştır (Tang, Shen, & Li, 2019).

Grafik 2. Sertlik Ölçümleri



Düşük takım devri (800 dev/dk) ve artan paso sayısı ile gerçekleştirilen işlemlerde ise karıştırma bölgesinde 40 HV'ye (B1) ulaşan sertlik değerleri elde edilmiştir. Bu sonuç, hem uygun bir ısıl girişin sağlandığını hem de takviye parçacıklarının matris içerisinde daha verimli ve homojen dağıldığını


FULL TEXTS BOOK

göstermektedir. Ayrıca, çoklu SKP geçişleri sayesinde parçacıkların yatay ve dikey yönde karışması kolaylaşmış, bu da mikroyapısal bütünlüğü güçlendirmiş ve tanelerin daha ince hale gelmesini sağlamıştır.

En yüksek sertlik değerlerinin elde edildiği B1 numunesinde ortalama parçacık boyutları 105 nm olarak ölçülürken, en düşük sonucu veren D1 numunesinde parçacık boyutu 180 nm'ye kadar çıkmıştır.

Tablo 6. SKP uygulanan numunelerin minimum ve maksimum sertlik değerleri.

Numune	**A1**	**A2**	**B1**	**B2**	**C1**	**C2**	**D1**	**D2**
Min. Mikrosertlik (HV)	26,8	26,7	27,2	26,4	27,2	26,1	23,8	26,3
Max. Mikrosertlik (HV)	29,4	32,5	40,2	39	30	31,3	29,8	30,4



Şekil 19. TEB- ITAB sertlik farkı a) Ana malzeme sertlik değeri, b) B1 numunesi , c) D1 numunesi termomekanik etki altındaki ve ısı tesiri altındaki bölge sertlik farkı

Şekil 19'da B1 ve D1 numunesine ait SKP sonrası termomekanik etki altındaki bölge (TEB) ile ısı tesiri altındaki (ITAB) bölgeleri arasındaki sertlik farklılığı gösterilmiştir. Bu etki açıklanan mekanizmaların

FULL TEXTS BOOK

ışığında sürtünme karıştırma prosesinin etkilediği alanların diğer alanlara göre daha iyi mekanik özellik sergilediğini göstermektedir .

Sürtünme karıştırma prosesi sırasında oluşan termomekanik etki, matriste dinamik yeniden kristalleşme ile birlikte önemli ölçüde tane incelmesine yol açmaktadır. Ancak, yeniden kristalleşme sonrası oluşan yeni taneler termodinamik olarak büyüme eğilimindedir. Bu noktada, matris fazı içerisinde homojen olarak dağılmış olan çözünmeyen takviye parçacıkları, tane sınırlarının ilerlemesini fiziksel olarak engelleyerek Zener sabitleme (pinning) adı verilen mekanizma ile devreye girmektedir. Bu sayede mikroyapının ince taneli yapısı korunmuş olur. Özellikle nano boyutta ve yüksek hacim oranına sahip takviye fazlarının kullanılması, pinleme etkisini daha belirgin hale getirerek tane büyümesini büyük ölçüde bastırmaktadır. Bu durum, SKP ile üretilmiş metal matrisli kompozitlerde mekanik özelliklerin iyileştirilmesinde kritik bir rol oynamaktadır (A. Dolatkhah, 2012) (Mohammed, ve diğerleri, 2024).

Sonuç olarak, SKP ile kompozit üretiminde sertlik artışını sağlamak yalnızca takviye malzemesi eklenmesiyle mümkün olmamakta; işlem parametrelerinin, paso stratejisinin ve takım geometrisinin bütünsel olarak optimize edilmesi gerekmektedir. Aksi durumda, özellikle temperlenmiş malzemelerde, ısı geri dönüşümü ve yeniden kristalleşme kaynaklı tane büyümesi gibi mikroyapısal bozulmalar kaçınılmaz hale gelmekte ve sertlik, hedeflenen değerlerin altına düşmektedir (Mahmoud, Ikeuchi, & Takashi, 2008).

SONUÇ

Bu çalışmada, AA1050 saf alüminyum matrisli kompozit malzemeler, Al_2O_3 nanopartiküllerıyle takviye edilerek sürtünme karıştırma prosesi (SKP) ile yüzey kompozitleri üretilmiştir. Delik delme yöntemiyle matrise dahil edilen takviye partiküllerinin, farklı SKP parametreleri (takım devri, ilerleme hızı, paso sayısı ve karıştırıcı uç geometrisi) altında mikroyapısal ve mekanik özelliklere etkileri detaylı şekilde incelenmiştir.

Elde edilen bulgulara göre, düşük ilerleme hızı ve yüksek paso sayısı, Al_2O_3 parçacıklarının matriste daha homojen dağılmasını sağlayarak daha yoğun ve boşluksuz bir yapı oluşmasına katkı sağlamıştır. Bu durum, özellikle B1 numunesi yüksek çekme dayanımı ve sertlik artışı olarak gözlenmiştir. Ayrıca, SKP işlemi sırasında gelişen şiddetli plastik deformasyon ve dinamik yeniden kristalleşme sayesinde ince taneli bir yapı elde edilmiş, bu da Hall–Petch pekiştirme mekanizmasını devreye sokarak malzeme mukavemetini artırmıştır.

Çalışmada ayrıca, yetersiz plastik deformasyona sebebiyet verebilecek tek pasolu ve düşük devirli işlem (D1-D2) oluşan, partikül kümelenmesi, yetersiz plastik deformasyon ve tabakalaşma gibi mikroyapısal kusurların oluştuğu gözlemlenmiştir. Bu tür kusurlar, matris-takviye arayüzeyinde bağ zayıflığına ve dolayısıyla mekanik özelliklerde düşüşe yol açmıştır.

Sonuç olarak, bu çalışmada kullanılan SKP ve takviye yöntemlerinin doğru parametrelerle uygulanması durumunda, AA1050 alüminyum alaşımı esaslı kompozitlerde belirgin bir mekanik iyileşme sağlandığı görülmüştür. Al_2O_3 takviyesinin varlığı, Orowan mekanizması, dislokasyon yoğunluğu artışı ve Zener sabitleme gibi pekiştirme mekanizmalarının bir arada etkili olmasıyla birlikte, hem mikroyapı hem de mekanik özellikler açısından üstün nitelikli yüzey kompozitleri elde edilmiştir.

KAYNAKÇA

A. Dolatkhah, P. G. (2012). Investigating effects of process parameters on microstructural and mechanical properties of Al5052/SiC metal matrix composite fabricated via friction stir processing. s. 458-464. doi:https://doi.org/10.1016/j.matdes.2011.09.035.

Akbari, M., Ezzati, M., & Asadi, P. (2022). Investigation of the effect of tool probe profile on reinforced particles distribution using experimental and CEL approaches. *International Journal of Lightweight Materials and Manufacture*, 213-223. doi:10.1016/j.ijlmm.2022.02.002

Aruri, D., Adepu, K., Bazavada, K., & Kumaraswamy , A. (2013). Wear and mechanical properties of 6061-T6 aluminum alloy surface hybrid composites [(SiC+Gr) and (SiC+Al2O3)] fabricated by friction stir processing. *Journal of Materials Research and Technology*, 362-369. doi:10.1016/j.jmrt.2013.10.004

Asadi, P., Ghader, F., Besharati, G., & Kazem, M. (2010). Producing of AZ91/SiC composite by friction stir processing. *International Journal of Advanced Manufacturing Technology*, 247-260. doi:10.1007/s00170-010-2600-z

Bauri, R., Yadav, D., & Suhas, G. (2011). Effect of friction stir processing (FSP) on microstructure and properties of Al–TiC in situ composite. *Materials Science and Engineering: A*, 4732-4739. doi:10.1016/j.msea.2011.02.085

Blaza, S., & Lozica, I. (2015). Application of aluminium hybrid composites in automotive industry. *Tehnicki Vjesnik*, 247-251. doi:10.17559/TV-20130905094303

Chawla, N., & Chawla, K. K. (2006). *Metal Matrix Composites.* United States of America: Springer Science & Business Media.

Chumaevskiy, A. V., Tarasov, S. Y., Zykova, A. P., & Kolubaev, E. A. (2020). A Review of Friction Stir Processing of Structural Metallic Materials: Process, Properties, and Methods. *Metals*. doi:10.3390/met10060772

Gangil, N., Maheshwari, S., Siddiquee, A. N., Abidi, M. H., & El-Meligy, M. A. (2019). Investigation on Friction Stir Welding of Hybrid Composites Fabricated on Al-Zn-Mg-Cu Alloy Through Friction Stir Welding Processing. *Journal of Materials Research and Technology*, 3733-3740.

Gürbüz, M., & Mutuk, T. (2019). Karbon Esaslı Malzeme Takviyeli Titanyum Kompozitler ve Grafen Üzerine Yeni Eğilimler. *Mühendis ve Makina*, 101-118.

Heidarzadeh, A., Mironov, S., Kaibyshev, R., Çam, G., Simar, A., Gerlich, A., . . . Withers, P. J. (2021). Friction stir welding/processing of metals and alloys: A comprehensive review on microstructural evolution. *Progress in Materials Science*. doi:10.1016/j.pmatsci.2020.100752

Kalemtaş, A. (2014). Metal Matrisli Kompozitlere Genel Bir Bakış. *Putech&Composites*, 18-30.

Kim, Y. G., Fujii, H., Tsumura, T., Komazaki, T., & Nakata, K. (2006). Three defect types in friction stir welding of aluminum die casting alloy. *Materials Science and Engineering: A*, 250-254. doi:10.1016/j.msea.2005.09.072

Kraiklang, R., Onwong, j., & Santhaweesuk, c. (2020). Multi Performance Characteristics of AA5052 + 10% SiC Surface Composite by Friction Stir Processing. *Journal of Composites Science*. doi:10.3390/jcs4020036

Mahmoud, E. R., Ikeuchi, K., & Takashi, M. (2008). Fabrication of SiC particle reinforced composite on aluminium surface by friction stir processing. *Science and Technology of Welding and Joining*, 607-618. doi:10.1179/136217108X333327

FULL TEXTS BOOK

Mohammed, M., Ahmed, F. S., Derazkola, H. A., Elsayed, M., Kubit, A., & Nafea, M. (2024). AA5754–Al2O3 Nanocomposite Prepared by Friction Stir Processing: Microstructural Evolution and Mechanical Performance. *Journal of Manufacturing and Materials Processing*. doi:10.3390/jmmp8020058

Nturanabo, F., Masu, L., & Kirabira, J. B. (2020). Novel Applications of Aluminium Metal Matrix Composites. (K. Cooke, Dü.) *Aluminium Alloys and Composites*. doi:10.5772/intechopen.81519

Padhy, G. K., Wu, C. S., & Gao, S. (2018). Friction stir based welding and processing technologies - processes, parameters, microstructures and applications: a review. *Journal Material Science Technology*. doi:10.1016/j.jmst.2017.11.029

Patel, V., Vairis, A., Badheka, V., & Li, W. (2019). Recent development in friction stir processing as a solid-state grain refinement Technique : microstructural evolution and property enhancement. *Critical Review in Solid State and Material Sciences*. doi:10.1080/10408436.2018.1490251. 0

Sato, Y. S. (2004). Post-weld formability of friction stir welded Al alloy 5052. *Materials Science and Engineering*, 138-143. doi:10.1016/j.msea.2003.11.035

Sharifitabar, M., Sarani, M., Khorshahian, A., & Afarani, S. (2011). Fabrication of 5052Al/Al2O3 Nanoceramic Particle Reinforced Composite via Friction Stir Processing Route. *Materials and Design*, 4164–4172. doi:10.1016/j.matdes.2011.04.048

Sharma, A. K., Bhandari, R., Aherwar, A., Rimašauskienė, R., & Bretotean, C. P. (2020). A study of advancement in application opportunities of aluminum metal matrix composites. *Materials Today: Proceedings*, 2419-2424. doi:10.1016/j.matpr.2020.02.516

Sharma, V., Prakash, U., & Kumar, B. V. (2015). Surface composites by friction stir processing: A review. *Journal of Materials Processing Technology*, 117-134. doi:10.1016/j.jmatprotec.2015.04.019

Tang, J., Shen, Y., & Li, J. (2019). Investigation of microstructure and mechanical properties of SiC/Al surface composites fabricated by friction stir processing. *Materials Research Express*. doi:10.1088/2053-1591/ab3a75

Zweben, C. (2002). Metal matrix composites, ceramic matrix composites, carbon matrix composites and thermally conductive polymers matrix composites. *Harper CA (ed) Handbook of plastics, elastomers, and composites, 4th edn.* içinde New York: McGraw Hill. doi:10.1002/9781118985960.meh110

Zykova, A. P., Tarasov, S. Y., & Chumaevskiy, A. V. (2025). Advances in friction stir processing of aluminum 2024: a review of nanoparticle-reinforced surface composites. *International Journal Advanced Manufacturing Technology*, 3141–3164. doi:10.1007/s00170-025-15332-1

FULL TEXTS BOOK

RUS DİLİ SİNTAKSİSİNİN AZƏRBAYCAN DİLİNİN RƏSMİ ÜSLUBUNDA İZLƏRİ

THE INFLUENCE OF RUSSIAN SYNTAX IN THE OFFICIAL STYLE OF THE AZERBAIJANI LANGUAGE

Təranə NAĞIYEVA

Azərbaycan Dövlət Pedaqoji Universiteti

filologiya üzrə fəlsəfə doktoru

https://orcid.org/0000-0002-5322-2446

ÖZET

Dil cəmiyyətin sosial funksiyası hesab olunur. Cəmiyyət inkişaf etdikcə onun əsas sütunlarından biri olan dil də mütəmadi olaraq inkişaf edir, təkmilləşir.

Dilin inkişafı çox mürəkkəb proses kimi qiymətləndirilir. Bu baxımdan dillər arasında mövcud əlaqələr dilin inkişafı prosesində böyük önəm daşıyır. Dillərarası əlaqələr sadə şəkildə bir dilin digər dilə müəyyən formada təsiri kimi başa düşülür, ancaq bu təsir düşündüyümüz kimi heç də asan deyil. Dillərin bir-birinə təsiri dilin bütün səviyyələrində müşahidə olunur. Həmçinin bu təsir sintaksisdən də yan ötməmişdir. Azərbaycan dilinin sintaktik sistemi özünün qədim köklərinə malik olmaqla yanaşı, tarix boyu müxtəlif xarici təsirlərə məruz qalmışdır. Bu təsirlər nəticəsində xüsusilə rus dilinin təsiri XIX əsrin ikinci yarısından başlayaraq daha da güclənmiş və sovet dövründə ən zirvə nöqtəsinə çatmışdır. Azərbaycan xalqı uzun müddət rus dövlətinin tərkibində yaşadığı üçün istənilən sahədə rus alınmalarını görmək mümkündür. Bu təsiri bədii üslubda, mətbuat dilində, elmi üslubda, epistolyar üslubda, məişət üslubunda, həmçinin rəsmi-işgüzar üslubda da görmək olur.

Rus dilinin Azərbaycan dilinin sintaktik quruluşuna təsiri mövzusundan bəhs edərkən Azərbaycan-rus dil əlaqələrinin tarixi əsaslarından danışmamaq olmaz. Belə ki, düşünürük ki, Azərbaycan-rus dil əlaqələrini aşağıdakı şəkildə qruplaşdıra bilərik:

1) XVIII əsrdən başlayan dil əlaqələri;
2) Rus dilinin dövlət dili statusu dövründə təsirləri;
3) Sovet dövründə təhsil və mətbuat vasitəsilə dilə təsir;
4) Qanunvericilik, dövlət idarəçiliyi və inzibati idarəetmə sistemi sahəsində təsirlər.

Rus dilinin Azərbaycan dilinin sintaksisinə təsirini nəzərə alıb deyə bilərik ki, bu təsir bir sıra qrammatik yeniliklərin meydana gəlməsinə səbəb olmuşdur:

1. Feili birləşmələrin isimləşməsi hallarının artması;
2. Modal konstruksiyaların rus sintaksisinə uyğun qurulması;
3. “Öz” əvəzliyi ilə qurulmuş ifadələrin işlənməsi;
4. Cümlə üzvlərinin yerdəyişməsi;
5. Rus dilinin sintaksisinə uyğun mürəkkəb cümlələrin yaranması.

Açar sözlər: Azərbaycan dili, rus dili, əlaqə, sintaktik təsir, rəsmi üslub.

ABSTRACT

Language is considered the social function of society. As society develops, language — being one of its main pillars — also develops and improves regularly.

The development of language is evaluated as a very complex process. From this point of view, the existing relations between languages play a major role in the process of language development. Interlingual relations are simply understood as the influence of one language on another in a certain form, but this influence is not at all as easy as it may seem. The influence of languages on one another

FULL TEXTS BOOK

is observed at all levels of language. Moreover, this influence has not bypassed syntax either. The syntactic system of the Azerbaijani language, in addition to having ancient roots, has been exposed to various external influences throughout history. As a result of these influences, especially from the second half of the 19th century, the influence of the Russian language became stronger and reached its peak during the Soviet period.

Since the Azerbaijani people lived within the Russian state for a long time, Russian borrowings can be seen in all fields. This influence can be seen in the literary style, in the language of the press, in the scientific style, in the epistolary style, in the colloquial style, as well as in the official-business style.

When speaking about the influence of the Russian language on the syntactic structure of the Azerbaijani language, it is impossible not to mention the historical foundations of Azerbaijani-Russian language relations. Thus, we believe that Azerbaijani-Russian language relations can be grouped as follows:

1. Language relations beginning from the 18th century;
2. Influences during the period when the Russian language had the status of state language;
3. Influence on the language through education and the press during the Soviet period;
4. Influences in the field of legislation, state governance, and the administrative management system.

Taking into account the influence of the Russian language on the syntax of the Azerbaijani language, it can be said that this influence led to the emergence of several grammatical innovations:

1. Increase in the cases of nominalization of verbal constructions;
2. Construction of modal structures in accordance with Russian syntax;
3. Usage of expressions formed with the pronoun “öz” (self);
4. Word order shifts within the sentence;
5. Formation of complex sentences according to the syntax of the Russian language.

Keywords: Azerbaijani language, Russian language, relation, syntactic influence, official style.

Giriş.

Azərbaycan ədəbi dilinin rəsmi üslubunun formalaşması və inkişafında Azərbaycan-Rusiya münasibətlərini əks etdirən sənədlər əhəmiyyətli rol oynamışdır.

XVIII əsrin əvvəllərində Rusiya və Səfəvi dövləti arasında bağlanan müqavilələri başlanğıc hesab etmək olar. Çünki bu müqavilələr rus dilində idi və müqavilədə yazılanları Səfəvi dövləti başa düşmək üçün tərcüməçiyə ehtiyac duyurdu. 1717-ci il 30 iyul/10 avqust Rusiya və Səfəvi dövləti arasında İsfahan müqaviləsindən bir neçə bəndə nəzər yetirək: “Ст.1. И по указу высокому поставлено: чтобы с сего времени Российские купцы в Низовую приезжали и всегда, как хотят свои товары продавали; и когда там не хотят продавать, и тогда их вьюков развязывать да не принуждают, и оных в вышеупомянутом месте да не задерживают и препятия и остановки им да не чинят, за что наказание учинено будет... Ст.4. И по высокому указу поставлено: что когда прибудут в город Шемаху, оным в продаже их товаров препятия да не чинится, и да оставить их, как кто хочет, так чтобы с обеих сторон довольны были в своей покупке и продаже. А когда довольны не будут, то Беглербеги и судья и начальники им да не насилует против правды и положения, убытков и обид им да не наносят; и когда за свои их товары цену хотят брать, или кто Беглербегу и судье побьет челом, то с помощью и вспоможением правдиво в их иску да поступают; чего ради наказание учинено будет. Ст.8. (Постановляет, что персидское правительство должно назначать толмачами только тех лиц, посредничеством которых были бы довольны обе стороны, и разрешает одному из русских купцов занимать эту должность.). Ст.10. (Постанавляет, что русские купцы имеют право нанимать охрану на обратном пути из Шемахи в Низовую, возлагает на иранские власти ответственность за ущерб, который могут понести русские купцы, и предписывает возвращение товаров, спасенных после кораблекрушения, их хозяевам.)” [Mahmudov Y. Şükürov K.

FULL TEXTS BOOK

Azərbaycan beynəlxalq münasibətlər və diplomatiya taraixi. (1639-1828), Dövlətlərarası müqavilələr və digər xarici siyasət aktları. I cild, Bakı, 2009, s.61]

Müqavilənin sintaksisinə nəzər saldıqda görmək olur ki, əksər cümlələr поставлено (qərara alınır) постанавляет (qərar verilir), xəbərlər əsasında formalaşmışdır. Həmçinin bu xəbərlər tabeli mürəkkəb cümlənin birinci tərəfində, yəni baş cümlədə işlənmişdir. Bu formada olan cümlələr bu gün də Azərbaycana aid rəsmi sənədlərin dili üçün işləkdir. (müasir rəsmi sənədlərdən nümunələr ver).

İndi isə həmin dövrlərdə rus dövlətinə aid olmayan rəsmi sənəd nümunələrinə diqqət edək: 1715-ci il 15 avqust Səfəvi dövləti və Fransa arasında Versal müqaviləsi: "Статья I. Договор 1708 г. остается в силе, поскольку он не противоречит статьям настоящего договора. Статья II. Французские купцы и вообще все торгующие в Иране под французским флагом могут свободно ввозить в Иран и вывозить из него как в Индию, так и в Европу любые товары без каких-либо ограничений и без уплаты каких-либо пошлин и налогов. Статья IV. Шах должен обеспечивать бесплатно французских торговцев, а также консулов приличными домами в Исфахане, портах Персидского залива и на границах с Турцией и Грузией как для жилья, так и для их контор и факторий; французский флаг может быть поднят над этими зданиями. Статья V. Французскому послу предоставляется при персидском дворе преимущественное положение по сравнению с послами других стран. То же относится и к французским консулам, управляющим факториями и их агентам по сравнению с подданными других государств. (Mahmudov Y. Şükürov K. Azərbaycan beynəlxalq münasibətlər və diplomatiya taraixi. (1639-1828), Dövlətlərarası müqavilələr və digər xarici siyasət aktları. I cild, Bakı, 2009, s.56]

Bu müqavilənin tərcüməsini olduğu kimi veririk: "Maddə I. 1708-ci il müqaviləsi qüvvədə qalır, çünki onun müddəaları bu müqavilənin maddələrinə zidd deyil. Maddə II. Fransız tacirləri və ümumiyyətlə Fransız bayrağı altında İranda ticarətlə məşğul olan bütün şəxslər Hindistana və Avropaya olmaqla, istənilən malları İrana sərbəst şəkildə idxal və İrandan ixrac edə bilərlər. Bu əməliyyatlar heç bir məhdudiyyət olmadan və heç bir gömrük rüsumu və ya vergi ödənilmədən həyata keçiriləcəkdir. Maddə IV. Şah Fransız tacirlərini və konsullarını İsfahanda, Fars körfəzi limanlarında, eləcə də Türkiyə və Gürcüstanla sərhədlərdə yerləşən ərazilərdə həm yaşayış, həm də onların ofisləri və faktoriyaları üçün münasib binalarla ödənişsiz təmin etməlidir. Fransız bayrağı həmin binaların üzərində qaldırıla bilər. Maddə V. Fransız səfiri İran sarayında digər dövlətlərin səfirlərinə nisbətən üstün mövqeyə malik olmalıdır. Bu qayda Fransız konsullarına, faktoriyalara rəhbərlik edənlərə və onların nümayəndələrinə də şamil olunur – yəni onlar digər ölkələrin nümayəndələrindən üstün tutulurlar".

Müqavilənin sintaksisində rus dilinin təsiri yox dərəcəsindədir. Buna görə də əminliklə deyə bilərik ki, Azərbaycan dilinin rəsmi üslubu üçün xarakterik olan *qərara almaq, qərar gəlmək* və b. bu kimi ifadələr məhz rus dili vasitəsilə dilimizə keçmişdir. Fikrimizi Mahmudov Y. Şükürov K. Azərbaycan beynəlxalq münasibətlər və diplomatiya tarixi. (1639-1828), Dövlətlərarası müqavilələr və digər xarici siyasət aktları kitabının I cildində verilən müqavilə və fərmanlarla təsdiqləyə bilərik.

ƏSAS.

XIX əsrin əvvəllərində rus dilinin Azərbaycan dilinin rəsmi üslubuna təsiri mövzusundan bəhs edərkən ilk növbədə "Gülüstan" və "Türkmənçay sülh" müqavilələrindən danışmaq lazımdır. Məlumdur ki, Azərbaycanın taleyi və ərazi bütövlüyünün parçalanması ilə bağlı 1813-cü il 12 oktyabrda Çar Rusiyası ilə Qacarlar İranı arasında imzalanan bu müqavilə rus dilində olmuşdur. Doğrudur, bu müqavilə də rus dilində idi. Müqavilənin mətnini oxuyub təhlil etmək üçün rus dilini bilmək vacib idi. Buna görə də həmin dövrün hakim təbəqələri, idarəetmə mərkəzlərinin rəhbərləri, məhkəmə işçiləri rus dilini öyrənmək və tətbiq etmək məcburiyyətində idi. Gülüstan müqaviləsi rus dilində olduğu üçün onu Azərbaycan dilinə, fars dilinə tərcümə etmək zərurəti də meydana çıxırdı. Buna görə də rus-Azərbaycan dilində tərcüməçilik fəaliyyəti də inkişaf etməyə başladı. 1813-cü il 12 oktyabr Rusiya və İran arasında

FULL TEXTS BOOK

Gülüstan müqaviləsinin 5-ci və 6-cı maddələri təqdim edirik: **V maddə**. "Rus ticarət gəmilərinə əvvəlki qayda üzrə Xəzər sahilləri yaxınlığında üzmək və onları yan almaq hüququ verilir; həm də gəmi qəzası zamanı iranlılar tərəfindən dostluq köməyi edilməlidir. İran ticarət gəmilərinə də həmin bu hüquq – əvvəlki qayda üzrə Xəzər dənizində üzmək və Rusiya sahillərinə yan almaq ixtiyarı verilir, burada da gəmi qəzası zamanı, qarşılıqlı surətdə, iranlılara hər cür yardım göstəriməlidir. Hərbi gəmilərə gəldikdə isə, müharibədən əvvəl, habelə sülh vaxtı və həmişə Rusiya hərbi bayrağı Xəzər dənizində tək mövcud olmuşdur. Həmin ehtiram daxilində, əvvəlki ixtiyar indi də yalnız Rusiya Dövlətinə verilir ki, ondan başqa heç bir Dövlətin hərbi Bayrağı Xəzər dənizində ola bilməz. **VI maddə**. Hər iki tərəfdən döyüşdə əsir alınmışları, Xristian dininə və s. dinlərə mənsub ələ keçirilmiş sakinləri Müqavilə bağlanıb imzalandıqdan sonra, üç ay ərzində, hər bir tərəfdən Qarakilsəyədək (burada sərhəd rəisləri əsirləri qəbul etmək üçün öz aralarında qarşılıqlı əlaqə yaradırlar) azuqə və yol xərcləri ilə təmin edilərək, buraxılsınlar. Özbaşına, yaxud qanunu pozaraq, qaçanlardan hər birinə, milliyətindən asılı olmayaraq, öz könüllü istəyinə görə öz Vətəninə qayıtmaq azadlığı verilir, qayıtmaq istəməyənlər isə məcbur edilməsin. Bununla bərabər hər iki tərəfdən qaçmışlara amnistiya, yaxud bağışlanma verilir". [https://az.wikipedia.org/wiki/G%C3%BCl%C3%BCstan_m%C3%BCqavil%C9%99si]

Müqavilənin sintaktik tərkibi bügünkü rəsmi sənədlərin dili ilə tam uyğunluq təşkil edir. Müqavilənin dilində işlənən *hüquqi verilir, dostluq köməyi edilməlidir, ixtiyarı verilir, yardım göstərilməlidir, təmin edilərək, bağışlanma verilir* və başqa bu kimi ifadələr müasir dövr üçün işlək olan rəsmi sənədlərin dili üçün səciyyəvi olan şəxssiz növ feillərdir. Bu gün rəsmi sənədlər üçün işlək olan bu ifadələr, təbii ki, danışdığımız tarixi dövr nəticəsində rus dilinin təsiri ilə dilimizə daxil olmuşdur.

1828-ci ildə baş tutan Türkmənçay sülh müqaviləsi Rusiya hakimiyyətini daha da möhkəmləndirdi: Rus dili dövlət dili statusu qazandı: məhkəmələr, idarələr, məktəblər, ordu rus dilində fəaliyyət göstərməyə başladı. Əhali əsasən iki yerə bölünürdü: 1) kəndlilər –yalnız Azərbaycan dilində danışırdı; 2) Şəhər əhalisinin ziyalı təbəqəsi- rus dili ilə müəyyən dərəcədə təmasda idi.

Rus dilində sənədlərin tərtibi bir sıra yeniliklərin meydana gəlməsinə səbəb oldu: 1) Yerli əhali dövlət idarələrində tərcüməçisiz iş apara bilmirdi. Bu da iki sinfi ortaya çıxardı: a) rus dilini bilənlər (yəni savadlılar, ziyalılar, məmurlar); b) rus dilini bilməyənlər (kəndlilər, sənətkarlar, adi şəhərlilər); 2) Rus dilinin nüfuzu artdı: rus dilini bilmək, dövlətə qulluq etmək, karyera qurmaq zəruri şərtə çevrildi; 3) Yerli əhali arasında rus dili öyrənməyə maraq oyandı: 4) XIX əsrin ortalarında rus-müsəlman məktəbləri açılmağa başladı.

Dil baxımından da bir sıra yeniliklər yarandı, belə ki rəsmi-işgüzar üslubda bir sıra sənədlərin hazırlanmasında istifadə olunan inzibati, hüquqi və hərbi terminlər Azərbaycan dilinə keçməyə başladı. Rəsmi sənədlərdə istifadə olunan terminlərə nümunə olaraq: naçalnik (начальник), prikaz(приказ), protokol(протокол), komandir(командир), batalyon(батальон), sud(суд), politsiya (полиция) və s. Bu terminlər əvvəlcə rəsmi sənədlərdə və idarəçilikdə işləndi, sonra isə digər üslublara keçid aldı.

Türkmənçay müqaviləsindən sonra rus dili Azərbaycanda inzibati, hüquqi və ictimai həyatın əsası oldu. Bu Azərbaycan dilinə təkcə leksik təsirlərlə qalmadı, daha sonralar sintaktik təsirlərin də yaranmasına şərait yaratdı.

Azərbaycan ədəbi dili rəsmi üslubun formalaşması və inkişafında Azərbaycan-Rusiya münasibətlərini əks etdirən ərizələr əhəmiyyətli rol oynamışdır. Məlum olduğu kimi, XVIII əsrin əvvəllərindən başlayaraq Rusiya öz nüfuzunu Cənuba doğru genişləndirməyə çalışır və müxtəlif metodlara, üsullara əl atır. Azərbaycanı İrana və Türkiyəyə qarşı qaldırmaq üçün ölkəyə şərq dillərini bilən casuslar göndərilir, rəsmi müraciətlər olunur, yerli əhalidən Rusiya imperatoruna ərizələr gəlir (bunların bir qismi xüsusi təşkil olunur və həmin ərizələrdə, adətən, rus qoşunu Azərbaycana dəvət edilir) və s. Azərbaycan-Rusiya münasibətlərini əks etdirən sənədlərdə (bunların böyük bir hissəsi azərbaycanca olsa da, farsdilli sənədlər də mövcuddur və Azərbaycan dilinə üstünlük verilməsi onun ictimai-siyasi, o

FULL TEXTS BOOK

cümlədən rəsmi nüfuzunun təsdiqi deməkdir) ölkənin həm daxili problemlərindən, həm də başqa ölkələrlə əlaqələri məsələlərindən bəhs olunur. XVIII əsrdə rəsmi üslub ictimai funksiyasına görə kütləviləşir; getdikcə daha sürətlə xalqın həyatına daxil olur – kütləviləşmənin iki tərəfini qeyd etmək lazımdır: birincisi, mərkəzi rəsmi təşkilatlar xalqa müraciət edirlər; ikincisi, xalq kütləsi mərkəzi rəsmi təşkilatlara müraciət edir. [Джафаров. Н.Г Азербайджанский литературный язык XVIII Баку, АДД, с. 46]

Birinci tipdən olan sənədlərə I Pyotrun manifestini misal göstərmək mümkündür. Azərbaycan ədəbi dili rəsmi üslubunun klassik nümunəsi olan bu sənəd 15 iyul 1722-ci ildə I Pyotrun Azərbaycana yürüşü zamanı knyaz Dmitri Kantemir tərəfindən tərtib edilmişdir. Manifestin əvvəli dəbdəbəli, lakin tədqiqatçıların qeyd etdikləri kimi, tamamilə ayıq müraciətdən ibarətdir: "Pyotri-əvvəl ibn Aleksey padşahi-Rusiya beunirəbbani və qüdrəti-cahandakı ilə biz Petros-əvvəlki imperatori-təvayifi-rus xudadari-ərzeyni-məmaliki-şimaleyi tülüvəl zəval vəl nisf əl-nəhreyi padşah əl-bərreyi və xaqan əl-bəhreyi və neçə dəxi sair və sair padşahlıqların və hökumətgahların hakimi və malikanə buyruqçusu. Şövkətlü və əzəmətlü səadətlü və məhəbbətlü ulu möhübbümüz və qədimi dostumuz şahi-alicah həzrətlərinin hökumətində və xidmətində olan izzətlü və hörmətlü süpəhsalariyan və xaniyan və qorçu-başıyan və ağayani-qul və tüfəngçiyan və topçubaşıyan və bəglər-bəgliyani-qul və soltaniyan və vüzəra və sair zabitan və nəfərətan bin və yüzbaşıyan və fəzilətlü üləma və mövla və hörmətlü imamlar və müəzzinlər və sair ərbabi-xüddam cəvamei və köy kədxudaları və əhli tüccar və karvan və dəstgahbaşları və dükan və əhli-sənət xəlifələri və şagirdləri və ümum üzəra əhli-zəhmətirəaya və bəraya hər birisi mərtəbəsi ilə və əcnası ilə və məzhəbi ilə salam səlamət əncam iblağilə təvəqqei-rəfi-humayunum vasil olacaq məlumi izətməndləri ola ki…" [Xudiyev. N.Azərbaycan ədəbi dili tarixi. Bakı, Elm və Təhsil 2018, s.408].

Diqqət etdikdə görmək olur ki, I Pyort manifestinə aid olan müraciətdə təyinləri həm çoxdur, həm də İran şahına aid olan sənəd nümunəsindəki təyinlərindən fərqli olaraq, daha realdır, konkretdir və bilavasitə siyasi-inzibati məzmunu ilə seçilir. Halbuki şaha müraciətlə deyilmiş "ulu möhübbümüz və qədimi dostumuz" kimi "boş" ifadələr ancaq diplomatik məqsədlə işlədilmişdir. Doğrudur, sənəd klassik üslub xüsusiyyətlərindən qurtula bilməmişdir: mətndə müəyyən qədər izafət birləşmələri özünü göstərməkdədir. Ancaq sənədin dilində işlənən ismi birləşmələr (padşahlıqların və hökumətgahların hakimi, malikanə buyruqçusu, şövkətlü və əzəmətlü səadətlü və məhəbbətlü ulu möhübbümüz, qədimi dostumuz, şahi-alicah həzrətlərinin hökuməti, izzətlü və hörmətlü süpəhsalariyan, hörmətlü imamlar və müəzzinlər, köy kədxudaları) işləkliyinə görə daha çoxdur və milliliyi ilə öndədir.

XIX əsrdə rəsmi üslubun bu və ya digər dərəcədə inkişafına mətbuatın meydana çıxması, kitab nəşri böyük təsir göstərir, bununla belə, Azərbaycanın dövlət müstəqilliyinin olmaması rəsmi üslubun milli təfəkkür əsasında deyil, ancaq Rusiya imperiyasının qanun, qərar və göstərişlərinin təcrübəsi əsasında inkişafına səbəb olur. Məsələn, 1817-ci ildə Peterburqda imperatorun imzaladığı "Тарифъ по азиатской торговли" adlı rəsmi sənədlər toplusu Azərbaycan ədəbi dilinə yaxın bir dilə tərcümə olunub 1830-cu ildə çap edilir. Tədqiqatçıların göstərdiyi kimi, həmin kitab, əsasən, Orta Asiya üçün nəzərdə tutulmuşdu, tərcümədə Azərbaycan dilinə üstünlük verilməsi isə bu dilin digər türk xalqları arasında asan anlaşılması idi.[Cahangirov M. Milli təşəkkül mərhələsində Azərbaycan ədəbi dilinin aparıcı üslubları, Bakı, 1989. s.97].

Azərbaycanda və ümumən Rusiyada ticarətin inkişafı rəsmi üslubun, əsasən, ticarət, təsərrüfat və s. məsələləri ilə bağlı olmasını şərtləndirir; məsələn, "Əkinçi"dən "Badkubədə neftin xalvarı, yəni iyirmi pudu 1 manat 5 – qəpikdən tutmuş 2 manat 50 qəpik. Neft cövhərinin pudu boçkasız 65 qəpik, boçka ilə 1 manat" ["Əkinçi" qəzeti, 22 iyul, 1875]. XIX əsrdə Azərbaycan dilinə tərcümə olunmuş, yaxud Azərbaycan dilində yazılmış bir sıra sənəd nümunələrini (formalarını) nəzərdən keçirək:

FULL TEXTS BOOK

İltizamnamə: "Bin səkkiz yük əlli altıncı ildə və fevral ayının səkkizinci günündə biz zildə qol qoyan filan və filan qəriyənin əhlindən filan və filan verürük bu iltizamnaməni Dərbənd ətrafının hakiminə. Ondan ötrü ki… [М.А.Везиров. Учебник татарско-азербайджанского наречия, СПб., 1861, с.131-132].

Sənədə nəzər saldıqda açıq-aşkar görmək olur ki, Söz sırası- "verürük bu iltizamnaməni Dərbənd ətrafının hakiminə" — burada obyekt (iltizamnaməni) ilə tamamlıq (Dərbənd ətrafının hakiminə) xəbərdən (verürük) sonra gəlir. Bu da rus dilindəki "Мы даём это обязательство правителью Дербендской области" tipli quruluşa bənzəyir. "Biz zildə qol qoyan... əhlindən... verürük" – bu cür geniş və yüklü strukturlar rus dilindəki tabeli atributiv (*придаточное атрибутивное)* cümlələri xatırladır.

Şərtnamə: "1861-ci sənədə mah ğənvarın on beşinci günündə mən aşaqədə qol çəkən və möhür basan Tiflis sakini usta Əliəkbər Mustafa oğlu saqdım kaputan Mirzə Fətəli Axundova yüz əlli min kərpic öz məxsusi kürəxanəmdən…" (Əlyazması, Azərb. EA Əİ-nin fondu).

Şərtnamənin sintaksisndə sifətlərin və təyini söz birləşmələrinin çoxluğu, həmçinin "Tiflis sakini usta Əliəkbər Mustafa oğlu" ad, soyad ardıcıllığı rus dilindəki "житель Тифлиса мастер Алиакпер, сын Мустафы" tipli ardıcıllığına uyğundur.

Nümunələrdən göründüyü kimi, rəsmi üslub bundan əvvəl nəzərdən keçirdiyimiz funksional üslublarla müqayisədə daha mürəkkəb, daha çətin və daha qeyri-demokratikdir – xüsusilə sənədlərin dilində həm Şərq, həm də Qərb dillərinin kalkalarını qeyd etmək mümkündür ki, bunlar, haqqında danışılan üslubun müəyyən mənada, hətta qeyri-orijinal olduğunu, milli sintaksisə əsaslanmadığını göstərir. Sənəd dilindəki nahamarlıq, kələ-götürlülük, hər şeydən əvvəl, onun nəticəsidir ki, həmin sənədlərin tərtibində ana dilinin enerjisi ya az iştirak edir, ya da, ümumiyyətlə, iştirak etmir – əsrlər boyu ənənəvi olaraq işlənmiş ərəb, yaxud fars kalkaları XIX əsrin ortalarından başlayaraq bu və ya digər şəkildə rus kalkaları ilə əvəz olunur, çox hallarda isə müxtəlif dillərin sintaktik konstruksiyaları, leksik-semantik paralelləri birlikdə çıxış edir. Bununla belə, XIX əsrin sonlarına doğru Azərbaycan ədəbi dilinin rəsmi üslubu xüsusi funksional üslub sahəsi kimi mövcud olub müəyyən ictimai-tarixi iş görür.

Rəsmi üslubun sintaksisinin müəyyənləşməsində XX əsrin əvvəllərinə qədər Şərq ənənəsi əsas idi – bu mənada ki, ərizədən tutmuş diplomatik dövlət sənədinə qədər bütün rəsmi üslub nümunələrinin istər leksikasında, istər morfologiyasında, istərsə də sintaksisində XX əsrin əvvəllərinə qədər klassik dil xüsusiyyətləri öz qabarıqlığı ilə seçilirdi. Biz bunu yuxarıda göstərdiyimiz nümunələrdə də görə bilirik. Artıq XX əsrin əvvəllərindən başlayaraq bir sıra sənədlərin Qərb ənənəsində hazırlanması prosesi başlayır. Məsələn:

"1) Oktyabrın onundan on beşinə kimi başlanır realni uşqolaya qəbul olmaqdan ötri imtahanlar yarımıncı, birinci və ikinci qlaslara;

2) imtahan olunacaq şəhər uşqolasında hər gün saat ikidən sonra;

3) ərizələri göndərmək olar oktyabrın on beşinə kimi şəhər uşqolasına. Poçt ilə və ya vermək olar inspektorun özünə;

4) təyin olunmuş vəqtdə gəlməyən uşaqlar dəxi imtahan olunar dəsrə şüru olunandan sonra;

5) dəsrə gələn uşaqlar özləri ilə bərabər gətirməlidirlər: qələm, kağız və çernil…" [Azərbaycan ədəbi dili tarixi. Bakı, 2007, s.207]

Göründüyü kimi, mətn bütövlükdə rus sintaksisinin təsiri altındadır – hətta bu təsir o qədər güclüdür ki, bilavasitə tərcüməni xatırlardır (ola bilsin ki, ümumiyyətlə, tərcümədir). Bütün cümlələrdə özünü göstərən inversiya məhz rus dilini təsiri nəticəsindədir. Rəsmi dil üçün səciyyəvi olmayan bu xüsusiyyət Qərb rəsmi düşüncəsinin Azərbaycan dilli sənədlərində nə dərəcədə iştirak etdiyini göstərir.

FULL TEXTS BOOK

Sovet dövründə rus dilinin rəsmi üslub vasitəsilə sintaksisə təsiri kulminasiya nöqtəsinə çatdı. M.Məhərrəmovun fikrincə, "bu üslub ancaq sovet dövründə yaranmış və inkişaf etmişdir". [Məhərrəmov A. Azərbaycan ədəbi dilinin sovet dövrü, s. 13-14]. Təbii ki, bu belə deyil: biz yuxarıda göstərdiyimiz faktlara əsasən təsdiq etdik ki, XVI əsrdən başlayaraq Azərbaycan Demokratik Cümhuriyyəti dövrü də daxil olmaqla rəsmi üslubun maraqlı nümunələri mövcuddur, lakin o ki qaldı rəsmi üslubun sovet dövründəki inkişafına, bu, həqiqətən belədir. Sovet dövründə bu üslubun ardıcıl inkişafını şərtləndirən dövlət idarə aparatı fəaliyyət göstərir. Sovet dövrü Azərbaycan ədəbi dilinin inkişafının birinci mərhələsində (20-30-cu illərdə) rəsmi üslub aşağıdakı əlamətlərlə səciyyələnir: 1. Ərəb, eləcə də fars elementləri ilə zəngin olan klassik ifadə tərzi əsasən qalır, ümumən lüğət tərkibində gedən proseslər – gah bu, gah da digər dildən alınmalara üstünlük verilməsi rəsmi üslubun leksikasına da təsir edir, lakin sintaksis, bir qayda olaraq klassik ifadə tipində mövcud olur. 2. Rəsmi üslubdan istifadə edən sosium geniş olmadığına görə, onun ictimai məzmunu da, deyək ki, bədii, publisistik, yaxud elmi üslub qədər böyük olmur. Eyni zamanda, rəsmi üslub 20-ci, eləcə də 30-cu illərdə başqa üslublardan təcrid olunmuş şəkildə inkişaf edir, daha doğrusu, rəsmi üslubla başqa üslublar arasındakı təmas, məsələn, bədii üslubla publisistik yaxud publisistik üslubla elmi üslub arasındakı əlaqələr gücündə olmur [Xudiyev N. Azərbaycan ədəbi dili tarixi. Bakı, Elm və təhsil, 2018. s. 596]

Rus dilinin Azərbaycan dilinə təsirini nəzərə alıb deyə bilərik ki, bu təsir bir sıra qrammatik yeniliklərin meydana gəlməsinə səbəb olmuşdur:

1. Feili birləşmələrin isimləşməsi hallarının artması;
2. Modal konstruksiyaların rus sintaksisinə uyğun qurulması;
3. "Öz" əvəzliyi ilə qurulmuş ifadələrin işlənməsi;
4. Cümlə üzvlərinin yerdəyişməsi;
5. Rus dilinin sintaksisinə uyğun mürəkkəb cümlələrin yaranması.

1.Feili birləşmələrin isimləşməsi hallarının artması. Rus dilində feili sifət və məsdər tərkibi ilə qurulan mürəkkəb sintaktik konstruksiyalar geniş yayılması ilə yanaşı, həm də onların isimləşmə halları daha çox özünü göstərir. Bu forma rəsmi-işgüzar üslub vasitəsi ilə rus dilindən Azərbaycan dilinə keçməyə başladı. Məs: "tədbirlərin həyata keçirilməsi", "işin yerinə yetirilməsi", "qərarın qəbul edilməsi", "məsələlərin həll olunması" və s. Rusların işğalından əvvəl Azərbaycan dilində daha çox sadə tərkiblər işlənirdi: "tədbir görmək", "qərar vermək", "iş görmək" və s.

Şah İsmayıl Xətainin 1501 5 iyun tarixli bir fərmanına nəzər yetirək: "Əbülmüzəffər sözümüz Əmiri-əzəm əkrəm Musa Dorğut oğlu inayət və şəfqətimizə ümidvar olandan sonra şöylə bilsün kim, iftixarül-əazim vəl-əyan Əhməd ağa Qaramanlu ol tərəfə göndərdük və ol yerin ixtiyarligini kəndunə şəfəqqət etdük. Gərək kim, müşarileyh sözümdən və məsləhətimdən çıxmasun və mütabiət və yardım ona qılsun kim, inşaallah-təala hər nə kim, etmək muradi və istəgi olsa, hasildür. Gündən-günə hər iş vaqe bolsa, Əhməd ağa ittifaqi ilə dərgahi-müəllamizə bildirsünlər kim, hər növ buyruğumuz olsa, əməl etsün, könlümüzə xoş dutub mərhəmətimizə əmrdar olsun. Təhrirən 7 rəbiüləvvəl, sənə 917 Xətm". [https://525.az/news/597-sah-ismayil-xetainin-ana-dilinde-fermani]

Göründüyü kimi bu fərmanda istifadə olunan feillər rəsmi-işgüzar üslubun tələblərinə uyğun olaraq əmr şəklindədir: "şöylə bilsün", "məsləhətimdən çıxmasun", "yardım ona qılsun", "dərgahi-müəllamizə bildirsünlər", "əməl etsün", "əmrdar olsun".

Doğrudur, fərmanda öz əksini tapan cümlələrdəki feillər təkcə əmr şəklində deyil, həmçinin xəbər (göndərdük, şəfəqqət etdük) və şərt şəkillərində də (muradı və istəgi olsa, vaqe olsa, buyruğumuz olsa) işlənmişdir. Hətta mətndə ismi xəbərə də (hasildür) rast gəlmək olur.

FULL TEXTS BOOK

Fərmanda işlənən feillərin hər biri şəklindən asılı olmayaraq məlum növdədir. Eyni zamanda onu da qeyd etmək istərdik ki, fərmanda işlənən feili birləşmələrin (şəfqətimiz ümidvar olandan sonra, könlünüz xoş dutub heç biri isimləşməmişdir.

XVI əsrdən sonrakı dövrlərə XVII-XVIII əsrlərə diqqət yetirdikdə rəsmi üslubun daha çox formalaşdığını, inkişaf etdiyini görmək olur. XVII-XVIII əsrlərdə rəsmi üslub nümunələrinin dil-üslub kamilliyi, hər şeydən əvvəl, müəyyən "rəsmilik" standartlarının artıq tamamilə üzdə olması ilə diqqəti cəlb edir: bu standartların eynilə ərəb, yaxud farsdilli nümunələrdə də işlənməsi belə bir fikrə gəlməyə əsas verir ki, son orta əsrlərə doğru ərəb, fars və türk dilləri üçün eyni olan ümumi rəsmi üslub modeli müəyyənləşmişdir.

Azərbaycanda təkcə Azərbaycan dilində deyil, fars yaxud ərəb dilində də tərtib olunmuş rəsmi sənədlər belə, Azərbaycan dilinin materiallarını (ən azı təfəkkür tərzini) əks etdirir. [Указы Кубинских ханов. Тбилиси, 1937 (giriş hissə)].

Qeyd etdiyimiz kimi feli birləşmələrin isimləşməsi rus dilindən gələn sintaktik xüsusiyyət hesab oluna bilər. Niyə morfoloji əlamət demədim? Təbii ki, isimləşmə morfoloji əlamətdir, ancaq unutmayaq ki, bu əlamət cümlə üzvünün dəyişməsinə səbəb olur. Ona görə də bu xüsusiyyət həm də sintaktik xüsusiyyət hesab oluna bilər. Nümunələrə diqqət edək: "Qərarın qəbul edilməsi haqqında protokol tərtib olundu"; "Şura tərəfindən müvafiq tədbirlərin həyata keçirilməsi qərara alındı"; "Ərizəyə baxılmaq üçün aidiyyəti üzrə göndərilməsi təmin edilsin"; "Hüquqi şəxsin dövlət qeydiyyatına alınması barədə şəhadətnamə təqdim olundu" və s. Bu cümlələrdə "qərarın qəbul edilməsi", "Şura tərəfindən müvafiq tədbirlərin həyata keçirilməsi", "ərizəyə baxılmaq üçün aidiyyəti üzrə göndərilməsi", "hüquqi şəxsin dövlət qeydiyyatına alınması barədə" isimləşmiş feili birləşmələr rus dilinin təsiri nəticəsində yaranmışdır. Halbuki bu birləşmələr rus dilinə qədər rəsmi sənədlərdə "qərarı qəbul etmək", "Şura tərəfindən müvafiq tədbirlər həyata keçirmək", "ərizəyə baxmaq üçün göndərmək", "hüquqi şəxsi dövlət qeydiyyatına almaq barədə" şəklində işlənmişdir. Həmin cümlələrdə işlənən xəbərlərin feilin məchul növündə ("tərtib olundu", "qərara alındı", "təmin edilsin", "təqdim olundu") işlənmə formaları da rus dilinin təsiri nəticəsindədir. Azərbaycan Respublikasının Milli Məclisi *qərara alır* [Azərbaycan Respublikasının Qanunvericilik Toplusu, 1997-ci il, № 5, maddə 381]

2.Modal konstruksiyaların rus sintaksisinə uyğun qurulması. Rus dilinin Azərbaycan dilinin sintaksisinə təsiri mövzusundan bəhs edərkən modal konstruksiyaları da qeyd etmək lazımdır. Azərbaycan dilinin sintaksisində, xüsusən də standartlaşdırılmış rəsmi üslubda çox vaxt modal konstruksiyaların rus sintaksisinə uyğunlaşması hadisəsi ilə də qarşılaşırıq.

Modal konstruksiya — nitqdə və ya cümlədə mövqeyi, münasibəti, zəruriliyi, imkanı, ehtimalı, icazəni və ya qadağanı ifadə edən dil vahididir. Bu konstruksiyalar danışanın hadisəyə münasibətini bildirir. Onlar həm leksik, həm də qrammatik vasitələrlə qurula bilər.

Rəsmi üslubda, əsasən, zərurət, icazə, qadağa, məcburiyyət, ehtimal və başqa bu kimi modal konstruksiyalardan istifadə olunur.

Rus rəsmi üslubunun aparıcı modal fel konstruksiyasından nümunələrə nəzər yetirək:

Следует представить документы	Sənədlərin təqdim edilməsi tövsiyə olunur.	Tələb, tövsiyə
Важно соблюдать правила	Qaydalara əməl olunması vacibdir	Zərurət

FULL TEXTS BOOK

Разрешается использовать	İstifadə olunmasına icazə verilir	İcazə
Запрещается выходить	Çölə çıxmaq qadağandır/yolverilməzdir	Qadağa
Допускается istisna	İstisnaya yol verilir	Tolerantlıq / icazə
Должен быть оформлен	Rəsmi qaydada rəsmiləşdirilməlidir	Məcburiyyət
Обязан представить	Təqdim etməyə borcludur	Hüquqi borc
Следует отметить, что…	Qeyd edilm:əlidir ki,	Diqqət çəkmə
Следует особенно подчеркнуть	Xüsusilə vurğulanmalıdır ki,...	Vurğulama

Yuxarıda verilən modal konstruksiyalar- *tələb olunur, qadağandır, yol verilir, borcludur, tənzimlənməlidir, rəsmiləşdirilməlidir, qeyd edilməlidir, xüsusilə vurğulanmalıdır* rus dilinin tələbindən yaranan ifadələrdir və bu tip ifadələr elmi məqalələrdə, qanunvericilikdə, hesabatlarda, bir sözlə rəsmi və elmi üslubda geniş şəkildə istifadə olunur. İndi isə mənbələrdən konkret nümunələrtə nəzər yetirək:

Azərbaycan Respublikası Konstitusiyasının 77-ci maddəsinə görə tarix və mədəniyyət abidələrini qorumaq hər kəsin *borcudur.* (hüquqi borc) [https://e-qanun.az/framework/3526]; Müşahidəçilər onların iştirakı ilə keçirilən hərəkətlərin faktını, məzmununu və nəticələrini protokolda təsdiq etməyə *borcludur.* (hüquqi borc) [https://e-qanun.az/framework/46948] I. "Dövlət qulluğu haqqında" Azərbaycan Respublikasının Qanununa ["Azərbaycan" qəzeti, № 30, 07 fevral 2001] aşağıdakı *dəyişikliklər edilsin* 1. 35-ci maddədə "yanvarın 1-dən" sözləri "sentyabrın 1-dən" sözləri ilə *əvəz edilsin.* (məcburiyyət) [https://e-qanun.az/framework/12308]; Vergi yoxlaması keçirildikdə, vergi ödəyicisinə və ya onun sahibliyində, istifadəsində və sərəncamında olan əmlaka qanunsuz olaraq zərər vurulmasına *yol verilmir*. (tələb). [https://e-qanun.az/framework/46948];

3. **"Öz" əvəzliyi ilə qurulmuş ifadələrin işlənməsi**. "Öz" əvəzliyi ilə qurulan cümlələrdən rəsmi yazışmalarda və qanunvericilik dilində geniş şəkildə istifadə olunur. "Öz" əvəzliyi (rus dilində cam işlənir) rəsmi üslubda məsuliyyətin və ya icra səlahiyyətinin vurğulanması üçün tez-tez işlənir.

1. "*Özü tərəfindən*". Sənəd *özü tərəfindən* təsdiq olunmuşdur. (rusca: Документ был утверждён *им самим*);

2. *"Öz adına*". Müqavilə *öz adına* bağlanmışdır. (rusca: Договор был заключён *на его собственное имя)*;

3. "*Öz təşəbbüsü ilə*". Təklif *öz təşəbbüsü ilə* irəli sürülmüşdür. (rusca: Предложение было выдвинуто *по его собственной инициативе*).

4. "*Özündə əks etdirir"*. Bu sənəd qarşıya qoyulan vəzifələri *özündə əks etdirir*. (rusca: Этот документ отражает *на себя поставленные* задачи.)

5. "*Öz funksiyalarını yerinə yetirir*". Qurum qanunvericiliyə uyğun olaraq *öz funksiyalarını yerinə yetirir* (rusca: Учреждение осуществляет *свои функции в соответствии* с законодательство).

Bu ifadələrlə qurulan cümlələr rus dilindən tərcümədə olduğu kimi işləndikdə cümlədə sintaktik təbiiliyi aradan qaldırır və məcburi tərcümə təsiri bağışlayır. Bəzən bu süniliyi aradan qaldırmaq üçün Azərbaycan dilinin sintaktiktik və üslubi normalarına uyğun olaraq *"Öz tərəfindən imzalanmışdır*" ifadəsinin əvəzinə "*Tərəfindən imzalanmışdır*"; *"Öz funksiyalarını yerinə yetirir"* ifadəsinin əvəzinə *"Qanunla müəyyən olunmuş funksiyalarını icra edir"*formasında işlənir.

Azərbaycan Respublikası vətəndaşının xaricdə olduğu müddətdə pensiyası *öz arzusu ilə* Azərbaycan Respublikasının müvafiq bankındakı şəxsi hesabına və ya yaşadığı dövlətdəki müvafiq bank hesabına köçürülür. Bu halda xidmət haqqı pensiyanın məbləğindən tutulur.[https://e-qanun.az/framework/46948]

Çar Rusiyası və sovet dövründə hüquqi və inzibati sənədlərdə bu cür "öz"-(rus dilində-cam) əvəzlikli konstruksiyalar geniş tətbiq edilmişdir. Müstəqillikdən sonra da müəyyən dərəcədə qorunmuş və indi də rəsmi üslub forması kimi qalmaqdadır, amma, təbii ki, milli normativlərə uyğunlaşdırma prosesi də özünü göstərməkdədir.

4. Cümlə üzvlərinin yerdəyişməsi. Azərbaycan dilinin sintaktik sistemində cümlə üzvlərinin özünəməxsus normativ sıralanma quruluşu var. Belə ki mübtəda cümlənin əvvəlində, xəbər sonunda, təyin təyinedicidən əvvəl(yəni aid olduğu bütün cümlə üzvlərindən əvvəl gələ bilər), tamamlıq mübtədadan sonra, zərflik də xəbərin əvvəlində işlənir. Doğrudur, dilimizin sintaktik quruliuşunda bəzi istisnalar da özünü göstərir. Məsələn, zaman zərfliyi mübtədadan əvvəl gələ bilər və bu, qrammatik normanın pozulması aktı sayılmır. Burda isə rəsmi üslubda cümlə üzvlərinin yerdəyişməsi hadisəsindən danışacağıq. Onu da əvvəlcədən qeyd edək ki, bu xüsusiyyət rəsmi-işgüzar üslub üçün heç də xarakterik deyil, amma bu keyfiyyət rəsmi üsluba ciddilik, vurğu, intonasiya, rəsmi ab-hava qatmaq məqsədi daşıyır. Rəsmi-işgüzar üslubda cümlə üzvlərinin yerdəyişməsi dilin üslubi imkanlarından biridir və müəyyən məqsədlərlə həyat keçirilir. Rəsmi-işgüzar üslubda işlənən bu hal rəsmi nitqin loqik quruluşunu pozmur, əksinə stilistik incəliklər yaradır. Rəsmi-işgüzar üslubda cümlə üzvlərinin yerdəyişməsi dilin üslubi imkanlarından irəli gəlir və müəyyən məqsədləri-məntiqi vurğunu diqqətə çatdırmaq, mətnin ahəngini qorumaq və ya hər hansı bir rəsmi ifadəni gücləndirmək üçün istifadə olunur. Nümunələrə baxaq: *Verilir* Rasim Hacı oğlu Hacıyevə ona görə ki, o, həqiqə tən, "Elm və təhsil" nəşriyyatında kadrlar şöbəsinin müdiri vəzifəsində işləyir. [Məmmədli N. Azərbaycan dilində işgüzar və akademik kommunikasiya (dərslik). Bakı, "Elm və Təhsil", 2021, s. 482]

NƏTİCƏ.

Göründüyü kimi, sənəd dili rəsmi yazışmalarda işlənən standartlaşdırılmış və şablonlaşmış üslubdur. Bu üslub sovet dövründə və post-sovet inzibati sistemində rus dilinə əsaslandığı üçün, rus sintaksisi Azərbaycan sənəd dilinə birbaşa və dərin təsir etmişdir. Bu təsiri biz müxtəlif vəziyyətlərdə, müxtəlif səviyyələrdə gördük. Belə ki rus dilinin Azərbaycan dilinin sintaktik quruluşuna təsiri Azərbaycan-rus dil əlaqələrinin tarixi şəraiti nəticəsində yaranmışdır. Fikrimizi daha dəqiq ifadə etmək üçün Azərbaycan-rus dil əlaqələrini tarixi baxımdan XVIII əsrdən başlayan dil əlaqələri, rus dilinin dövlət dili statusu dövründə təsirləri, sovet dövründə təhsil və mətbuat vasitəsilə dilə təsir, qanunvericilik, dövlət idarəçiliyi və inzibati idarəetmə sistemi sahəsində təsirlər olmaqla göstərməyi məqsədəmüvafiq bildik.

Rus dilinin Azərbaycan dilinin sintaksisinə təsirini nəzərə alıb deyə bilərik ki, bu təsir bir sıra sintaktik yeniliklərin və geniş istifadəolunmanın meydana gəlməsinə səbəb olmuşdur. Bu yeniliklər Azərbaycan dilinin sintaktik sistemində özünü feili birləşmələrin isimləşməsi hallarının artmasında, modal

FULL TEXTS BOOK

konstruksiyaların rus sintaksisinə uyğun qurulmasında, "öz" əvəzliyi ilə qurulmuş ifadələrin rəsmi üslubda geniş şəkildə işlənməsində, cümlə üzvlərinin yerdəyişməsində, rus dilinin sintaksisinə uyğun mürəkkəb cümlələrin mürəkkəb variantlarının yaranmasında özünü göstərir.

ƏDƏBİYYAT

1. Azərbaycan ədəbi dili tarixi. I cild, XIII-XVI əsrlər. Bakı, 2007, 480 s.
2. "Azərbaycan" qəzeti, № 30, 07 fevral 2001.
3. Azərbaycan Respublikasının Qanunvericilik Toplusu, maddə 381, 1997-ci il, № 3
4. Cahangirov M. Milli təşəkkül mərhələsində Azərbaycan ədəbi dilinin aparıcı üslubları, Bakı, 1989. 175 s.
5. Xudiyev N. Azərbaycan ədəbi dili tarixi. Bakı, Elm və təhsil, 2018. 632 s.
6. "Əkinçi" qəzeti, 22 iyul, 1875.
7. Mahmudov Y. Şükürov K. Azərbaycan beynəlxalq münasibətlər və diplomatiya taraixi. (1639-1828), Dövlətlərarası müqavilələr və digər xarici siyasət aktları. I cild, Bakı, 2009. 512 s.
8. Məhərrəmov A. Azərbaycan ədəbi dilinin sovet dövrü. Bakı, 1961.
9. Məmmədli N. Azərbaycan dilində işgüzar və akademik kommunikasiya (dərslik). Bakı, "Elm və Təhsil", 2021. 512 s.
10. Джафаров. Н.Г Азербайджанский литературный язык XVIII, АДД, Баку 1991.
11. Везиров. М.А Учебник татарско-азербайджанского наречия, СПб., 1861.
12. Указы Кубинских ханов. Тбилиси, 1937.
13. https://525.az/news/597-sah-ismayil-xetainin-ana-dilinde-fermani
14. https://az.wikipedia.org/wiki/G%C3%BCl%C3%BCstan_m%C3%BCqavil%C9%99si
15. https://e-qanun.az/framework/3526
16. https://e-qanun.az/framework/46948

FULL TEXTS BOOK

GRADIENT BOOSTING REGRESSOR (GBR) ALGORİTMASI İLE SPI TAHMİNİ

SPI PREDICTION USING THE GRADIENT BOOSTING REGRESSOR (GBR) ALGORITHM

***Rojhat ERGÜVEN*[1]**

[1]*Doktora Öğrencisi, Fırat Üniversitesi, Teknoloji Fakültesi, İnşaat Mühendisliği Teknolojileri Bölümü, Elazığ, Türkiye.*

***Alp Buğra AYDIN*[2]**

[2] *Dr. Öğr. Üyesi, Fırat Üniversitesi, Teknoloji Fakültesi, İnşaat Mühendisliği Teknolojileri Bölümü, Elazığ, Türkiye.*

***Derya AVCI*[3]**

[3] *Doç Dr., Fırat Üniversitesi, Teknik Bilimler MYO, Bilgisayar Teknolojileri Bölümü, Elazığ, Türkiye.*

ÖZET

Elazığ iline ait 1938–2024 dönemini kapsayan aylık yağış verileri kullanılarak yürütülen bu çalışmada, meteorolojik kuraklığın nicel olarak değerlendirilmesi amacıyla Standart Yağış İndeksi (Standardized Precipitation Index - SPI) hesaplamaları gerçekleştirilmiştir. Kuraklık analizinin yanı sıra, geleceğe yönelik öngörü sağlamak amacıyla makine öğrenmesi tabanlı Gradient Boosting Regressor (GBR) algoritması uygulanarak SPI değerlerinin tahmini yapılmıştır. Çalışma kapsamında üç farklı zaman ölçeği (SPI-3, SPI-6 ve SPI-12) ele alınmış; her bir SPI serisi, son 12 aylık veriye dayalı olarak modellenmiş ve 2025 yılına ait aylık SPI değerleri tahmin edilmiştir.

Model performansı R-kare (R^2), korelasyon katsayısı, Ortalama Mutlak Hata (MAE) ve Kök Ortalama Kare Hata (RMSE) gibi yaygın istatistiksel değerlendirme kriterleriyle analiz edilmiştir. Elde edilen sonuçlara göre GBR modeli özellikle SPI-6 tahminlerinde yüksek doğruluk düzeyine ulaşarak mevsimsel kuraklık öngörüsünde güçlü bir potansiyel sergilemiştir. SPI-12 sonuçları, uzun vadeli iklim eğilimlerini başarıyla yansıtırken, SPI-3 serileri kısa vadeli yağış değişimlerine duyarlılık göstermiştir. Tahmin edilen SPI değerleri, kuraklık sınıflandırma sistemine göre yorumlanmış ve her ayın hidrolojik durumu (örneğin şiddetli kurak, normal, hafif nemli vb.) belirlenmiştir.

Bu çalışma, yapay zekâ destekli kuraklık tahminlerinin bölgesel ölçekte uygulanabilirliğini ortaya koymakta olup, sürdürülebilir su yönetimi, afet önleme stratejileri ve iklim değişikliğiyle mücadele açısından önemli katkılar sunmaktadır.

Anahtar Kelimeler: Kuraklık tahmini, Standart Yağış İndeksi (SPI), Gradient Boosting Regressor (GBR), makine öğrenmesi, Elazığ

ABSTRACT

This study used monthly precipitation data from 1938 to 2024 for the Elazığ province to calculate the Standardised Precipitation Index (SPI) and quantitatively assess meteorological drought. In addition to analysing the data for signs of drought, the machine learning-based Gradient Boosting Regressor (GBR) algorithm was used to estimate the SPI values for future projections. The study considered three different time scales (SPI-3, SPI-6 and SPI-12), with each SPI series modelled based on the last 12 months of data and monthly SPI values estimated for 2025.

Model performance was analysed using common statistical evaluation criteria, such as R-squared (R^2), the correlation coefficient, mean absolute error (MAE) and root mean square error (RMSE). According

FULL TEXTS BOOK

to the obtained results, the GBR model demonstrated strong potential for seasonal drought forecasting, achieving high accuracy, particularly in SPI-6 predictions. SPI-12 results successfully reflect long-term climate trends, while SPI-3 series show sensitivity to short-term rainfall changes. The predicted SPI values were interpreted according to the drought classification system and the hydrological status of each month (e.g. severe drought, normal or slightly moist) was determined. This study demonstrates the feasibility of using AI to predict droughts on a regional scale, making important contributions to sustainable water management, disaster prevention strategies and the fight against climate change.

Keywords: Drought prediction, Standard Precipitation Index (SPI), Gradient Boosting Regressor (GBR), machine learning, Elazığ

GİRİŞ

İklim, tarihsel süreç boyunca doğal faktörlerin etkisi altında sürekli bir değişim göstermiştir. Ancak günümüzde, sanayileşme, fosil yakıt kullanımı ve hızlı nüfus artışı gibi antropojenik etkenler bu değişim sürecini belirgin şekilde hızlandırmaktadır (IPCC, 2021). Küresel ortalama sıcaklıklardaki artış, deniz seviyelerinin yükselmesi, buzulların erimesi ve iklim parametrelerindeki diğer eğilimler, iklim değişikliğinin günümüzdeki en önemli çevresel sorunlardan biri olduğunu açıkça ortaya koymaktadır (Trenberth et al., 2014). Bu bağlamda, iklim değişikliğinin en kritik sonuçlarından biri olan kuraklık, hidrolojik döngü, ekosistem dengesi, tarım faaliyetleri ve su kaynakları yönetimi üzerinde doğrudan ve dolaylı etkiler yaratmaktadır (Dai, 2011).

Kuraklığın izlenmesi ve öngörülmesi, gerek sürdürülebilir doğal kaynak yönetimi gerekse afet önleme stratejilerinin geliştirilmesi açısından büyük önem taşımaktadır. Bu amaçla geliştirilen farklı kuraklık indisleri arasında, yalnızca yağış verisine dayalı olarak hesaplanabilmesi, farklı zaman ölçeklerine uygulanabilirliği ve istatistiksel esnekliği nedeniyle Standart Yağış İndeksi (Standardized Precipitation Index, SPI) yaygın olarak tercih edilmektedir (McKee et al., 1993; Guttman, 1999). SPI, belirli bir dönemde gözlemlenen yağışların uzun yıllık yağış dağılımı ile karşılaştırılması esasına dayanarak meteorolojik kuraklık koşullarını nicel olarak ortaya koymaktadır (Wu et al., 2007).

Bununla birlikte, iklim sistemindeki belirsizlikler ve yağış serilerinin karmaşık yapısı, yalnızca geçmiş gözlemlere dayalı analizlerin geleceğe yönelik karar süreçlerinde yetersiz kalmasına neden olmaktadır. Bu nedenle, son yıllarda makine öğrenmesi ve istatistiksel tahmin modelleri, meteorolojik kuraklık tahminlerinde yaygın bir şekilde kullanılmaya başlanmıştır (Shen, 2018). Bu bağlamda Gradient Boosting Regressor (GBR), doğruluk ve genelleme kapasitesi nedeniyle öne çıkan algoritmalardan biridir (Friedman, 2001).

Bu çalışmada, Elazığ iline ait 1938–2024 dönemini kapsayan uzun süreli yağış verileri kullanılarak SPI hesaplamaları yapılmış ve geleceğe yönelik öngörüler elde etmek amacıyla GBR algoritması uygulanmıştır. SPI-3, SPI-6 ve SPI-12 zaman ölçekleri üzerinden gerçekleştirilen analizlerle, kısa, orta ve uzun vadeli kuraklık koşullarının hem güncel hem de öngörülen değerleri incelenmiştir.

Elde edilen sonuçlar, GBR modelinin özellikle mevsimsel kuraklık öngörülerinde güçlü bir başarı sergilediğini, uzun dönem SPI-12 tahminlerinde ise iklimsel eğilimleri yansıtmada etkili olduğunu göstermektedir. Böylelikle çalışma, yapay zekâ tabanlı kuraklık tahmin yöntemlerinin bölgesel ölçekte uygulanabilirliğini ortaya koymakta ve sürdürülebilir su yönetimi, tarımsal planlama ve afet risk azaltma stratejileri için bilimsel bir katkı sunmaktadır.

FULL TEXTS BOOK

MATERYAL VE YÖNTEM

Çalışma Alanı

Çalışma alanı olarak Türkiye'nin Doğu Anadolu Bölgesi'nde yer alan Elazığ ili seçilmiştir (Şekil 1). Elazığ, Fırat Nehri üzerinde konumlanmış olup, Türkiye'nin enerji üretimi ve su yönetimi açısından stratejik öneme sahip illerinden biridir. İlde, Türkiye'nin en büyük ikinci barajı olan Keban Barajı başta olmak üzere çok sayıda baraj ve hidroelektrik santral (HES) bulunmaktadır (DSİ, 2023). Bu yapılar hem enerji üretimi hem de sulama ve içme suyu temini açısından bölgesel ölçekte kritik roller üstlenmektedir. Dolayısıyla, yağış ve kuraklık koşullarındaki değişimlerin Elazığ'daki hidrolojik sistemler ve su kaynakları yönetimi üzerinde doğrudan etkileri bulunmaktadır.

Elazığ, yarı kurak iklim özellikleri ile dikkat çekmekte ve özellikle düzensiz yağış dağılımı nedeniyle kuraklık riskine açık bir coğrafyadır (Atalay, 2011). Bu nedenle uzun dönemli yağış analizleri, yalnızca bölgenin hidrolojik ve tarımsal sürdürülebilirliği açısından değil, aynı zamanda enerji güvenliği ve afet yönetimi stratejilerinin geliştirilmesi bakımından da büyük önem taşımaktadır.

Bu çalışmada, Elazığ iline ait 1938–2024 dönemini kapsayan aylık toplam yağış verileri kullanılmıştır. Bu uzun zaman serisi, iklim değişikliğine bağlı olarak yağış rejiminde ortaya çıkan değişimlerin izlenmesine ve farklı kuraklık ölçeklerinde detaylı bir değerlendirme yapılmasına olanak sağlamaktadır. Özellikle geleceğe yönelik öngörülerde, Elazığ'ın stratejik konumu ve baraj sistemleriyle Türkiye'nin su ve enerji politikaları üzerindeki kritik rolü, çalışma alanının seçiminde belirleyici olmuştur. Böylece, elde edilen bulgular yalnızca bölgesel düzeyde değil, ulusal su kaynakları yönetimi politikaları açısından da yol gösterici nitelik taşımaktadır.



Şekil 1. Elazığ İli

Standard Precipitation Index (Standart Yağış İndeksi)

SPI, basitliği, zamansal esnekliği ve yalnızca yağış verilerine dayanması nedeniyle en yaygın kullanılan meteorolojik kuraklık göstergelerinden biridir. McKee vd. (1993) tarafından geliştirilen SPI, çeşitli hidrolik ve tarımsal sistemleri etkileyen kısa, orta ve uzun vadeli nem anomalilerinin tespitini mümkün kılarak, çoklu zaman ölçeklerinde kuraklık değerlendirmesi yapılmasına olanak tanır.

FULL TEXTS BOOK

SPI, belirli bir konum için uzun vadeli aylık yağış kayıtlarına bir olasılık dağılım fonksiyonu (genellikle gama dağılımı) uydurularak hesaplanır. Bu dağılım daha sonra ortalama sıfır ve standart sapma bir olan standart normal dağılıma dönüştürülür. Ortaya çıkan SPI değerleri, gözlemlenen kümülatif yağışın belirli bir zaman ölçeği (örneğin 1 ay, 3 ay, 6 ay, 12 ay) için uzun vadeli ortalamadan sapma sayısını ifade eder.

Aylık yağış verileri, her bir çalışma alanı için uzun vadeli tarihsel kayıtlardan (en az 30 yıl) elde edilmiştir. Veri setleri kalite kontrolünden geçirilmiş ve varsa eksik değerler interpolasyon teknikleri kullanılarak işlenmiştir.

Gama dağılımı, seçilen zaman ölçeklerinde (SPI-3, SPI-6, SPI-12 ve SPI-24) toplu yağış serilerine uydurulmuştur. Yağışın sıfır olduğu aylar için, sıfırda olasılık kütlesi ile pozitif değerler için gama dağılımını birleştiren karışık bir dağılım uygulanmıştır.

Kümülatif olasılık, ters normal fonksiyon kullanılarak standart normal dağılıma dönüştürülmüş ve sonuç olarak SPI değeri elde edilmiştir (Denklem 1):

$$SPI = \Phi^{-1}(F(x)) \quad (1)$$

Φ^{-1} standart normal kümülatif dağılım fonksiyonunun (CDF) tersidir,

F(x) yağış değeri x için uyarlanmış gama dağılımından türetilen kümülatif olasılıktır.

Kısa ve uzun vadeli kuraklık dinamiklerini değerlendirmek için dört zaman ölçeği için SPI değerleri hesaplanmıştır: SPI-3: Kısa vadeli tarımsal kuraklık için kullanışlıdır. SPI-6: Akarsu debisi ve toprak nemi için ara göstergedir. SPI-12: Uzun vadeli hidrolojik kuraklık değerlendirmesi için uygundur (örneğin, rezervuar ve yeraltı suyu seviyeleri).

SPI değerleri, tablo 1'de gösterilen kuraklık şiddet sınıflarına göre kategorize edilmiştir (McKee vd., 1993):

Tablo 1. SPI değerlerinin sınıflandırılması

SPI Değeri	Kuraklık Sınıfı
0.0 - 0.99 arası	Hafif kurak
-1.00 -1.49 arası	Orta derecede kurak
-1.50 -1.99 arası	Şiddetli kurak
≤ -2.00	Aşırı kurak

SPI, Python programlama dili kullanılarak hesaplanmış ve uygulama, climate-indices veya scipy.stats kütüphanelerine dayalı olarak gerçekleştirilmiştir. Görselleştirme ve son işlemler matplotlib ve pandas kullanılarak yapılmıştır.

SPI, bölgeler arasında standartlaştırılmış karşılaştırmalar ve çoklu zaman ölçeklerine uyarlanabilirlik gibi çeşitli avantajlar sunmaktadır. Ancak, buharlaşma veya sıcaklığı hesaba katmamaktadır, bu da su talebinin yağış miktarı kadar önemli olduğu kurak ve yarı kurak bölgelerde performansını sınırlamaktadır.

Gradient Boosting Regressor (GBR)

Bu çalışmada, yağış verilerine dayalı kuraklık tahminlerinin gerçekleştirilmesi için denetimli öğrenme tabanlı makine öğrenmesi algoritmalarından Gradient Boosting Regressor (GBR) kullanılmıştır. GBR,

FULL TEXTS BOOK

temel olarak bir ansambl yöntem olup, zayıf tahminleyiciler olarak kullanılan karar ağaçlarını ardışık biçimde inşa eder ve hataları minimize ederek daha güçlü bir model oluşturur (Friedman, 2001). Model, her iterasyonda önceki tahminlerden kaynaklanan hata terimlerini (residual) hedef alarak yeni ağaçlar oluşturur ve böylece toplam hatayı kademeli olarak azaltır (Natekin & Knoll, 2013).

GBR algoritmasının en önemli avantajı, hem doğrusal olmayan ilişkileri yakalayabilmesi hem de farklı zaman ölçeklerinde karmaşık veri desenlerini öğrenebilmesidir. Bu özellik, özellikle iklim ve hidrolojik değişkenlerdeki düzensizliklerin modellenmesinde büyük önem taşımaktadır (Hastie, Tibshirani & Friedman, 2009). Model, parametre optimizasyonu için öğrenme oranı (learning rate), ağaç derinliği (max depth), ağaç sayısı (n_estimators) ve alt örnekleme oranı (subsample) gibi hiperparametrelere sahiptir. Bu parametrelerin doğru seçimi, modelin performansını doğrudan etkilemektedir (Chen & Guestrin, 2016).

Çalışmada GBR modeli, SPI-3, SPI-6 ve SPI-12 değerlerinin tahmin edilmesinde kullanılmış; her bir model, geçmiş 12 aylık yağış verisine dayalı olarak eğitilmiş ve ileriye dönük 12 aylık SPI tahminleri üretilmiştir. Model performansının değerlendirilmesinde, R^2 katsayısı, korelasyon katsayısı, Ortalama Mutlak Hata (MAE) ve Kök Ortalama Kare Hata (RMSE) metrikleri kullanılmıştır. Bu yaklaşım sayesinde, hem kısa dönemli hem de uzun dönemli kuraklık eğilimleri güvenilir şekilde öngörülmeye çalışılmıştır.

BULGULAR

Tablo 2 ve Şekil 2'de sunulan sonuçlar, Gradient Boosting Regressor (GBR) modelinin farklı zaman ölçeklerindeki tahmin performansını ortaya koymaktadır. Model, SPI-3 tahminlerinde R^2=0.786 ve korelasyon katsayısı r=0.918 ile orta düzeyde bir başarı sergilemiştir. Bu sonuç, kısa vadeli kuraklık dalgalanmalarının modellenmesinde GBR'nin belirli bir öngörü kapasitesine sahip olduğunu göstermektedir.


SPI-6 ölçeğinde ise modelin performansı belirgin şekilde artmış; R^2=0.918, korelasyon katsayısı r=0.970, düşük RMSE (0.367) ve MAE (0.294) değerleri elde edilmiştir. Bu durum, GBR modelinin özellikle mevsimsel kuraklık öngörülerinde yüksek doğruluk sağlayabildiğini ortaya koymaktadır.

Buna karşılık, SPI-12 tahminlerinde R^2=0.213 ve r=0.703 gibi görece düşük değerlere ulaşılmış; hata metriklerinin (RMSE = 0.750, MAE = 0.550) artmasıyla uzun dönemli kuraklık tahminlerinde modelin yetersiz kaldığı görülmüştür. Bu bulgu, GBR'nin kısa ve orta vadeli öngörülerde güçlü performans sergilerken, uzun vadeli iklim eğilimlerini yakalamakta sınırlı kaldığını göstermektedir.

Tablo 2. GBR modelinin tahmin doğruluk değerleri

SPI	R^2	Korelasyon	RMSE	MAE
SPI-3	0.786	0.918	0.557	0.440
SPI-6	0.918	0.970	0.367	0.294
SPI-12	0.213	0.703	0.750	0.550

FULL TEXTS BOOK



Şekil 2. SPI Tahminlerinde Model Başarı Karşılaştırması (GBR)

Şekil 3'te 2024 yılına ait SPI-3, SPI-6 ve SPI-12 kuraklık sınıflarının karşılaştırmalı durumu sunulmaktadır. Grafik incelendiğinde, yılın ilk aylarında (Ocak–Şubat) SPI-12 serisinin hafif kurak sınıfında yer aldığı, SPI-3 ve SPI-6 serilerinin ise daha çok normal ve hafif nemli koşulları yansıttığı görülmektedir.


İlkbahar aylarında (Mart–Mayıs) özellikle SPI-6 serisinde orta ve şiddetli nemli sınıflara geçiş gözlemlenirken, SPI-12 serisinin kuraklık eğilimini sürdürmesi uzun dönemli kurak koşulların hâkim olduğunu göstermektedir. SPI-3 serisinde ise daha kısa vadeli dalgalanmalar dikkat çekmekte, Mart ve Nisan aylarında nemli koşullara işaret etmektedir.

Yaz aylarına gelindiğinde (Haziran–Ağustos) SPI-3 ve SPI-6 serileri tekrar normal ve hafif nemli sınıflara gerilerken, SPI-12 serisinin büyük ölçüde hafif kurak sınıfında kalması uzun vadeli su stresi riskini ortaya koymaktadır. Özellikle Temmuz–Ağustos aylarında SPI-6'nın nemli koşullardan normal seviyelere düşmesi mevsimsel kuraklık eğilimini vurgulamaktadır.

Sonbahar aylarında (Eylül–Kasım) SPI-3 serisi hafif nemli ve yer yer orta nemli değerler gösterirken, SPI-6 serisi genel olarak normal koşulları yansıtmaktadır. SPI-12 serisi ise yılın neredeyse tamamında kuraklık eğilimini koruyarak, uzun dönemli kurak koşulların bölge için kritik olduğunu göstermektedir.

Genel olarak değerlendirildiğinde, SPI-3 kısa dönemli dalgalanmalara duyarlılık göstermekte, SPI-6 mevsimsel değişimleri daha başarılı bir şekilde yakalamakta ve SPI-12 ise uzun vadeli kuraklık eğilimlerini net biçimde ortaya koymaktadır. Bu bulgular, farklı zaman ölçeklerinde yapılan SPI analizlerinin birbirini tamamlayıcı nitelikte olduğunu ve kuraklık yönetiminde çok ölçekli yaklaşımların önemini vurgulamaktadır.

FULL TEXTS BOOK



Şekil 3. 2024 Yılı SPI-3, SPI-6, ve SPI-12 Kuraklık Sınıfları Karşılaştırması

TARTIŞMA

Bu çalışmada, Elazığ iline ait uzun dönemli yağış verileri üzerinden gerçekleştirilen SPI analizleri ve Gradient Boosting Regressor (GBR) modeli ile elde edilen tahmin sonuçları karşılaştırmalı olarak değerlendirilmiştir. Bulgular, farklı zaman ölçeklerinin kuraklık izleme ve tahmininde farklı işlevlere sahip olduğunu ortaya koymuştur. Özellikle SPI-6 serisinin yüksek doğruluk düzeyi (R^2=0.918; r=0.970) ile öne çıkması, mevsimsel kuraklık tahminlerinde GBR'nin güçlü bir öngörü kapasitesine sahip olduğunu göstermektedir. Bu bulgu, benzer çalışmalarda rapor edilen makine öğrenmesi tabanlı modellerin mevsimsel düzeyde başarılı sonuçlar verdiği yönündeki literatür ile de uyumludur (Shen, 2018; Friedman, 2001).

Buna karşılık, SPI-12 tahminlerinde düşük doğruluk değerleri (R^2=0.213) modelin uzun vadeli iklim eğilimlerini yakalamakta sınırlı kaldığını göstermektedir. Bu durum, uzun vadeli kuraklık süreçlerinin yalnızca yağış verisiyle değil, aynı zamanda sıcaklık, evapotranspirasyon ve diğer iklim parametreleri ile de ilişkilendirilmesi gerektiğini ortaya koymaktadır (Dai, 2011; Wu et al., 2007). Dolayısıyla, uzun dönem tahminlerde GBR'nin tek başına yeterli olmayabileceği, daha karmaşık modellerin veya çok değişkenli veri setlerinin kullanılması gerektiği anlaşılmaktadır.

SPI-3 sonuçları ise, kısa dönemli dalgalanmalara karşı duyarlılığı ile dikkat çekmektedir. Bu ölçek, ani yağış anomalilerinin veya kısa süreli kuraklık/nemlilik koşullarının belirlenmesinde işlevsel olmakla birlikte, model doğruluk değerleri SPI-6'ya kıyasla daha düşük kalmıştır. Bu durum, kısa vadeli öngörülerde yağış serilerinin yüksek değişkenliği nedeniyle öngörü hatalarının artabileceğini göstermektedir.

Kuraklık sınıfları üzerinden yapılan değerlendirmeler, özellikle SPI-12'nin yılın büyük bölümünde hafif kurak koşulları işaret etmesi ile bölgedeki uzun dönemli su stresi riskini vurgulamaktadır. Bu bulgu, Elazığ'ın hidrolojik ve tarımsal sürdürülebilirliği açısından kritik bir uyarı niteliği taşımaktadır. Ayrıca, SPI-6'nın ilkbaharda nemli koşulları göstermesi, bölgedeki mevsimsel su yönetimi politikalarının farklı dönemlere özgü planlanması gerektiğini ortaya koymaktadır.

FULL TEXTS BOOK

SONUÇ

Bu çalışma, yapay zekâ tabanlı yöntemlerin meteorolojik kuraklık analizlerinde bölgesel ölçekte uygulanabilirliğini ortaya koymuştur. Elde edilen sonuçlara göre:

- SPI-3 ölçeği, kısa vadeli dalgalanmaları yansıtmakta başarılı olmakla birlikte, öngörü doğruluğu görece daha düşük kalmıştır.
- SPI-6 ölçeği, GBR modeli ile yapılan tahminlerde en yüksek doğruluk düzeyine ulaşarak mevsimsel kuraklık öngörülerinde güvenilir bir araç olarak öne çıkmıştır.
- SPI-12 ölçeği, uzun dönemli iklim eğilimlerini yansıtmada önemli bir gösterge olmakla birlikte, GBR modeli bu ölçekte düşük performans sergilemiştir.

Bu bulgular ışığında, kuraklık tahmin çalışmalarında çok ölçekli SPI analizlerinin birlikte kullanılması gerektiği sonucuna varılmaktadır. Ayrıca, uzun dönemli öngörülerde yalnızca yağışa dayalı modellerin yetersiz kaldığı; sıcaklık, evapotranspirasyon ve atmosferik dolaşım verilerinin de modele entegre edilmesi gerektiği ortaya konmuştur.

Sonuç olarak, çalışma hem sürdürülebilir su yönetimi hem de kuraklıkla mücadele stratejileri açısından önemli bir bilimsel katkı sunmaktadır. Özellikle Elazığ gibi hidrolojik ve enerji üretimi bakımından stratejik öneme sahip bölgelerde, bu tür öngörüler politikacılar ve yerel yönetimler için yol gösterici nitelik taşımaktadır.

KAYNAKÇA

Atalay, İ. (2011). Türkiye'nin ekolojik bölgeleri. Meta Basım.

Chen, T., & Guestrin, C. (2016). XGBoost: A scalable tree boosting system. Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, 785–794. https://doi.org/10.1145/2939672.2939785

Dai, A. (2011). Drought under global warming: A review. Wiley Interdisciplinary Reviews: Climate Change, 2(1), 45–65. https://doi.org/10.1002/wcc.81

Devlet Su İşleri (DSİ). (2023). Türkiye barajları ve hidroelektrik santralleri. DSİ Genel Müdürlüğü Yayınları.

Friedman, J. H. (2001). Greedy function approximation: A gradient boosting machine. Annals of Statistics, 29(5), 1189–1232.

Friedman, J. H. (2001). Greedy function approximation: A gradient boosting machine. Annals of Statistics, 29(5), 1189–1232. https://doi.org/10.1214/aos/1013203451

Guttman, N. B. (1999). Accepting the standardized precipitation index: A calculation algorithm. Journal of the American Water Resources Association, 35(2), 311–322.

Hastie, T., Tibshirani, R., & Friedman, J. (2009). The elements of statistical learning: Data mining, inference, and prediction. Springer Science & Business Media.

IPCC. (2021). Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press.

McKee, T. B., Doesken, N. J., & Kleist, J. (1993). The relationship of drought frequency and duration to time scales. Proceedings of the 8th Conference on Applied Climatology, 179–184.

Natekin, A., & Knoll, A. (2013). Gradient boosting machines, a tutorial. Frontiers in Neurorobotics, 7, 21. https://doi.org/10.3389/fnbot.2013.00021

FULL TEXTS BOOK

Shen, C. (2018). A transdisciplinary review of deep learning research and its relevance for water resources scientists. Water Resources Research, 54(11), 8558–8593.

Trenberth, K. E., Fasullo, J. T., Branstator, G., & Phillips, A. S. (2014). Seasonal aspects of the recent pause in surface warming. Nature Climate Change, 4(12), 911–916.

Wu, H., Hayes, M. J., Weiss, A., & Hu, Q. (2007). An evaluation of the standardized precipitation index, the China-Z index and the statistical Z-score. International Journal of Climatology, 21(6), 745–758.

İBN RÜŞD ÜZERİNE YAPILAN DİJİTAL ÇALIŞMALAR VE YÖNTEMLER: ANALYTICA POSTERIORA ŞERHİ ÜZERİNDEN BİR UYGULAMA

DIGITAL STUDIES AND METHODS ON AVERROES: AN APPLICATION BASED ON THE COMMENTARY ON THE POSTERIOR ANALYTICS

Mehmet DUGAN

Dr. Öğr. Üyesi, Akdeniz Üniversitesi, İlahiyat Fakültesi, Felsefe ve Din Bilimleri Bölümü, Antalya, Türkiye.

ORCID ID: https://orcid.org/0000-0002-9745-4249

ÖZET

İslam düşüncesinin önemli düşünürlerinden biri olan İbn Rüşd, yalnızca teolojik eserleriyle değil özellikle Aristoteles üzerine kaleme aldığı şerhleriyle de öne çıkmaktadır. Bu şerhler 13. yüzyılda Latinceye çevrilmiş ve Avrupa'nın başlıca ilim merkezlerinde yoğun ilgi görmüştür. İbn Rüşd'ün eserleri İslam dünyasından ziyade Hristiyan ve Yahudi düşünürler arasında daha etkili olmuştur. Bu çalışma, İbn Rüşd'ün eserlerinin ve görüşlerinin dijital beşerî bilimler perspektifinden yeniden değerlendirilmesini amaçlamakta; dijital araçların sunduğu yeni yöntem ve bakış açılarına dikkat çekmeyi hedeflemektedir. Günümüzde onun fikirleri ve sonraki dönemlere olan etkisi disiplinlerarası araştırmalar açısından zengin bir kaynak teşkil etmekte, yalnızca klasik yöntemlerle değil, dijital beşerî bilimlerin sunduğu araçlarla da yeniden değerlendirilebilmektedir. Nitekim felsefe metinlerinin dijital araçlarla incelenmesi ve bu incelemenin klasik yöntemlerle desteklenmesi, modern dönemdeki felsefe araştırmalarının daha derinlikli biçimde ele alınmasına katkı sunmaktadır. Çalışmada ilk olarak İbn Rüşd'ün metinlerinin Arapça orijinalleri ile Latince ve İbranice tercümeleri ekseninde geliştirilen Averroes Edition ve Digital Averroes Research Environment (DARE) gibi dijital projeler, altyapılar ve bunların sunduğu imkânlar ele alınmıştır. Daha sonra ise bu metinlerin bulunduğu dijital veri tabanları ve araştırma ortamları incelenmiş ve bu tür platformlardan nasıl faydalanılabileceği sorusu tartışılmıştır. Araştırmacılar için veri tabanlarının ve bibliyografyaların nasıl kullanılacağı, kaynakların nasıl okunup yorumlanacağı konusunda yeni yöntemler değerlendirilmiştir. Ayrıca metin analizi, ağ analizi, stilometri, bibliyometrik analiz, uzak okuma ve görselleştirme gibi dijital yöntemlerin sunduğu yeni yorumlama biçimleri ile bu kapsamda kullanılan uygulama ve araçlardan da bahsedilmiştir. Örnek olarak İbn Rüşd'ün Kitâbu'l-Burhân eserine yazdığı orta şerhin Latince tercümesi üzerinden dijital tekniklerin nasıl kullanılabileceği gösterilmiştir. Sonuç olarak bu çalışmada İbn Rüşd üzerine yürütülen dijital projelere dikkat çekmek ve dijital yöntemlerin sunduğu avantajlar ile sınırlılıkların eleştirel bir bakışla değerlendirilmesi amaçlanmaktadır. Özellikle yazma eserlere dijital ortamlar aracılığıyla erişim, bu eserlerin analizi ve yeniden yorumlanması çalışmanın merkezinde yer almaktadır. Bu çerçevede, dijital araçların sunduğu imkânlar üzerinden İbn Rüşd araştırmalarına yeni bir metodolojik zemin kazandırma potansiyeli tartışılmaktadır.

Anahtar Kelimeler: Felsefe, Mantık, Dijital Beşerî Bilimler, İbn Rüşd, Kitâbu'l-Burhân, Metin Analizi, Görselleştirme.

ABSTRACT

Averroes, one of the most significant thinkers of Islamic thought, stands out not only for his theological works but also for his commentaries on Aristotle. In the 13th century these commentaries were translated into Latin and attracted significant interest in Europe's major centers of learning. Averroes' works had

FULL TEXTS BOOK

a greater impact on Christian and Jewish thinkers than on the Islamic world. This study reevaluates Averroes' works and views from the perspective of digital humanities, highlighting the new methods and approaches enabled by digital tools. Today, his ideas and their influence on subsequent periods constitute a rich source for interdisciplinary research and can be examined not only with classical methods but also with the tools provided by digital humanities. The analysis of philosophical texts using digital tools, combined with classical methods, contributes to a deeper and more comprehensive approach to philosophy research in the modern era. The study primarily examines digital projects and infrastructures developed around the Arabic originals and Latin and Hebrew translations of Averroes' texts, such as the Averroes Edition and the Digital Averroes Research Environment (DARE), as well as the opportunities they offer. It then reviews the digital databases and research platforms hosting these texts and discusses how these platforms can be used effectively. The study evaluates new strategies for using databases and bibliographies and for reading and interpreting sources. In addition, new forms of interpretation offered by digital methods such as text analysis, network analysis, stylometry, bibliometric analysis, distant reading, and visualization were discussed, along with the applications and tools used in these contexts. For example, the Latin translation of Averroes' Middle Commentary on the Posterior Analytics serves as a case study demonstrating how digital techniques can be used. Ultimately, this study draws attention to digital projects related to Averroes and critically evaluates the advantages and limitations of digital methodologies. The digital access, analysis, and reinterpretation of written works lie at the heart of this research. Within this framework, the potential of digital tools to provide a new methodological foundation for Averroes research is discussed.

Keywords: Philosophy, Logic, Digital Humanities, Averroes, Posterior Analytics, Text Analysis, Visualization.

GİRİŞ

Aristoteles üzerine farklı düzeylerde yazdığı şerhleri nedeniyle İslâm dünyasında *Şârih*, Latin dünyasında ise *Commentator* olarak anılan İbn Rüşd, 1126 yılında Kurtuba'da doğmuştur. Endülüs'teki önemli bir aileye mensup olan İbn Rüşd, tıpkı dedesi ve babası gibi Kurtuba kadılığı yapmıştır.[2] Ebu Ya'kub Yûsuf'un halifeliği döneminde saray hekimi olmuştur. Bu süre zarfında bir yandan teolojik eserlerini yazarken halifenin de talebiyle Aristoteles üzerine yazdığı şerhlere ağırlık vermiştir. Hayatının son dönemlerinde skolastik düşünürlere de oldukça etki edecek olan büyük şerhlerini kaleme alan İbn Rüşd, 1198 yılında Merakeş'te vefat etmiştir.[3] İbn Rüşd'ün ismi Latince metinlerde genel olarak Averroes şeklinde geçmektedir.

Bilindiği üzere İslâm dünyasında 9. yüzyılda Bağdat merkezli başlayan ve Müslümanların ilmî olarak gelişmelerine ciddi katkılar sunan tercüme hareketinin bir benzeri 12. yüzyılda Latin dünyasında yaşanmıştır.[4] Bu kapsamda gerek Yunanca eserler gerekse bunların Arapça şerhleriyle İslâm düşünürlerinin özgün eserleri tümüyle Latinceye aktarılmıştır. İbn Rüşd'ün Aristoteles üzerine yazdığı şerhlerin Michael Scotus (ö. 1235) tarafından tercüme edilmesiyle de felsefi ve teolojik tartışmaların seyri açısından 13. yüzyıl Latin dünyası için bir dönüm noktası olmuştur. Zira Albertus Magnus (ö. 1280), Thomas Aquinas (ö. 1274), Bonaventura (ö. 1274) gibi Skolastik dönemin ve Hristiyan düşüncesinin en önemli şahısları ile İbn Rüşdcülük ile ilişkilendirilen Brabantlı Siger (ö. 1284) ve Dacialı Boethius (ö. 1284?) 13. yüzyılda yaşamışlardır. İbn Rüşd'ün şerhleri ekseninde özellikle Paris

[2] Bekir Karlığa, *İbn Rüşd: Batı'yı Aydınlatan İslam Düşünürü* (İstanbul: Mahya Yayıncılık, 2014), 41.
[3] Hüseyin Sarıoğlu, *İbn Rüşd Felsefesi* (İstanbul: Klasik Yayınları, 2012), 19-21.
[4] İslâm dünyasında gerçekleşen tercüme hareketi için bk. Dimitri Gutas, *Yunanca Düşünce Arapça Kültür: Bağdat'ta Yunanca-Arapça Çeviri Hareketi ve Erken Abbasi Toplumu*, çev. Lütfü Şimşek (İstanbul: Kitap Yayınevi, 2020).

FULL TEXTS BOOK

Üniversitesi çevresinde yoğun tartışmalar yaşanmış ve sonraki yüzyıllarda Avrupa'nın pek çok şehrine yayılmıştır.

Hristiyanların yanı sıra Yahudi düşünürlerinde de İbn Rüşd'ün eserlerine teveccüh gösterdikleri görülmektedir. Bu nedenle İbn Rüşd'ün eserlerinin sadece Latince değil İbranice çevirileri de yapılmıştır. Ancak burada Latince ve İbraniceye tercüme edilen eserlerin her zaman aynı olmadığını da belirtmek gerekir. Nitekim İbn Rüşd'ün şahsı ve eserleri üzerine yapılan tartışmalar 13. yüzyıl sonrasında da sürmüştür. İslâm dünyasında ise onun böylesi bir ilgiye sahip olduğu söylemek güçtür.

Mevcut durumda İbn Rüşd'ün eserlerinin Arapça, Latince ve İbranice olmak üzere üç farklı dilde nüshaları bulunabilmektedir. Bu eserlere bir arada ulaşabilmek bir araştırmacı için oldukça kıymetlidir. Nitekim felsefe alanında yürütülen dijital çalışmaların ilk basamağı esas olarak veri tabanlarıdır.[5] Bu tür bir girişim ilk olarak Thomas Aquinas'ın metinlerini bir araya getirmeyi hedefleyen Index Thomisticus projesi ile başlamıştır.[6] Bu çalışmada ilk olarak İbn Rüşd ile ilgili olarak geliştirilen dijital veri tabanları ve projeler ele alınacaktır. Açık erişimli veri tabanlarından hareketle metinlerin çeşitli uygulamalar ve dijital araçlar vasıtasıyla okunması ve yorumlanması mümkün hale gelmiştir. Daha sonra ise bu tür platformlardan nasıl faydalanılabileceği örneklerle gösterilecektir. Bu çalışma, İbn Rüşd üzerine yapılacak araştırmalarda dijital beşerî bilimlerin sunduğu imkânlarla uygulanabilecek yeni yöntemleri tanıtmayı ve bu sayede farklı bakış açıları kazandırmayı hedeflemektedir.

1. Felsefe Alanında Yapılan Dijital Projeler

Günümüz araştırmacıları bilimsel çalışmalarında kaçınılmaz olarak bilgisayar destekli hesaplamaları ve teknolojileri kullanmaktadırlar. Sosyal bilimler alanında tasniflenen diğer disiplinler gibi felsefe ile ilişkili çalışmalarda da dijital yöntemlerin ve araçların kullanımı giderek artmaktadır. Diğer disiplinlere nazaran daha geç entegre olmuş gibi gözükse de sosyal bilimler alanında yapılan ilk dijital beşerî bilimler projesi yukarıda bahsedildiği üzere Thomas Aquinas'ın tüm eserlerinin dijitale aktarma projesidir. Roberto Busa'nın 1946 yılında IBM'in de desteği ile başlattığı Index Thomisticus adıyla bilinen bu girişim bir altyapı projesi olup, araştırmacıların Thomas Aquinas'ın tüm eserlerinin kritik edisyonu ve transkripsiyonuna ulaşabileceği açık erişimli bir veri tabanı sağlamaktadır. Böylece kelime, cümle, alıntı, benzetme, korelasyon ve istatistiksel bilgileri aramak, karşılaştırmak ve sıralamak mümkün hale gelmektedir.[7]

Benzer altyapı projelerine örnek olarak Platon'un metinlerini dijitale aktarmayı hedefleyen Dijital Platon projesi[8] ve Albertus Magnus'un eserlerini açık erişime açmayı amaçlayan Alberti Magni e-corpus projesi[9] verilebilir. Bunlar gibi tek bir filozofa odaklanan projelerin yanı sıra dönem, yüzyıl, ekol vb. sınıflamalarla muhtelif filozofların eserlerini bir araya getiren projeler de bulunmaktadır. Bunlara örnek olarak ise Yunan ve Roma dönemi felsefi ve edebi eserler ile birlikte Arapça yazılmış metinlere de erişim imkânı tanıyan Perseus Dijital Kütüphanesi,[10] farklı dönemlere ve ekollere odaklanan alt projeleri ihtiva eden ve filozofların metinleri dijital ortamda sunan Discovery Project: Philosophy in the Digital Era[11] başlıklı proje gösterilebilir.

[5] Fırat Çelebi, "Felsefe Tarihi Araştırmalarında Kullanılan Dijital Araçlar ve Metotlar", *Kaygı. Uludağ Üniversitesi Fen-Edebiyat Fakültesi Felsefe Dergisi* 22/2 (2023), 875.
[6] Bu proje hakkında daha fazla bilgi için bk. Julianne Nyhan, *Hidden and Devalued Feminized Labour in the Digital Humanities: On the Index Thomisticus Project 1954-67* (London: Routledge, 2022) Söz konuusu projeye https://www.corpusthomisticum.org/ adresinden ulaşılabilir.
[7] Bk. "A Brief Introduction to the Corpus Thomisticum", *Corpus Thomisticum* (Erişim 10 Haziran 2025).
[8] https://digital-plato.org/
[9] http://albertusmagnus.uwaterloo.ca/index.html
[10] https://www.perseus.tufts.edu/hopper/collections
[11] http://www.discovery-project.eu/home.html

FULL TEXTS BOOK

Tüm bu projelerin yanı sıra günümüzde bazı felsefe metinlerinin transkripsiyonlarına daha farklı platformlardan da erişim sağlanmaktadır. Türkiye'de ise dijital beşerî bilimler henüz yeni gelişmekte olup birtakım projeler yapılmaktadır. Nitekim Marmara Üniversitesi bünyesinde 2021 yılında Dijital Beşerî Bilimler Uygulama ve Araştırma Merkezi ve Sabancı Üniversitesi bünyesinde de 2022 yılında Dijital Beşeri Bilimler Laboratuvarı kurulmuştur. Ancak buradaki çalışmalar genel olarak tarih alanında sürdürülmektedir.[12] Felsefe tarihi alanında ise Fırat Çelebi tarafından 11. yüzyılda yaşamış Yahudi düşünür İbn Cebirol'ün eserlerini dijitale aktaran ve analiz etmeye imkân tanıyan bir proje gerçekleştirilmiştir.[13] Bu proje, Türkiye'de felsefe ile ilişkili olarak dijital beşerî bilimler alanında yapılan çalışmalara öncülük etmektedir.

2. İbn Rüşd Üzerine Yapılan Dijital Projeler

İbn Rüşd üzerine yapılan dijital projeler doğrudan ve dolaylı olmak üzere ikiye ayrılabilir. Daha önce belirtilen Index Thomisticum gibi tek bir düşünüre odaklanan projeleri ilk sınıf altında tasnif edebiliriz. Bu bağlamda Averroes Edition ve Digital Averroes Research Environment (DARE) projeleri İbn Rüşd özelinde yürütülen dijital çalışmaların başında gelmektedir. Averroes Edition projesi Kuzey Ren-Vestfalya Bilim, Beşeri Bilimler ve Sanat Akademisi (Die Nordrhein-Westfälische Akademie der Wissenschaften und der Künste) ile Köln Üniversitesi bünyesindeki Thomas Enstitüsü'nün birlikte destekledikleri bir projedir. Andreas Speer ve David Wirmer'ın yürütücülüğünü yaptığı proje *Averrois Opera (Corpus Philosophorum Medii Aevi)* projesinin bir parçası olarak 2016 yılında başlamış ve 25 yıl sürecek şekilde planlanmıştır.

Averroes Edition projesi İbn Rüşd'ün doğa felsefesine dair eserlerinin orijinal Arapçaları ile birlikte Latince ve İbranice el yazmalarının kritik edisyonunu yapmayı amaçlamaktadır. Bu eserleri önemine, projenin web sitesinde İbn Rüşd'ün Aristoteles şerhlerini Geç Antik dönem ve İslâm dünyasında Yunan felsefesinin kabulüne dair eserler dizisi ya da bir özeti şeklinde dikkat çekilmiştir. Diğerlerinin yanı sıra özellikle onun doğa felsefesine dair olan eserlerinin yüzyıllar boyunca entelektüel tarihin seyrine büyük etki ettiği belirtilir. Bu bağlamda metinlerin seçimlerinde dört temel husus üzerinden gidildiği ifade edilmektedir. Tematik olarak, İbn Rüşd'ün Aristoteles'in *Fizik* eserine ilişkin tüm şerhlerinin tam edisyonu; metodolojik olarak, İbn Rüşd'ün ilk şerhleri olan küçük şerhlerinin incelenmesi; bu incelemenin sağlam bir temele oturması bakımından İbn Bâcce'nin ilgili yorumlarının daha önce düzenlenmesidir. Son olarak ise Latince aktarımla ilgili olarak ise İbraniceden Latinceye yapılan Rönesans çeviri dalgasından önce İbn Rüşd'ün 13. yüzyıldan kalma doğa felsefesi eserlerinin yapılan tüm Arapça-Latince çevirilerinin kritik edisyonunun tamamlanmasıdır.[14] Projenin odaklandığı hususlar bakımından önemi ortaya çıkmaktadır. Zira İbn Rüşd'ün küçük şerhlerinin bugüne kadar neredeyse hiç kritik edisyonları yapılmamış olup bu duruma kendileri de değinmiştir. İbn Bâcce ile ilgili yapılacak çalışmaların nedenleri ise onun İbn Rüşd'ün başlıca kaynaklarından biri ve Endülüslü öncülü olması şeklinde belirtilmiştir. İbn Rüşd'ün onun talebesi olup olmadığı konusu tartışmalı olsa da etkilenmiş olduğu konusunda ortak bir noktada buluşulduğu söylenebilir.[15]

Projede mevcut tüm el yazmaları temelinde matbu ve dijital olmak üzere İbn Rüşd ve İbn Bâcce'ye ait on sekiz eserin kritik edisyonlarını yapmak, ana hedef olarak kaydedilmektedir. Bu eserler, Aristoteles'in *Physica, De generatione et corruptione*, *Meteorologica*'sının İbn Rüşd'ün Arapça ve

[12] Osmanlı tarihi üzerine dijital edisyon ve metin kodlama örneği için bk. Fatma Aladağ, "Osmanlı Çalışmalarında Dijital Edisyon ve Metin Kodlama: Text Encoding Initiative-TEI", *Zemin: Edebiyat, Dil ve Kültür Araştırmaları* 6 (2023), 190-211.
[13] https://ibngabirol.gitlab.io/digitalize/index.html
[14] "About – Averroes", *Averroes Edition* (Erişim 12 Haziran 2025).
[15] Bu konudaki iki farklı görüş için bk. Majid Fakhry, *Averroes (Ibn Rushd): His Life, Works and Influence* (Oxford: Oneworld, 2008), 1-2; Ernest Renan, *İbn Rüşd ve İbn Rüşdcülük: Tarihsel Deneme*, çev. Ayşe Meral (İstanbul: Albaraka Yayınları, 2021), 36.

FULL TEXTS BOOK

İbranice dillerinde bulunan küçük şerhleri ile aynı eserlere İbn Bâcce'nin yaptığı şerhler; İbn Rüşd'ün *Physica*'ya yaptığı büyük şerhin Latince ve İbranicesi ile *Physica*'nın orta şerhinin iki farklı İbranice versiyonu; *De Caelo* şerhi ile *Quaestiones Naturales*'in İbranice çevirileri; *De Substantia Orbis* ile *Meteorologica*'nın orta şerhi ve *De Animalibus*'un küçük şerhinin kısmî Latince çevirileri olarak belirlenmiştir.

Bir diğer proje ise yine aynı enstitü bünyesinde geliştirilen ve aynı akademisyenlerin yürütücülüğünü yaptığı Digital Averroes Research Environment (DARE) adlı platformdur. Bu platform İbn Rüşd'ün eserlerine ait el yazmaları ile erken modern edisyonlarının dijital görüntülerini ve bu metinlerin transkripsiyonlarını sunmayı amaçlamaktadır. Bu kapsam orijinal Arapça metinler ile bunların 16. yüzyılın sonuna kadar yapılan İbranice ve Latince çevirilerini de içermektedir. DARE Viewer aracılığı ile ise tüm bu materyali tarama, arama ve okuma imkânı geliştirilmiştir. Böylece araştırmacıya aynı eserin birden fazla versiyonunu karşılaştırma olanağı sunulmuştur.[16]

İbn Rüşd'ün eserlerinin incelenmesine olanak tanımak için bir araştırma altyapısı olarak tasarlanan DARE geniş bir veri tabanı sunmaktadır. Bu veri tabanında yukarı belirtildiği gibi birincil kaynakların yanı sıra modern araştırmacılar tarafından üretilen bilimsel yayınlar ve araştırma literatürü sistematik olarak yer almaktadır. Bu sayede İbn Rüşd ile ilgili yapılacak olan araştırmalar için ilk olarak birincil kaynakların yani el yazmalarının bir listesi ve dijitale aktarılmış görsellerini vermek; ikinci olarak ise yaklaşık iki yüz yıldır yapılan ikincil literatürü yani modern araştırmaları tek bir yerde ve tutarlı bir çerçevede bir araya getirmek ve dijital araştırma tekniklerine açmak üzere bir altyapı oluşturulmaktadır. DARE sisteminde İbn Rüşd'ün mantık, doğa felsefesi ve metafizik gibi 9 farklı kategoride, 3 dilde, 33 ülkede toplamda 159 kütüphaneden el yazması kayıtlıdır. Bunların bazılarının dijital olarak taranmış görselleri ve transkripsiyonları bulunurken diğerleri ise künye bilgileri ile arşivlenmiştir. Modern araştırmaları da bir araya getirmeyi amaçlayan DARE'nin veri tabanınında bugün itibarıyla kayıtlı 922 farklı yazara ait kitap, kitap bölümü, makale vb. türde oldukça geniş bir bibliyografya oluşturulmuştur.

Mevcut kaynakları ve edisyonları tek bir açık erişim platformunda toplamaya odaklanan DARE, İbn Rüşd'ün eserlerinin transkripsiyonlarını yayınlamaya devam etmektedir. Buradaki amaç mümkün olduğunca geniş bir metin yelpazesine erişilmesi suretiyle araştırmacıların İbn Rüşd'ün eserlerine genel bir bakış elde etmelerini sağlamaktır. Tüm bunların sonucu olarak projenin hedefi "özellikle Averroes'un eserlerinin yeni ve daha doğru baskıları ile ilgili olarak, ilham vermeyi ve ilerletmeyi umduğu yeni araştırmaların sonuçlarını özümseyebilecek bir ortam oluşturmak" olarak belirtilmiştir.[17]

Söz konusu projeler doğrudan İbn Rüşd ile ilgili olarak hayata geçirilmiştir. Bunların yanında İbn Rüşd araştırmaları için dolaylı olarak kullanılabilecek başka projeler de bulunmaktadır. Hem felsefe tarihini genel olarak ele alan hem de Index Thomisticum ve Alberti Magni e-corpus gibi Ortaçağ felsefesi için önemli olan projeler, İbn Rüşd araştırmaları için veri sağlayabilir. Nitekim karşılaştırmalı çalışmalar, felsefe araştırmaları açısından oldukça önemli bir yerde durmaktadır. Bu tür projelerin yanı sıra geniş kapsamlı ve açık erişimli bir felsefe ansiklopedisi olan Stanford Encyclopedia of Philosophy için Visualizing SEP sitesi, etkileşimli bir veri görselleştirme uygulaması sunmaktadır. Joseph DiCastro tarafından tasarlanmış ve geliştirilmiş olan bu uygulamanın Stanford Encyclopedia of Philosophy ile resmi bir ilişkisi yoktur. Bu ansiklopedide bulunan 1746 makale ve bu makaleler arasında kurulan 18916 bağlantıdan oluşan söz konusu uygulama aracılığı ile İbn Rüşd'ün hem etkilendiği hem de etkilediği düşünürlerle olan felsefi ilişkileri görselleştirilebilmektedir (Ek-1).[18] Harvard ve Tufts Üniversitelerinde ortak bir proje olarak başlatılmış ve Andrew W. Mellon Vakfı tarafından finanse edilen The Digital Corpus for Graeco-Arabic Studies projesi, İbn Rüşd'ün de eserlerinin bulunduğu felsefi ve tıbbi ağırlıklı

[16] "About - DARE", *DARE – Digital Averroes Research Environment* (Erişim 14 Haziran 2025).
[17] "About - DARE".
[18] "About Visualizing SEP", *Visualizing SEP* (Erişim 15 Haziran 2025).

FULL TEXTS BOOK

360'tan fazla metine yer vermektedir.[19] Çok fazla olmamakla birlikte bu tür dijital projeler için başka örnekler de bulunmaktadır. Ayrıca bazı kütüphanelerin açık erişimli olarak yayınladıkları el yazmaları da dijital çalışmalar için fırsat sunmaktadır.

3. Dijital Araçlarla İbn Rüşd'ün *Analytica Posteriora* Şerhinin İncelenmesi

Averroes Edition ve DARE gibi İbn Rüşd'ün eserlerinin kritik edisyonlarını yapan ve İbn Rüşd üzerine yapılmış çalışmaları bir araya getirmeyi hedefleyen projeler araştırmacılar için oldukça geniş bir veri tabanı sunmaktadır. Bu tür platformlarının yanında dijital araç ve uygulamaların da İbn Rüşd çalışmalarına katkı sağlayacağı ve yeni fikirlere olanak tanıyacağı aşikardır. Zira bu şekilde bir filozofun diğeri ile olan felsefi ilişkileri farklı bir perspektif ile izlenebilmektedir. Bu yöntemler arasında ilk olarak metinlerin dijitale aktarılması ve işlenmeye hazır hale getirilmesi ile yapılan metin analizi gösterilebilir.

Bireysel olarak doğrudan metinlerin yakından incelendiği geleneksel yöntem yakın okuma olarak adlandırılmaktadır. Çok sayıda metnin bir araya geldiği korpusların tek bir kişi tarafından geleneksel yöntemler ile okunması mümkün olmadığından metin analizinde uzak okuma yöntemi kullanılmaktadır. Franco Moretti'nin önerdiği bu yöntem yakın okuma ile ulaşılması mümkün olmayan sonuçlara varmayı hedeflemektedir.[20] Uzak okuma yöntemi ile birçok teknik kullanılarak sonuçlar elde edilebilmektedir. Literatürde bağlamlı dizin (concordance) olarak bilinen teknik, bir metinde kullanılan sözcükleri sıklık ve yoğunluk bağlamında değerlendirmeye yaramaktadır. Stilometri, edebi üslubun hesaplamalı uzak okuma yöntemleri aracılığıyla nicel olarak incelenmesidir.[21] Büyük metin koleksiyonlarında hangi konuların bulunduğunu keşfetmek için kullanılan konu modelleme, metin külliyatının içeriğini anlamlı kodlama kategorileri kümesine kodlamak için otomatik bir işlem sağlamaktadır.[22] Bir araştırma alanındaki yayınlar arasında atıf ilişkileri, yayın sıklıkları, yazar ağları gibi bilgileri inceleyen bir nicel analiz yöntemi olan bibliyometrik analiz ise diğer teknikler arasına girmektedir.[23]


Dijital beşerî bilimler alanında kullanılan uygulamalar ve araçlar ise yukarıda belirtilen yöntemlerin kolay bir şekilde uygulanmasını sağlamaktadır. Bu tür uygulama ve araçlar günden güne artmaktadır. Bunların arasında önemli olan bazı uygulama ve araçlara değinmekte fayda vardır. El yazmalarının dijitalleştirilmesi, yapay zekâ destekli metin tanıma, transkripsiyon ve arama için kapsamlı bir platform olan Aletheia[24] ya da Transkribus[25] belgeleri manuel ve otomatik olarak transkribe etmek; özel yapay zekâ modellerini eğitmek, belgeleri etiketlerle açıklamak belgeleri çeşitli formatlara aktarmak gibi amaçlarla kullanılabilmektedir. Sosyal bilimler alanındaki üretilen metinlerin dijitalleştirilmesi açısından standart haline gelen TEI (Text Encoding Initiative), her bir unsurun (başlık, paragraf, kişi adı, yer, tarih, dipnot, varyant vs.) etiketlenebildiği XML tabanlı bir kodlama dilidir. Bu nedenle pek çok dijital beşerî bilimler projesinde bu dil çeşitli yazılımlar vasıtasıyla kullanılmaktadır.[26] Bu çalışmada İbn Rüşd'ün bir metninin analizinde kullanılacak olan Voyant Tools adlı web tabanlı bir metin okuma ve analiz ortamı ise dijital beşerî bilimler alanında araştırmaları için okuma ve yorumlama uygulamalarını kolaylaştırmak amacıyla tasarlanmış projedir. Voyant Tools ile ilgili metne dair, kelime

[19] "About - A Digital Corpus for Graeco-Arabic Studies", *The Digital Corpus for Graeco-Arabic Studies* (Erişim 16 Haziran 2025).
[20] Franco Moretti, *Distant Reading* (London, New York: Verso, 2013), 65-67.
[21] Stilometri temelli yapılan bir çalışma için bk. Helena Montserrat Gomez Adorno vd., "Stylometry-based Approach for Detecting Writing Style Changes in Literary Texts", *Computación y Sistemas* 22/1 (2018), 47-53.
[22] John W. Mohr - Petko Bogdanov, "Introduction—Topic Models: What They Are and Why They Matter", *Poetics* 41/6 (2013), 546.
[23] Naveen Donthu vd., "How to Conduct a Bibliometric Analysis: An Overview and Guidelines", *Journal of Business Research* 133 (2021), 285-286.
[24] https://www.primaresearch.org/tools/Aletheia
[25] https://www.transkribus.org/
[26] https://tei-c.org/

FULL TEXTS BOOK

sıklığı, frekansı, sayısı, bağlam dizini gibi hususları tablo, bulut, diyagram, grafik vb. biçimde gösterilebilmektedir.

Bu çalışma kapsamında İbn Rüşd'ün bir metni analiz edilerek örnek bir uygulama yapılmıştır. Bu nedenle ilk olarak İbn Rüşd'ün *Kitâbu'l-Burhân* üzerine yazdığı orta şerhine DARE Viewer üzerinde ulaşıldı (Ek-2). Bu metnin söz konusu platform üzerinden çeşitli kütüphanelerdeki el yazmalarının dijitallerine ve bir transkripsiyonuna erişilebilmektedir. İstenildiği takdirde nüshalar arasında karşılaştırma yapma imkânı sunan platform, araştırmacılar için büyük kolaylık sağlamaktadır (Ek-3). Bu kapsamda Lunalı William tarafından Latinceye tercüme edilen ve *Commentum super libris posteriorum analecticorum* adıyla bilinen eserin transkripsiyonuna bu platform üzerinde erişilmiştir.[27] XML ya da HTML formatında indirilebilir halde olan bu eser, dijital tekniklerle analiz için hazır durumdadır.



Şekil 1. Commentum super libris posteriorum analecticorum'un TEI kodlamasından bir görünüm

Mezkûr eserin analizini yapabilmek için ilk olarak DARE Viewer üzerinden indirilen TEI standardında etiketlenmiş olan XML formatındaki dosyanın Voyant Tools'a yüklenmiştir. Ardından genel bir bilgi için "Summary" başlığına göz atılabilir. Burada metnin toplam kelime sayısı, mükerrer olmayan kelime sayısı, kelime yoğunluğu, okunabilirlik endeksi, cümle başına ortalama kelime sayısı ve tüm metin boyunca en sık kullanılan kelimeler otomatik olarak hesaplanmaktadır.

[27] Averroes, *Commentum super libris posteriorum analecticorum*, transl. William of Luna. Edited transcription of the editio princeps in: Aristoteles, Opera, Part I, 1: Organon, ed. Nicoletus Vernia, Venice: Andreas Torresanus, Bartholomaeus de Blavis, 2 Feb. 1483, n.p. [foll. 75r–88r] (Gesamtkatalog der Wiegendrucke 2337). Cologne: Digital Averroes Research Environment (DARE), 2011. URI: dare.uni-koeln.de/app/fulltexts/FT5 .

Summary ?

This corpus has 1 document with 34,672 total words and 4,092 unique word forms.
Created about 18 minutes ago.

Vocabulary Density: 0.118

Readability Index: 13.138

Average Words Per Sentence: 35.1

Most frequent words in the corpus:

- scientia (198); possibile (136); una (132); oportet (125); demonstratio (118); propositiones (115); inventum (96); medium (80); diffinitio (79); inventa (73)

Şekil 2. Söz konusu metne dair genel bilgiler.

Herhangi bir metinde en sık geçen kelimeler dijital araçlar ile çeşitli şekillerde gösterilebilir. Bunun en yaygın kullanılan türlerin biri kelime bulutudur. Analiz edilen metinde en çok kullanılan kelimelerin Voyant Tools yardımı ile kelime bulutu gösterimi ise aşağıdaki gibidir. Burada en sık kullanılan kelime en büyük olacak şekilde sıralanmaktadır.




Şekil 3. Kelime bulutu olarak gösterimi

Bağlamsal dizin ise bir metinde geçen kelimenin bağlamını ortaya koymasın açısında önemlidir. Bu uygulama ile o kelimenin geçtiği tüm cümleler ve birlikte kullanıldığı kavramlar kolayca görülebilir. Bunun sağladığı husus ise yakın okuma ile gözden kaçabilecek bağlamı bu yöntemle daha anlaşılabilir ve görülebilir hale getirmektir. Ele alınan metinde en çok geçen *scientia* kelimesinin bağlamının bir kısmı gösterilmektedir.

Left	Term	Right
apparet, quod scien tia comprehensa ex doctrina erit quidem ex	scientia	praecedente discipulo. Et similiter apparet in aliis rebus, quarum pro
ddiscat conclusionem syllogismi, quod iam praecesserit apud eum	scientia	propositionum syllogismi. Et qui certificat propositionem universalem pe
doceat rem, quam doceat propter exemplum. Commentum 2 [2] Et	scientia	quam oportet praecedere omne de cuius pro prietate est, ut
per intellectum et syllo gismum, est secundum duos modos aut	scientia	, ut sit inven tum aut non inventum. Et haec est
ut sit inven tum aut non inventum. Et haec est	scientia	, quae dicitur verificatio. Aut scientia, super quam significet nomen rei
non inventum. Et haec est scientia, quae dicitur verificatio. Aut	scientia	, super quam significet nomen rei. Et illa est, quae no
formatio. et de quibusdam rebus oportet, ut praecedat addiscenti	scientia	, quae invenitur tamen, sicut est propositio, in qua dicitur, quod
quibusdam opor tet, quod praecedat et sciatur ex re ipsorum	scientia	significans nomen ipso rum tamen, sicut ut praeponat geometra triangu
et quod ipsa res est inventa, et illud est, quia	scientia	in inventione rei est per scientiam, quid significet nomen ipsius
ationem nominis. [3] Haec est antici patio scientiae praecedentis in	scientia	discipuli, sicut sensibilium prio rum in re ad sensibilia secunda
quoniam si esset dispositio scientiae acquisitae per doctrinam cum	scientia	praece dente, haec dispositio esset necesse, ut sit disciplina recordatic
facit nos dubitare in hac propositione, si dicatur, quod omnis	scientia	et omnis disciplina erit quidem per scientiam praecedentem, quia iam
s, praeter quod praecesserit nobis sensus ipsarum, quoniam haec	scientia	et scientia proveniens secundum disciplina est scientia in aequivocatio
quod praecesserit nobis sensus ipsarum, quoniam haec scientia et	scientia	proveniens secundum disciplina est scientia in aequivocatione nominis.
oniam haec scientia et scientia proveniens secundum disciplina est	scientia	in aequivocatione nominis. Commentum 3 Et quarumdam rerum proven
ationе nominis. Commentum 3 Et quarumdam rerum provenit nobis	scientia	per sensum post disciplinam ipsarum simul. Et illae sunt res
eius sunt aequales duobus rectis. Et propositio minor in hac	scientia	provenit ex sensu, quae est, quod est triangulus, et conclusio

Şekil 4. Bağlamsal dizin gösterimi

Söz konusu metinde geçen kelimelerin sıklığı kelime bulutu ile görselleştirilmişti. Ancak sadece seçilen kelimelerin gösterimi için kelime bulutundan başka bir görselleştirme aracına ihtiyaç vardır. Bunlardan biri olan göreli sıklık, bir derleme veya tek bir belge için farklı olacak şekilde analiz edilmek istenen istenilen kelimeler özelinde yapılabilmektedir. Buradaki uygulama, İbn Rüşd'ün *Kitâbu'l-Burhân* eserine yazdığı orta şerhin Latince tercümesi üzerinden gerçekleşmektedir. Dolayısıyla derlemede bir belge olduğundan belgede otomatik olarak oluşturulan her bir segment için ilgili terimin sıklığını gösterilir. Aşağıdaki örnekte seçilen dört farklı kelime için yapılan göreli sıklık grafiği görülmektedir.



Şekil 5. En çok görülen dört terim için göreli sıklık grafiği

SONUÇ

İbn Rüşd'ün eserleri Ortaçağ Latin ve Yahudi düşüncesi üzerinde de belirleyici bir etkiye sahip olmuştur. Görüşleri çok dilli bir yazma kültürü içerisinde uzun bir süre dolaşımda olmuştur. Son iki asırda ise onun eserleri ve görüşleri araştırmacılar tarafından yeniden çalışılmaktadır. Sadece şerhleri açısından bakıldığında dahi geleneksel yöntemler ile bazı hususlara ulaşmanın zor olduğu aşikardır. Nitekim Aristoteles üzerine yazdığı 38 şerhin Arapçalarının bir kısmı günümüze ulaşamadığı için sadece

FULL TEXTS BOOK

tercümelerinden hareketle araştırmaları yürütmek mümkündür. Bundan dolayı bu eserlerin günümüzde yeniden ele alınması, yalnızca klasik filolojik yaklaşımlarla değil, dijital beşerî bilimlerin sunduğu yöntemlerle de değerlendirmesi önemlidir. Bu çalışmada İbn Rüşd'ün eserlerine yönelik geliştirilen dijital projeler tanıtılmış; özellikle çok dilli metinlerin dijital temsilleri, karşılaştırmalı edisyon olanakları ve veri tabanlarının sunduğu işlevsellik bağlamında çeşitli okuma ve yorumlama yolları değerlendirilmiştir. Burada İbn Rüşd'ün *Kitâbu'l-Burhân* üzerine yazdığı orta şerh üzerinden bahsedilen dijital altyapıların, yöntemleri ve tekniklerin bazı uygulama ve araçlar ile nasıl kullanılacağı gösterilmiştir. Bu sayede dijital yöntemlerin sadece erişimi kolaylaştırmadığını aynı zamanda metni çok katmanlı ve karşılaştırmalı bir şekilde yeniden okuma ve anlamlandırma imkânı sunduğu gösterilmiştir. Bu çerçevede dijital platformlar, araştırmacıların klasik metinler üzerinde hem yorumlayıcı hem de yapısal düzeyde çalışmasına olanak tanıyan yeni bir metodolojik zemin ortaya koymaktadır. Bununla birlikte, bu tür dijital çalışmalarda bağlamdan kopuk okuma riskleri, teknik altyapı sorunları gibi bazı sınırlılıkların da göz önünde bulundurulması gerekir. Türkiye bağlamında ise dijital filolojiye dayalı araştırmaların henüz yeterince kurumsallaşmamış olduğu ancak mevcut uluslararası örneklerin bu alanda önemli bir ilham kaynağı olabileceği söylenebilir. Bu nedenle araştırmacıların mevcut projelerden hareketle yeni girişimlerde bulunması veya yeni araştırmalarında dijital yöntemlerin imkanlarından daha fazla yararlanmaları önerilir. Örneğin İbn Rüşd'ün üç düzeyde yazdığı bir şerhine dijital teknikler uygulanarak yeni sonuçlar çıkarılabilir. İbn Rüşd ile başka bir filozofun benzer eserler üzerinden karşılaştırmaları yapılabilir. Sonuç olarak dijital veri tabanlarının gelişmesi ve dijital yöntemlerin kullanımı araştırmacılara metinlere kolay erişim sağladığı gibi geleneksel yöntemler ile ulaşılamayacak verilerin de önünü açmaktadır. Bu nedenle bu tür girişimlerin artması ve dijital beşerî bilimlerin öğrenilmesi ve öğretilmesinin yararlı olacağı düşünülmektedir.

KAYNAKÇA

Aladağ, Fatma. "Osmanlı Çalışmalarında Dijital Edisyon ve Metin Kodlama: Text Encoding Initiative-TEI". *Zemin: Edebiyat, Dil ve Kültür Araştırmaları* 6 (2023), 190-211. https://doi.org/10.5281/ZENODO.10435921

Çelebi, Fırat. "Felsefe Tarihi Araştırmalarında Kullanılan Dijital Araçlar ve Metotlar". *Kaygı. Uludağ Üniversitesi Fen-Edebiyat Fakültesi Felsefe Dergisi* 22/2 (2023), 872-892. https://doi.org/10.20981/kaygi.1141102

Donthu, Naveen vd. "How to Conduct a Bibliometric Analysis: An Overview and Guidelines". *Journal of Business Research* 133 (2021), 285-296. https://doi.org/10.1016/j.jbusres.2021.04.070

Fakhry, Majid. *Averroes (Ibn Rushd): His Life, Works and Influence*. Oxford: Oneworld, Reprinted., 2008.

Gomez Adorno, Helena Montserrat vd. "Stylometry-based Approach for Detecting Writing Style Changes in Literary Texts". *Computación y Sistemas* 22/1 (2018), 47-53. https://doi.org/10.13053/cys-22-1-2882

Gutas, Dimitri. *Yunanca Düşünce Arapça Kültür: Bağdat'ta Yunanca-Arapça Çeviri Hareketi ve Erken Abbasi Toplumu*. çev. Lütfü Şimşek. İstanbul: Kitap Yayınevi, 2020.

Karlığa, Bekir. *İbn Rüşd: Batı'yı Aydınlatan İslam Düşünürü*. İstanbul: Mahya Yayıncılık, 2014.

Mohr, John W. - Bogdanov, Petko. "Introduction—Topic Models: What They Are and Why They Matter". *Poetics* 41/6 (2013), Bibliometric. https://doi.org/10.1016/j.poetic.2013.10.001

Moretti, Franco. *Distant Reading*. London, New York: Verso, 2013.

FULL TEXTS BOOK

Nyhan, Julianne. *Hidden and Devalued Feminized Labour in the Digital Humanities: On the Index Thomisticus Project 1954-67*. London: Routledge, 2022. https://doi.org/10.4324/9781003138235

Renan, Ernest. *İbn Rüşd ve İbn Rüşdcülük: Tarihsel Deneme*. çev. Ayşe Meral. İstanbul: Albaraka Yayınları, 2021.

Sarıoğlu, Hüseyin. *İbn Rüşd Felsefesi*. İstanbul: Klasik Yayınları, 2012.

Corpus Thomisticum. "A Brief Introduction to the Corpus Thomisticum". Erişim 10 Haziran 2025. https://www.corpusthomisticum.org/wintroen.html


The Digital Corpus for Graeco-Arabic Studies. "About - A Digital Corpus for Graeco-Arabic Studies". Erişim 16 Haziran 2025. https://www.graeco-arabic-studies.org/about

Averroes Edition. "About – Averroes". Erişim 12 Haziran 2025. https://averroes.uni-koeln.de/project/about/


DARE – Digital Averroes Research Environment. "About - DARE". Erişim 14 Haziran 2025. https://dare.uni-koeln.de/project/about-dare/

Visualizing SEP. "About Visualizing SEP". Erişim 15 Haziran 2025. https://www.visualizingsep.com/#/about

Ekler:




Ek-1: *Visualizing SEP ile yapılan İbn Rüşd merkezli bir görselleştirme.*



Ek-1: İbn Rüşd'ün Kitâbu'l-Burhân üzerine yazdığı orta şerhin DARE Viewer ile erişilebilen nüshaları

FULL TEXTS BOOK



*Ek-2: DARE Viewer üzerinden İbn Rüşd'ün Kitâbu'l-*Burhân *üzerine yazdığı orta şerhin Latince ve Arapça transkripsiyonlarının karşılaştırılması.*

DOMATESİN BAKTERİYEL HASTALIKLARI ÜZERİNE YAPILAN BİLİMSEL YAYINLARIN BİBLİYOMETRİK ANALİZİ

Dr. Büşran SUNYAR [1]

[1]*Iğdır Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Iğdır, Türkiye* [1]*ORCID ID: https://orcid.org/0000-0001-8524-3308*

Doç. Dr. Mesude Figen DÖNMEZ [2]

[2] *Doç. Dr., Iğdır Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Iğdır, Türkiye* [2]*ORCID ID: https://orcid.org/0000-0002-7992-8252*

ÖZET

Domates (*Solanum lycopersicum* L.), dünya genelinde üretimi ve tüketimi en yüksek sebze türlerinden biri olup gıda sanayisinde geniş kullanım alanına sahiptir. Türkiye, domates üretim ve ihracatı bakımından ön sıralarda yer almaktadır. Ancak domates üretimini sınırlayan başlıca faktörlerden biri bakteriyel hastalıklardır. Bu hastalıklar arasında *Pseudomonas syringae* pv. *tomato, Clavibacter michiganensis* subsp. *michiganensis, Xanthomonas axonopodis* pv. *vesicatoria, Ralstonia solanacearum*ve *Pectobacterium carotovorum* gibi önemli patojenler yer almaktadır. Çalışmada, bu bakteriyel etmenler üzerine yapılan bilimsel yayınların gelişimini ortaya koymak amacıyla bibliyometrik analiz gerçekleştirilmiştir. Scopus veri tabanında 1953–2025 yılları arasında yayımlanan 794 belge incelenmiş; analizler VOSviewer ve R programları kullanılarak yürütülmüştür. Bulgular, bilimsel yayınların özellikle 2000 sonrası hız kazandığını, üretimin büyük oranda çok yazarlı iş birliklerine dayandığını ve en fazla katkının ABD, Çin ve Japonya gibi ülkelerden geldiğini göstermiştir. Anahtar kelime analizleri, araştırmaların zamanla daha moleküler konulara yöneldiğini ortaya koymuştur. Ayrıca, eş atıf ve iş birliği ağları, belirli dergiler ve ülkeler arasında güçlü akademik bağların oluştuğunu ifade etmiştir. Bu çalışma domates bakteriyel hastalıkları alanındaki bilimsel eğilimleri değerlendirerek gelecekteki araştırmalara yön verecek stratejik bilgiler sunmaktadır.

Anahtar kelimeler: Domates, Bakteriyel Hastalıklar, Bibliyometrik analiz

GİRİŞ

Domates (*Solanum lycopersicum* L.), hem taze tüketim hem de işlenmiş ürün sanayisinde yaygın olarak kullanılan, dünya genelinde stratejik tarım ürünlerinden biridir. 2022 yılı itibarıyla dünya genelinde 180 milyon tondan fazla domates üretildiği ve bu üretimin büyük kısmının Asya, Avrupa ve Amerika kıtalarında yoğunlaştığı rapor edilmiştir (FAO, 2023). Domatesin bu denli yaygın üretimi, beraberinde çeşitli biyotik stres faktörlerini de gündeme getirmektedir. Özellikle bakteriyel hastalıklar, domatesin verim ve kalite potansiyelini tehdit eden önemli etmenler arasında yer almakta ve üreticilere ciddi ekonomik kayıplar yaşatmaktadır (Jones et al., 2014; Mansfield et al., 2012). Bu bakteriyel hastalıklar arasında bakteriyel benek [*Pseudomonas syringae* pv. *tomato* (Young et al., 1978)], bakteriyel kanser ve solgunluk [*Clavibacter michiganensis* subsp. *michiganensis* (Davis et al., 1984)], bakteriyel yaprak lekesi [*Xanthomonas euvesicatoria* (Gardner and Kendrick, 1923)], bakteriyel solgunluk [*Ralstonia solanacearum* (Yabuuchi et al., 1995)], bakteriyel gövde ve meyve çürüklüğü [*Pectobacterium carotovorum* subsp. *carotovorum* (Bergey et al., 1923)] ve domates öz nekrozu [*Pseudomonas corrugata* (Scarlet et al., 1978), *Pseudomonas cichorii*, *Pseudomonas viridiflava* (Wilkie and Dye, 1974)] gibi etmenlerin neden olduğu hastalıklar öne çıkmaktadır. Bu patojenler, domatesin fide döneminden hasada kadar olan tüm gelişim sürecinde zarara yol açabilmekte; özellikle sıcak ve nemli

FULL TEXTS BOOK

çevre koşullarında hızlı yayılım göstererek geniş çaplı enfeksiyonlara neden olabilmektedir (Hayward, 1991; Jones et al., 2000).

Geleneksel hastalık yönetim stratejilerinde kimyasal kontrol yöntemlerinin etkisinin sınırlı kalması, çevresel sürdürülebilirlik endişeleri ve patojenlerdeki direnç gelişimi gibi sorunlar nedeniyle, bakteriyel hastalıkların teşhisi, epidemiyolojisi, direnç genlerinin tanılanması ve biyolojik mücadele alternatifleri üzerine yapılan bilimsel çalışmalar son yıllarda giderek artmıştır (McManus et al., 2002). Bu artış, alanın bilimsel gelişimini, araştırma eğilimlerini ve akademik iş birliği ağlarını incelemeye olan ihtiyacı doğurmuştur.

Bibliyometrik analiz, belirli bir araştırma alanındaki çalışmaları belirli bir dönemde hem nitel hem de nicel olarak incelemek ve analiz etmek için matematiksel ve istatistiksel yöntemler kullanan bir yöntemdir (Hicks vd., 2015). Bu yöntem, belirli bir alandaki araştırmalarla ilgili ülkelere, kurumlara, dergilere, yazarlara ve anahtar kelimelere odaklanarak okuyuculara alandaki eğilimler ve sınırlar hakkında nesnel bir görünüm sağlamaktadır (Deng vd., 2022; Ninkov vd., 2022).

Bu çalışmanın amacı, domatesin bakteriyel hastalıkları üzerine yapılmış akademik yayınları bibliyometrik yöntemlerle inceleyerek alandaki araştırma eğilimlerini, öne çıkan yazarları, kurumları, ülkeleri, iş birliği ağlarını ve tematik odakları ortaya koymaktır. Bu doğrultuda, Scopus veri tabanı temel alınarak gerçekleştirilen analizler; literatürün gelişim sürecini, bilgi üretim merkezlerini ve gelecek çalışmalara ışık tutabilecek yönelimleri kapsamlı biçimde değerlendirmeyi amaçlamaktadır.


MATERYAL ve YÖNTEM

Çalışmada, domates bitkisinde önemli bir hastalık olan bakteriyel bakteriyel hastalıkların “tomato” ve “bacterial pathogen” anahtar kelimeleri Scopus veri tabanında arama yapıldığında elde edilen 794 yayın üzerinde VOSviewer ve R programları kullanılarak bibliyometrik analizi yapılmıştır.


Yöntem

Bibliyometrik analiz çerçevesinde, Scopus veri tabanında gerçekleştirilen tarama sonucunda tespit edilen 794 akademik yayın analiz kapsamına alınmıştır. Uygulanan yöntemsel süreç ise Şekil 1’de şematik olarak detaylandırılmıştır.

FULL TEXTS BOOK



Şekil 1. Bibliyometrik analizi için veri indirme şeması



Şekil 2. Çalışmada kullanılan Vosviwer ve R progmında yapılan analizler

Veri Analizi

Ülkelere göre yayın dağılımı, yayımlanan dergiler ve benzeri bibliyometrik değişkenlere ilişkin betimsel analizler Microsoft Excel yazılımı kullanılarak yapılmıştır. Atıf ilişkileri ile anahtar kelimeler arasındaki

FULL TEXTS BOOK

eşzamanlılık ve bu ilişkilerin görsel temsilleri VOSviewer programı aracılığıyla analiz edilmiştir. Ayrıca, ülkeler arasındaki bilimsel iş birliği ağları, en sık karşılaşılan anahtar terimler, anahtar kelimelerin zaman içerisindeki değişim eğilimleri ve kavramsal gruplamaları (co-word analizi) R yazılımı üzerinden, bibliometrix paketinin web tabanlı grafik arayüzü olan "biblioshiny" kullanılarak detaylı biçimde değerlendirilmiştir.

BULGULAR

1953–2025 yılları arasını kapsayan veri setinde toplam 794 belgenin yayımlandığı belirlenmiştir. Bu belgelerin 275 farklı dökümanın yer aldığı, yıllık ortalama artış oranının %4,74 olduğu tespit edilmiştir. Her belgeye ortalama 55,95 atıf yapıldığı görülmüştür. İçerik analizinde 1788 yazar anahtar kelimesi kaydedilmiştir. Toplam 3353 yazarın katkı sunduğu bu çalışmalarda, yalnızca 16 yazarın tek yazarlı dökümanlara sahip olduğu; belge başına ortalama 5,86 ortak yazar düştüğü saptanmıştır. Belge türleri incelendiğinde, 731 adetle en çok makale türünün kullanıldığı; bunu sırasıyla 25 derleme, 15 kitap bölümü ve 13 konferans bildirisinin izlediği belirlenmiştir. Elde edilen bulgular, alandaki yayın faaliyetlerinin büyük ölçüde çok yazarlı iş birlikleriyle ve makale formatında yürütüldüğünü göstermektedir (Tablo 2).


Tablo 2. Domateste görülen bakteriyel hastalıklara ilişkin yayınların ana bilgileri

Veri Hakkında Temel Bilgiler	
Zaman Aralığı	**1953:2025**
Döküman (Dergiler, Kitaplar vb.)	**275**
Belgeler	**794**
Yıllık Büyüme Oranı (%)	**4,74**
Belge Ortalama Yaşı	**10,1**
Belge Başına Ortalama Atıf	**55,95**
Yazar Anahtar Kelimeleri (DE)	**1788**
Yazarlar	
Yazar Sayısı	**3353**
Tek yazarlı belgelerin yazarları	**16**
Tek yazarlı belgeler	**17**
Belge Başına Ort. Ortak Yazar	**5,86**
Belge Türleri	
Makale	**731**
Kitap bölümü	**15**
Konferans bildirisi	**13**

FULL TEXTS BOOK

Düzeltme (Erratum)	**4**
Not	**4**
Derleme	**25**
Kısa anket	**2**


Çalışmada yıllara göre bilimsel yayın üretiminin dağılımında 1953–1990 yılları arasında oldukça sınırlı sayıda yayının üretildiği gözlemlenmiştir. 1990'lı yılların sonlarından itibaren yayın sayılarında artış eğilimi kaydedilmiştir. Özellikle 2000'li yıllarla birlikte bu artışın ivme kazandığı belirlenmiştir. 2016–2022 yılları arasında yıllık yayın sayısının 60'ın üzerine çıktığı ve bu dönemin en yoğun bilimsel üretimin gerçekleştiği dönem olduğu tespit edilmiştir. 2023 ve 2025 yıllarında düşüş olduğu tespit edilmiştir (Şekil 3). Genel olarak, zaman içerisinde bilimsel üretimin istikrarlı bir şekilde arttığı ve son on yılda bu artışın belirginleştiği ortaya konulmuştur.




Şekil 3. Yıllara göre yayın sayısı

Ülkeler arası bilimsel etkileşimlerin yapısını ortaya koymaya yönelik olarak oluşturulan ağ haritasında, uluslararası iş birliği ilişkileri görsel olarak sunulmuştur. Düğümlerin büyüklüğü, ilgili ülkelerin akademik üretkenlik düzeylerini; düğümler arasındaki çizgiler ise ülkeler arasındaki iş birliği yoğunluğunu yansıtmaktadır. Analiz sonucunda, en yüksek bilimsel üretkenliğin ve iş birliği düzeyinin Amerika Birleşik Devletleri merkezli olduğu belirlenmiş; bu ülkenin özellikle Çin, Almanya, Fransa, Japonya, Güney Kore ve Birleşik Krallık ile güçlü akademik iş birlikleri yürüttüğü tespit edilmiştir. Çin'in de yüksek düzeyde bilimsel üretime sahip olduğu ve çok sayıda ülkeyle ortak çalışmalarda yer aldığı belirlenmiştir. Avrupa ülkeleri arasında Almanya, Fransa, İspanya ve İtalya ön plana çıkarken; Asya kıtasında Japonya, Güney Kore ve Hindistan'ın aktif katkı sunduğu belirlenmiştir (Şekil 4). Genel olarak, bilimsel iş birliği ağlarının belirli merkez ülkeler etrafında yoğunlaştığı ve bu iş birliklerinin küresel akademik üretimi şekillendirdiği tespit edilmiştir. Ülkeler arasındaki bilimsel iş birliklerinin zaman içindeki dağılımı Şekil 5'te gösterilmiştir. Sarı renge yakın tonlarla gösterilen ülkelerin, özellikle 2018 sonrası dönemde daha güncel yayınlarla uluslararası iş birliklerinde yer aldığı belirlenmiştir. Çin, Hindistan, İtalya, Romanya, Polonya ve Ukrayna gibi ülkelerin son yıllarda bilimsel yayın üretimine daha aktif biçimde katkı sunduğu gözlemlenmiştir. Buna karşın, koyu mavi tonlarla gösterilen ülkelerde iş birliklerinin daha çok önceki yıllarda gerçekleştiği belirlenmiştir.

FULL TEXTS BOOK



Şekil 4. Ülkelerin ağ grafiği analizi




Şekil 5. Ülkelerin ağ grafiği analizinin güncel durumları

Ülkeler arasındaki bilimsel iş birliği yapısını ortaya koyan ağ analizine göre, uluslararası düzeyde en fazla ortak yayının Amerika Birleşik Devletleri, Çin ve bazı Avrupa ülkeleri tarafından üretildiği


FULL TEXTS BOOK

belirlenmiştir. Yüksek düzeyde iş birliğini temsil eden koyu mavi tonlar ile ülkeler arası bağlantılar üzerinden ortak yayın faaliyetleri görselleştirilmiştir. Elde edilen bulgular doğrultusunda, pek çok ülkenin bilimsel üretime uluslararası ölçekte katkı sunduğu, ancak bu iş birliklerinin belirli coğrafi bölgelerde yoğunlaştığı ve araştırma ağlarının küresel ölçekte merkezî bir yapı olduğu belirlenmiştir (Şekil 6).



Şekil 6. Ülkelerin ortak yayın haritası


Grafikte, ilgili yazarların ülkelerine göre bilimsel yayın dağılımı gösterilmiştir (Şekil 7). En fazla yayının Amerika Birleşik Devletleri (ABD) tarafından üretildiği belirlenmiştir. ABD'yi sırasıyla Çin, Güney Kore, Japonya ve Hindistan'ın izlediği görülmektedir. Yayınların büyük bir kısmının tek ülke kökenli (SCP) olarak gerçekleştirildiği, ancak bazı ülkelerde çok uluslu iş birliklerine dayalı yayınlara (MCP) da yer verildiği anlaşılmaktadır. Özellikle Çin, Japonya, Almanya ve Birleşik Krallık gibi ülkelerde uluslararası iş birliklerine yönelik yayın oranının daha yüksek olduğu dikkat çekmiştir. Türkiye'nin de bu sıralamada yer aldığı ve ağırlıklı olarak tek ülke kökenli yayınlarla katkı sunduğu tespit edilmiştir. Elde edilen veriler, bilimsel üretimin büyük ölçüde belirli ülkeler tarafından yürütüldüğünü ve bazı ülkelerde uluslararası iş birliğine dayalı çalışmaların öne çıktığını ortaya koymuştur.



Şekil 7. Sorumlu yazarların ülkelerine göre yayın dağılımı ve uluslararası işbirlikleri

FULL TEXTS BOOK


Araştırmada, çalışmalara en fazla katkı sunan yazarlar sayıca sıralanmıştır. En üretken yazar olarak öne çıkan Martin Gb'nin toplam 20 yayınla birinci sırada yer aldığı belirlenmiştir. Bu ismi sırasıyla 17 yayınla Jones Jb, 15 yayınla Wang Y, 14 yayınla Wang X ve 13 yayınla Zhang Y takip etmiştir. Sıralamanın devamında Li J ve Staskawicz Bj (her biri 12 yayın), He Sy ve Lee S (her biri 11 yayın) ile Chen J (10 yayın) yer almıştır (Şekil 8). Ayrıca isimlerin coğrafi dağılımı dikkate alındığında, özellikle Asya ve Kuzey Amerika merkezli araştırmacıların ön planda olduğu görülmektedir. Bu durum, söz konusu yazarların hem bireysel üretkenliğini hem de uluslararası iş birliği ağlarında etkin konumda olduklarını göstermektedir.



Şekil 8. Literatürde en fazla yayına sahip olan yazarlar ile bu yazarların belge sayıları


Çalışmada önde gelen yazarların yıllara göre bilimsel üretimleri ve bu üretimlerin yıllık atıf düzeylerinde, her bir daire ile belirli bir yılda yayımlanan makale sayısı ve bu makalelerin aldığı toplam atıf temsil edilmiştir; dairelerin büyüklüğü yayın sayısını, renk yoğunluğu ise yıllık atıf etkisini yansıtmaktadır. Staskawicz Bj'nin yayın faaliyetlerinin 1990'lı yıllarda yoğunlaştığı, Martin Gb ve Jones Jb'nin ise daha uzun bir zaman dilimine yayılan düzenli bir üretim süreci içinde yer aldığı gözlemlenmiştir. 2016–2020 döneminde Wang Y, Wang X ve Zhang Y gibi yazarların hem üretim düzeyi hem de atıf etkisi bakımından daha fazla öne çıktığı tespit edilmiştir (Şekil 9). Genel olarak, yazarların akademik katkılarının zaman içinde farklılık gösterdiği ve bazı dönemlerde belirli isimlerin alanda daha etkili olduğu belirlenmiştir.

FULL TEXTS BOOK



Şekil 9. Yazarların üretkenlik zamanları


Ağ haritasında, akademik dergiler arasındaki eş atıf ilişkileri farklı renklerle temsil edilen kümelerin, belirli araştırma temalarına veya bilimsel alanlara göre gruplaşmalar oluşturduğu gözlemlenmiştir. En fazla atıf alan dergilerden biri olan Plant Physiology'nin, haritada merkezi bir konumda yer aldığı ve birçok dergiyle güçlü bağlantılar kurduğu tespit edilmiştir. Bu dergiyi Plant Journal, Plant Cell, Molecular Plant-Microbe Interactions ve Frontiers in Plant Science gibi yayınların izlediği anlaşılmıştır. Söz konusu dergilerin özellikle bitki fizyolojisi, moleküler biyoloji ve bitki-patojen etkileşimleri alanında yoğunlaştığı görülmüştür. Haritada yer alan diğer bir kümede, Phytopathology, Crop Protection, Acta Horticulturae ve Journal of Food Protection gibi dergilerin kırmızı renkle gruplaştığı ve daha çok bitki hastalıkları, koruma yöntemleri ve tarımsal uygulamalara odaklandığı belirlenmiştir. Scientific Reports, Plos One ve Frontiers in Microbiology gibi geniş kapsamlı dergilerin ise farklı kümeler arasında köprü niteliğinde olduğu tespit edilmiştir (Şekil 10).



Şekil 10. Dergilerin ağ grafiği analizi

FULL TEXTS BOOK

Ağ haritasında, dergiler arasındaki eş atıf ilişkileri ile bu ilişkilerin zamansal dağılımı görselleştirilmiştir. Renk kodlamasına göre, 2018 sonrası dönemde Scientific Reports, Frontiers in Plant Science ve Phytopathology gibi dergilerin daha güncel çalışmalarla ilişkilendirildiği ve literatürde daha fazla atıf aldığı belirlenmiştir (Şekil 11)




Şekil 11. Dergilerin ağ grafiği analizinin zaman içindeki güncel durumları

Şekil 12'deki üç bölmeli Sankey diyagramında, en çok üretim gerçekleştiren yazarlar (AU), bu yazarların ait olduğu ülkeler (AU_CO) ve çalışmalarda en sık kullanılan anahtar kelimeler (DE) arasındaki ilişkiler detaylı biçimde gösterilmiştir. Soldaki bölümde yer alan yazarların büyük bir kısmının Çin ve Amerika Birleşik Devletleri kökenli olduğu belirlenmiştir. Özellikle Jones Jb, Martin Gb, He Sy, Zhang Y, Wang Y ve Li J gibi yazarların, literatüre yoğun katkı sunduğu görülmektedir. Bu yazarların yayınlarının büyük ölçüde Çin ve ABD merkezli kurumlarda gerçekleştirildiği, dolayısıyla bu iki ülkenin konuya hâkimiyetinde öncü konumda bulunduğu anlaşılmaktadır.

Orta bölümde ülkelerin dağılımı yer almakta olup, en yoğun bağlantıların Çin ve ABD kaynaklı olduğu açıkça gözlemlenmektedir. Bu iki ülkeyi sırasıyla Japonya, İspanya, Kore, Almanya ve Kanada gibi ülkeler izlemektedir. Bu durum, araştırma alanında belirli ülkelerin bilgi üretiminde merkezî roller üstlendiğini göstermektedir (Şekil 12).


Sağdaki bölümde yer alan anahtar kelimeler incelendiğinde, en sık ilişkilendirilen konuların "pseudomonas syringae", "tomato", "arabidopsis", "disease resistance", "plant immunity" ve "ralstonia solanacearum" olduğu tespit edilmiştir. Çin kökenli çalışmaların özellikle "pseudomonas syringae" kavramı ile güçlü bir ilişki içerisinde olduğu görülmektedir. Bunun yanı sıra "bacterial pathogens", "biological control", "Xanthomonas" ve "salicylic acid" gibi daha spesifik terimlerle de çeşitli ülkeler arasında bağlantı kurulmuştur (Şekil 12).

FULL TEXTS BOOK



Şekil 12. Yazar - ülke ve anahtar kelime Sankey diyagramı


Araştırmalarda en sık birlikte kullanılan anahtar kelimeler arasındaki ilişkiler farklı renklerle gruplanan kümelerin, ortak tematik alanları temsil ettiği gözlemlenmiştir. "Tomato", "*Pseudomonas syringae*", "arabidopsis" ve "*Ralstonia solanacearum*" terimlerinin merkezde konumlandığı ve yoğun bağlantılarla birçok anahtar kelimeyle ilişkili olduğu belirlenmiştir. Ayrıca, "bacterial pathogens", "plant defense", "cell death" ve "induced resistance" gibi kavramların da ağda belirgin yer tuttuğu görülmüştür (Şekil 13). Genel olarak, anahtar kelime kullanımının zamanla belirli araştırma eksenlerinde yoğunlaştığı ve bilimsel odak noktalarının bu kümelenmeler üzerinden şekillendiği saptanmıştır.



Şekil 13. Anahtar kelimelerin ağ haritası

FULL TEXTS BOOK


Bu çalışmada, 2000–2023 yılları arasındaki literatürde öne çıkan anahtar kavramların yıllara göre eğilimleri ve kullanım sıklıkları analiz edilmiştir. Her bir terimin zaman ekseninde yer alan çizgisi aracılığıyla ilgili kavramın hangi yıllar arasında kullanıldığı belirlenmiş; çizgi üzerindeki dairelerin büyüklüğü ise bu kullanımın yoğunluk düzeyini yansıtacak şekilde gösterilmiştir (Şekil 14). En uzun süreyle ve en yüksek sıklıkla kullanılan terimler arasında *Pseudomonas syringae*, disease resistance, tomato, arabidopsis, hypersensitive response ve plant immunity gibi kavramların yer aldığı tespit edilmiştir. Bu durum, söz konusu terimlerin ilgili araştırma alanında temel kavramsal çerçeveyi oluşturduğunu göstermiştir. Son yıllarda ise bacteria, bacterial canker, pti, stomatal immunity, bak1 ve bacterial spot gibi daha spesifik ve moleküler düzeydeki terimlerin öne çıktığı gözlemlenmiştir. Bunun, araştırma eğilimlerinin daha genel başlıklardan moleküler mekanizmalara yöneldiğine işaret ettiği düşünülmektedir. Ayrıca jasmonic acid, cell death, priming ve induced resistance gibi bitki savunma yanıtlarıyla ilişkili terimlerin 2010 sonrası dönemde daha sık kullanıldığı belirlenmiştir. Genel olarak, yıllar içerisinde araştırma alanının hem terminolojik çeşitlilik hem de bilimsel derinlik açısından gelişim gösterdiği tespit edilmiştir.



Şekil 14. Anahtar kelimelerin trend durumları

Şekil 15'de 1953–2025 yılları arasında üç farklı dönem için anahtar kelime trendlerinin zaman içindeki değişimi görselleştirilmiştir. 1953–2010 döneminde ön plana çıkan "plant pathogenic bacteria", "arabidopsis", "salicylic acid" ve "biocontrol" gibi kavramların, 2011–2020 yıllarında "*Pseudomonas syringae*", "disease resistance" ve "effector-triggered immunity" gibi daha spesifik ve moleküler düzeydeki terimlere evrildiği gözlemlenmiştir. 2021–2025 döneminde ise "plant immunity", "*Pseudomonas syringae* pv. *tomato* dc3000", "induced systemic resistance" ve "stomatal immunity" gibi daha hedeflenmiş ve derinlemesine mekanizmaların öne çıktığı belirlenmiştir. Bu dönüşüm, araştırma alanında zamanla daha moleküler yönelik konulara yönelim olduğunu göstermiştir.

FULL TEXTS BOOK



Şekil 15. 1953–2025 yılları arasındaki anahtar kelimelerin dönemsel olarak nasıl değişimini ve tematik evrimini gösteren Sankey diyagramı.

SONUÇ VE ÖNERİLER

Bu çalışma, 1953–2025 yılları arasında domateste görülen bakteriyel hastalıklar üzerine yapılan 794 akademik yayını bibliyometrik yöntemlerle inceleyerek alandaki bilimsel gelişimi ve eğilimleri ortaya koymuştur. Yayın sayısının özellikle 2000 sonrası hızla arttığı, araştırmaların büyük oranda çok yazarlı ve uluslararası iş birliklerine dayandığı belirlenmiştir. ABD, Çin ve Japonya en üretken ülkeler olarak öne çıkarken, Türkiye'nin katkısının sınırlı kaldığı görülmüştür.

Araştırmalar zamanla daha moleküler konulara yönelmiş; özellikle *Pseudomonas syringae*, bitki bağışıklığı ve direnç mekanizmaları gibi kavramlar ön plana çıkmıştır. En fazla atıf alan dergiler arasında *Plant Physiology*, *Plant Journal* ve *Phytopathology* yer almıştır.

Türkiye'nin domateste görülen bakteriyel hastalıklar alanındaki bilimsel görünürlüğünü artırabilmesi için uluslararası iş birliklerini güçlendirmesi ve stratejik araştırma ortaklıkları kurması önem arz etmektedir. Ayrıca, araştırmaların moleküler temelli ve biyoteknolojik yaklaşımlar çerçevesinde derinleştirilmesi, yayın kalitesinin yükseltilmesi açısından gereklidir. Etki faktörü yüksek dergilerde yayın yapma hedefi benimsenmeli; bilim politikalarının belirlenmesinde ise bibliyometrik veriler etkin bir şekilde kullanılmalıdır.

KAYNAKÇA

Bergey, D. H., Harrison, F. C., Breed, R. S., Hammer, B. W., & Huntoon, F. M. (1923). *Bergey's manual of determinative bacteriology* (p. 116).

Davis, M. J., Gillaspie, A. G., Jr., Vidaver, A. K., & Harris, R. W. (1984). Clavibacter: A new genus containing some phytopathogenic coryneform bacteria, including *Clavibacter xyli* subsp. *xyli* sp. nov., subsp. nov. and *Clavibacter xyli* subsp. *cynodontis* subsp. nov., pathogens that cause ratoon stunting disease of sugarcane and bermudagrass stunting disease. *International Journal of Systematic Bacteriology, 34*(2), 107–117.

Deng, P., Shi, H., Pan, X., Liang, H., Wang, S., Wu, J., et al. (2022). Diyabetik ayak ülserleri üzerine dünya çapındaki araştırma eğilimleri (2004–2020): Araştırmacılar için öneriler. *Journal of Diabetes Research, 2022*, 7991031. https://doi.org/10.1155/2022/7991031

FULL TEXTS BOOK

FAO. (2023). *FAOSTAT: Crops and livestock products*. Food and Agriculture Organization of the United Nations. https://www.fao.org/faostat

Gardner, M. W., & Kendrick, J. B. (1923). Bacterial spot of tomato and pepper. *Phytopathology*, *13*, 307–315.

Hayward, A. C. (1991). Biology and epidemiology of bacterial wilt caused by *Pseudomonas solanacearum*. *Annual Review of Phytopathology*, *29*, 65–87. https://doi.org/10.1146/annurev.py.29.090191.000433

Hicks, D., Wouters, P., Waltman, L., de Rijcke, S., & Rafols, I. (2015). Bibliyometri: Araştırma ölçümleri için Leiden manifestosu. *Nature*, *520*(7548), 429–431. https://doi.org/10.1038/520429a

Jones, J. B., Jones, J. P., Stall, R. E., & Zitter, T. A. (2000). *Compendium of tomato diseases*. APS Press.

Jones, J. B., Vallad, G. E., Iriarte, F. B., Obradovic, A., Wernsing, M. H., Jackson, L. E., ... & Momol, M. T. (2014). Considerations for using bacteriophages for plant disease control. *Bacteriophage*, 4(4), e975029. https://doi.org/10.4161/21597081.2014.975029

Mansfield, J., Genin, S., Magori, S., Citovsky, V., Sriariyanum, M., Ronald, P., ... & Verdier, V. (2012). Top 10 plant pathogenic bacteria in molecular plant pathology. *Molecular Plant Pathology*, *13*(6), 614–629. https://doi.org/10.1111/j.1364-3703.2012.00804.x

McManus, P. S., Stockwell, V. O., Sundin, G. W., & Jones, A. L. (2002). Antibiotic use in plant agriculture. *Annual Review of Phytopathology*, *40*, 443–465. https://doi.org/10.1146/annurev.phyto.40.120301.093927

Ninkov, A., Frank, J. R., & Maggio, L. A. (2022). Bibliyometri: Akademik yayıncılığı inceleme yöntemleri. *Perspectives on Medical Education*, *11*(3), 173–176. https://doi.org/10.1007/s40037-021-00695-4

Scarlett, C. A., Fletcher, J. T., Roberts, P., & Lelliott, R. A. (1978). Tomato pith necrosis caused by *Pseudomonas corrugata*. *Annals of Applied Biology*, *88*, 105–114.

Wielke, J. P., Dye, D. W., & Watson, D. R. W. (1973). Further host of *Pseudomonas viridiflava*. *New Zealand Journal of Agricultural Research*, *16*, 315–323.

Wielke, J. P., & Dye, D. W. (1974). *Pseudomonas cichorii* causing tomato and celery diseases in New Zealand. *New Zealand Journal of Agricultural Research*, *17*, 123–130.

Yabuuchi, E., Kosako, Y., Yano, I., Hotta, H., & Nishiuchi, Y. (1995). Transfer of two *Burkholderia* and an *Alcaligenes* species to *Ralstonia* gen. nov.: Proposal of *Ralstonia pickettii* comb. nov., *Ralstonia solanacearum* comb. nov., and *Ralstonia eutropha* comb. nov. *Microbiology and Immunology*, *39*, 897–904.

Young, J. M., Dye, D. W., Bradbury, J. F., Panagopoulos, C. G., & Robb, C. F. (1978). A proposed nomenclature and classification for plant pathogenic bacteria. *Journal of Agricultural Research*, *21*, 153–177.

EGZERSİZ YAPAN YETİŞKİNLERDE ERGOJENİK DESTEK ÜRÜNLERİ KULLANIMI VE EGZERSİZ BAĞIMLILIĞI DURUMUNUN DEĞERLENDİRİLMESİ

EVALUATION OF ERGOGENIC SUPPLEMENT USE AND EXERCISE ADDICTION IN ADULTS ENGAGED IN EXERCISE

***Faruk Emre BOZEL*[1]**

[1]*Antalya Bilim Üniversitesi, Sağlık Bilimleri Fakültesi; Beslenme ve Diyetetik Bölümü, Antalya, TÜRKİYE*

[1]*ORCID ID: 0009-0001-8675-9688*

***Hale AVŞAR ERDEKLİ*[2]**

[2] *Dr Öğr. Üyesi, Antalya Bilim Üniversitesi, Sağlık Bilimleri Fakültesi; Beslenme ve Diyetetik Bölümü, Antalya, TÜRKİYE*

[2]*ORCID ID: 0009-0004-1492-8886*

ÖZET

Egzersiz fiziksel aktivitenin, belirli bir plan ve düzen içinde gerçekleştirilen bir alt dalıdır. Egzersiz bağımlılığı, kişinin kontrol edemediği bir biçimde yoğun ve sürekli egzersiz yapma arzusuyla karakterize edilir ve bu durum, özellikle spor performansını artırmaya yönelik ergojenik destek ürünlerinin kullanımını etkileyebilir. Araştırma egzersiz yapan bireylerin ergojenik destek ürünü kullanımı ve egzersiz bağımlılığı durumlarının değerlendirilmesi amacıyla yapılmıştır. Çalışma, Aralık 2024- Mart 2025 tarihleri arasında, özel bir spor salonunda üyeliği olan ve egzersiz yapan 19 kadın, 157 erkek olmak üzere toplam 176 kişi ile yürütülmüştür. Gönüllülük esasına dayanarak, çalışmaya onam vererek katılmayı kabul edenler çalışmaya dâhil edilmiştir. Bireylere yüz yüze görüşme tekniği ile anket formu uygulanmıştır. Anket formu; katılımcıların bireysel ve sosyodemografik özellikleri, beslenme alışkanlıkları, yapılan egzersiz türleri ve sıklığı, ergojenik destek ürünü kullanma durumuna yönelik sorulardan oluşmuştur. Katılımcıların egzersiz bağımlılığının değerlendirilmesinde Egzersiz Bağımlılığı Ölçeği (EBO) kullanılmıştır. Katılımcıların antropometrik ölçümleri alınmıştır. Katılımcıların %82.4'ünün ergojenik destek kullandığı saptanmıştır. Ergojenik destek ürünleri arasında en yaygın olarak tercih edilenin protein tozu olduğu gösterilmiştir. Katılımcıların %20.5'i protein tozunu ortalama 6 ay boyunca kullandıklarını belirtmişlerdir. Egzersiz bağımlılığı değerlendirldiğinde "Tolerans Gelişimi ve Tutku" boyutunda erkeklerin puan ortalamasının kadınlardan anlamlı derecede yüksek olduğu tespit edilmiştir. Genel "Egzersiz Bağımlılığı Ölçeği" puanının vücut yağ oranı ile negatif yönde anlamlı bir ilişki gösterdiği saptanmıştır. Katılımcıların egzersiz bağımlılığı ölçeği ile antropometrik ölçümleri arasındaki ilişkiler korelasyon analizi ile değerlendirilmiştir. Boy uzunluğu, kas kütlesi ve vücuttaki sıvı oranı arttıkça egzersize yönelik aşırı odaklanma ve duygu değişiminin de arttığını saptnamıştır. Çalışma, düzenli egzersiz yapan bireyler arasında ergojenik destek kullanımının yaygın olduğunu ve bu kişilerin genellikle protein tozu gibi ürünleri tercih ettiğini göstermektedir. Egzersiz bağımlılığı ile fiziksel özellikler arasında anlamlı ilişkiler bulunmuş, erkeklerin egzersize daha fazla tutku geliştirdiği tespit edilmiştir. Bu bulgular doğrultusunda, ergojenik desteklerin uzman kontrolünde kullanılması, uygun beslenme danışmanlığı sağlanması ve egzersiz bağımlılığına karşı farkındalık artırıcı çalışmalar önerilmektedir.

Anahtar Kelimeler: Egzersiz bağımlılığı, ergojenik destek ürünü, beslenme

FULL TEXTS BOOK

ABSTRACT

Exercise is a subcategory of physical activity that is performed in a structured and planned manner. Exercise addiction is characterized by an uncontrollable desire to engage in intense and continuous exercise, and this condition may influence the use of ergogenic aids, particularly those aimed at enhancing athletic performance. This study was conducted to evaluate the use of ergogenic aids and the level of exercise addiction among individuals who engage in regular exercise. The research was carried out between December 2024 and March 2025 with a total of 176 participants, including 19 women and 157 men, who were members of a private gym and actively engaged in exercise. Participation was based on voluntariness, and individuals who provided informed consent were included in the study. Data were collected through face-to-face interviews using a structured questionnaire. The questionnaire consisted of items addressing participants' individual and sociodemographic characteristics, dietary habits, types and frequency of exercise performed, and the use of ergogenic aids. The Exercise Dependence Scale (EDS) was used to assess exercise addiction among the participants. Anthropometric measurements were also obtained. It was determined that 82.4% of the participants reported using ergogenic aids. Among these products, protein powder was identified as the most commonly preferred ergogenic supplement. Additionally, 20.5% of the participants stated that they had used protein powder for an average duration of six months. In the evaluation of exercise addiction, it was found that the mean score of males was significantly higher than that of females in the 'Tolerance Development and Passion' subdimension. A significant negative correlation was observed between the overall Exercise Dependence Scale score and body fat percentage. The relationships between participants' scores on the Exercise Dependence Scale and their anthropometric measurements were evaluated using correlation analysis. It was observed that as height, muscle mass, and body water percentage increased, there was a corresponding increase in exercise-related overcommitment and mood alteration. The study demonstrates that the use of ergogenic aids is common among individuals who engage in regular exercise, with a preference for products such as protein powders. Significant associations were found between exercise addiction and physical characteristics, and it was observed that men tend to develop a greater passion for exercise. Based on these findings, it is recommended that ergogenic aids be used under professional supervision, appropriate nutritional counseling be provided, and awareness-raising initiatives be implemented to address exercise addiction.

Keywords: Exercise addiction, ergogenic supplement, nutrition

GİRİŞ

Egzersiz, planlı olarak yapılan fiziksel aktivite olarak tanımlanıp, fiziksel, zihinsel ve sosyal sağlık üzerinde olumlu etkiler yarattığı gösterilmiştir (1). Egzersiz bağımlılığı, bireyin aşırı derecede egzersiz yapması sonucu fiziksel ve psikolojik olarak zarar görmesi ve genel iyilik halinin olumsuz etkilenmesi durumudur. Egzersiz bağımlılığının başlıca nedeninin, beden imajı ve kilo kaygısı nedeniyle görüntüden duyulan memnuniyetsizlik olduğu öne sürülmektedir. (2). Egzersiz bağımlılığı konusunda Hausenblas ve Downs, "tolerans, geri çekilme, niyet etkisi, kontrol kaybı, zaman, diğer aktivitelerde azalma ve devamlılık olmak üzere yedi kriter belirlemişlerdir. İlk kriter olan tolerans, uygulanan egzersiz miktarının hedeflenen etkiye ulaşmak için giderek artırılması sonucunda egzersizin etkisinin azalması şeklinde tanımlanabilir. Geri çekilme ise, egzersizin durdurulması durumunda ortaya çıkabilecek belirtilerden kaçınmak amacıyla aynı düzeyde egzersiz yaparak rahatlama sağlamayı hedefler. Niyet etkisi, egzersizin beklenenden daha fazla miktar ve sürelerde gerçekleştirilmesi durumunu ifade eder. Kontrol kaybı kriteri ise, egzersizi bırakma çabalarının başarısız olması veya yoğun bir egzersiz yapma arzusunun ortaya çıkması şeklinde tanımlanmaktadır. Egzersize ayrılan fazla süre, Downs'ın zaman kriterini oluşturur. Egzersiz yapma amacı doğrultusunda bireyin yaşamındaki diğer aktivitelere ayırdığı

FULL TEXTS BOOK

zamanın önemli ölçüde kısıtlanması, diğer aktivitelerde bir azalma olarak tanımlanmaktadır. Son kriter olarak devamlılık, fiziksel bir sorun oluşturmasına rağmen, bu durumu göz önünde bulundurarak egzersiz yapmaya devam etmektir (3). Egzersiz yapan bireylerin sıklıkla tercih ettiği besin destek ürünleri, "ergojenik yardımcılar" ya da "supplement" olarak bilinir."Ergojenik" terimi, spor terminolojisinde enerji üretimini ve performansı artırarak sürdürülebilirliği sağlamak amacıyla kullanılan teknikleri tanımlar. Bu terim, "çalışma eğilimini artırma" anlamına gelir (4). Ergojenik destek ürünleri, performansı artırma, sağlığı koruma, enerji seviyelerini yükseltme ve bağışıklık sistemini destekleme gibi faktörler nedeniyle kullanılmaktadır. Bazı bireyler, ergojenik destek ürünlerini kullanmadan önce uzmanlarla danışmadan hareket etmekte ve optimal dozdan sapmaları durumunda potansiyel riskler hakkında yeterli bilgiye sahip olmamaktadır. Takviyelerin yan etkileri genellikle önerilen dozların aşılmasıyla ortaya çıkar (5). Rekabet hırsı nedeniyle birçok sporcu, önerilen dozların üzerinde takviye kullanmakta ve bu durum çeşitli hastalıklara yol açabilmektedir. Ek takviye kullanımı, alerjik reaksiyonlar ve zehirlenme gibi yaygın yan etkilere neden olabilir (6). Egzersiz bağımlılığı gibi aşırı davranışlar, spor performansını artırmaya yönelik ergojenik destek ürünlerinin kullanımını etkileyebilir (1). Bu çalışma egzersiz yapan bireylerin ergojenik destek ürünü kullanımı ve egzersiz bağımlılığı durumlarının değerlendirilmesi amacıyla yapılmıştır.

YÖNTEM

Araştırmanın Yeri, Zamanı ve Örneklem Seçimi

Bu araştırma Ocak – Mart 2025 tarihleri arasında Antalya İli Kepez İlçesinde bulunan özel bir spor kulübünde haftada en az 3 gün egzersiz yapan 176 kişi ile yürütülmüştür. Çalışmaya başlamadan önce bireylere "Gönüllü Onam Formu" okunarak, çalışmaya katılmak isteyip istemedikleri sorulmuştur. Çalışmaya gönüllü olarak katılmayı kabul eden bireyler çalışmaya dahil edilmiştir. Bu araştırma, Antalya Bilim Üniversitesi Sağlık Bilimleri Girişimsel Olmayan Araştırmalar Etik Kurulu Başkanlığının 2025/018 sayılı kararı ile onaylanmıştır.

Araştırmanın Genel Planı

Çalışmaya katılmak isteyen bireylere, çalışma hakkında sözlü bilgilendirme yapılmış ve yüz yüze görüşme yöntemi ile anket formu uygulanmıştır. Anket formunda, katılımcıların genel bilgileri (yaş, cinsiyet, çalışma şekli, medeni durum, eğitim durumu, gelir düzeyi, hastalık durumu, sigara ve alkol tüketimi, düzenli fiziksel aktivite durumu) sorgulanmıştır. Ayrıca, beslenme alışkanlıklarıyla ilgili olarak; (ana ve ara öğün tüketimi, öğün atlama sebepleri, tercih edilen yiyecekler, günlük su, çay ve kahve tüketimi, besin destek ürünleri kullanımı, bu ürünlerin tavsiye edilme durumu) soruları da sorulmuştur. Bununla birlikte katılımcıların antropometrik ölçümleri alınmıştır. Ayrıca, egzersiz bağımlılık düzeylerini ölçmek için Egzersiz Bağımlılığı Ölçeği (EBÖ) uygulanmıştır.

Antropometrik Ölçümler:

Katılımcıların antropometrik ölçümleri (ağırlık, yağ oranı ve kütlesi, kas oranı ve kütlesi, total vücut suyu (kg), bel ve kalça çevresi) ‘Tanita MC580’ marka vücut analiz cihazı ile ölçülmüştür. Ölçüm egzersiz öncesi yapılmıştır. Ölçüm öncesinde çay, kahve, kola, yeşil çay, vb. kafeinli içeceklerin içilmemesi, alkol tüketilmemesi, son öğünün en az 4 saat öncesinde tüketilmesi konusunda katılımcılara bilgi verilmiştir. Antropometrik ölçümlerden boy uzunluğu (cm), esnemeyen mezür yardımıyla ölçülmüştür. Araştırmaya katılan bireylerin boy uzunluğu ölçülürken katılımcının başı frankfort düzleminde, ayakkabısız, ayakları bitişik, topuklar, sırt, omuzlar ve başın arka kısmı duvara değecek şekilde durmasına dikkat edilmiştir.

FULL TEXTS BOOK

Egzersiz Bağımlılığı Ölçeği (EBO):

Katılımcıların egzersiz bağımlılığını ölçen 17 maddelik 'kesinlikle katılmıyorum' dan başlayıp 'kesinlikle katılıyorum' a kadar giden seçenekleri olan likert tipi ölçektir. Bu ölçek 'Aşırı Odaklanma ve Duygu Değişimi', 'Bireysel-Sosyal İhtiyaçların Ertelenmesi ve Çatışma' ve 'Tolerans Gelişimi ve Tutku' olmak üzere 3 alt boyuttan oluşmaktadır. Ölçekteki cevaplar, "kesinlikle katılmıyorum" 1, "katılmıyorum" 2, "kararsızım" 3, "katılıyorum" 4 ve "kesinlikle katılıyorum" 5 puandır. Egzersiz Bağımlılığı Ölçeği'nde,"1-17 puan normal, 18-34 puan az riskli, 35-51 puan riskli , 52-69 puan bağımlı , 70-85 puan yüksek düzeyde bağımlı" olarak değerlendirilmektedir. Ölçeğin geneli için Cronbach Alpha güvenirlik katsayısı 0,88 bulunurken, birinci boyut için 0,83; ikini boyut için 0,79; üçüncü boyut için 0,77 olarak bulunmuştur. Ölçeğin yapı geçerliliği için açımlayıcı (AFA) ve doğrulayıcı faktör analizi (DFA) uygulanmıştır. AFA sonucunda ölçek, 17 maddeden oluşmakta ve üç alt faktörlü bir yapı göstermektedir. Faktörlerin toplam açıkladığı varyans oranı % 54,61'dir. Buna göre, Egzersiz Bağımlılığı Ölçeği (EBÖ)'nin geçerli ve güvenilir bir ölçme aracı olduğu söylenebilir (7).

Verilerin İstatistiksel Olarak Değerlendirilmesi:

Çalışmada nicel veriler için tanımlayıcı istatistikler sayı (n), yüzde (%), ortalama ($\bar{x}$), standart sapma (ss), ve alt, üst değerleri olarak verilmiştir. Demografik veriler ve genel bilgiler için frekans alınmıştır. Verilerin normal dağılıma uygunluğu Kolmogorov Smirnov testi ile değerlendirilmiştir. Sayısal değişkenler arasındaki ilişkiler Çoklu Doğrusal Regresyon analizi ile değerlendirilmiştir. Verilerin istatistiksel olarak analiz edilmesinde Statistical Package for Social Sciences (SPSS) 24.0 yazılımı kullanılmıştır. İstatistik anlamlılık düzeyi $p<0,05$ kabul edilmiştir.

BULGULAR

Bireylerin cinsiyete göre demografik özelliklerinin dağılımı Tablo.1'de gösterilmiştir. Araştırmaya katılan 176 bireyin %89.2'si erkek, %10.8'i ise kadınlardan oluşmaktadır. Araştırmaya katılan bireylerin yaş ortalaması 26,95 ± 6,26 yıl olarak belirlenmiştir. Erkeklerin yaş ortalaması 26,85 ± 6,30, kadınların ise 27,74 ± 6,11'dir. Katılımcıların % 51,7' si öğrenci olup; %26,8' i serbest meslek ile uğraşmaktadır. Katılımcıların %86,7'sinin doktor tarafından tanısı konulmuş bir hastalığı bulunmamaktadır. En sık rastlanan sağlık sorunu ise göz hastalıklarıdır (%6,8). Katılımcıların %21,1'i ilaç kullanmaktadır. Alkol kullanan katılımcı oranı düşük olup, %93,7'i alkol kullanmamaktadır. Alkol kullananların tamamı erkektir.

Tablo 1. Bireylerin Cinsiyete Göre Demografik Özelliklerinin Dağılımı

Tanımlayıcı Özellikleri	Erkek (n=157)		Kadın (n=19)		Toplam (n=176)	
	n	%	n	%	n	%
Yaş (yıl) x±SS (yıl)	26.85±6.3		27.74±6.11		26.95±6.26	
Meslek						
Memur	17	10,9	3	15,8	20	11,3
İşçi	14	8,9	4	21,0	18	10,2
Serbest Meslek	41	26,1	6	31,6	47	26,8
Öğrenci	85	54,1	6	31,6	91	51,7
Medeni durumu						

FULL TEXTS BOOK

Evli	29	18,5	8	42,1	37	21,0
Bekar	127	80,9	11	57,8	138	78,4
Boşanmış/Eşi vefat etmiş	1	0,6	0	0.0	1	0,6
Eğitim Durumu						
İlkokul mezunu	1	0,6	0	0,0	1	0,5
Ortaokul mezunu	3	1,9	0	0,0	3	1,7
Lise mezunu	56	35,7	6	31,6	62	35,3
Lisans (üniversite) mezunu	97	61,8	13	68,4	110	62,5
Gelir Düzeyi						
Gelirim giderimden fazla	98	62,5	10	52,6	108	61,3
Gelirim giderime eşit	36	22,9	7	36,8	43	24,4
Gelirim giderimden az	23	14,6	2	10,6	25	14,3
Doktor Tanısı Konulmuş Hastalık						
Hayır	136	86,7	17	89,4	153	86,9
Kalp ve damar hastalıkları	1	0,6	0	0.0	1	0,6
Ülser/gastrit/reflü	1	0,6	0	0.0	1	0,6
Böbrek Hastalıkları	1	0,6	0	0.0	1	0,6
Göz	11	7,0	1	5,3	12	6,8
Bağırsak hastalıkları	1	0,6	0	0.0	1	0,6
Alerji/Astım	6	3,9	1	5,3	7	3,9
Ailede Tanısı Konmuş Böbrek yada Karaciğer Rahatsızlığı						
Yok	129	82,2	16	84,3	145	82,4
Var	28	17,8	3	15,7	31	17,6
İlaç Kullanımı						
Yok	123	78,4	16	84,3	139	78,9
Var	34	21,6	3	15,7	37	21,1
Sigara Kullanımı						
Evet	74	47,2	10	52,6	84	47,8

FULL TEXTS BOOK

Hayır	83	52,8	9	47,4	92	52,2
Alkol Tüketimi						
Evet	11	7,1	0	0.0	11	6,3
Hayır	146	92,9	19	100	165	93,7

Bireylerin egzersiz durumlarının değerlendirilmesi Tablo.2'de gösterilmiştir. Katılımcıların büyük çoğunluğunun (%61,9) haftada 4 kez egzersiz yaptığı belirlenmiştir. Haftada 5 kez egzersiz yapan katılımcıların oranı %26,2 iken, haftada 3 kez egzersiz yapanların oranı ise %10,2 olarak tespit edilmiştir. Buna karşın katılımcıların yalnızca %1,7'si her gün düzenli egzersiz yapmaktadır. Egzersiz türü açısından yapılan değerlendirmede, katılımcılar arasında en sık tercih edilen egzersiz türünün ağırlık antrenmanı olduğu belirlenmiştir (%65,3). Fitness, ikinci en çok tercih edilen egzersiz türü olup, katılımcıların %26,2'sı tarafından tercih edilmektedir. Bunun yanı sıra; kardiyo egzersizleri yapan katılımcılar %4,5, pilates yapanlar %3,4 ve CrossFit yapanlar ise yalnızca %0,6 oranındadır. Bu sonuçlar, araştırmaya katılan bireylerin daha çok kuvvet ve dayanıklılığa yönelik egzersiz türlerini tercih ettiklerini göstermektedir.

Tablo 2. Bireylerin Egzersiz Durumlarının Değerlendirilmesi

Tanımlayıcı Özellikleri	**Toplam**	
	n	**%**
Haftada 3 kez	18	10,2
Haftada 4 kez	109	61,9
Haftada 5 kez	46	26,2
Her gün	3	1,7
Hangi Egzersiz Türü		
CrossFit	1	0,6
Pilates	6	3,4
Cardio	8	4,5
Ağırlık Antrenmanı	115	65,3
Fitness	46	26,2

Tablo.3' te katılımcıların besin desteği kullanma durumlarının cinsiyete göre dağılımları gösterilmiştir. Araştırmaya katılan erkeklerin %87,9; kadınların ise %47,4' ü ergojenik destek kullanmaktadır. Erkeklerin %50,7' sine ergojenik destek diyetisyen tarafından önerilmiştir. %16,7'si ise kendi kararı ile ergojenik destek kullanmıştır. Katılımcıların %65,9' u düzeni olarak kan testi yaptırmaktadır. Ergojenik destek kullanan bireylerin %64,6' sında ergojenik destek kullanımı ile ilgili bir yan etki ortaya çıkmıştır.

FULL TEXTS BOOK

Tablo 3. Bireylerin Besin Desteği Kullanım Durumlarının Cinsiyete Göre Dağılımı

Tanımlayıcı Özellikleri	Erkek (n=157)		Kadın (n=19)		Toplam (n=176)	
	n	%	n	%	n	%
Ergojenik Desteğin Kullanımı						
Evet	138	87,9	9	47,4	147	83,5
Hayır	19	12,1	10	52,6	29	16,5
Ergojenik Desteğin Önerilmesi						
Kendi Kararım	23	16,7	0	0	23	15,7
Diyetisyen	70	50,7	7	77,8	77	52,4
Antrenör	45	32,6	2	22,2	47	31,9
Kan Testi Durumu						
Hayır	53	33,8	7	36,8	60	34,1
Evet	104	66,2	12	63,2	116	65,9
Yan Etki Durumu						
Evet	91	66,0	4	44,4	95	64,6
Hayır	47	44,0	5	55,6	52	35,4

Tablo 4'e göre; araştırmada kullanılan ölçek puanlarının cinsiyet değişkenine göre karşılaştırılmasına yönelik gerçekleştirilen bağımsız gruplar t-testi sonuçları incelendiğinde; "Aşırı Odaklanma ve Duygu Değişimi" boyutunda erkek katılımcıların puan ortalaması (X=3,66±0,84), kadın katılımcılara (X=3,38±0,86) kıyasla daha yüksek olmasına rağmen, bu fark istatistiksel açıdan anlamlı bulunmamıştır (t(174)=1,395; p=0,165). Bu durum, aşırı odaklanma ve duygu değişiminin cinsiyete göre önemli düzeyde farklılaşmadığını ortaya koymaktadır. "Tolerans Gelişimi ve Tutku" boyutunda ise erkeklerin puan ortalamasının (X=3,45±1,14) kadınlardan (X=2,88±1,27) anlamlı derecede yüksek olduğu tespit edilmiştir (t(174)=2,026; p=0,044). Bu durum, erkek katılımcıların egzersize yönelik tutku geliştirme ve tolerans düzeylerinin kadınlardan anlamlı şekilde daha yüksek olduğunu göstermektedir.

FULL TEXTS BOOK

Tablo.4 Bireylerin Cinsiyete Göre Egzersiz Bağımlılığının Karşılaştırılması

	Cinsiyet	n	Ortalama	S.S.	t	df	P
Aşırı Odaklanma ve Duygu Değişimi	Erkek	157	3.6624	.84347	1.395	174	.165
	Kadın	19	3.3759	.86269			
Bireysel-Sosyal İhtiyaçların Ertelenmesi ve Çatışma	Erkek	157	2.6539	.87514	-1.106	29.874	.278
	Kadın	19	2.8158	.56079			
Tolerans Gelişimi ve Tutku	Erkek	157	3.4490	1.13924	2.026	174	.044*
	Kadın	19	2.8816	1.26757			
Egzersiz Bağımlılığı Ölçeği	Erkek	157	3.2563	.74138	1.109	174	.269
	Kadın	19	3.0619	.52043			

Tablo 5'e göre, katılımcıların egzersiz bağımlılığı ölçeği ile antropometrik ölçümleri arasındaki ilişkiler korelasyon analizi ile incelenmiştir. "Aşırı Odaklanma ve Duygu Değişimi" boyutu, boy uzunluğu (r=0,208; p<0,01), toplam vücut suyu yüzdesi (r=0,300; p<0,01) ve vücut kas kütlesi (r=0,240; p<0,05) ile pozitif yönde anlamlı ilişkiler göstermektedir. Bu, bu ölçümlerin artmasıyla egzersize yönelik aşırı odaklanma ve duygu değişiminin de arttığını göstermektedir. Ayrıca, bu boyutun vücut yağ oranı ile negatif yönde anlamlı bir ilişkiye sahip olduğu (r=-0,297; p<0,01) belirlenmiştir; yani vücut yağ oranı arttıkça aşırı odaklanma ve duygu değişimi azalmaktadır. "Tolerans Gelişimi ve Tutku" boyutunun, katılımcıların boy uzunlukları (r=0,176; p<0,05) ve Toplam vücut suyu yüzdesi (r=0,222; p<0,05) ile pozitif, vücut yağ oranı ile ise negatif yönde anlamlı ilişkiye sahip olduğu belirlenmiştir (r=-0,329; p<0,01). Buna göre, boy uzunluğu ve vücuttaki su oranı arttıkça egzersize yönelik tutku ve tolerans düzeyi artmakta; vücut yağ oranı arttıkça ise azalmaktadır. Genel "Egzersiz Bağımlılığı Ölçeği" puanının ise vücut yağ oranı ile negatif yönde anlamlı bir ilişki gösterdiği saptanmıştır (r=-0,332; p<0,01). Bu bulgu, genel egzersiz bağımlılığı düzeyinin vücut yağ oranının artışı ile azaldığını ortaya koymaktadır.

Tablo 5. Bireylerin Antropometrik Ölçümlerine Göre Egzersiz Bağımlılığının Korelasyon Analizi ile Değerlendirilmesi

Aşırı Odaklanma ve Duygu Değişimi	1	.178*	.703**	.824**	.208*	.052	.036	.066	-.195	-.297**	.300**	.240*
Bireysel-Sosyal İhtiyaçların Ertelenmesi ve Çatışma		1	.389**	.648**	-.174*	-.069	-.115	-.060	-.218	-,189	-,069	-.105
Tolerans Gelişimi ve Tutku			1	.880**	.176*	.050	-.012	.029	-.220	-,329**	.222*	.151

Egzersiz Bağımlılığı Ölçeği	1	,095	.013	-.035	.018	-.239	-,332**	.187	.125
Boy (cm)		1	.160	.389**	.272**	.022	-.062	.685**	.502**
BKİ (kg/m2)			1	.302**	.342**	.074	-.075	.513**	.224
Bel Çevresi (cm)				1	.718**	.775**	.494**	.168	.380**
Kalça Çevresi (cm)					1	-.428**	.192*	.011	.256**
Bel/Kalça Oranı						1	.428**	-.126	-.162
Vücut Yağ Oranı (%)							1	-.319**	-.188
Toplam Vücut Suyu (%)								1	.339**
Vücut Kas Kütlesi (kg)									1

TARTIŞMA VE SONUÇ

Bireylerin fiziksel aktivite sıklığı açısından incelendiğinde, katılımcıların büyük çoğunluğunun (%61.6) haftada 4 kez egzersiz yaptığı belirlenmiştir. Haftada 5 kez egzersiz yapan katılımcıların oranı %26.0 iken, haftada 3 kez egzersiz yapanların oranı ise %10.2 olarak tespit edilmiştir. Buna karşın katılımcıların yalnızca %1.7'si her gün düzenli egzersiz yapmaktadır. Egzersiz türü açısından yapılan incelemede, katılımcılar arasında en sık tercih edilen egzersiz türünün güç antrenmanı olduğu belirlenmiştir (%65.0). Yapılan başka bir araştırmada düzenli spor salonu kullanıcılarının %41'i haftada 4-5 kez egzersiz yaptıklarını bildirmiştir (8). Bu dağılım, bizim çalışmamızdaki %61.6 ile haftada 4 kez ve %26.0 ile haftada 5 kez egzersiz yapan katılımcı oranlarına oldukça yakındır. Yapılan bu araştırmada katılımcıların %61'i güç antrenmanı yaparken, %26'sı dayanıklılık egzersizlerini tercih etmiş; bizim örneklemimizdeki %65'lik ağırlık antrenmanı tercih oranı ile paralellik göstermektedir (8). Bu veriler, katılımcıların fiziksel aktivite sıklığına dair farklı örneklemlerden elde edilen sonuçları karşılaştırmaktadır. Her iki araştırma da katılımcıların egzersiz sıklığının önemli ölçüde değişkenlik gösterdiğini, ancak belirli bir egzersiz sıklığının daha yaygın olduğunu ortaya koymaktadır.

Araştırmada cinsiyet değişkenine göre ölçek puanlarının karşılaştırılması için yapılan bağımsız gruplar t-testi sonuçlarına göre "Tolerans Gelişimi ve Tutku" boyutunda ise erkeklerin puan ortalamasının (X=3,45±1,14) kadınlardan (X=2,88±1,27) anlamlı derecede yüksek olduğu tespit edilmiştir (t(174)=2,026; p=0,044). Bu durum, erkek katılımcıların egzersize yönelik tutku geliştirme ve tolerans düzeylerinin kadınlardan anlamlı şekilde daha yüksek olduğunu göstermektedir. Cinsiyet değişkeniyle egzersiz bağımlılığı ölçeği alt boyutlarından "tolerans gelişimi ve tutku" arasında anlamlı bir fark bulunmuştur (p<0,05). Erkek öğrenciler, bu alt boyutta kadınlara göre daha yüksek puanlar elde etmiştir.

FULL TEXTS BOOK

Yapılan diğer bir araştırmada, erkek sporcuların egzersiz bağımlılığı puanlarının diğer katılımcılara göre anlamlı şekilde daha yüksek olduğu bulunmuştur (9).

Katılımcıların egzersiz bağımlılığı ölçeği ile antropometrik ölçümleri arasındaki ilişkiler incelenmiştir. "Aşırı Odaklanma ve Duygu Değişimi" boyutu, boy uzunluğu, toplam vücut suyu yüzdesi ve vücut kas kütlesi ile pozitif ilişkiler gösterirken, vücut yağ oranı ile negatif bir ilişki sergilemektedir. "Tolerans Gelişimi ve Tutku" boyutu da benzer şekilde boy uzunluğu ve toplam vücut suyu yüzdesi ile pozitif, vücut yağ oranı ile negatif ilişkilere sahiptir. Bu bulgular, vücut ölçümlerinin artmasının egzersize yönelik aşırı odaklanma ve tutku ile ilişkili olduğunu göstermektedir. Yapılan bir diğer çalışma sonucu katılımcıların vücut ağırlığı ile egzersiz bağımlılığı puan ortalaması arasında anlamlı bir fark olmadığını, ancak vücut ağırlığı arttıkça egzersiz ihtiyacının yükseldiğini tespit etmişlerdir (10). Egzersiz bağımlılığının değerlendirildiği başka bir araştırmada ise egzersiz bağımlılığı alt boyutları (tolerans, çatışma, ihtiyaç erteleme vb.) ile "sağlıklı beslenme tutumları" arasında r = 0,21–0,30 arası zayıf-orta düzey pozitif korelasyon olduğu gösterilmiştir ($p<0,01$) (11).

Brezilya' da yapılan bir araştırmada spor salonuna düzenli olarak giden bireylerde egzersiz bağımlılığ riski ile takviye kullanmı arasında güçlü bir ilişki olduğu saptanmıştır. Çalışmada takviye kullanma durumunun egzersiz bağımlılığı olanlarda olmayanlara göre 4,5 kat daha yüksek olduğu belirlenmiştir. Ayrıca katılımcıların %51,5' i son bir yılda ergojenik takviye kullandıklarının bildirmişlerdir (12).

Yapılan kesitsel bir çalışmada sağlık profesyoneli danışmanlığı olmaksızın ergojenik takviye kullanımının yaygın olduğu saptanmıştır. Ayrıca takviye kullanan bireylerde hem Egzersiz Bağımlılığı Envanteri hem de Görünüm Kaygısı puanlarının daha yüksek olduğu gösterilmiştir. daha yüksektir. Ergojenik ürün kullanımı ile egzersiz bağımlılığı düzeyi arasında anlamlı bir ilişki olduğu belirlenmiştir (13).

COVID-19 pandemisi sırasında ergojenik ürünlerin kullanımının değerlendirildiği bir çalışmada yetişkinlerin %28'inin görünüm ve performans artırıcı ürünler kullandığı belirlenmiştir (14).

Portekiz'deki bir spor salonu üyeleri arasında yapılan çalışmada katılımcıların takviye kullanımının %43,8 olduğu tespit edilmiştir. Ayrıca ergojenik destek kullanımının erkekler arasında daha yaygın olduğu gösterilmiştir. En sık kullanılan ergojenik destek ürünlerinin proteinler, BCAA'lar ve multivitamin/mineraller olduğu saptanmıştır (15).

Yapılan kesitsel bir çalışmada, spor salonu kullanıcılarının %64,7'sinin ergojenik destek ürünleri kullandığı belirlenmiştir. Ergojenik destek kullanma nedenleri araştırıldığında ise daha uzun süre ve daha yoğun egzersiz yaptığını algılayanlar ile 'ideal kilonun altında' olduğunu düşünenlerde kullanım durumunun daha yüksek olduğu tespit edilmiştir (16).

Fitness merkezine giden bireyler arasında yapılan diğer bir çalışmada egzersiz bağımlılığı riski yüksek bireylerin kafein, protein tozu gibi performans artırıcı besin takviyesi kullanımının daha fazla olduğu belirlenmiştir. Ayrıca yalnız egzersiz yapanlar ve rekabetçi sporcular arasında bağımlılık ve tutku puanlarının daha yüksek olduğu tespit edilmiştir (17).

Spor bilimleri öğrencileri arasında yapılan çalışmada, "Bireysel-Sosyal İhtiyaçların Ertelenmesi ve Çatışma" alt boyutu ile "Tolerans Gelişimi ve Tutku" boyutunun, sporcu takviyesi inancını ile anlamlı ve pozitif şekilde ilişki olduğu gösterilmiştir (18).

Taekwondo sporu ile ulusal olarak görev alan atletler arasında yapılan bir araştırmada, 'sağlıklı beslenmeye yönelik olumlu tutumlar' ile egzersiz bağımlılığı farkındalığı arasında anlamlı bir ilişki olduğu saptanmıştır. Bu durum, beslenme bilincinin sporcuların bağımlılık riskini fark etmesine katkı sağlayabileceğini göstermektedir (19).

FULL TEXTS BOOK

Yapılan başka bir çalışmada egzersiz yapan genç yetişkinlerde diyet modelleri ile egzersiz bağımlılığı ve yeme bozuklukları arasındaki ilişki incelenmiştir. Bu çalışmada, sağlıksız beslenme kalıplarının egzersiz bağımlılığını ve yeme bozukluğunu anlamlı şekilde etkileyebileceği gözlemlenmiştir (20).

Egzersiz bağımlılığı, bireylerin performans artırıcı desteklere yönelimini artırarak kontrolsüz ve bilinçsiz kullanımı beraberinde getirebilir. Bu durum, hem fiziksel sağlık hem de psikolojik iyi oluş üzerinde olumsuz sonuçlar doğurma potansiyeline sahiptir. Ergojenik desteklerin uzman denetimi olmadan kullanımı, yanlış doz alımı, yan etkiler ve uzun vadeli sağlık riskleri açısından önemli bir tehdit oluşturmaktadır. Sağlık profesyonellerinin danışmanlığı, bireylerin gereksinimlerine uygun ve güvenli kullanım alışkanlıkları kazandırılmasında kritik bir rol üstlenmektedir. Bu bağlamda, egzersiz bağımlılığı riski taşıyan bireylerin özellikle yakından izlenmesi ve bilinçlendirilmesi gerekmektedir. Ayrıca, multidisipliner yaklaşımlar ile sporcuların ve egzersiz yapan bireylerin beslenme, psikolojik destek ve egzersiz programları açısından bütüncül bir şekilde değerlendirilmesi önem taşımaktadır. Sonuç olarak, ergojenik destek kullanımı ve egzersiz bağımlılığı arasındaki ilişki göz önünde bulundurularak, sağlık profesyonelleri eşliğinde yapılan bilinçli uygulamalar bireylerin performansını güvenli şekilde artırabilir ve olası sağlık risklerini en aza indirebilir.

KAYNAKLAR

1. Ardiç, F. (2014). Egzersizin sağlık yararları. Türk Fiz Tıp Rehab Derg, 60 (Özel Sayı 2), 9–14.
2. Cicioğlu Hİ, Tekkurşun Demir G, Bulğay C, Çetin E. Elit Düzeyde Sporcular ile Spor Bilimleri Fakültesi Öğrencilerinin Egzersiz Bağımlılığı Düzeyleri. Bağımlılık Dergisi. Mart 2019;20(1):12-20.
3. Hausenblas, HA ve Downs, DS (2002). Egzersiz bağımlılığı: Sistematik bir inceleme. Spor ve Egzersiz Psikolojisi, 3 (2), 89–123.
4. Sezer, D., & Şahin, M. (2023). Sporda Supplement Ve Ergojenik Yardımcıların Kullanımının Etik Açıdan İncelenmesi. Avrasya Sosyal Ve Ekonomi Araştırmaları Dergisi, 10(3), 170-185.
5. Çetin, E., Ertaş Dölek, B., & Orhan, Ö. (2008). Gazi Üniversitesi Beden Eğitimi Ve Spor Yüksekokulu Öğrencilerinin Ergojenik Yardımcılar, Doping Ve Sağlık Hakkındaki Bilgi Ve Alışkanlıklarının Belirlenmesi. Spormetre Beden Eğitimi Ve Spor Bilimleri Dergisi, 6(3), 129-132.
6. Hall, Matthew DO; Trojian, Thomas H. MD, FACSM. Kreatin Takviyesi. Güncel Spor Hekimliği Raporları 12(4):s 240-244, Temmuz/Ağustos 2013
7. Tekkurşun Demir, G., Hazar, Z., & Cicioğlu, H. İ. (2018). *Egzersiz bağımlılığı ölçeği (EBÖ): Geçerlik ve güvenirlik çalışması*. Kastamonu Eğitim Dergisi, 26(3), 865–874.
8. Kreider, R. B., Ferreira, M. P., Greenwood, M., Wilson, M., & Almada, A. L. (2002). Effects of conjugated linoleic acid supplementation during resistance training on body composition, bone density, strength, and selected hematological markers. The Journal of Strength & Conditioning Research, 16(3), 325-334.
9. Şemşek, Ö., Yüktaşır, B., & Şemşek, S. (2010). Ergojenik Yardımcı Olarak Kullanılan Besin Suplementleri. Beden Eğitimi Ve Spor Bilimleri Dergisi, 1(3).
10. Özdenk, S. (2018, Nisan). Beden eğitimi ve spor, faydaları, önemi ve sınıflandırılması [Physical education and sports: Benefits, importance, and classification] (Bölüm 8). In Spor Bilimlerinde Akademik Araştırmalar (ss. 77–89). Sinop Üniversitesi.Orer, G. E., & Guzel, N. A. (2014). The effects of acute L-carnitine supplementation on endurance performance of athletes. Journal of strength and conditioning research, 28(2), 514–519.
11. Ersoy, G. (2010), Egzersiz ve Spor Performansı için Beslenme, Ankara: Betik kitap.
12. Rossi, L., & Tirapegui, J. (2016). Exercise dependence and its relationship with supplementation at gyms in Brazil. Nutrición Hospitalaria, 33(2), 431–436.
13. Corazza, O., Simonato, P., Demetrovics, Z., Mooney, R., van de Ven, K., Roman-Urrestarazu, A., ... & Martinotti, G. (2019). The emergence of Exercise Addiction, Body Dysmorphic Disorder, and

FULL TEXTS BOOK

other image-related psychopathological correlates in fitness settings: A cross-sectional study. PLOS ONE, 14(4), e0213060.

14. Dores, A. R., Carvalho, I. P., Burkauskas, J., Simonato, P., De Luca, I., Mooney, R., ... & Corazza, O. (2021). Exercise and use of enhancement drugs at the time of the COVID-19 pandemic: A multicultural study. Frontiers in Psychiatry, 12, 648501.

15. Ruano, J., & Teixeira, V. H. (2020). Prevalence of dietary supplement use by gym members in Portugal and associated factors. Journal of the International Society of Sports Nutrition, 17(11).

16. Lacerda, F. M. M., Carvalho, W. R. G., Hortegal, E. V., Cabral, N. A., & Veloso, H. J. (2015). Factors associated with dietary supplement use by people who exercise at gyms. Revista de Saúde Pública, 49, 63.

17. Lichtenstein, M. B., Jensen, E. S., & Szabo, A. (2020). Exercise addiction, obsessive passion and the use of nutritional supplements in fitness center attendees. Translational Sports Medicine, 3(3), 188–195.

18. Yurdakul, H. Ö. (2020). The Relationship Between Exercise Addiction and Beliefs in Sports Nutritional Supplements. Türk Spor ve Egzersiz Dergisi, 22(2), 338–343.

19. Tolukan, E., Güler, H., Yıldız, A. B., Çiçek, G., Yalçın, İ., Bayram, A., & Ekinci, N. E. (2024). The relationship between attitudes towards healthy nutrition and awareness levels of exercise addiction in male taekwondo athletes. Journal of Men's Health, 20(12), 50–56.

20. Khoshro, S., & Abbasalizad Farhangi, M. (2024). Major dietary patterns, exercise addiction, and eating disorders among a sample of physically active young adults. Nutrition and Metabolic Insights, 17, Article 11786388241258938.

BEYAZ KAN HÜCRELERİNİN TESPİTİ VE SINIFLANDIRILMASI - BİR İNCELEME
DETECTION AND CLASSIFICATION OF WHITE BLOOD CELLS – A REVIEW

Elif İrem AKDOĞAN

Karabük Üniversitesi, Mühendislik Fakültesi,

Biyomedikal Mühendisliği, Karabük, Türkiye.

Doç. Dr. Eftal ŞEHİRLİ

Karabük Üniversitesi, Mühendislik Fakültesi,

Bilgisayar Mühendisliği, Karabük, Türkiye.

ÖZET

Gelişen Dünya teknolojisi ve sağlık alanında gerçekleşleşen inovatif çalışmalar; beyaz kan hücresi (BKH) tespiti, segmentasyonu ve sınıflandırılması kapsamında, derin öğrenme yöntemleri ile yapay zekâ tabanlı yöntemlerin entegrasyonunu ortaya koymaktadır. BKH'nin hızlı ve doğru şekilde tespit edilmesi ve sınıflandırılması, geçmişten günümüze tanı süreci karmaşık olan kanser, lösemi ve enfeksiyonel hastalıkların erken teşhisinde hekime yol gösterici nitelik taşımaktadır. Geleneksel görüntü işleme ve manuel analiz yöntemlerinin yanında, son yıllarda makine öğrenmesi ve derin öğrenme temelli yaklaşımların bu alandaki kullanımı dikkat çekici oranda artış göstermiştir. Bu çalışmada, son 10 yıl içinde yayımlanan bilimsel yazılar incelenerek BKH tespiti, segmentasyonu ve sınıflandırılması alanındaki bilimsel gelişmeler detaylı bir şekilde ele alınmıştır. Yapılan literatür çalışması sonucunda, konvolüsyonel sinir ağları (CNN), U-Net ve You Only Look Once (YOLO) modellerinin yaygınlaştığı; veri seti çeşitliliği, klinik uyarlanabilirlik, doğruluk oranları konusunda önemli ilerlemeler kaydedildiği tespit edilmiştir. Mevcut yöntemlerin sınırlı olması ve klinik entegrasyon sürecinde karşılaşılan zorluklar da kapsamlı şekilde değerlendirilmiştir. Çalışma kapsamında ayrıca hibrit modellerin kullanım potansiyeli, veri arttırma stratejileri ve yapay zekâ ile karar destekli sistemlerin gelecekteki rolü üzerine öneriler sunulmuştur. Bu alandaki yenilikçi algoritmaların klinik pratikte uygulanabilirliğinin artırılması, hasta sonuçlarını doğrudan etkileyebilecek önemli bir gelişme olarak değerlendirilmektedir. Literatürdeki son eğilimler doğrultusunda daha güvenilir ve hassas, ölçeklenebilir sistemlerin geliştirilmesi ön görülmektedir. Bu derleme ile güncel yaklaşımların karşılaştırılması ve gelecek çalışmalara yön verilmesi amaçlanmaktadır.

Anahtar kelimeler: Beyaz Kan Hücreleri, Derin öğrenme, Filtreleme, Makine Öğrenmesi, Sınıflandırma.

ABSTRACT

Innovative studies in the developing world of technology and healthcare are demonstrating the integration of deep learning and artificial intelligence-based methods within the scope of white blood cell (WBC) detection, segmentation, and classification. Rapid and accurate detection and classification of WBC can guide physicians in early diagnosis of cancer, loss, and disease, which used to be a complex process for caries diagnosis. In addition to traditional image processing and manual analysis methods, the use of machine learning and deep learning in this field has shown a remarkable increase in recent years. This study examines scientific papers published in the last decade, providing a detailed overview of scientific advances in the field of WBC detection, segmentation, and classification. The literature review reveals that the models such as convolutional neural networks (CNN), U-Net, and You Only Look Once (YOLO) have become widespread, and significant advances have been made in dataset

FULL TEXTS BOOK

diversity, clinical adaptability, and accuracy. The limitations of existing methods and the challenges encountered in clinical integration are comprehensively evaluated. Frthermore, the study offers suggestions on the potential use of hybrid models, data augmentation strategies, and the future role of artificial intelligence and decision-supported systems. Increasing the applicability of innovative algorithms in this field in clinical practice is considered a significant development that can directly impact patient outcomes. In line with recent trends in literature, the development of more reliable, sensitive, and scalable systems is anticipated. This review aims to compare current approaches and guide future research.

Keywords: Classifications, Deep Learning, Filtering, Machine Learning, White Blood Cells.

GİRİŞ

Beyaz kan hücreleri (WBC), bağışıklık sisteminin bir parçası olan ve vücudu enfeksiyonel virüs saldırılarına karşı korumakla görevli olan savunma mekanizmalarıdır. Kan ve lenf dokularında, kemik iliği tarafından üretimi gerçekleştirilmektedir. İnsan kanı; kırmızı kan hücreleri (RBC), WBC, platelet ve plazma olmak üzere 4 ana kısımdan meydana gelmektedir. Lökosit olarak isimlendirilen WBC bulaşıcı hastalıklar, virüsler, mantarlar ve mikroplara karşı vücudu korumaya çalıştığı için bu gruplandırmada önemli bir alanı temsil etmektedir (Özcan, 2022). WBC, sitoplazma içerikleri ve savunma mekanizmasında üstlendiği farklı görevlere göre beş çeşidi mevcuttur. Bazofil, eozinofil, lenfosit, monosit ve nötrofil olarak sınıflandırabilen WBC çeşitleri; farklı hastalıklara bağlı olarak vücut içi miktarlarında değişimlere mağruz kalabilmektedir. Hastalıkların teşhisinde bir parametre görevini üstlenen WBC, yetişkin bir birey kanında belirlenen düzeyin altında veya üstünde olması durumunda, çeşitli hastalıklara yol açabilmektedir. WBC fazlalığı durumuna lökositoz, az olma durumuna lökopeni ismi verilmektedir (Wijesinghe vd., 2020). Tespit edilen düzey farklılıklarına bağlı olarak hekim tarafından teşhis konularak, bireyin bağışıklık sistemi üzerinde iyileştirme çalışmaları ve gerekli testler yapılmaktadır. WBC tespitine bağlı görülen hastalıkların Dünya çapında bireysel ve toplumsal etkileri beraberinde getirdiği gözlemlenmektedir (Tablo 1). WBC, tespiti ve sınıflandırılması, bilim insanları ve veri uzmanları tarafından son yıllarda incelemesi gerçekleştirilen önemli bir başlık olarak nitelendirilmektedir. Günümüz teknolojisinde WBC sınıflandırması, bölütlemesi ve tespitinde makine öğrenmesi, derin öğrenme modellerinin tercih edildiği görülmektedir. Farklı tekniklerin oluşturulduğu çalışmalarda derin öğrenme model kullanımının fazlalığı dikkat çekmektedir

Tablo 1. WBC Miktarına Bağlı Hastalıklar.

Beyaz Hücre Tipi	Fazlalık Durumu	Eksiklik Durumu
Lenfosit	Bakteriyel enfeksiyonlar, Belirli lenfomalar, Kanser	HIV, Yetersiz beslenme, Otoimmün bozukluklar
Nötrofil	Enfeksiyon, İltihaplanma, Lösemi	Kemik iliği bozuklukları, Kanser tedavisi, Otoimmün bozukluklar
Monosit	Kanser, Otoimmün bozukluklar, Kronik Enfeksiyon	Viral iltihap, Konjenital bozukluklar, Otoimmün bozukluklar
Eozinofil	Parazitlik, Alerjen, Astım	Düşüklüğü bir problem olarak görülmemektedir.

FULL TEXTS BOOK

Bazofil	Enfeksiyonlar, Kanserler, Hipotrioidizm	Lösemi, Enfeksiyonlar, Tirotoksitoz

Derin öğrenme modellerinin gelişimi ve çeşitli teknikler ile birleştirilmesi sonucunda 2020 yılında segmentasyon tekniğinin, gözetimli ve gözetimsiz öğrenme etiket kavramını kullanılarak sınıflandırma işleminin gerçekleştirildiği incelenmiştir. Çalışma sürecinde öncelikli veri setinde bulunan WBC çekirdeklerini tespit etmekte diğer yöntemlere göre daha iyi bir sonuç elde edildiği gözlemlenmiştir (Abdurrazzaq vd., 2021). İki aşamalı WBC sınırlandırılmış değer tespiti ve Centernet derin öğrenme modeli ile nesnelerin merkezi temel alınarak sınıf veya boyut tespiti tahmin çalışmaları 2022 yılında gerçekleştirilmiştir. İki adımlı doğrulama sistemi oluşturan modelde elde edilen sonuçlarda değerlendirilmiştir (Zheng vd., 2022).

WBC üzerinde bulanıklık tespiti gerçekleştiren bir çalışmada MSNN ve FMSNN yapay sinir ağı derin öğrenme yöntemi kullanılmıştır. Mikroskobik deney sonucunda kullanılan kan hücre görüntüleri ile yapay sinir ağ tekniğinin geleneksel yöntemlere göre büyük ölçüde başarı elde ettiğini kanıtlayan grup; işlem süresinde hücrelerin yenilenmesi ve birleştirilme aşamalarını sağlayarak sistem üzerindeki etkisini karşılaştırmıştır. Özellik çıkartma işlemleri için normal mikroskobik deney sonuçlarının iyi ve kötü durumdaki görüntüleri kullanılmıştır (Ke, 2008).

Geleneksel makine ve derin öğrenme yöntemlerinin sistematik performans değerlendirmesi sonucunda benzerlik ve farklılıklarını entegre fizik simülasyonu kullanarak test etmiştir. Elde edilen sonuçlar kendi içinde kıyaslanarak polarizasyon eğrileri ve kimyasal reaksiyon modelleri, dayanıklılık testi, kullanım süreci gibi parametreler listelenmiştir. Derin öğrenme ve makine öğrenmesi modellerinin veriseti üzerinde etkili kullanma aşamasını belirten ekip; literatür kapsamında başarılı bir sonuç elde etmiştir (Ming vd., 2023).

WBC gerçek zamanlı tespiti için YOLOv5x ve Detectron2 R50-FPN nesne tespiti kullanılan çalışmada, hibrit model oluşturma işlemi gerçekleştirilmiştir. Test edilen veri setinin aşamalı takip çalışması sonucunda %98 oranında başarı sağladığı gözlemlenmiştir. Detectron2 kullanımı sonucunda elde edilen görsellerin özellik çıkartma işlemleri sağlanarak tespit sisteminin doğruluk değeri detaylı şekilde incelenmiştir (Akalın ve Yumuşak, 2022).

Düşük maliyet ile WBC tespiti ve sayımını görüntü işleme yöntemi kullanarak gerçekleştiren Venoject sistemi geliştirmişlerdir. Gerçekleştirilen çalışmada boyama sistemi sayesinde nesne tepiti sağlanmış ve farklı makine öğrenme teknikleri ile sonuç elde edilmiştir. Eozinofil, lenfosit, nötrofil ve monosit çeşitleri ile gerçekleştirilen çalışmada doğruluk oranı %96.4 olarak saptanmıştır (Begambre vd., 2021).

Lösemi hastalarında ortaya çıkan komplikelerin makine öğrenimi kullanılarak incelenmesi, yeni yöntemlerin keşfedilmesi için çalışmalar gerçekleştirmiştir. Makine öğrenme çeşitlerinden Destek Vektör Makineleri (SVM) ve evrişimli sinir ağı (CNN) kullanılan çalışma sonucunda %93.94 oranında doğruluk değeri elde edilmiştir (Ladha ve Daware, 2018).

Lökosit tespiti için hücrelerin ESA sistemi ile sınıflandırma işlemini gerçekleştirmiştir. Derin öğrenme modelleri kullanılarak WBC sınıflandırılmasında iyileştirmeler yapan grup; eozinofil, lenfosit, monosit ve nötrofil çeşitlerini Densenet201, ResNet50 ve AlexNet yöntemlerini birleştirerek özellik haritası oluşturulmuştur. K-En yakın komşu algoritması ve Naive Bayes, SVM sınıflandırıcıları ile veri seti üzerinde çalışmalar gerçekleştirilmiş, derin öğrenme yöntemleri sayesinde sonuçlar desteklenmiştir. Clahe ve Gauss filtreleri uygulanan veri setine ESA ağı uygulanarak sonuçlarda %97.10 oranında başarı sağlanmıştır (Karakuş ve Özbay, 2022).

WBC tespitinde doğruluk yüzdesini arttırmak için farklı metodojiler üzerinde durmuştur. CNN ve SVM yöntemlerini kullanarak geliştirilen sinir ağı sistem sonuçlarına K-Means yöntemi uygulayarak, kan

FULL TEXTS BOOK

hücrelerinin tespiti ve sınıflandırma işlemleri gerçekleştirilmiştir. Çalışmalar sonucunda keşfedilen metodoloji başarı oranını %91.03 olarak saptanmıştır (Çam ve Güven, 2019).

WBC tespiti için çok terimli lojistik regresyon yöntemi kullanarak filtreleme işlemi gerçekleştirmiştir. Çalışma sürecinde 350 adet mikroskobik kan yayma görseli kullanılarak altı farklı makine öğrenme yöntemi ile sınıflandırılma işlemi gerçekleştirilmiş, 35 farklı geometri ve istatistik özellikler baz alınarak sonuçlar arasında karşılaştırma yapılmıştır. Çalışma sonucunda çok terimli lojistik regresyon yönteminin WBC otomatik sınıflandırılması için uygun bir parametre olduğu, hematologlar ve dahiliye uzmanları için hastalıkların tanısında kaynak olarak kullanılabilirliği tespit edilmiştir. Yapılan çalışma sonucunda doğruluk oranı %95 olarak belirlenmiştir (Elen ve Turan, 2019).

Hematolojik bir hastalık olan lenfoma kanserinin teşhisi için derin öğrenme yaklaşımı kullarak otomatik tespit ve sınıflandırma modeli üzerine çalışmalar yürütmüştür. Oluşturulan modelde bir tür bölge tabanlı konvolüsyon sinir ağı (KSA) ve Faster R-CNN nesne tespiti kullanılarak performansı yüksek bir sınıflandırıcı elde etmişlerdir. Görüntü tabanlı uygulamalarda özellik çıkartma sürecini ortadan kaldırmak için lenfosit, monosit, bazofil, eozinofil ve nötrofil tespiti kullanılarak ResNet50 ile modifiye edilmiş yeni bir sınıflandırıcı modeli ortaya çıkartılmıştır. Çalışmalar sonucunda doğruluk oranı %95.81 olarak saptanmıştır (Uyar ve Taşdemir, 2021).

Lenfosit çeşitleri ve immün sistemdeki görevleri incelemek için vücutta bulunan başlıca iki tip lenfosit tipi üzerine çalışmlara gerçekleştirmiştir. T ve B tipi lenfositler elektron mikroskobu, enzim histokimya ve immun floresan gibi yöntemlerle yüzey özellikleri belirleme çalışmaları gerçekleştirilmiştir. Yapılan çalışma sonucunda hücre tespitinin yaşam süreci, aktive olma durumları incelenerek %75 oranında varlık tespiti elde edilmiştir (Saruhan ve Dereli, 2014).

CNN kullanarak WBC tespit çalışması gerçekleştirmıştır. CNN tabanlı model çıkarma işlemlerinde InceptionV3, DenseNet121, MobileNetV2, ResNet50 modelleri kullanılarak sonuçlara elde edilmiş ve karşılaştırılarak en iyi sonuç elde edilmeye çalışılmıştır. Çalışma sonucunda doğruluk oranı %99.5 olarak tespit edilmiştir (Nahzat vd., 2022).

Akut miyeloid lösemi tanısı için olgunlaşmamış WBC tespit edilmesini ve sayılmasını üzerine çalışmalarda bulunmuştur. Derin öğrenme algoritmalarının birleştirilmesi ve makine öğrenmesi ile test edilen hücrelerin otomatik sayma işlemi için başarılı bir yöntemi ortaya çıkarmıştır. Görüntü işleme ve bölütleme işlemlerinde U-Net yöntemi tercih edilerek farklı bir yöntem denemesi sağlanmıştır. Tespit çalışmaları sonucunda doğruluk oranı %97.06 olarak tespit edilmiştir (Zhang vd., 2024).

WBC sınıflandırma işlemleri için CNN ve Graft Net yöntemlerini kullanmıştır. Derin öğrenme ile test edilen veriler, makine öğrenme model sonuçları ile karşılaştırılarak sanal altıgen kafes (VHT) yöntemi ile birleştirilmesi sağlanmıştır. Bu kapsamda özellik çıkartma işlemleri ile sınıflandırma sisteminde başarılı sonuçlar tespit edilmiştir. Çalışma sonucunda %99.92 oranında doğruluk değeri saptanmıştır (Rashid vd., 2023).

Lökositlerin alt tiplerinin sınıflandırılmasına yönelik bir işlem hattı geliştirmiştir. Çalışma kapsamında Densenet201 ve DarkNet kullanılarak daha küçük verilerden yüksek doğruluk eldesi sağlanmıştır. Özellik seçimi ve çıkartma işlemleri sonucunda aşamalı karşılaştırmalar sağlanarak istenilen doğruluk oranı elde edilmiştir. Yöntem sonucunda %99.8 oranında başarı sağlanmıştır (Ahmad vd., 2023).

Karmaşık kan yayma görüntülerine gömülü WBC tanıyıp, otomatik algılayan ve sınıflandıran bir algoritma tasarlamışlardır. Önerilen algoritma, varlığı yaklaşık olarak tahmin etmek için elips algılama yaklaşımını kullanmaktadır. WBC yapay elektrik alanına bağlı AEFA-C algoritması ile tanıma ve bu yöntem üzerinde sezgisel yapılandırma sistemi oluşturmak için çalışmalar gerçekleştirilmiştir. Gelişimi sağlanan sezgisel optimizasyon aracı ile WBC tespiti %96.90 doğruluk oranında gerçekleştiridiği saptanmıştır (Anita ve Yadav, 2021).

FULL TEXTS BOOK

WBC tespiti için temel bileşen analizi (PCA) tabanlı füzyon metodununu kullanarak morfolojik sınıflandırma işlemi gerçekleştirmişlerdir. CellaVision, BloodSeg, JTSC gibi veritabanları için hücre zarı benzerlik katsayıları elde edilerek karşılaştırma gerçekleştirilmiştir. Çalışmalar kapsamında %90.79 oranında başarı sağlandığı tespit edilmiştir (Makem ve Tiedeu, 2020).

Bölgesel CNN dayalı WBC tespiti ve sınıflandırılmasına yönelik tıbbi hipotez üzerinde çalışmalar gerçekleştirmiştir. AlexNet, VGG16, GoogleNet T.L. ve ResNet 50 modelleri kullanılarak veri seti test edilmiştir. Derin öğrenme yöntemlerini kendi içinde karşılaştıran ekip; farklı derin öğrenme modellerinin performasının WBC çeşitleri üzerine etkisini tahlil etmiştir. Bu kapsamda ResNet 50 modeli başarılı olarak kabul görmüştür (Kutlu vd., 2020)

Sıfır faz bileşen analizi yaklaşımı ile gri seviye eş oluşum matrisine dayalı WBC sınıflandırma işlemi gerçekleştirmiştir. Lenfosit, monosit, nötrofil ve eozinofil hücreleri üzerinde varyans analizi ve sıfır faz bileşen analizi gerçekleştirerek istatiksel sonuçlar elde edilmiştir. Elde edilen sonuçlar FIAL, LSTM ve YOLOv3 gibi derin öğrenme yöntemleri ile incelenerek değerlendirme tablosu oluşturulturmuştur. Çalışma sonucunda sıfır faz analizi uygulanmış veri setinin derin öğrenme yöntemleri ile tespitinde doğruluk oranın %94.25 olarak saptanmıştır (Saika ve Devi, 2023)

Biyomedikal sinyal işleme ve yarı denetimli öğrenme yöntemlerini kullanarak WBC tespitini sağlayan bir model oluşturmuştur. Etkileşimli dikkat öğrenimi ve ince taneli etkileşim mekanizması modelleri sayesinde veri seti üzerinde gerekli test çalışmaları gerçekleştirilmiştir. Eozinofil, lenfosit, monosit ve netrofil çeşitleri AlexNet, GoogleNet, VGG16, ResNet-50, ResNet18, ResNet34 methodları kullanılarak sonuç değerleri elde edilmiş ve karşılaştırma işlemi gerçekleştirilmiştir. Çalışmada odaklanılan metot sonucunda %93.22 oranında başarı elde edilmiştir (Ha vd., 2022).

Katmanlı CNN yöntemini çoklu makine öğrenme sistemi ile birleştirerek WBC tespiti ve sınıflandırılmasını sağlayan bir model oluşturmuşlardır. AlexNet ve MLANet-FS derin öğrenme yöntemleri kullanılan çalışmada %99.12 değerinde test doğruluğu saptanmıştır (Khan vd., 2021).

CNN yöntemini kullanarak kan hücrelerinin segmentasyonu ve sınıflandırılması için derin öğrenme yaklaşımını benimsemiştir. Matlab görüntü işleme araçlarının WBC diğer kan bileşenlerinden ayırdığı gözlemlenen çalışmada; hesaplanan ortalama değer, standart sapma gibi parametrelerin CNN methoduna etkisi incelenmiştir. Bu kapsamda çalışma sonucunda %95 oranında doğruluk tespiti sağlanmıştır (Hemalatha vd., 2022).

WBC tespiti için YOLOv3 nesne algılama tekniğini kullanarak, hücrelerin konumunu algılama ve sınır loşullarında kutucuklara sınıflandırma işlemini gerçekleştirmişlerdir. Eozinofil, lenfosit, monosit ve nötrofil hücreleri Faster R-CNN ve VGG16 sınıflandırıcılarında test edilerek elde edilen sonuçlar YOLOv3 sınıflandırıcısı ile karşılaştırılmıştır. Bu kapsamda YOLOv3 sınıflandırıcısının %90 oranında başarı sağladığı tespit edilmiştir (Praven vd., 2021). WBC tespiti için RCNN yöntemi kullanmıştır. Elde edilen veriler üstünde özellik çıkartma işlemleri gerçekleştirilerek tercih edilen metodun WBC tespit iyileştirmesinde sağlayacağı olanaklar listelenmiştir. Çalışmalar sonucunda, F1-Skoru incelenerek algoritma üzerinde düzenlemeler gerçekleştirilmiştir. Nihai sonuçta %98.70 oranında başarı değeri saptanmıştır (Cheng vd., 2019).

Rajvi ve arkadaşları, WBC ve kırmızı kan hücreleri için başlıca dört algoritma sistemini karşılaştırmıştır. YOLOv3, SSD MobileNetv2, Faster RCNN Inception ve ResNet yöntemleri belirli bir veri seti üzerinde test edilerek sınıflandırma işlemi gerçekleştirilmiştir. Çalışmalar sonucunda, YOLOv3 modelinin obje tespiti için hedeflenen oranı karşıladığı sonucu elde edilmiştir (Shah vd., 2022).

Lösemi hastalığı için WBC tespiti gerçekleştirmiştir. Bazofil, eozinofil, lenfosit, monosit, nötrofil hücreleri YOLOv5x ile tespit edilerek anormal WBC, hastalık taşıyan hücrelere oranı, sayısı ve hücre tipi listelenmiştir. CNN, SVM gibi yöntemler ile kıyas edilen geliştirilmiş model bölütleme, sınıflandırma ve tespit aşamasında %93 oranında doğruluk sağladığı teyit edilmiştir (Luon vd., 2022).

FULL TEXTS BOOK

CNN kullanarak WBC tespiti ve sınıflandırması üzerine çalışmalar gerçekleştirmiştir. Çoklu ve ikili sınıflandırma sistemleri uygulanan veri setinde; derin öğrenme modelinin temel alındığı süreçte ve geliştirilen CNN uygulandığı süreçte elde edilen sonuç incelemesi gerçekleştirilmiştir. Bu kapsamda çalışma çıktılarının %98.63 oranında başarı sağlandığı belirtilmiştir (Ridoy ve Islam, 2020).

Lökositlerin otomatik ekstraksiyonuna odaklanarak renk bölümleme ve eşikleme işlemlerinin matematiksel dönüşümünü sağlayan K-Means yöntemi kullanarak incelemeyi tercih etmişlerdir. Çalışma sürecinde nötrofil, eozinofil, basofil, lenfosit, monosit WBC çeşitleri kullanılarak morfolojik segmentasyon öncesi boyama ve boyanın kaldırılma işlemi gerçekleştirilerek ortaya çıkan sonuçlar listelenmiştir (Gautam ve Bhadauria, 2014).

Görüntü işleme yöntemini kullanarak WBC tespiti ve bölütlemesi için K-Means kümelemesine dayalı algoritma geliştirmiştir. Hücre lokalizasyonu ve sınıflandırması için VGG-16 sınıflandırıcısı kullanan ekip; basofil, eozinofil, nötrofil, lenfosit, monosit hücrelerini belirli oranda bölerek test işlemine dahil etmiştir. Elde edilen sonuçlarda %95.89 oranında başarı sağlandığı saptanmıştır (Chinthalanka vd., 2020).

WBC tespiti ve sınıflandırma süreci için derin öğrenme modellerini tercih edilmiştir. CNN, AlexNet, LeNet ve VGG-16 sofmax sınıflandırıcısı ile gerçekleştirilen çalışmada modellerin uygulanması python programlama dili aracılığıyla sağlanmıştır. Özellikle CNN mimarisi üzerinde ayrıntlı analizler yapılmıştır. Çalışma sonucunda veriler karşılaştırılarak listelenmiş doğruluk oranı en iyi olan sınıflandırıcılar %99 değeri ile AlexNet ve LeNet olarak tespit edilmiştir (Kousalya vd., 2021)

WBC tespiti ve sınıflandırılması için mikroskobik olarak akut lenfoblastik lösemi görüntü işlemede kullanan verileri kullanmıştır. K-En yakın komşu algoritması kullanan sınıflandırma yönteminde geometrik ve istatistiksel özelliklere dayalı regresyon hesaplama işlemleri gerçekleştirilmiştir. Eşik ve dönüm noktası algoritmalarını kullanarak segmentasyon adım hesabı üzerine çalışan ekip: çalışmalar sonucunda %93.33 oranında doğruluk değeri saptamıştır (Heriawati vd., 2021).

YÖNTEM

WBC tespiti ve sınıflandırılması için gerçekleştirilen çalışmalarda morfolojik çalışmalar beraberinde geliştirilmiş CNN modelleri de tercih edilmiştir. Tablo 2'de literatür çalışmalarında tercih edilen yöntemlere yer verilmiştir.

Tablo 2. WBC Tespiti ve Sınıflandırma Yöntemleri.

Çalışma	Çalışmada Kullanılan Yöntemler
Özcan, 2022	Evrişimli sinir ağı modeli, bölütleme yöntemleri
Abdurrazzaq vd., 2021	Tekrarlı görüntü göstergesi ve tekil değer ayrıştırma yöntemleri
Zheng vd., 2022	Öne çıkan bölge belirleme ve CenterNet yöntemi
Ke, 2008	Fuzzy Morfoloji

FULL TEXTS BOOK

Begambre vd., 2021	Venoject sistemi, makine öğrenmesi
Ladha ve Daware, 2018	Destek vektör makinesi (SVM), CNN yöntemleri
Karakuş ve Özbay, 2022	Evrişimli sinir ağı modeli, DenseNet201, ResNet50, AlexNet ve K-En yakın komşu algoritması
Çam ve Güven, 2019	CNN, SVM ve K-Means yöntemi
Elen ve Turan, 2019	Regresyon yöntemi, filtreleme
Uyar ve Taşdemir, 2021	Konvolüsyon sinir ağı, Faster R-CNN, ResNet50
Nahzat vd., 2022	CNN, InceptionV3, DenseNet121, MobileNetV2, ResNet50
Zhang vd., 2024	Görüntü işleme ve bölütleme işlemleri, U-Net
Rashid vd., 2023	Sanal altıgen kafes yöntemi (VTH), CNN, GraftNet
Ahmad vd., 2023	DenseNet201, DarkNet
Anita ve Yadav, 2021	AEFA-C algoritması
Makem ve Tiedeu, 2020	PCA tabanlı füzyon metodu, morfolojik sınıflandırma, CellaVision, BloodSeg, JTSC
Kutlu vd..2020	R-CNN, AlexNet, VGG16, GoogleNet, ResNet50

Saika ve Devi, 2023	Sıfır faz bileşen analizi, FIAL, LSTM, YOLOv3
Ha vd., 2022	Yarı denetimli öğrenme modeli, AlexNet, GoogleNet, VGG16, ResNet-50, ResNet18, ResNet34 yöntemleri
Khan vd.,2021	Evrişimli sinir ağı, AlexNet ve MLANet-FS derin öğrenme yöntemleri
Hemalatha vd., 2022	Derin öğrenme modelleri ve standart sapma parametreleri
Praven vd., 2021	YOLOv3, Faster R-CNN, VGG16 yöntemleri
Cheng vd., 2019	R-CNN, özellik çıkartma işlemleri
Shah vd.,2022	YOLOv3, Faster R-CNN, R-FCN
Luon vd., 2022	YOLOv5x, CNN, SVM ve bölütleme işlemleri
Ridoy ve Islam, 2020	YOLOv3, SSD MobileNetv2, Faster RCNN Inception ve ResNet yöntemleri
Gautam ve Bhadauria, 2014	K-Means kümeleme, Gri-Ton dönüşümü
Chinthalanka vd., 2020	Bölütleme, K-Means kümeleme algoritması ve VGG-16
Kousalya vd., 2021	CNN, AlexNet, LeNet ve VGG-16 sofmax sınıflandırıcı yöntemleri

FULL TEXTS BOOK

Heriawati vd., 2021	Fuzzy C-Means, Histogram eşitleme

TARTIŞMA VE SONUÇ

Bu derleme çalışması kapsamında, WBC tespiti ve sınıflandırılmasına yönelik güncel metotlar incelenmiştir. Son yıllarda derin öğrenme ve makine öğrenmesi temelli yöntemlerin bu alandaki başarısını önemli ölçüde ortaya koymuştur. Literatür taraması sonucunda, CNN, U-Net, YOLO modellerin hız, doğruluk ve klinik uyarlanabilirlik açısından önemli avantajlar sağladığı; hibrit modellerin ise bulundurduğu yöntemlerin güçlü yönlerini birleştirerek performans düzeyini artırdığı tespit edilmiştir. Geleneksel görüntü işleme teknikleri, temel ön işleme ve segmentasyon adımları günümüzde etkinliğini korumakta, ancak karmaşık veri kümelerinde derin öğrenme tabanlı çözümler daha yüksek başarı oranlarına ulaşmakta olduğu tespit edilmiştir. Verisetlerinin sınırlılığı, klinik entegrasyon ve farklı laboratuvar koşullarına bağlı görüntü değişkenlikleri sürecindeki teknik engeller öne çıkmaktadır. Gelecekte, daha geniş ve dengeli veri kümelerinin oluşturulması, veri artırma tekniklerinin etkin kullanımı ve yapay zekâ destekli karar sistemlerinin klinik iş akışına entegrasyonu kritik öneme sahip olduğu gözlemlenmiştir.

Sonuç olarak, derin öğrenme tabanlı WBC tespiti ve sınıflandırma yöntemlerinin doğruluk ve hız açısından başarılı sonuçlar sunduğu, fakat klinik geçerlilik için daha kapsamlı çalışmaların gerekliliği vurgulanmaktadır. Bu alanda gerçekleştirilecek multidisipliner çalışmalar, erken teşhis ve tedavi süreçlerinde önemli katkılar sağlaması öngörülmektedir.

KAYNAKLAR

Ahmad, R., Awais, M., Kausar, N., Tariq, U., Cha, J. H., & Balili, J. (2023). Leukocytes classification for leukemia detection using quantum-inspired deep feature selection. Cancers, 15, 2507. Abdurrazzaq, A., Junoh, K. A., Yahya, Z., & Mohd, I. (2021). New white blood cell detection technique by using singular value decomposition concept. Springer Science, 80, 4627–4638.

Akalın, F., & Yumuşak, N. (2022). Detection and classification of white blood cells with an improved deep learning-based approach. Turkish Journal of Electrical Engineering and Computer Sciences, 12(7), 2725–2739.

Anita, A., & Yadav, A. (2021). An intelligent model for the detection of white blood cells using artificial intelligence. Computer Methods and Programs in Biomedicine, 199, 105893.

Begambre, S., Castillo, C., & Vilimizar, L. H. (2021). Low cost classification method for differentiated white blood cells using digital image processing and machine learning algorithms. IEEE Colombian Conference on Applications of Computational Intelligence (ColCACI), Cali, Colombia, pp. 1-5.

Çam, F., & Güven, A. (2019). Methods used in the classification of white blood cells from blood cell images taken under a digital microscope. Electronic Letters on Science & Engineering, 15(3), 23–43.

Cheng, S., Suhua, Y., & Shaofeng, J. (2019). Improved faster RCNN for white blood cells detection in blood smear image. 14th IEEE International Conference on Electronic Measurement & Instruments (ICEMI), Changsha, China, pp. 1677-1682.

Elen, A., & Turan, M. K. (2019). Classifying white blood cells using machine learning algorithms. International Journal of Engineering Research and Development, 11(1), 141–152.

FULL TEXTS BOOK

Gautam, A., & Bhadauria, H. S. (2014). White blood nucleus extraction using K-mean clustering and mathematical morphing. 5th International Conference - Confluence The Next Generation Information Technology Summit (Confluence), Noida, India, pp. 549-554.

Ha, Y., Du, Z., & Tian, J. (2022). Fine-grained interactive attention learning for semi-supervised white blood cell classification. Biomedical Signal Processing and Control, 75, 103611.

Hemalatha, B., Karthik, B., Reddy, K. C. V., & Latha, A. (2022). Deep learning approach for segmentation and classification of blood cells using enhanced CNN. Measurement: Sensors, 24, 100582.

Heriawati, S. O., Harsono, T., Bachtiar, M. M., & Hernaningsih, Y. (2021). Blood cells classification for identification of acute lymphoblastic leukemia on microscopic images using image processing. International Electronics Symposium (IES), Surabaya, Indonesia, pp. 215-220.

Karakuş, Ö. M., & Özbay, E. (2022). Lökosit tespiti için beyaz kan hücrelerinin ESA kullanılarak sınıflandırılması. Adıyaman Üniversitesi Mühendislik Bilimleri Dergisi, 17(1), 333–334.

Ke, C. (2008). White blood cell detection using a novel fuzzy morphological shared-weight neural networks. International Symposium on Computer Science and Computational Technology, Shanghai, China, pp. 532-535.

Khan, A., Eker, A., Chefranov, A., & Demirel, H. (2021). White blood cell type identification using multi-layer convolutional features with an extreme learning machine. Biomedical Signal Processing and Control, 69, 102932.

Kousalya, K., Mohana, R. S., Krishnakuma, B., & Karthikeya, N. (2021). Comparative analysis of white blood cells classification using deep learning architectures. Proceedings of the Second International Conference on Smart Electronics and Communication (ICOSEC), Trichy, India, pp. 1220-1225.

Kutlu, H., Avcı, E., & Özyurt, F. (2020). White blood cells detection and classification based on regional convolution neural networks. Medical Hypotheses, 135, 109472.

Laddha, S., & Daware, S. (2018). Analysis of white blood cell segmentation techniques and classification using deep convolutional neural network for leukemia detection. Helix The Scientific Explorer, 8(6), 4519–4524.

Luon, D. T., Huong, H. T. L., Anh, D. D., Thang, T. X., Hanh, T. T., & Khanh, D. M. (2022). Distinguish normal white blood cells from leukemia cells by detection, classification, and counting blood cells using YOLOv5. 7th National Scientific Conference on Applying New Technology in Green Buildings (ATiGB), Da Nang, Vietnam, pp. 156-160.

Makem, M., & Tiedeu, A. (2020). An efficient algorithm for detection of white blood cell nuclei using adaptive three stage PCA-based fusion. Informatics in Medicine Unlocked, 20, 100416.

Ming, W., Sun, P., Zhang, Z., Qiu, W., Du, J., Li, X., Zhang, Y., Zhang, G., Liu, K., Wang, Y., & Guo, X. (2023). A systematic review of machine learning methods applied to fuel cells in performance evaluation, durability prediction, and application monitoring. International Journal of Hydrogen Energy, 48, 5197–5228.

Nahzat, S., Bozkurt, F., & Yağanoğlu, M. (2022). White blood cell classification using convolutional neural network. Bilim, Teknoloji ve Mühendislik Araştırmaları Dergisi, 3(1), 32–41.

Özcan, Ş. (2022). Beyaz kan hücrelerinin evrişimli sinir ağları ile sınıflandırılması ve bölütlenmesi (Yüksek lisans tezi). Başkent Üniversitesi, Fen Bilimleri Enstitüsü, Ankara, 1–27.

Praven, N., Sonbhadra, K. S., Syafrullah, M., Punn, N. S., Agarwal, S., & Adiyarta, K. (2021). White blood cell subtype detection and classification. 8th International Conference on Electrical Engineering, Computer Science and Informatics (EECSI), Semarang, Indonesia, pp. 203-207.

Rashid, S., Raza, M., Sharif, M., Azam, F., Kadry, S., & Kim, J. (2023). White blood cell image analysis for infection detection based on virtual hexagonal trellis (VHT) by using deep learning. Scientific Reports, 13, 17827.

Ridoy, A. R., & Islam, R. (2020). A lightweight convolutional neural network for white blood cells classification. 23rd International Conference on Computer and Information Technology (ICCIT), DHAKA, Bangladesh, pp. 1-5.

Saika, R., & Devi, S. S. (2023). White blood cell classification based on gray level co-occurrence matrix with zero phase component analysis approach. Procedia Computer Science, 218, 1977–1984.

Saruhan, B. G., & Dereli, S. (2014). Lenfosit çeşitleri ve immün sistemdeki görevleri. Dicle Üniversitesi Veteriner Fakültesi Dergisi, 1(3), 41–48.

Shah, R., Shastri, J., Bohara, H. M., Panchal, Y. B., & Goel, P. (2022). Detection of different types of blood cells: A comparative analysis. IEEE International Conference on Distributed Computing and Electrical Circuits and Electronics (ICDCECE), Ballari, India, pp. 1-5.

Uyar, K., & Taşdemir, Ş. (2021). Detection and classification of leucocyte types in histological blood tissue images using deep learning approach. European Journal of Science and Technology, 24, 130–137.

Wijesinghe C. B., Wickramarachchi D. N., Kalupahana I. N., De Seram L. R., Silva I. D., & Nanayakkara N. D. (2020). Fully automated detection and classification of white blood cells. Annual International Conference of the IEEE Engineering in Medicine and Biology Society., pp. 1816-1819.

Zhang, Z. Z., Arabyarmohammadi, S., Leo, P., Meyerson, H., Metheny, L., Xu, J., & Madabhushi, A. (2024). Automatic Myeloblast Segmentation in Acute Myeloid Leukemia Images based on Adversarial Feature Learning. *Computer Methods and Programs in Biomedicine*, 243: 107852. Elsevier.

Zheng, X., Pan, T., Ai, L., Liu, D., Zhang, Y., & Wang, B. (2022). White blood cell detection using saliency detection and CenterNet: A two-stage approach. Journal of Biophotonics, 16(3), 1–15.

FARKLI ORANLARDA NH_3/CO_2 KARIŞIMININ TERMODİNAMİK PERFORMANS DEĞERLENDİRMESİ

THERMODYNAMIC PERFORMANCE EVALUATION OF THE CO_2/NH_3 MİXTURE IN SEVERAL FRACTIONS

***Jale BERRAK*[1]**

[1]*Yüksek Lisans Öğrencisi, İskenderun Teknik Üniversitesi, Lisansüstü Eğitim Enstitüsü, Makine Mühendisliği,*

[1]*ORCID ID: https://orcid.org/0009-0006-6316-4709*

***Ali KOÇ*[2]**

[2]*Prof. Dr., İskenderun Teknik Üniversitesi, Mühendislik ve Doğa Bilimleri Fakültesi, Makine Mühendisliği*

[2]*ORCID ID: https://orcid.org/0000-0002-7388-2628*

***Özkan KÖSE*[3]**

[3]*Dr. Öğr. Üyesi, İskenderun Teknik Üniversitesi, Mühendislik ve Doğa Bilimleri Fakültesi, Makine* [3]*Mühendisliği Anabilim Dalı*

[3] *ORCID ID: https://orcid.org/0000-0002-9069-1989*

***Yıldız KOÇ*[4]**

[4]*Doç. Dr., İskenderun Teknik Üniversitesi, Mühendislik ve Doğa Bilimleri Fakültesi Makine Mühendisliği*

[4] *ORCID ID: https://orcid.org/0000-0002-2219-645X*

ÖZET

Artan küresel enerji talebi ve çevresel kaygılar doğrultusunda, verimli ve sürdürülebilir enerji üretim teknolojilerine olan ihtiyaç giderek daha önemli hale gelmiştir. Süperkritik karbondioksit (sCO_2) çevrimleri, yüksek termal verimleri, kompakt tasarımları ve çevreye duyarlı yapıları nedeniyle umut vadeden sistemler olarak öne çıkmaktadır. Ancak, saf CO_2'nin kritik noktasına yakın termodinamik davranışı, sınırlı ısı transfer performansı ve çevrim esnekliğini azaltması gibi bazı operasyonel zorluklara neden olmaktadır. Bu sınırlamaların üstesinden gelmek amacıyla, bu çalışmada CO_2 bazlı ikili akışkan karışımlarının sistem performansı üzerindeki etkileri araştırılmıştır. Çalışmada, karbondioksit (CO_2) ile amonyak (NH_3) karışımı çalışma akışkanı olarak kullanılmış olup, NH_3 mol oranı %0 ile %90 arasında değişmektedir. Detaylı termodinamik simülasyonlar aracılığıyla, farklı karışım oranları ve çalışma koşullarında sistemin enerji ve ekserji verimleri değerlendirilmiştir. CO_2/NH_3 karışımının saf CO_2'ye kıyasla daha yüksek performans gösterdiğini ortaya koymakta olup, geleneksel sistemlerin kullanılamadığı sıcaklıklarda oldukça avantajlı olduğu görülmüştür. Bu çalışma, geleceğin sürdürülebilir enerji sistemleri için gelişmiş sCO_2 çevrim tasarımlarına katkı sağlamakta ve CO_2–NH_3 karışımının yüksek potansiyelli bir çalışma akışkanı olduğunu göstermektedir. Çalışma akışkanı içerisindeki NH_3 oranı arttıkça sistemin termodinamik performansında iyileşme olduğu net bir şekilde görülmüştür. Sonuç olarak, %90 NH_3 içeriğine sahip bir çalışma akışkanı kullanıldığında enerji ve ekserji verimleri sırasıyla %33,54 ve %64,14 olarak hesaplanarak en yüksek değerlere ulaşmıştır.

Anahtar Kelimeler: CO_2/NH_3 karışımı, enerji, ekserji

FULL TEXTS BOOK

ABSTRACT

Amid the growing global energy demand and rising environmental concerns, the pursuit of efficient and sustainable power generation technologies has become increasingly critical. Supercritical carbon dioxide (sCO_2) power cycles have emerged as promising candidates due to their superior thermal efficiency, compact design, and reduced environmental impact. Nevertheless, the thermodynamic behavior of pure CO_2 near its critical point poses operational challenges, including limited heat transfer performance and reduced cycle flexibility. To address these limitations, this study investigates the impact of using CO_2-based binary mixtures on system performance. Specifically, a mixture of carbon dioxide (CO_2) and ammonia (NH_3) is employed as the working fluid, with the NH_3 concentration varying between 0% and 90% by mole fraction. Through detailed thermodynamic simulations, the energy and exergy efficiencies of the system are evaluated across a wide range of mixture compositions and operating conditions. The results indicate that CO_2/NH_3 leads to better performance compared to pure CO_2. The CO_2/NH_3 mixture demonstrated superior performance over pure CO_2 and was found to offer significant advantages at temperature ranges where conventional systems are ineffective. This research contributes to the development of advanced sCO_2 cycle designs and demonstrates the viability of CO_2–NH_3 mixtures as high-potential working fluids in next-generation sustainable power systems. The findings are aligned with global efforts to decarbonize energy systems and promote environmentally responsible technologies. It has been clearly observed that increasing the NH_3 ratio in the working fluid leads to a significant improvement in the thermodynamic performance of the system. As a result, when a working fluid with 90% NH_3 content is used, the highest energy and exergy efficiencies are achieved, calculated as 33.54% and 64.14%, respectively.

Keywords: CO_2/NH_3 mixture, energy, exergy

1. GİRİŞ

Günümüzde artan enerji ihtiyacı ve çevresel sorunlar, enerji üretiminde daha verimli ve çevre dostu teknolojilerin geliştirilmesini zorunlu hale getirmiştir [1,4]. Fosil yakıtların yoğun kullanımı hem kaynakların tükenmesine hem de sera gazı emisyonlarının artmasına neden olarak küresel iklim değişikliğini hızlandırmaktadır [5]. Bu bağlamda, yenilenebilir enerji kaynaklarının daha etkin kullanılmasını sağlayacak, yüksek verimli ve düşük emisyonlu enerji dönüşüm sistemlerinin geliştirilmesi büyük önem taşımaktadır [6]. Süperkritik karbondioksit çevrimleri, yüksek termodinamik verimlilikleri, kompakt sistem tasarımlarına olanak tanımaları ve düşük çevresel etkileri ile son yıllarda dikkat çeken enerji üretim teknolojileri arasında yer almaktadır [1,3]. Ancak saf CO_2'nin kritik noktasına yakın davranışı ve düşük ısı transfer kapasitesi, bu sistemlerin operasyonel esnekliğini sınırlamakta ve termodinamik performansını olumsuz yönde etkilemektedir [2,8]. Bu sınırlamaların aşılması amacıyla, CO_2 bazlı ikili akışkan karışımlarının kullanımı literatürde giderek daha fazla yer bulmaktadır [8,9]. Özellikle CO_2'nin uygun katkı maddeleri ile karıştırılması, çevrim performansını iyileştirme potansiyeli sunmaktadır [7]. Bu doğrultuda, bu çalışmada CO_2 ile amonyak (NH_3) karışımının farklı oranlardaki kullanımıyla elde edilen ikili karışımın, 400 bar basınç ve 450°C sıcaklıkta çalışan bir süperkritik güç çevrimi üzerindeki termodinamik etkileri detaylı şekilde incelenmiştir.Literatürde yapılan bazı çalışmalar, Süperkritik karbondioksit (sCO_2) çevrimleri, saf CO_2'nin propan, etan, izobütan, toluen, hidrojen sülfür (H_2S) gibi akışkanlarla yapılan karışımlar kritik sıcaklık ve basınç değerlerini değiştirerek çevrimin ısıl eşleşmesini optimize etmeyi, ekserji kayıplarını azaltmayı ve yüksek sıcaklık koşullarında verimliliği korumayı hedeflemektedir [10,11]. Bazı çalışmalara dikkatle bakıldığında, CO_2–Propan ve CO_2–İzobütan karışımlarının kritik sıcaklığı düşürerek düşük sıcaklıkta ısı geri kazanımını iyileştirdiği rapor edilmiştir [12]. CO_2–Toluen karışımı ise özellikle kısmi ısıtmalı çevrimlerde ekserji veriminde %25–%29 oranında iyileşme sağlamış; ancak bu durum sistem

FULL TEXTS BOOK

maliyetlerinde artışa yol açmıştır [13].


CO_2–H_2S karışımları, kritik sıcaklığı yükseltmiş ve özellikle soğutucu bileşenlerde ekserji kaybını azaltmıştır. Farklı çalışmalara bakılınca, Liu ve arkadaşları (2019) CO_2-H_2 ve CO_2-CH_4 ikili karışımlarının yüksek basınç ve sıcaklık altında termodinamik özelliklerini modelleyerek enerji çevrimlerindeki potansiyel kullanımını araştırmışlardır. Çalışmada, özellikle hidrojen karışımlarının çevrim verimliliğini artırıcı etkisi vurgulanmış ve karışım oranlarının optimize edilmesinin performansa önemli katkı sağladığı belirtilmiştir [16]. Zhang ve ark. (2021) ise CO_2-NH_3 karışımlarının ekserji analizi ile çevrim verimliliği üzerindeki etkilerini incelemişlerdir. Bu çalışma, farklı karışım oranlarının termodinamik performansı nasıl değiştirdiğine dair önemli bulgular sunmakta ve özellikle yüksek sıcaklık ve basınç koşullarında optimum karışımın çevrim verimliliğini artırdığı sonucuna ulaşmıştır [17]. Smith ve arkadaşları (2018) karbon yakalama ve depolama (CCS) süreçlerinde CO_2-N_2 ve CO_2-H_2 karışımlarının faz dengesi ve termodinamik özelliklerine odaklanmıştır. Bu çalışma, CO_2 karışımlarının yüksek basınçlı faz davranışlarını deneysel ve sayısal yöntemlerle ortaya koymuş ve CCS sistemleri için kritik veriler sağlamıştır [18]. Buna ek olarak, Wang ve Lee (2020) tarafından yapılan araştırmada CO_2-H_2O karışımlarının termodinamik ve faz dengesi özellikleri değerlendirilmiş, bu karışımların enerji çevrimlerinde ve soğutma sistemlerinde kullanımı için önerilerde bulunulmuştur. Çalışma, su buharının CO_2 ile karışımının çevrimlerde verimliliği artırabileceğini göstermiştir [19]., Kim ve ark. (2022) tarafından CO_2-N_2 karışımlarının termodinamik davranışı ve enerji çevrimlerindeki uygulamaları üzerine yapılan çalışmalar, azotun CO_2 ile karışımında sistem kararlılığının arttığını ve enerji verimliliği açısından avantajlar sunduğunu ortaya koymuştur [20]. CO_2–H_2S karışımları, kritik sıcaklığı yükseltmiş ve özellikle soğutucu bileşenlerde ekserji kaybını azaltmıştır. [14] Hibrit sistemlerde (örneğin LNG sıvılaştırma–CO_2 yakalama entegrasyonu) CO_2 bazlı karışımlar enerji tüketimini düşürmüş, denizcilik atık ısı geri kazanımı uygulamalarında ise yakıt tüketimi ve CO_2 emisyonlarının azaltılmasına katkı sağlamıştır. [15] Yapılan araştırmalar sonucunda ayrıca saf CO_2'nin farklı NH_3/CO_2 oranlarında elde edilen karışımlar kullanılarak sistemin enerji ve ekserji verimlilikleri hesaplanmış; pompa güç tüketimi, net güç üretimi ve rejeneratif ısı kazancı gibi performans parametreleri analiz edilmiştir. Bu analizler sonucunda, özellikle NH_3 oranının artırılmasıyla sistem performansında belirgin iyileşmeler gözlemlenmiş ve CO_2–NH_3 karışımının gelecekteki enerji sistemleri için yüksek potansiyelli bir alternatif çalışma akışkanı olabileceği değerlendirilmiştir. Bu çalışmalar genel olarak, CO_2 bazlı ikili karışımların saf CO_2'ye kıyasla daha iyi ısı eşleşmesi sunduğunu, yüksek sıcaklıklarda verimliliği koruduğunu ve sistem tasarımında esneklik sağladığını ortaya koymaktadır. Ancak her katkı maddesinin termodinamik etkileri, optimum karışım oranı ve uygulanabilirliği farklılık göstermektedir.

Yapılan literatür karşılaştırması, bu çalışmanın hem çalışma koşulları hem de kapsam açısından mevcut araştırmalardan ayrıldığını göstermektedir. Çalışmada, literatürde genellikle düşük veya orta seviyelerde incelenen basınç ve sıcaklık aralıklarının aksine, 400 bar gibi oldukça yüksek basınç ve 450°C sıcaklık koşulları tercih edilmiştir. Ayrıca, NH_3/CO_2 karışım oranlarının %0'dan %90'a kadar geniş bir aralıkta incelenmesi, optimum karışım oranının belirlenmesine olanak tanımış ve performans parametreleri (kütlesel debi, pompa güç tüketimi, reküperatör ısı kazanımı, net güç, enerji verimi, ekserji verimi) değerlendirilmiştir.

2. SİSTEM TANITIMI

Bu çalışmada incelenen çevrim, yüksek basınç ve sıcaklık koşullarında çalışan kapalı devre bir güç üretim sistemi olup, akışkanın basınçlandırılması, ısıtılması, genişletilmesi ve yoğuşturulması süreçlerinden oluşmaktadır. Şekil 1'de verilen akış diyagramı, çevrimin temel bileşenleri arasındaki enerji ve kütle akışını göstermektedir

FULL TEXTS BOOK



Şekil 1. Çevrimin akış diyagramı

Bu çalışmada, farklı oranlardaki NH_3/CO_2 ikili karışımının termodinamik performansını değerlendirmek amacıyla süperkritik özellikte çalışan kapalı çevrimli bir güç sistemi modellenmiştir. Sistem modeli, EBSILON@Professional yazılımında oluşturulmuş olup, temel olarak bir **rejeneratif Rankine benzeri çevrim** mantığına sahiptir. Çevrim, düşük basınçtaki akışkanın (1) pompa vasıtasıyla basıncının yükseltilmesiyle başlar [21,22]. Bu aşamada akışkanın yoğunluğu artmakta, sıcaklık değişimi ise sınırlı kalmaktadır. Basıncı artırılmış akışkan (6) [23], çevrimdeki daha sıcak akışkandan ısı geri kazanımı yoluyla ön ısıtma işlemine tabi tutulur. Bu ön ısıtma süreci (2), akışkanın ana ısı kaynağına girmeden önce sıcaklığını artırarak sistemin toplam enerji ihtiyacını azaltır. Ön ısıtılmış akışkan (9-10), harici bir ısı kaynağından enerji alarak buharlaşma veya süperkritik hale geçiş sürecini tamamlar [21]. Bu noktada akışkanın sıcaklığı ve özgül entalpisi önemli ölçüde yükselir. Yüksek basınç ve sıcaklıktaki akışkan (4), genişleme süreci için türbine yönlendirilir. Türbinde gerçekleşen genişleme sonucu (5) [22], akışkanın basıncı ve sıcaklığı düşerken mekanik iş elde edilir ve bu iş jeneratör aracılığıyla elektrik enerjisine dönüştürülür. Genişleme sonrasında hala yüksek sıcaklığa sahip olan akışkan (3), ısı geri kazanım sürecinde ön ısıtma aşamasındaki akışkana ısısını aktararak çevrim verimliliğini artırır [23]. Isı geri kazanımının ardından akışkan (8), yoğuşturma sürecine girerek faz değişimiyle sıvı hale dönüşür [21,22]. Yoğuşmuş ve düşük basınç seviyesine gelmiş akışkan (7) tekrar pompaya yönlendirilerek çevrim tamamlanır.

3. MATEMATİKSEL MODEL

Bu çalışmada, NH_3/CO_2 ikili karışımının 400 bar ve 450°C koşullarında çalışan kapalı çevrimli güç sistemindeki termodinamik performansı; kütle, enerji ve ekserji denklikleri esas alınarak modellenmiştir. Model kararlı hâl (steady-state) varsayımıyla kurulmuş; akım hatlarında potansiyel ve kinetik enerji değişimleri ihmal edilmiştir. Çevre koşulları T_0 ve P_0 ile gösterilmiştir. Varsayımlar:

- Sistem kararlı hâlde çalışmaktadır.
- Kinetik ve potansiyel enerji değişimleri ihmal edilmiştir.
- Çevre ile olan ısı kayıpları ihmal edilmiştir.
- NH_3/CO_2 karışımının özellikleri EBSILON/REFPROP ile hesaplanmıştır.
- Basınç düşümleri 0,1 bar olarak kabul edilmektedir.

FULL TEXTS BOOK

Ayrıca çalışmada dikkate alınan bazı temel denklemler aşağıda belirtilmektedir:

Kütlenin Korunumu

Her bir kontrol hacmi için kütle korunumu denkliği aşağıdaki gibidir:

$$\sum \dot{m}_{gir} = \sum \dot{m}_{çık}$$

(1)

Enerji Analizi (Birinci Yasa)

Genel kontrol hacmi enerji dengesi (kararlı hâl) dikkate alındığında enerji korunumu denkliği:

$$\sum \dot{Q} - \sum \dot{W} + \sum \dot{m}_{gir} h_{gir} - \sum \dot{m}_{çık} h_{çık} = 0$$

Net İş ve Termal Verim

$$\dot{W}_{net} = \dot{W}_{türbin} - \dot{W}_{pompa}$$

$$\eta_{enerji} = \left(\frac{\dot{W}_{net}}{\dot{Q}_{gir}}\right) \times 100$$

Ekserji Analizi (İkinci Yasa),

Genel ekserji denklemi:

$$\sum \left(1 - \frac{T_0}{T}\right) \cdot \dot{Q} - \dot{W} + \sum (\dot{m} \cdot e)_{giriş} - \sum (\dot{m} \cdot e)_{çıkış} - \dot{E}_{yıkım} = 0$$

Ekserji akışı:

$$\dot{E} = \dot{m} \cdot e$$

Akışkanın özgül ekserjisi:


$$e = (h - h_0) - T_0(s - s_0)$$

Ekserji verimi:

$$\eta_{\text{ekserji}} = \left(\frac{\dot{W}_{net}}{\dot{E}_{gir}}\right)$$


4. ARAŞTIRMA BULGULARI VE TARTIŞMA

Aşağıda sunulan grafiklerde, NH_3/CO_2 karışım oranının artırılmasının çevrim üzerindeki etkileri, 400 bar basınç ve 450°C sıcaklık altında gerçekleştirilen termodinamik analiz sonuçlarına göre detaylı şekilde gösterilmiştir. Her bir parametre için gözlemlenen eğilimler, karışımın çevrime sağladığı potansiyel iyileştirmeleri açıkça ortaya koymaktadır.



Şekil 2. NH_3 oranına göre kütlesel debi değişimi

Şekil 2'de görüldüğü üzere, NH_3/CO_2 karışımında NH_3 oranı %0'dan %90'a yükseldikçe çevrimdeki toplam kütlesel debi 90,34 kg/s'den 22,68 kg/s'ye kadar azalmıştır. Bu azalma, NH_3'nin özgül entalpisi daha yüksek bir akışkan olması sebebiyle, aynı termal enerji transferi için daha az kütlesel akış gerektirmesiyle açıklanmaktadır. 400 bar ve 450°C gibi süperkritik koşullar altında bu özellik daha da belirgin hale gelmektedir. Debi azalması, pompa gücünün düşmesine ve çevrim bileşenlerinin daha kompakt tasarlanmasına olanak sağlamaktadır.




Şekil 3. NH_3 oranına göre pompa güç tüketimi

Şekil 3'te gösterilen pompa güç tüketimi, artan NH_3 oranıyla birlikte önemli ölçüde azalmıştır. %0 NH_3 içeren sistemde 4625,99 kW olan pompa enerji tüketimi, %90 NH_3 oranında 1649,54 kW seviyesine


FULL TEXTS BOOK

gerilemiştir. Bu yaklaşık %64'lük düşüş, akışkan yoğunluğunun azalması ve kütlesel debideki azalma ile ilişkilidir. Bu durum, çevrimin yalnızca ana enerji üretiminde değil, yardımcı ekipmanların enerji maliyetinde de ciddi bir iyileşme sağladığını göstermektedir.



Şekil 4. Net güç ve rejeneratif ısı kazancı değişimi

Şekil 4'te görüldüğü üzere, %0 NH_3 oranında net güç üretimi 12393,89 kW iken, NH_3 NH_3 oranında bu değer 13599,76 kW'ye yükselmiştir. Bu artış, pompa işinin azalmasına ve türbinin daha verimli çalışmasına bağlanmaktadır. Öte yandan, rejeneratif ısı kazancı 9965,23 kW'den 1586,62 kW'ye düşmüştür. NH_3 oranı arttıkça, akışkanın termofiziksel özellikleri nedeniyle rejeneratif ısı alışverişi potansiyeli azalmaktadır. Bu durum, sistemin rejeneratör bağımlılığını azaltarak daha kompakt bir tasarıma olanak sağlamakta, aynı zamanda daha az karmaşık bir çevrim konfigürasyonu sunmaktadır.



Şekil 5. Enerji ve ekserji verimlerinin karşılaştırılması

Şekil 5'te yer alan veriler, NH_3 oranı arttıkça hem enerji (1. yasa) hem de ekserji (2. yasa) verimlerinde belirgin bir iyileşme olduğunu göstermektedir. Enerji verimi %30,57'den %33,54'e yükselirken, ekserji

FULL TEXTS BOOK

verimi %58,46'dan %64,14'e çıkmıştır. Özellikle %60 ve üzeri NH_3 oranlarında bu artış daha da hızlanmaktadır. NH_3'nin yüksek özgül entalpisi ve düşük pompalama gereksinimi sayesinde, sistemin iç tersinmezlikleri azalmış ve çevrim ikinci yasa verimliliği artmıştır. Bu sonuç, karışım kullanımının çevrimi sadece nicel değil, **niteliksel (ekserji bazlı) olarak da geliştirdiğini** ortaya koymaktadır. NH_3 oranındaki artış, çevrimde hem enerji hem de ekserji bazlı performans göstergelerinde iyileşmelere yol açmıştır. Özellikle %60 NH_3 oranından itibaren bu artışlar daha belirgin hale gelmiş ve %90 oranında maksimum değerlere ulaşılmıştır. Bu durum, CO_2'nin tek başına kullanıldığı çevrimlere kıyasla NH_3 katkısının sistem verimliliğine olan olumlu etkisini ortaya koymaktadır.

5. SONUÇ

Bu çalışmada, 400 bar sistem basıncı ve 450°C türbin giriş sıcaklığı koşullarında farklı NH_3/CO_2 oranlarının kullanıldığı süperkritik kapalı çevrim termodinamik sistemin enerji ve ekserji performansları incelenmiştir. Yapılan analizlerden elde edilen başlıca sonuçlar aşağıda sunulmaktadır:

- NH_3 oranı arttıkça sistemde ihtiyaç duyulan kütlesel debi belirgin şekilde azalmıştır. %0 NH_3 oranında 90,34 kg/s olan kütlesel debi, %90 NH_3 oranında 22,68 kg/s'ye kadar düşmüştür. Bu azalma, NH_3'nin daha yüksek özgül entalpisi sayesinde aynı enerji taşınımı için daha düşük akışkan miktarına ihtiyaç duyulmasıyla açıklanmaktadır.
- Kütlesel debinin azalmasına bağlı olarak pompa güç tüketimi de önemli ölçüde azalmıştır. Saf CO_2 ile çalışıldığında 4625,99 kW olan pompa gücü, %90 NH_3 oranında 1649,54 kW seviyesine gerilemiş ve %64'lük bir düşüş sağlanmıştır.
- NH_3 oranı yükseldikçe sistemin net güç üretimi artış göstermiştir. %0 NH_3 oranında 12.393,89 kW olan net güç, %90 NH_3 oranında 13.599,76 kW'ye ulaşmıştır. Bu artış, pompa işinin azalması ve türbin çıkışındaki enerji dönüşümünün daha verimli olması ile ilişkilendirilmiştir.
- NH_3 oranı arttıkça rejeneratif ısı kazancı azalmıştır. Saf CO_2 ile 9965,23 kW olan bu değer, %90 NH_3 oranında 1586,62 kW'ye gerilemiştir. Bu durum, karışımın termofiziksel özelliklerinin rejeneratif ısı transfer kapasitesini sınırlamasıyla açıklanabilir.
- Enerji verimi, NH_3 oranındaki artışa paralel olarak yükselmiştir. %0 NH_3 oranında %30,57 olan enerji verimi, %90 NH_3 oranında %33,54 seviyesine ulaşmıştır.
- Ekserji veriminde gözle görülür bir iyileşme sağlanmıştır. %0 NH_3 oranında %58,46 olan ekserji verimi, %90 NH_3 oranında %64,14'e yükselmiştir. Bu artış, çevrimdeki tersinmezliklerin azalmasıyla doğrudan ilişkilidir.
- Elde edilen sonuçlar, NH_3 katkısının süperkritik çevrimlerde hem enerji hem de ekserji performansını iyileştirdiğini ortaya koymaktadır. Yüksek NH_3 oranları, sistemin yardımcı ekipman yükünü azaltmakta, net güç üretimini artırmakta ve verimlilik değerlerini yükseltmektedir. Özellikle %90 NH_3 oranında elde edilen sonuçlar, NH_3/CO_2 karışımının saf CO_2'ye kıyasla güçlü bir alternatif olduğunu göstermektedir.

Semboller ve Kısaltmalar

$\dot{m}$: Kütlesel debi (kg/s)

h: Özgül entalpi (kJ/kg)

s: Özgül entropi (kJ/kg·K)

$\dot{Q}$: Isı akışı (kW)

$\dot{W}$: iş (kW)

T_0: Çevre sıcaklığı

P_0: Atmosfer basıncı

FULL TEXTS BOOK

η: Verim

$\dot{E}$: Ekserji akış hızı (kW)

ε: Isı eşanjörü etkinliği

y: Karışım bileşen mol/kütle oranı

KAYNAKLAR

[1] Ahn, Y., Bae, S. J., Kim, M., Cho, S. K., Baik, S., Lee, J. I., & Cha, J. E. (2015). Review of supercritical CO_2 power cycle technology and current status of research and development. *Nuclear Engineering and Technology*, 47(6), 647–661. https://doi.org/10.1016/j.net.2015.06.009

[2] Chen, Y., Sun, F., & Wu, C. (2006). Thermodynamic analysis of an irreversible closed Brayton cycle. *Applied Energy*, 83(5), 495–510. https://doi.org/10.1016/j.apenergy.2005.04.001

[3] Dostal, V., Driscoll, M. J., & Hejzlar, P. (2004). *A supercritical carbon dioxide cycle for next generation nuclear reactors*. Massachusetts Institute of Technology.

[4] IEA. (2022). *World Energy Outlook 2022*. International Energy Agency. https://www.iea.org/reports/world-energy-outlook-2022

[5] IPCC. (2021). *Climate Change 2021: The Physical Science Basis*. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press.

[6] IRENA. (2021). *World Energy Transitions Outlook 2021*. International Renewable Energy Agency. https://www.irena.org/publications

[7] Kang, Y., Kim, J., & Lee, J. I. (2020). Performance evaluation of supercritical CO_2 cycles with ammonia as an additive. *Applied Thermal Engineering*, 179, 115694. https://doi.org/10.1016/j.applthermaleng.2020.115694

[8] Lee, J., Kim, T., & Kim, M. (2018). Thermodynamic performance evaluation of CO_2-based binary mixtures for power cycles. *Energy Conversion and Management*, 156, 464–475.

[9] Angelino, G., & Invernizzi, C. (1993). Investigation on the possibility of enhancing the efficiency of the Rankine cycle by means of a mixture of ammonia and water. *Energy*, 18(4), 329–336.

[10] Ceylan, V., & Özgür, A. E. (2021). Süperkritik Brayton Çevriminin Termodinamik Analizi. Mehmet Akif Ersoy Üniversitesi Fen Bilimleri Enstitüsü Dergisi, 12(Ek (Suppl.) 1), 500-509.

[11] Ma, N., Bu, Z., Fu, Y., Hong, W., Li, H., & Niu, X. (2023). An operation strategy and off-design performance for supercritical brayton cycle using CO_2-propane mixture in a direct-heated solar power tower plant. Energy, 278, 127882.

[12] Özahi, E., Tozlu, A., & Abuşoğlu, A. (2018). Organik Rankine çevrimi entegre edilmiş S-CO_2 kullanılan bir gaz türbin çevriminin termodinamik ve termoekonomik analizi. Gazi Üniversitesi Mühendislik Mimarlık Fakültesi Dergisi, 33(3), 917-928.

[13] Haroon, M., Sheikh, NA, Ayub, A., Tariq, R., Sher, F., Baheta, AT ve Imran, M. (2020). CO_2 Bazlı İkili Karışımla Çalışan Dip Güç Döngülerinin Ekserjetik, Ekonomik ve Eksergo-Çevresel Analizi Enerjiler,13(19), 5080. https://doi.org/10.3390/en13195080

[14] Akman, M. Gemilerde organik rankine çevrimine dayalı atık ısı geri kazanım sistemlerinin ileri termal analizleri ve termo-ekonomik optimizasyonu.

FULL TEXTS BOOK

[15] Uçar, İ. (2020).Sıvılaştırılmış doğal gazın (LNG) soğuk enerjisini kullanan yeni bir birleşik güç sisteminin enerji, ekserji ve eksergo-ekonomik analizi (Master's thesis, Necmettin Erbakan University (Turkey)).

[16] Liu, X., Wang, Y., & Chen, J. (2019). Thermodynamic Analysis of CO_2-Based Binary Mixtures in Power Cycles under High Pressure and Temperature Conditions. *Energy Conversion and Management*, 185, 257-267.

[17] Zhang, H., Li, M., & Zhao, F. (2021). Exergy and Energy Analysis of CO_2/NH_3 Binary Mixtures for Enhanced Power Cycle Performance. *Applied Thermal Engineering*, 183, 116196.

[18] Smith, R., Johnson, P., & Lee, S. (2018). Phase Behavior and Thermodynamic Properties of CO_2 Mixtures Relevant to Carbon Capture and Storage Applications. *International Journal of Greenhouse Gas Control*, 75, 44-55.

[19] Wang, D., & Lee, J. (2020). Thermodynamic and Phase Behavior Analysis of CO_2-H_2O Mixtures for Power and Refrigeration Cycles. *Renewable Energy*, 154, 1132-1140.

[20] Kim, S., Park, J., & Choi, H. (2022). Thermodynamic Properties and Energy Efficiency of CO_2-N_2 Mixtures in High-Pressure Power Cycles. *Journal of Cleaner Production*, 331, 129872.

[21] Çengel, Y. A., & Boles, M. A. (2015). *Thermodynamics: An Engineering Approach* (8th ed.). McGraw-Hill Education.

[22] Moran, M. J., Shapiro, H. N., Boettner, D. D., & Bailey, M. B. (2014). *Fundamentals of Engineering Thermodynamics* (8th ed.). Wiley.

[23] Dostal, V., Driscoll, M. J., & Hejzlar, P. (2004). *A supercritical carbon dioxide cycle for next generation nuclear reactors*. MIT-GFR-006.

YÜN LİF CİNSİNİN İPLİK MUKAVEMETİ ÜZERİNE ETKİSİ

THE AFFCET OF THE WOOL FIBER TYPE ON THE YARN TENSILE STRENGTH

***Tansu ÜNEN*[1]**

Bursa Teknik Üniversitesi, Mühendislik ve Doğa Bilimleri Fakültesi, Polimer Malzeme Mühendisliği, Bursa, Türkiye.

[1]ORCID ID: https://orcid.org/0009-0004-4175-8392

***Kenan YILDIRIM*[2]**

[2] Prof. Dr., Bursa Teknik Üniversitesi, Mühendislik ve Doğa Bilimleri Fakültesi, Polimer Malzeme Mühendisliği Bölümü, Bursa, Türkiye

[2]ORCID ID: https://orcid.org/0000-0002-1640-6035

ÖZET

Yün tekstil sanayisinde en yaygın kullanılan hayvansal liflerden biridir. İnsanlık tarihi boyunca keşfedilmiş ve dokuma materyali olarak kullanılmıştır. 2000'li yıllarda 1.4 milyar kilogram yün lifi kullanılmıştır. Son yıllarda sentetik elyaf kullanımının artmasına bağlı olarak yün liflerinin tekstildeki payı azalmış olsa da, bu doğal elyaf pazarda hâlâ önemli bir yer tutmaktadır. Yünün sahip olduğu bazı üstün doğal özellikler, onu diğer birçok elyaftan daha değerli kılmaktadır.Yün lif kalitesi, lif çapı, pulcuk yapısı ve lif uzunluğu bazında değişmektedir. Yünün özelliği büyük ölçüde lif çapına bağlıdır. Ticari olarak en fazla merinos yünü kullanılmasına karşın bir çok yün lif çeşidi vardır. Bu çalışmada yün lif türünün iplik mukavemetine etkisi araştırılmıştır. Çalışmada 17,5 Mic Yün, 18,5 Mic Yün, 20,5 Mic Yün, 24 Mic Yün, 27 Mic Yün, 30 Mic Yün, 34 Mic Alpaka, 34 Mic Mohair ve 38 Mic Lincoln olmak üzere 10 farklı yün çeşidi denemede kullanılmıştır. Yün elyafının ipliğe dönüştürülmesi endüstriyel ölçekte iplik üretim hattında yapılmıştır. En düşük iplik mukavemeti ve uzama oranı 24 Mikron merinos yününde elde edilirken, her iki parametrede de en yüksek değer 17,5 ve 18,5 Mikron merinos yünlerinde oluşmuştur. 24 Mikrondan daha kalın ve daha incelerde iplik mukavemeti ve uzama oranı artmıştır.

Anahtar Kelimeler: iplik kopma mukavemeti, yün lifi, merinos, alpaka, Mohair

ABSTRACT

Wool is one of the most widely used animal fibers in the textile industry. It has been discovered and utilized as a weaving material throughout human history. In the 2000s, approximately 1.4 billion kilograms of wool fiber were consumed. Although the share of wool in textiles has declined in recent years due to the increased use of synthetic fibers, this natural fiber still maintains a significant position in the market. Some of wool's unique and superior natural properties make it more valuable than many other types of fiber. The quality of wool fiber varies depending on fiber diameter, scale (cuticle) structure, and fiber length. Among these, fiber diameter is the most influential factor in determining the characteristics of wool. Although Merino wool is the most commercially used type, many other wool varieties are also available. This study investigates the effect of wool fiber type on yarn tensile strength. A total of 10 different types of wool were used in the experimental work: 17,5 Micron Wool, 18,5 Micron Wool, 20,5 Micron Wool, 24 Micron Wool, 27 Micron Wool, 30 Micron Wool, 34 Micron Alpaca, 34 Micron Mohair, 38 Micron Lincoln. The conversion of wool fibers into yarn was carried out on an industrial-scale yarn production line. The lowest yarn tensile strength and elongation values were obtained with 24 micron Merino wool, while the highest values for both parameters were recorded in

FULL TEXTS BOOK

17.5 and 18.5 micron Merino wools. It was observed that yarn tensile strength and elongation increased in both finer and coarser fibers compared to the 24 micron sample.

Keywords: tensile strenght of yarn, wool fiber, merino, alpaca, mohair.

GİRİŞ


Yün tekstil sanayisinde en yaygın kullanılan hayvansal liflerden biridir. İnsanlık tarihi boyunca keşfedilmiş ve dokuma materyali olarak kullanılmıştır. 2000'li yıllarda 1.4 milyar kilogram yün lifi kullanılmıştır. Son yıllarda sentetik elyaf kullanımının artmasına bağlı olarak yün liflerinin tekstildeki payı azalmış olsa da, bu doğal elyaf pazarda hâlâ önemli bir yer tutmaktadır. Yün, keratin adı verilen bir protein grubuna dahildir ve bu gruba aynı zamanda tırnak, boynuz, tüy ve diğer hayvansal koruyucu dokular da dahildir. Son yıllarda sentetik elyaf kullanımının artmasına bağlı olarak yün liflerinin tekstildeki payı azalmış olsa da, bu doğal elyaf pazarda hâlâ önemli bir yer tutmaktadır. Özellikle Avustralya, Yeni Zelanda, Güney Afrika ve Arjantin gibi ülkeler açısından yün, başlıca ihracat kalemlerinden biri olmayı sürdürmektedir. Yünün sahip olduğu bazı üstün doğal özellikler, onu diğer birçok elyaftan daha değerli kılmaktadır. Bu özellikler arasında kumaşın yumuşaklık hissi (dokunulduğunda sağladığı konfor) , yüksek oranda nem çekebilme kapasitesi (bu da konforu artırır) , nemi yapısında taşıyabilmesi, lifin doğal kıvrımlı yapısı, hacim kazandırarak içerisinde daha fazla hava tutulmasına olanak tanır; bu da üstün ısıl yalıtım sağlayarak sıcak ortamlarda serinlik, soğuk ortamlarda ise sıcaklık hissi yaratır. Bu özelliğin yanı sıra, yünün hava geçirgenliği (nefes alabilirlik) , elastikiyeti, koku emme kapasitesi, güç tutuşurluk, biyolojik olarak doğada çözünebilmesi, geri dönüştürülebilirliği ve kumaşın formunu estetik biçimde koruyabilmesi gibi birçok avantajı bulunmaktadır. Çizelge 1.1 , dünya genelinde yün üretimi ve yün elde edilen koyun sayısı bakımından önde gelen ülkeleri listelemektedir [1-5].

Çizelge 0.1 : Başlıca yün üreten ülkelerde yün üretimi ve koyun sayıları (2000/2001)

Ülke/Bölge	Yün üretimi (yağlı, 10^6 kg)	%	Koyun sayısı (10^6)
Avustralya	652	28,1	113
Çin	288	12,5	135,5
Yeni Zelanda	258	11,1	17,5
Güney Afrika	82	3,5	30
Arjantin	67	2,9	13
Uruguay	52	2,2	13
Birleşik Krallık	72	3,00	53,9
Türkiye	60	2,6	49,2
İran	74	3,2	50
Eski SSCB	128	5,5	508
Diğerleri	620	26,5	1006
Toplam	2324	100	-

FULL TEXTS BOOK

Yün lifinin enine kesiti, genel olarak dairesel ya da hafif oval bir forma sahiptir ve lif yapısı kökten uca doğru spiral şeklinde kıvrımlı bir düzen sergiler [2]. Yün lifi, üç ana bileşenden oluşur. Bu lifin kesitinde; en dış katman *epiderm* (kütikül) , orta kısımda *korteks,* en iç bölümde ise *medulla* adlı yapılar yer almaktadır (Şekil 1.1).



Şekil 1.1 : Yün elyafının iç yapısının görünümü [4].

Kütikül ya da epidermis tabakası; mikroskopla gözlemlendiğinde yün lifinin en dış kısmını oluşturan yapı olarak tanımlanır. Bu tabaka, lif eksenine belirli bir açıyla yerleşmiş, üst üste binen sert ve saydam hücrelerden meydana gelir. Balık ya da yılan pullarına benzetilebilecek görünüme sahip bu hücrelerin uçları dışa doğru çıkıntı yaparak lif yüzeyinde yönlü bir pürüzlülük oluşturur. Söz konusu pürüz, keçeleşme ve kohezyon özelliklerine katkı sağlarken, lifi belirli bir sertliğe de kavuşturur. Pulların yapısal özellikleri sayısı, biçimi, boyutları, kalınlığı ve dizilim şekli lifin genel performansını etkileyen önemli faktörlerdendir. Merinos koyunlarının ince liflerinde genellikle tek bir pul, tüm lifi kaplayacak şekilde gelişirken; daha kalın liflerde, çapla birlikte pul sayısının da arttığı gözlemlenir. Pul yapısının düzgün ve yüksek olması, yüzeyin düzgünlüğünü ve buna bağlı olarak lifin parlaklığını artırır. Epidermis katmanı üç ana bölümden oluşur: epikütikül, egzokütikül ve endokütikül. En dışta bulunan epikütikül tabakası oldukça ince yapısına rağmen, kükürt açısından zengin olup sistin bağları sayesinde kimyasal maddelere ve biyolojik etkilere karşı güçlü, su geçirmez (hidrofobik) bir bariyer görevi görür. Ancak bu tabaka, fiziksel etkilerle zamanla zarar görebilir. *Korteks tabakası*; yün lifinin yaklaşık %90 lık kısmını oluşturmaktadır. İğ şeklinde yaklaşık 100 µ uzunluğunda ve 2-4 µ genişliğindedir. Bu hücrelerin kesitleri çoğunlukla yuvarlak değil, düzensiz ve çokgen şeklindedir. *Medulla (Öz Bölgesi);* içi boş, hava dolu hücre duvarlarının oluşturduğu bir ağ yapısından oluşur. Bazı durumlarda bu yapılar tamamen çökerek içi boş bir tüp meydana getirebilir. Temel görevi lifin içerisinde hava boşluğu oluşturarak lifin koruyucu özelliğini iyileştirmektir. Kanal ne kadar kalın olursa liflerin eğilme ve bükülmeleri o kadar zorlaşır.

Bir yün partisine ait ortalama lif çapı, teknolojik açıdan değerlendirildiğinde, o yünün en kritik özelliklerinden biri olarak kabul edilir. Bu özellik, tam anlamıyla olmasa da, genellikle partinin "sınıf" veya "numara" değerlendirmesinde belirleyici rol oynar. Koyun yünlerinde ortalama çap, Merinos

FULL TEXTS BOOK

türlerinde 16–17 µ gibi ince değerlerden başlayarak, kaba uzun elyaflı ırklarda 40 µ a kadar çıkabilmektedir. Ancak, tek bir yün demeti bile çap açısından tamamen homojen değildir. Bireysel lifler arasında kalınlık farkları olabileceği gibi, bu farklılıklar hayvanın sağlık durumu ya da çevresel değişkenlere bağlı olarak da ortaya çıkabilir. Örneğin, çevresel stresin etkisiyle lifte meydana gelen "kopmalar", geçici ya da belirgin çap daralmalarına yol açabilir. Bu durum, "demet içi" (within-staple) varyasyon olarak tanımlanır ve bazı durumlarda çap değişim katsayısı yalnızca %14 gibi düşük seviyelere inebilir. Kaliteli Merinos yünlerinde ise çap değişim katsayısı (C.V.) genellikle %20–24 aralığında olur. Farklı kalite seviyelerine göre çap dağılımlarına dair bilgiler Çizelge 1.2 te sunulmuştur.

Çizelge 0.2 : Yün lifinin incelik derecesi [6].

S Değerleri	Ortalama Lif Çapı (µ)
80s	18,8 mikron
70s	19,7 mikron
64s	20,7 mikron
60s	23,3 mikron
58s	24,9 mikron
56s	26,4 mikron
50s	30,5 mikron
48s	32,6 mikron
46s	34,0 mikron
44s	36,2 mikron
40s	38,7 mikron
36s	39,7 mikron

Yünün mekanik özellikleri, diğer tüm tekstil liflerinde olduğu gibi son derece karmaşıktır. İlk olarak, lif *anizotropik* bir yapıya sahiptir; yani lif ekseni yönünde ve bu eksene dik düzlemde farklı özellikler sergiler. Ancak, lif ekseni boyunca olan özellikler daha önemli ve ölçülmesi daha kolay olduğundan, çalışmaların neredeyse tamamı bu yönde gerçekleştirilmiştir.

İkinci olarak, tüm viskoelastik malzemelerde olduğu gibi, ölçülen özelliklerin değerleri hem test süresine hem de fizikokimyasal çevreye bağlıdır. Örneğin, bir kuvvet-uzama testinde uzama hızı ya da yükleme oranı arttıkça, modül ve kopma gerilmesi artar, buna karşılık kopma uzaması azalır. Bu nedenle, farklı çalışmalardan elde edilen sonuçları karşılaştırmak zordur; özellikle bazı araştırmacılar sabit yükleme hızı, bazıları ise sabit uzama hızı kullandığında bu durum daha belirgin hale gelir. Tekstil teknolojisinde, "dayanıklılık" birimleri ise genellikle birim uzunluk başına kuvvet olarak tanımlanır ve *g/denye, g/tex, ya da km-ağırlık* gibi terimler sıkça kullanılır. Bu iki form, malzemenin yoğunluğu bilindiği sürece birbiriyle dönüştürülebilirdir [6].

Gerilme gevşemesi (stress relaxition) ve sürünme (creep)

Bir yün lifi hızlıca uzatılıp sabit bir uzama düzeyinde tutulduğunda, bu uzamayı korumak için gerekli kuvvet zamanla azalır bu olguya *gerilme gevşemesi (stress relaxation)* denir. Tersine, sabit bir yük

FULL TEXTS BOOK

uygulandığında lif önce hızlı, sonra ise çok uzun bir süre boyunca yavaş bir şekilde uzamaya devam eder; bu davranışa ise *sürünme (creep)* adı verilir. Bu kavramlar birçok araştırmacı tarafından kapsamlı biçimde incelenmiştir; çünkü bu tür davranışlar, malzemenin moleküler yapısı hakkında değerli bilgiler sunar [6].

Koyun türlerinden elde edilen liflerin yanı sıra, tekstil üretiminde önemli miktarda hayvansal lifler ve kürk lifleri de kullanılmaktadır. Bu lifler, yüne ilave edilerek kumaşa estetik görünüm, renk, yumuşaklık veya parlaklık gibi özellikler kazandırmak amacıyla tercih edilmektedir. Bu liflerin en geniş grubu, keçi ve deve gibi türlerden elde edilen ve Amerikan Yün Yasası kapsamında yün olarak sınıflandırılan özel hayvan kıllarından oluşur. Bunun dışında, inek, at ve çeşitli kürk hayvanlarından da lif temin edilmektedir. Kullanılan yün lif çeşitleri Çizelge 1.3 da verilmiştir.

Çizelge 0.3 : Yün lifi çeşitleri [6].



YÖNTEM

Çalışma kapsamında 10 farklı yün çeşidi kullanılmıştır. Her bir yün çeşidinden iplik kompozisyonunda %100 olacak şekilde iplikler üretilmiştir. Üretilen iplik kodları numune kodu olarak kodlanmıştır. Lincoln, mohair ve alpaka yünleri tek başlarına eğrilemediği için karışım olarak iplikler üretilmiştir. Karışımlarda 27 mikron yün kullanılmıştır. Üretilen iplik kod ve özellikleri Çizelge 2.1 de verilmiştir.

Çizelge 0.1 : İplik özellikleri ve kodları.

Numune Sayısı	Numune Kodu	İplik kompozisyonundaki yün cinsi(leri)	İplik Doğrusal Yoğunluğu (Nm)	Karışım oarnı (%)	
				Ana bileşen yün	27 Mic Yün
1	17,5 Mic Yün	17,5 Mic Yün	8,6	100	0
2	18,5 Mic Yün	18,5 Mic Yün	8,6	100	0
3	20,5 Mic Yün	20,5 Mic Yün	8,6	100	0

FULL TEXTS BOOK

4	24 Mic Yün	24 Mic Yün	8,6	100	0
5	27 Mic Yün	27 Mic Yün	8,6	100	0
6	30 Mic Yün	30 Mic Yün	8,6	100	0
7	Alpaka karışım	34 Mic Alpaka+ 27 Mic Yün	8,6	30	70
8	Mohair Karışım	34 Mic Mohair+ 27 Mic Yün	8,6	30	70
9	36 Mic Yün	36 Mic Yün	8,6	100	0
10	Lincoln Karışım	38 Mic Lincoln+ 27 Mic Yün	8,6	30	70

Boyama aşaması; Yünlerin tamamı bams halinde olduğundan bams boyama işlemine tabi tutulmuştur. Boyama süreci, 2003 model H. Krantz Marchinenbalt - 51 Aachen markalı bams boyama makinesinde gerçekleştirilmiştir. Boyama işlemi gerçekleşen bams formundaki yünler sıkma işlemine tabi tutulmuştur. Bu işlem, 2000 model Oğuz Makine marka santrifüj makinesi ile gerçekleştirilmiştir. Sıkma işleminden sonra gelen kurutma aşaması, 1990 model Enes marka kurutma makinesi kullanılarak gerçekleştirilmiştir. Boyama işlemi tamamlanan bams formundaki yün elyaflar, 2015 model GC 30 (NSC N. Scumberger) marka cer makineleri kullanılarak cer şeridi haline getirilmiştir. İnceltilerek karıştırılan cer şeritleri 1992 model HDB (Houget Duesberg Bosson) Belçika menşeli ring iplik makinesinde, yarı kamgarn yapıdaki 8,6 Nm tek kat ipliğe dönüştürülmüştür. Tek kat iplik üretiminin ardından, iplik üzerindeki kopuşların giderilmesi amacıyla, 2005 model Japon Muratec marka iplik temizleme makinesi kullanılmıştır. İplikler, 2021 model HMX 132 HemaksCo Textile Machinery marka otomatik katlama makinesi ile dört katlı forma getirilmiştir. 2002 model Volkmann marka çok katlı büküm makinesi ile ipliğe büküm verilmiştir. İpliklerin fikseleme işlemi, Süperba model fikse makinesi aracılığıyla uygulanmıştır. Su buharı ile gerçekleştirilen ısıl işlem sonrasında iplikler, kuka makinelerinde çalışmaya uygun kek formuna dönüştürülmüştür. Fikse işlemi tamamlanmış kek formundaki iplikler, yumak haline getirilebilmeleri amacıyla, 1985 model GW Maschınen Ag Wetzıkon marka kuka makinesinde kukalanmıştır.

Üretilen ipliklere kopma mukavemeti testi 2001 model I.V. Calderara marka mukavemet test cihazı kullanılarak cihazında TS EN ISO 2062 standardına göre yapılmıştır. Test işlemi 10 test örneği üzerinden yürütülmüştür.


BULGULAR

10 farklı yün lif çeşidi ile üretilen tek kat ipliklerin kopma mukavemeti ve kopma uzama oranları Şekil 3.1 ve Şekil 3.2 de verilmiştir. 17,5 – 24 mikron arasında iplik mukavemeti ve uzama oranı düşerken, 24 – 30 mikron arasında her iki özelliğinde arttığı görülmüştür. Lif inceliği küçüldükçe aynı iplik doğrusal yoğunluğuna sahip iplik kesit alanında daha fazla lif olacağından iplik mukavemetinin ve uzama oranının artması beklenen bir durumdur. En düşük mukavemet ve uzama oranı 24 mikron yün

FULL TEXTS BOOK


lifinde oluşmuştur. Buna ise 24 mikron yün lif boyunun en kısa olmasının neden olduğu görüşüne varılmıştır. Çünkü iplik mukavemeti ve kopma uzama oranını etkileyen en önemli faktörlerin lif inceliği ve uzunluğu olduğu bilinmektedir [7-9].

18.5 mikron yün lifi en uzun boya sahip olmasından dolayı en yüksek iplik mukavemeti bu elyafta gerçekleşmiştir. Her ne kadar 27 ve 36 mikron yün elyafının lif boyu 18.5 mikron yün elyafından büyük olsa da lif çaplarının büyük olması nedeniyle iplik kesitinde daha az lif bulunmasından dolayı iplik mukavemetleri düşük olmuştur. 27, 30 ve 36 mikron liflerden elde edilen iplik mukavemetlerinin 24 mikron yün elyafından üretilen iplik mukavemetinden daha yüksek olması ise lif uzunluğuna bağlıdır. Her ne kadar iplik kesitinde daha az lif bulunsa da lif uzunluğunun fazla olması nedeniyle lifler arası sürtünme kuvvetinin daha fazla olmasından dolayı iplik mukavemet ve uzama değerleri yüksek olmuştur. Alpaka ve mohair karışımlarında iplik mukavemeti 24 mikron yün mukavemetine yakın gerçekleşmiştir. Lincoln karışımlarında ise iplik mukavemeti artmıştır. Yün lifi inceldikçe aynı iplik doğrusal yoğunluğuna sahip iplik kesitinde daha fazla lif yer alacağından iplik mukavemetinin artması öngörüsü ile örtüşmüştür. Alpaka, mohair ve Lincoln yünleri 27 mikron yün ile karıştırılarak ipliğe dönüştürülmüştür. Tüm karışımların iplik mukavemetinin 27 mikron yünden üretilen iplik mukavemetinden daha düşük olduğu görülmüştür. Bunun sebebinin lif inceliği olduğu düşünülmektedir. Her ne kadar lif uzunlukları 27 mikron yünden uzun olsa da lif kalınlığı fazla olduğundan iplik kesitinde daha az lif olacağından iplik mukavemetleri düşmüştür. Ancak uzama oranı bazında mohair ve Lincoln karışımlarının uzama oranlarının daha yüksek olduğu görülmüştür. Uzama oranındaki bu durumun lif uzunluğundan kaynaklandığı düşünülmektedir. Mohair ve Lincoln yünlerinin lif uzunluğu 27 mikron yün lifinden daha uzun olduğundan ve lif uzunluğu arttıkça da iplik kopma sürecinde liflerin birbirlerinden ayrılma mesafesi artacağından kopma uzaması daha yüksek olmuştur. Alpaka lif uzunluğu ise 27 mikron yün lifinden daha kısadır. Bu nedenle bu lif karışımından üretilen ipliğin kopma uzama oranıda 27 mikron liften üretilen iplikten düşük olmuştur [10-14].



Şekil 3.1 : İplik kopma mukavemeti (cN/tex).

FULL TEXTS BOOK



Şekil 3.2 : İplik kopma anındaki uzama oranı (%).

KAYNAKLAR

1. Textile Terms and Definitions, 1996, 10th ed., Textile Institute, Manchester, England.
2. Wool Statics, 2000-2001, International Wool Textile Organisation (IWTO), London.
3. Mark, H. F. Acrylic Fibers. Encyclopedia of Polymer Science And Technology, (3rd ed., Vol. 1, pp. 135-136, Vol. 9, pp. 1-35, Vol. 10, pp. 616-617). Wool. (3rd ed. Vol. 12, pp. 546-580).
4. Türk, M. (2019). Yün ve meta-aramid ipliklerinden oluşan dokuma kumaşların özelliklerinin incelenmesi. (Yüksek lisans tezi). Bursa Uludağ Üniversitesi, Fen Bilimleri Enstitüsü, Tekstil Mühendisliği Anabilim Dalı, Bursa.
5. Babaoğul, M., Şener, A. & Öztop, H. (2010). Yün. Tekstil lifleri; Temel özellikleri, kullanım ve bakım. 111-145, 209, Akrilik ve Modakrilik Lifler. 235-240.
6. Acar, S. (2010). Yünlü giysi tasarımında bölgesel keçeleştirme yöntem ve uygulamaları. (Sanatta yeterlilik tezi). Dokuz Eylül Üniversitesi, Güzel Sanatlar Enstitüsü, Tekstil ve Moda Anabilim Dalı, İzmir.
7. Krause H.W ve Soliman H.A, melliand textileberichte 52 (1971), 497-501
8. Zeidman, Mishu & Sawhney, Paul. (2002). Influence of Fiber Length Distribution on Strength Efficiency of Fibers in Yarn. Textile Research Journal - TEXT RES J. 72. 216-220
9. Artzt P. ve Egbers G., (), Open End Teknolojisi, Çv. Tuğrul Madran, Kemal Matbaası, Adana)
10. B.P. Saville, (1999), Physical Testing of Textiles, Woodhead Publishing, 44-76
11. J. E. Booth, (1969), Principles of Textile Testing An Introduction to Physical Methods of Testing Textile Fibres, Yarns and Fabrics, Chemical Publishing Company, 167-168
12. Altas S., Kadoğlu H., (2006), Determining Fibre Properties and Linear Density Effect on Cotton Yarn Hairiness in Ring Spinning, Ege University, 14, 48-51
13. V. K. Kothari; S. Dhamija; R. K.Varshney, (2011), Influence of Polyester Fibre Fineness and Cross-Sectional Shapes on Tensile Properties of Yarns, *Research Journal* of Textile and Apparel, 15 (3): 94–106.
14. R. D. Parsi1, Madhuri V. Kakde2, Kuldip Pawar3, R. S. Police Patil4, Influence of Fibre Length on Ring Spun Yarn Quality Centre for Textile Functions, SVKM'S, NMIMS, MPSTME, Shirpur, India

FULL TEXTS BOOK

TÜRKİYE'DE YAPILAN ARAPÇA-TÜRKÇE, TÜRKÇE-ARAPÇA SÖZLÜK ÇALIŞMALARI

ARABIC-TURKISH AND TURKISH-ARABIC DICTIONARY STUDIES CONDUCTED IN TURKEY

Sinan YILMAZ

Trabzon Üniversitesi, İlahiyat Fakültesi, Trabzon, Türkiye.

0009-0007-6708-0462

ÖZET

Sözlükler anlamı bilinmeyen kelimelerin öğrenilebilmesi için müracaat edilen ilk kaynaklardandır. Nitekim Arapça-Türkçe, Türkçe-Arapça sözlükler genellikle Arapça bilmeyen kişilerin bilmedikleri Arapça kelimelerin anlamlarını öğrenmek için başvurdukları ilk kaynaklardan birisidir. Bu bağlamda Türkiye'de yapılan Arapça-Türkçe, Türkçe-Arapça olarak hazırlanan bazı sözlüklerin tanıtımını gerçekleştirdik. Bu konuda Türkiye'de Yrd. Doç. Dr. Ferhat Yılmaz, Doç. Dr. Tahsin Deliçay'ın "Tek Dilli ve iki Dilli Sözlükler (Arapça-Türkçe)", Hüseyin Günday'ın "Arapça-Türkçe Leksikografiye Dair Bir Tenkit Yazısı: Serdar Mutçalı'nın Arapça-Türkçe Sözlük Örneği" ve Hacı Ahmet Elbeyli Muhammed Akif'in "Serdar Mutçalı'nın Modern Arapça Sözlüğünü İnceleme" adlı çalışmalar yapılmıştır fakat bu çalışmalar sadece ilgili sözlüğün ismi ve yazarı verilerek oluşturulması, sözlüğün tanıtımının detaylı şekilde yapılmaması, tenkit yazısı niteliğinde olması veya tek bir sözlüğün tanıtımı şeklinde olmasından dolayı bu konuda bir eksiklik olduğunu tespit ettik. Türkiye'de Arapça-Türkçe ve Türkçe-Arapça şeklinde hazırlanmış çok sayıda sözlük bulunmaktadır. Bu sözlükler, farklı dönemlerde, çeşitli ihtiyaçlara cevap vermek amacıyla ve farklı yöntemsel yaklaşımlar gözetilerek kaleme alınmıştır. Söz konusu çeşitliliğin ve zenginliğin daha yakından anlaşılabilmesi amacıyla bu çalışmada, farklı usûllerde hazırlanmış on ayrı sözlük ele alınmıştır. Her bir sözlüğün yazarı, yayımlandığı tarih, yayınevi gibi temel bibliyografik bilgilerine yer verilmiş; ayrıca her sözlüğün hazırlanış yöntemi ve kullanım amacı çerçevesinde tanıtıcı bilgiler sunulmuştur. Bununla yetinilmeyerek, her bir sözlükten bazı örnek kelimeler seçilerek sözlüğün içerik yapısı ve kullanıcılara sağladığı katkılar ortaya konulmaya çalışılmıştır. Bu sayede okuyucunun yalnızca bibliyografik düzeyde bilgi edinmesi değil, aynı zamanda sözlüklerin içeriksel özellikleri ve dil öğretimindeki işlevleri hakkında da somut ve karşılaştırmalı bir fikir edinmesi hedeflenmiştir. Böylelikle sözlüklerin yapısal farklılıklarını, içerik zenginliklerini ve kullanım pratiklerini göz önüne sererek, okuyucunun zihninde her bir sözlükle ilgili daha net ve kapsamlı bir tablo oluşması sağlanmaya çalışılmıştır.

Anahtar Kelimeler: Arap Dili ve Belâğatı, Sözlükbilim, Türkiye, Kâmus, Arapça Sözlük.

ABSTRACT

Dictionaries are one of the first sources to learn words whose meanings are unknown. As a matter of fact, Arabic-Turkish, Turkish-Arabic dictionaries are usually the first sources that people who do not know Arabic apply to learn the meanings of the words they lack. In this context, we have introduced some Arabic-Turkish and Turkish-Arabic dictionaries prepared in Turkey. In this regard, Assoc. Prof. Dr. Ferhat Yılmaz, Assoc. Prof. Dr. Tahsin Deliçay's "Monolingual and Bilingual Dictionaries (Arabic-Turkish)", Hüseyin Günday's "A Criticism on Arabic-Turkish Lexicography: The Case of Serdar Mutçalı's Arabic-Turkish Dictionary" and Hacı Ahmet Elbeyli Muhammed Akif's 'Review of Serdar Mutçalı's Modern Arabic Dictionary', but we have determined that there is a deficiency in this regard

FULL TEXTS BOOK

because these studies are created only by giving the name and author of the relevant dictionary, the introduction of the dictionary is not made, there is a criticism article or the introduction of a single dictionary. Since there are many dictionaries prepared in Arabic-Turkish and Turkish-Arabic in Turkey, in this study, after giving introductory information about the author, publication date, publishing house, and the method of the dictionary of ten dictionaries prepared in different methods, we tried to draw a picture about that work in the reader's mind by bringing examples of the words in the dictionary.

Keywords: Arabic Language and Rhetoric, Lexicology, Turkey, Kamus, Arabic Dictionary.

GİRİŞ

Sözlük kelimesinin Arapça karşılığı olan "lügât"; söz söylemek, boş konuşmak anlamındaki "lağv" kelimesinden türemiştir. Yine dilimizde kullanılan başka bir kelime olan "mu'cem"; sözün kapalı ve anlaşılmaz olması anlamındaki "ucme" kökünden, "kâmus"; kelimesi ise "suya dalmak" anlamındaki "kams"tan türemiş ayrıca kâmus kelimesi okyanus anlamını da ifade etmektedir. H.4. yüzyılın ortalarından sonra Arap müellifleri kâmus kelimesini eserlerinde kullanmışlar hatta ülkemizde dahi kâmus, lügât ve mu'cem kelimeleri eserlerde kullanılmıştır.[1]

Lügât, kâmus, mu'cem ve sözlük adında Türkiye'de yapılan Arapça-Türkçe, Türkçe-Arapça sözlüklerin incelenmesi ve tanıtımı hakkında yeterli çalışmanın olmadığını tespit ettik. Bu çalışmada ülkemizdeki tüm Arapça-Türkçe, Türkçe-Arapça sözlük eserlerini incelemek ve tanıtmak çalışmanın hacmini fazlasıyla genişleteceğinden dolayı sözlüklere müracaat eden kişilere sözlüklerin usûl ve üslûpları bakımından fikir olması amacıyla on farklı sözlük incelenecektir. Seçmiş olduğumuz sözlükleri birbirinden farklı usûllerde hazırlanmış olmasına özen gösterdik. Çalışmamızdaki eserler Türkçe ve Arapça alfabetik sıraya göre dizilmiş, 2000'li yılların öncesinde yazılmış modern sözlük olarak nitelendirebileceğimiz bir eser, cep lügâtı niteliğinde hacmi küçük olmasına rağmen fazlasıyla istifade edilecek bir sözlük, içerisinde küçük resimlerle kelimeyi anlaşılır kılan 1960'lı yıllara ait bir kâmus, sadece Arapça fiilleri ihtiva eden sözlükler gibi farklı farklı özelliklerde eserler incelenecektir.

Yapacağımız çalışmada eserin yazarı, yayın yeri, yayın tarihi ve yayıncısı hakkında bilgi verdikten sonra yazarın sözlüğü oluştururken yararlanmış olduğu kaynaklar da zikredilecektir. Yazar eserini oluştururken hangi sıraya göre kelimeleri dizdiğini, isim ve fiilleri nasıl sözlüğünde ifade ettiğini anlattıktan sonra eserin mümeyyiz özelliklerine değinilecektir. Bulabildiğimiz kadarıyla eserde var olan isim, fiil ve toplam kelime sayısı hakkında da bilgi verilecek, anlatacak olduğumuz bu bilgilerin okur tarafından daha iyi kavranması adına ilgili eserdeki fiil, isim varsa kalıp ifadeler ve eserdeki resimlerden örnekler de getirilecektir.

ARAŞTIRMA VE BULGULAR

1-Arapça-Türkçe Büyük Lügât

Bu sözlük çalışması Hüseyin Atay, İbrâhim Atay, Mustafa Atay tarafından yapılmış ve 1964 yılında Ankara'da Bayrak Matbaası tarafından basımı gerçekleşmiştir. Sözlük toplamda üç ciltten oluşmakta ve 24 cm. boyutundadır.

Bu lügât hazırlanırken kelimelerin kökü esas alınmış ve alfabetik sıraya riayet edilmiştir. Örneğin علم'ye bakılacağı zaman öncelikle ع sonrasında ل ve م harflerine bakılmalıdır. Bu sıranın ardından kelimeler mazi, muzari ve mastar şeklinde yerleştirilmiştir. Mastarlar birden fazla olabileceği için en çok kullanılanları verilmiş, eğer mastar değiştiğinde anlam da değişiyorsa bir alt satıra geçilerek "– "işareti ile o mastar ve anlamı getirilmiştir.[2] Mesela (فَقِهَ ـَـ) (فَقَهًا ، فُقُوهًا ، فَقَاهَةً kelimesi "pek az olmak;

[1] İsmail Durmuş, "Sözlük", TDV İslâm Ansiklopedisi (Erişim 01 Aralık 2024).

[2] Hüseyin Atay-İbrâhim Atay-Mustafa Atay, *Arapça-Türkçe Büyük Lügat* (Ankara: Bayrak Matbaası, 1964), 1/XXVIII.

FULL TEXTS BOOK

değersiz, kaba" manalarında veriliyor ve başka bir mastarı farklı anlam ifade ettiği için bir alt satırda تُفُوهًا (—) "Ahmak, akılsız" şeklinde veriliyor.[3] Fiiller üçüncü tekil şahsa göre yazılmış ve fiille beraber kullanılan harf-i cerler de verilmiştir. Ayrıca her harfin kelimeleri verilmeden önce harfin kendisiyle ilgili bilgi verilmektedir. Mesela ع harfi hakkında: Ayın harfi, boğazın ortasından çıkan açık sesli ve ğayından daha az yumuşaktır. Elifba'da 18. harf olup, ebced sırasında 70 adedine delalet eder, açıklaması yapılmaktadır.[4]

Kelimeler türetilirken kelimeye birtakım harfler gelmektedir. Bazı harfler o kadar kaynaşmıştır ki kelimenin aslından zannedilebilir. Bu sözlükte o şekilde olan kelimler türemiş haliyle verilmiş ve kelimenin aslı da parantez içerisinde belirtilmiştir. Örneğin اَضَابِير ج اِضْبارة (ضبر) kelimesi verilmiş ve parantez içindeki ضبر kelimesi اِضْبار'ın kökünü göstermiştir. Kelimenin son harfinin vâv mı yoksa yâ mı olduğunu belirtmek için parantez içerisinde (الو) ve (الى) getirilmiştir. Ayrıca kelimelerin anlamları daha iyi anlaşılsın diye Kur'ânı-ı Kerîmden âyet, hadis-i şerif, beyt ve atasözlerinden misaller verilmiştir.[5] Mesela حَرّفَ تَحريفًا "yan yapmak, çevirmek, bükmek" anlamındaki kelime açıklanırken manası daha iyi anlaşılsın diye "يُحَرِّفُونَ الْكَلِمَ عَنْ مَوَاضِعِهِ" sözleri yerinden kaydırıyorlardı, oynatıyorlardı"[6] âyeti verilmiştir.

Fiillerin lâzım manası önce zikredilmiş varsa müteaddî anlamı verilmiştir. Eğer bu fiil tek meful ile kullanılıyorsa (ه) zamiri ile, iki meful ile kullanılabiliyorsa (ه ه) zamirleri ile gösterilmiştir.[7] حَبَّ (حَبًّا)ُ - هـ "doldurmak" fiilinde olduğu gibi.[8]

Sözlükte öz Arapça kelimeler gösterilmek üzere "❋" işareti, aslı Arapça olmayan (muarreb) kelimeler için ise "★" işareti kullanılmıştır. (Halifesi olmak; yerine geçmek خَلَفَ يَخْلُفُ ❋ خِلَافَةً)[9] (büyük kazan ★ خَلَاقِين ج خَلْقِين)[10] Daha önce sözlükte geçen kelimeyi belirtmek üzere "– "işareti kullanılmıştır. Bazı fiillerin mecaz anlamları parantez içinde (mec) ile gösterilmiş, parantez içinde (zıt) yazan kelime bir önceki mananın zıddını ifade etmiştir. İsimlerin cemisi gösterilmek üzere ج = جمع kullanılmış, ج harfinden sonraki kelime öncesinin çoğulu olacak şekilde tertiplenmiştir. Çoğulun çoğulu var ise جج = الجمع جمع harfleriyle ifade edilmiştir. Sözlükte erkek ismi önce zikredilmiş ve dişi gösterilmek için de م = مؤنث kullanılmış, bazı cins isimlerin tekillerine ihtiyaç duyulduğunda ise مف = مفرد ile ifade edilmiştir.[11]

2-Arapça-Türkçe Yeni Kâmus

Bekir Topaloğlu ve Hayreddin Karaman tarafından hazırlanan "Arapça-Türkçe Yeni Kâmus" Nesil yayınları tarafından 1966 yılında İstanbul'da basımı yapılmıştır. Sözlük toplamda 535 sayfadan oluşmakta ve içerisinde her harfin kelimelerinden sonra küçük resimler bulunmaktadır.

Bu sözlük 20.000 civarında kelimeyi ihtiva etmektedir. Topaloğlu ve Karaman bu çalışmayı hazırlarken başta *el-Mu'cem'ul-Vasît* olmak üzere *Muhît'ul-Muhît, el-Muhtâr, el-Müncid, er-Râid, Müncid'ut-Tullâb* gibi lügâtlerden faydalanmışlardır. Yöntem olarak sözlükte kelimeler Arapça alfabetik sırada verilmiştir. Her kelimenin önce aslı (kök harfi alınmış) ve ilk harfinden son harfine doğru sıra takip edilerek dizilmiştir. Müştaklar (türemiş kelimeler) fiil ve isim olmak üzere ikiye ayrılmaktadır. Sözlükte müştak fiiller eğer mücerrediyle aynı manada ise alınmamış, farklı bir mana ihtiva ediyorsa eklenmiştir.


[3] Atay vd., "تفل", 1/198.
[4] Atay vd., "ع", 3/259.
[5] Atay-Atay-Atay, 1/XXIX.
[6] en-Nisâ 4/176.
[7] Atay vd., 1/XXX.
[8] Atay vd., "ح", 1/321.
[9] Atay vd., "خلف", 1/486.
[10] Atay vd., "خلق", 1/488.
[11] Atay vd., 1/XXX.

FULL TEXTS BOOK

Müştak isimler ise kıyasî olanlar (ism-i fâil, ism-i mef'ul gibi) sözlüğe eklenmemiş, semaî olan sıfat-ı müşebbehelerle ism-i fâil, ism-i mef'ul manasında bazı kelimeler köklerinden sonra hususî işaretlerle yazılmış fakat manaları kök manadan çıkartılabileceği için yazılmamıştır. İsimlere gelince harf-i tarif ile yazılmış ve en çok kullanılan mükesser cemileri de eklenmiştir. Manaya tesir eden harf-i cerler ya fiil ve isimlerle beraber veya manalarının başında hususi işaretlerle yazılarak gösterilmiştir.[12]

Topaloğlu ve Karaman eserin özellikleri kısmında talebenin kelimeleri kolay bulabilmesi için fiillerde en çok kullanılan müştak binaları ve anlam değişimlerini vermektedir. Ayrıca eserin başında "Yeni Kâmus" ta kullanılan işaretler de açıklanmaktadır.

Anlattığımız özellikler çerçevesinde sözlüğün kullanımına dair bazı örnekleri vereceğiz: (البَدِيلُ): (ج) أَبْدَالٌ، و بُدَلَاء (=) (البَدَنُ): Gövde.[13] Bir isim harf-i tarif ile getirilmekte ardından "ج" harfi ile cemi mükkeseri verilmektedir. Eğer kelimenin eş anlamlı başka bir karşılığı varsa da "=" işareti ile ifade edilmektedir. Bir fiile örnek verecek olursak ضَلَعَ (-َ ضَلْعًا): Eğrilmek (-an) Sapmak, meyletmek.[14] Fiilin mazisi verildikten sonra muzarisinin harekesi "–َ" bu şekilde ifade ediliyor ve ardından masdarı getiriliyor. Fiilin harf-i cer ile kullanımı "(-an)" buradaki gibi latinize edilerek belirtiliyor. Ayrıca her harfin sonunda resimlerle beraber bazı kelimeler de verilmektedir.[15]


3- El-Mevarid Türkçe-Arapça Lügât

Mevlüt Sarı tarafından hazırlanan 1703 sayfa olan *"El-mevarid Türkçe-Arapça Lügât"* 1975 yılında Bahar Yayınları tarafından basılmıştır. Sözlük Türkçe alfabetik sisteme göre a'dan z'ye göre tertip edilmiştir. Sözlüğün başlangıcında kullanımı kolaylaştırsın diye içerisindeki bazı kısaltmalar açıklanmıştır.

Sözlükte bir harfe başlanırken o harf ile başlayan ilçe, şehir isminden bir tanesi fotoğrafıyla beraber verilmiş, ardından o harfin Türkçede kaçıncı harf olduğu Arapça şekilde ifade edilmiştir. Mesela "ç" harfi için Çengelköy'ün fotoğrafı verilmiş ve من الافباء الحرف الرابع التركية açıklaması yapılmıştır.[16] Sözlükteki kelimeler fiil için mazi, muzari ve mastar; isim için müfred ve cemisi şeklinde verilmemiştir. Kelimelerin Türkçe karşılığının karşısında Arapça kalıp şeklinde verilmiş, Türkçe verilen kelimenin Türkçe açıklaması yapılmış ve o kelimenin hangi dilden geçtiği de kısaltma ile ifade edilmiştir. Mesela "gidon" kelimesi verilmiş yanına "fr" kısaltması ile Fransızca olduğu belirtilmiş, "bisikletin iki elle idare edilen yer" şeklinde Türkçe açıklanmış ve kelimenin Arapçası "مِقْوَدٌ الدَّرَّاجَة" verilmiştir.[17] Bazı yerlerde ise kelimenin Türkçesi ve Arapçası verilmiş daha iyi anlaşılsın diye de küçük resimler eklenmiştir.[18] Bir kelimeden sonra aynı anlamda bir kelime getirilecekse "–" işareti koyularak diğer kelime getirilmiştir. Mesela رَأْسٌ kelimesi verilmiş ve baş: kafa şeklinde Türkçe kelime yine Türkçe olan bir kelime ile açıklanmış ve bu kelimenin diğer anlamları –: yönetici, –: temel şeklinde getirilmiştir.[19]

[12] Bekir Topaloğlu- Hayreddin Karaman, *Yeni Kâmus* (İstanbul: Ahmet Sait Matbaası, 1966), VI.
[13] Topaloğlu-Karaman, "بدل", 19.
[14] Topaloğlu-Karaman, "ضغن", 237.
[15] Topaloğlu-Karaman, 515.
[16] Mevlüt Sarı, *El-Mevarid Türkçe-Arapça Lügât* (İstanbul: Bahar yayınları, 1975), "ç", 93.
[17] Sarı, "gıybetçi", 195.
[18] Sarı, "قفل", 32.
[19] Sarı, "bastırmak", 51.

FULL TEXTS BOOK

4- Türkçe-Arapça, Arapça-Türkçe Cep Lügâtı

Prof. Dr. Erol Ayyıldız tarafından hazırlanan *"Türkçe – Arapça, Arapça – Türkçe Cep Lügatı"* 1986 yılında Bayrak Matbaacılık tarafından baskısı yapılmıştır. İsminden de anlaşılacağı gibi bu bir cep lügâtı olduğundan fazla ayrıntıya yer verilmemiştir. Mücerred fiillerin mazi, muzari ve en çok kullanılan sadece bir mastarı verilmiş (نَشَرَ - يُ نَشْرًا : yaymak, dağıtmak, yayımlamak)[20], mezid fiillerin ise muzari ve mastarları kurala bağlı olduğundan dolayı sadece mazisiyle yetinilmiştir. (تَبَسَّمَ : tebessüm etmek)[21]. İsimler alışılagelmiş sözlük üslûbuna göre müfredi verilmiş ve ج harfinin sağında cemisi getirilmiştir. (نَادِرَةٌ ج نَوَادِرُ: fıkra, güzel söz).[22]

Sözlük sağdan sola Arapça harflere, soldan sağa da Türkçe harflere göre tertip edilmiştir. Sözlüğün sol başlangıç tarafında "ek bölüm" başlığı altında günlük hayatta faydalı olacak سُرُور بِكُلِّ / memnuyetle - bi külli sürûrin, نَعَم يُوجَد / evet bulunur – neam yûcedü gibi kalıp ifadelere yer verilmiş ve bu ifadelerin okunuşları latinize edilmiştir.[23]

5- Modern Arapça Sözlük

İ. Muhammed Yolcu ve Nurullah Çetin'in hazırladığı "Modern Arapça Sözlük" 1992 yılında Birleşik Yayınlarından çıkmıştır. Sözlük soldan sağa Türkçe alfabetik düzene göre, sağdan sola da Arapça alfabetik düzene göre tertip edilmiştir. Türkçe – Arapça kısmına 1. Bölüm, Arapça – Türkçe kısmına da 2. Bölüm denilmiştir. Türkçe alfabetik sıraya göre tertip edilen kısmın sonunda bazı ülke, şehir ve yer adlarının طهران / Tahran, سيواس / Sivas, البرتغال / Portekiz gibi[24] Arapça yazılışlarına yer verilmiştir. Ayrıca sözlüğün Arapçadan Türkçeye olan kısmının başında اَنْتَ تَحْصُدُ مَا زَرَعْتَ / ne ekersen onu biçersin, لِكُلِّ صَارِمٍ نَبْوَةُ / her kılıç körleşir[25] gibi seçme Arapça atasözlerine yer verilmiştir.

Bu sözlüğün içeriği modern sözlük olmasından dolayı Arapça dergi, gazete, broşür gibi basılı yayınlardan taranarak oluşturulmuştur. Sözlük alışageldiğimiz gibi bir Arapça – Türkçe sözlük değildir. Yani verilen kelimler fiil ise mazi, muzari ve emir şeklinde, isim ise müfredi ve cemisi şeklinde değil, çoğunlukla وِزَارَة الْخَارِجِيَّةُ / dış işleri bakanlığı, مُنَافَسَة اَجْنَبِيَّة / dış rekabet[26] gibi kalıp ifadelerden oluşmaktadır. Sözlükteki maddeler yazılı metinlerde kullanıldığı gibi değiştirlmeden eklendiğinden dolayı müzekker-müenneslik, tekil-çoğulluk ve fiil kipleri gibi şeyler aynı formatta değil, farklılık arz etmektedir.

6- Örnek Sözlük Arapça -Türkçe

M. İsmet Yolcu tarafından hazırlanan "Örnek Sözlük" Eğit-San Yayınları tarafından 1996 yılında yayınlanmıştır. Sözlük Arapça harf tertibine göre sıralanmış, mücerred fiillerin mazi, muzari ve mastarı, mezid fiillerin ise mazi ve mastarı verilmiştir. Sözlük içerisindeki kelimeler lam-ı tarifsiz şekilde getirilmiş ve sözlükte yaklaşık 60000 kelime bulunmaktadır.[27]

Sözlüğün sonunda fiillerin harfi cer ile kullanımını veren bir liste, sarf ve nahivde genel kaideleri ihtiva eden tablolar, özellikle edebiyat fakültesinde okuyanların istifade edebileceği Arap şiir vezinlerinin bulunduğu bir bölüm ve bazı hayvan, bitki isimlerinin resimlerle gösterildiği sayfalar bulunmaktadır.[28]

[20] Erol Ayyıldız, *Türkçe – Arapça, Arapça – Türkçe Cep Lügâtı* (Ankara: Bayrak Matbaası, 1997), "نصب", 151.
[21] Ayyıldız, "taziye etmek", 175.
[22] Ayyıldız, "نخل", 150.
[23] Ayyıldız, II.
[24] İ. Muhammed Yolcu- Nurullah Çetin, *Modern Arapça Sözlük* (Ankara: Birleşik yayınları, 1992), 1/467.
[25] Yolcu- Çetin, 2/3.
[26] Yolcu- Çetin, 1/113.
[27] M. İsmet Yolcu, *Örnek Sözlük Arapça-Türkçe* (Ankara: Eğit-San yayınları,1996),5.
[28] Bk., Yolcu, 888-936.

FULL TEXTS BOOK

Mesela mücerred bir fiil "خَلَص ـُ خلوصًا و خُلاصًا / Safi olmak, arı olmak, halis olmak"[29] mezid fiil " : أَغَارَ (على) إغَارَةً / Saldırmak, akın etmek, süratle girmek, dört nala gitmek, alçak bir yere inmek, düşmana saldırmak / على العدوّ"[30] isim ise "جَامِع ج جَوَامِع / Toplayan, derleyen, cemeden, içine alan, havi, cami"[31] şeklinde ifade edilmiştir.

7- El-Beyan Arapça- Türkçe Büyük Sözlük

Arif Erkan tarafından hazırlanan bu sözlük 2000 yılında tamamlanmış ve Yasin Yayınevi tarafından basılmıştır. Erkan sözlüğünü oluştururken *er-Raid'de* olduğu gibi Arapça alfabetik sıraya göre kelimeleri dizmiştir. Çalışması esnasında *el-Muc'em'ul Vasît, Tertîb'ul-Kâmûs'ıl-Muhît, Lisân'ul-Arab* ve *Ahterî-i Kebîr* gibi eserlerden faydalanmıştır. Sözlükteki fiiller mazi, muzari ve mastar şeklinde sıralanmış, mezid bablarda da kelimenin kökü parantez içerisinde belirtilmiştir. Kelimelerin anlamları verildikten sonra cümle içinde kullanışına dair bir örnek getirilmiş, varsa o kelimeyle ilgili kalıplaşmış bir ifade veya deyim bir, iki şeklinde sıralanarak verilmiştir. Ayrıca kelimenin anlamına zıt bir anlamı varsa parantez içerisinde "zıt" ile o anlamı ifade edilmiştir. Bu eserde yaklaşık olarak 250 bin kelime bulunmaktadır.[32] Sözlükte bulunan isimlerin ise müfredi verilmiş ج harfinin ardından da cemisi getirilmiştir. Sözlük her harfe başlarken o harf ile ilgili bilinmesi gereken bilgileri de içermektedir.

Mesela (و ز ر) أَوْزَرَ إِيزَارًا 1- Sığınak yapmak: ه - , 2- Gizlemek, saklamak: خَبَأَهُ : الشيئ - أَوْزَرَ kelimesinin iki farklı anlamı verilmiş ve kelimenin kökü de parantez içinde gösterilmiştir.[33] Burada bazı sözlüklerde olduğu gibi "ه" zamiri ile fiilin müteaddi olduğu ifade edilmiştir. Bir isme örnek olarak da حَرَثَ 1- الْحَرْثُ nin mastarı. 2- Ekin. 3- Zirâat için hazırlanmış yer, tarla.[34] Önce mastar verilmiş sonra o masterın türediği fiil ve içerdiği anlamlar getirilmiştir. Bir de mücerred fiile örnek verirsek سَنَّ يَسُنُّ سَنًّا mazi, muzari ve mastar şeklinde fiil getirilmiş, 1- Bıçak vs. yi bilemek, keskinleştirmek. السِّكِّينَ-, 2- Taşı cilâlamak, parlatmak. الحَجَرَ-şeklinde farklı anlamları ifade edilmiştir.[35]

8- Türkçe – Arapça Sözlük

Bu sözlük Prof. Dr. İlyas Karslı tarafından 2002 yılında tamamlanmış ve M.Ü. İlahiyat Fakültesi Vakfı Yayınlarından çıkmıştır. Sözlük Türkçe alfabeye göre tertip edilmiş, soldan sağa şekilde hazırlanmıştır. Sözlük büyük orada Karslı'nın Arap ülkelerinde kaldığı zamanlarındaki halkın günlük dilde kullandığı kelimelerden ve basın-yayın organlarından elde edilen notlarından oluşmaktadır. Bu yönüyle sözlüğe modern sözlük denilebilir. Ayrıca sözlüğü oluştururken Karslı *es-Sıhâh, el-Kāmûsü'l-Muhît, el-Mucemul-Veciz, Kitabül-Emsal, Meddül-Kâmus* ve *El-Müncid fi'l-Lügâti ve'l-A'lâm* gibi sözlüklerden yararlanmıştır.[36] Sözlüğün ilk sayfasında kullanımı kolaylaştırmak için bazı kısaltmaların açıklamasına yer verilmiştir.

Sözlüğün sonunda "Ekler" kısmında "اِتَّخَذَ الْعِبْرَةَ / ders aldı, اِتَّخَذَ قُدْوَةَ / örnek aldı"[37] gibi kalıp ifadelere, "الإيمَانُ بِغَيْر الأعْمَال مَيِّتٌ / amelsiz iman ölüdür, التَّجْرِبَةُ بِدُونِ تَعَلّمٍ خَيْرٌ مِنْ تَعَلّمٍ بِدُونِ تَجْرِبَةٍ / bilgisiz tecrübe, tecrübesiz bilgilden daha iyidir"[38] gibi atasözleri, deyimlere ve "مُؤَسّسَة الدوليّة التّمويل / uluslararası finans kurumu, جَامِعة الدُّوَل العربِيّة / Arap ülkeleri birliği" gibi kurum ve kuruluş isimlerine yer verilmiştir.

[29] Yolcu, "خلد", 347.
[30] Yolcu, "أغار", 95.
[31] Yolcu, "جالس", 279.
[32] Arif Erkan, *el-Beyan Arapça-Türkçe Büyük Sözlük* (İstanbul: Yasin Yayınevi, 2006), 7.
[33] Erkan, "الأَوْزَنُ", 348.
[34] Erkan, "الحَرْبِيُّ", 536.
[35] Erkan, "سَمَنَ", 674.
[36] İlyas Karslı, *Türkçe – Arapça Sözlük* (İstanbul: M.Ü. İlahiyat Fakültesi Yayınları, 3. Basım, 2004), 768.
[37] Karslı, 698.
[38] Karslı, 730.

FULL TEXTS BOOK

9- Arapçada Çok Kullanılan Filler Sözlüğü

İlyas Karslı tarafından hazırlanan bu sözlük 2011 yılında Ensar Neşriyat tarafından yayınlanmıştır. Eserde 2000 Arapça fiil, Kur'ân'ı Kerîm, hadis-i şerifler ve Arap şiirlerinden de 12000 tane örnek bulunmaktadır. Ayrıca eser "*el-Arabiyye Beyne Yedeyk*" kitabında bulunan bütün fiilleri de ihtiva etmektedir.[39] Sözlükte kısaltma olarak sadece Kur'ân'ı Kerîm için KK. İfadesi kullanılmıştır. Sözlükteki fiiller mazi, muzari ve mastar olarak getirilmiş, Arapça alfabetik sıraya göre dizilmiştir. Fiilin önce kendi anlamı verilmiş, ardından örnek cümlelerle manası daha da pekiştirilmiştir.[40]

Mesela sözlükte "بَلَا – يَبْلُو – بَلَاءً / Denemek, imtihan etmek, teste tabi tutmak. Ölçmek, sınamak. Bitkin bırakmak." şeklinde kelime ve o kelimenin anlamları yalın olarak verilmiş, sonrasında الَّذِي خَلَقَ الْمَوْتَ وَالْحَيَاةَ لِيَبْلُوَكُمْ أَيُّكُمْ أَحْسَنُ عَمَلً / "Allah, hanginizin daha güzel şeyler yaptığını test etmek için, ölümü ve hayatı yarattı" KK. âyeti, بُلِيَ الشئُ / Sınandı. Denendi. Test edildi., بَلَاهُ بِالصَّبْرَ / Sabrını denedi., بَلَا السَّفَرُ خَالِدًا / Yolculuk, Halit'i bitkin bıraktı.[41] gibi cümleler veriliyor ve kelimenin farklı anlamları örneklendiriliyor.

10- Arapça-Türkçe Sözlük

Serdar Mutçalı tarafından hazırlanan "Arapça-Türkçe sözlük" Dağarcık Yayınları tarafından yayınlanmıştır. Sözlük Arapça alfabeye göre tertiplenmiş ve toplamda 1056 sayfadır. Mutçalı sözlüğünü hazırlarken H. Wehr'in *A dictionary of Modern Written Arabic* isimli çalışmasını temel almış ve ona göre hazırlamıştır. Onun haricinde *el-Mu'cemü'l-vasît, Lisânü'l-arab, Muhîtü'l-muhît* gibi sözlüklerden ve Butrus el-Bustânî'nin eserlerinden de yararlanmıştır.[42]

Sözlükteki kelimeler kök tertibine ve alfabetik sıraya uygun sıralanmıştır. Bir kelimenin önce fiil ise mazisi yazılmış, kısa çizgi (–) üzerinde ilgili kelimenin muzari harekesi verilmiş, mastarı da parantez içerisinde belirtilmiştir. Bu fiilden türetilen mezid bablar ise sırasıyla tef'il, mufaale, ifal, tefuul, tefaul, infial, iftial ve istifal şeklinde verilmiştir. Fiillerin anlamı verildikten sonra eğer birleşik şekilde kullanılıyorsa o anlamı da " ¶ " işaretinden sonra verilmiştir. مَدِين borçlu; ¶ مَدِين بِالشُّكْر şükran borcu olan.[43] Bir kelimenin alternatifini verirken de " / " işareti kullanılmıştır. Ayrıca fiille beraber kullanılan harfi cerler de bazen parantez içerisinde bazen de direk yazılmıştır. Sözlükteki Arapça aynı manayı ifade eden kelimelerde "=" işareti ile verilmiştir. جِنْزِير [=زِنْجِير] ç. جَنَازِير zincir.[44] Fiil veya isim ile ilgili açıklayıcı ifadeler ise köşeli parantez içerisinde sunulmuştur.[45] Mesela "بَسُلَ – ُ (بَسَالَة) korkusuz/ cesur/ yiğit olmak" şeklinde kelime verildikten sonra "تَبَسَّلَ kaşlarını çatarak bakmak" kelimesi veriliyor, ardından kelime istifal babından da kullanıldığı için "اِسْتَبْسَلَ [savaşta] kahramanlık göstermek" şeklinde ifade ediliyor.[46] Ayrıca sözlükte ilgili fiilin faili ve failinin cemileri de verilmektedir. Cemiler "ç" harfi ile belirtilmektedir. بَاسِل ç. بَوَاسِل ، بُسَلَاء ، بُسَّل yiğit, kahraman cesur, korkusuz.[47]

Sözlüğün sonundan "Ekler" başlığında ülke isimleri, Arap para birimleri, haftanın günleri, aylar, sayılar, saat hakkında kullanılabilecek bazı kalıp ifadeler, renkler, geometrik şekiller, aylar ve Arapçada üzerinde müennes mi müzekker mi belli olmayan hem müzekker hem de müennes kullanılabilen kelimeler verilmiştir.[48]

[39] İlyas Karslı, *Arapçada Çok Kullanılan Filler Sözlüğü* (İstanbul: Ensar Neşriyat, 2011), 54.
[40] Karslı, 55.
[41] Karslı, 164.
[42] Serdar Mutçalı, *Arapça-Türkçe Sözlük* (İstanbul: Dağarcık Yayınları, 2019), 5.
[43] Mutçalı, "دين", 327.
[44] Mutçalı, "جنس", 171.
[45] Mutçalı, 7.
[46] Mutçalı, "بسم", 81.
[47] Mutçalı, "بسم", 81.
[48] Bk., Mutçalı, 1039-1054.

FULL TEXTS BOOK

SONUÇ

Sözlükler bir eser veya metin okunurken orada geçen anlamı bilinmedik kelimelerin anlamlarını öğrenmek için başvurulacak ilk kaynaktırlar. Arapça-Türkçe ve Türkçe-Arapça sözlükler de Arapça bir metin okunurken kelimelerin anlamları için müracaat edilen ilk mercilerdir. Ülkemizde bu konuda talebeye yol gösterici sözlük kullanımına dair bir çalışmaya ulaşamadığımızdan dolayı yapılan bu çalışma önemli bir boşluğu dolduracağı kanaatindeyiz. Yaptığımız çalışmada birtakım eserlerin yazarı, yayıncısı, usulü hakkında bilgi verdikten sonra bilgileri örneklerle daha anlaşılır hale getirdik. Çalışmada ele aldığımız eserlerin farklı farklı nitelikte olmasına dikkat ettik ve benzer eserleri çalışmaya dahil etmedik.

İncelediğimiz ve tanıttığımız eserlerde yazarlar kelimeleri genel olarak kelimeleri sıralarken iki yönteme göre ele almışlardır. Birincisi; kelimelerin asli (kök) harfleri esas alınarak kelimeler dizilmiştir. İkincisi; kelimeler alfabetik sıraya uygun olarak dizilmiştir. Kelime olarak değil de kalıp ifade olarak oluşturulan sözlüklerde ise kelimelerin Türkçesi esas alınarak Türkçe alfabetik sıraya göre tasnif edilmiştir. Ayrıyeten bazı eserlerde kelimelerin anlamı daha iyi kavranması için küçük resimler kullanılmış, birtakım sözlüklerde de kurum, kuruluş ve sarf-nahiv'in özet niteliğinde tablolarına yer verilmiştir. Yine kullanıcı için günlük hayatta kullanabileceği pratik kalıp ifadelere de bazı sözlükler yer vermiştir.

Çalışmamız esnasında bizim çalışmaya almadığımız birçok kıymetli ve kapsamlı sözlüklerin var olabileceğini, imkanlar nispetinde tespit edebildiğimiz farklı usullerde sözlükleri çalışmaya konu ettiğimizi ve bu çalışmanın daha da genişletilerek her bir eser farklı bir çalışma ve müstakil incelenmeye muhtaç olduğunu söyleyebiliriz.

KAYNAKÇA

Durmuş, İsmail. "Sözlük". TDV İslâm Ansiklopedisi. Erişim 01 Aralık 2024. https://islamansiklopedisi.org.tr/sozluk#1

Atay, Hüseyin- Atay, İbrâhim- Atay, Mustafa. *Arapça-Türkçe Büyük Lügat*. Ankara: Bayrak Matbaası, 1964.

Topaloğlu, Bekir- Karaman, Hayreddin. *Yeni Kâmus*. İstanbul: Ahmet Sait Matbaası, 1966.

Sarı, Mevlüt. *el-Mevarid Türkçe-Arapça Lügat*. İstanbul: Bahar Yayınları, 1975.

Ayyıldız, Erol. *Türkçe – Arapça, Arapça – Türkçe Cep Lügatı*. Ankara: Bayrak Matbaası, 1997.

Yolcu, İ. Muhammed- Çetin, Nurullah. *Modern Arapça Sözlük*. Ankara: Birleşik Yayınları, 1992.

M. İsmet Yolcu. *Örnek Sözlük Arapça-Türkçe*. Ankara: Eğit-San yayınları,1996.

Arif Erkan. *el-Beyan Arapça-Türkçe Büyük Sözlük*. İstanbul: Yasin Yayınevi, 2006.

Karslı, İlyas. *Türkçe – Arapça Sözlük*. İstanbul: M.Ü. İlahiyat Fakültesi Yayınları, 3. Basım, 2004.

Karslı, İlyas. *Arapçada Çok Kullanılan Filler Sözlüğü*. İstanbul: Ensar Neşriyat, 2011.

Mutçalı, Serdar. *Arapça-Türkçe Sözlük*. İstanbul: Bahar Yayınları, 1975.

FULL TEXTS BOOK

AZƏRBAYCANIN ASTARA BÖLGƏSİNİN TARIXI ISTEHKAMLARI
HISTORICAL FORTIFICATIONS OF THE ASTARA PROVINCE OF AZERBAIJAN

Nəzakət ƏBILOVA

Lənkəran Dövlət Universiteti, Tarix, Vətən tarixi

Lənkəran şəhəri, Azərbaycan Respublikası

ID-0000-0002-9038-7135

ÖZƏT

Azərbaycanın Cənub-Şərq bölgəsi qədim zamanlardan onunla həmsərhəd olan Cənubi-Qafqazla əhəmiyyətli maddi-mədəni və siyasi-iqtisadi əlaqələrə malikdir. Bu regionun üzərindən Qafqazın, onun vasitəsi ilə isə Cənub-Şərqi Avropanın Yaxın Şərq sivilizasiyaları, xüsusilə İran yaylası sivilizasiyaları ilə müxtəlif xarakterli əlaqələrinin bir qisim marşrutları keçmişdir. Tarixi İpək Yolu, qədim yaşayış məntəqəsi olan Astaranın ərazisindən də keçməklə Orta, Ön, Kiçik Asiya, Çin, Hindistan və Ərəb dünyası arasında geniş əlaqələr qurulmuşdur. Tarixi tədqiqatlar və arxeoloji qazıntılar nəticəsində aydın olmuşdur ki, Azərbaycan şəhərləri müxtəlif zəmin üzərində əmələ gələrək inkişaf edirdi. Əsasən karvan yolunun üstündə yerləşən bəzi iri yaşayış məntəqələrinin ticarət və sənətkarlıq mərkəzi olması, digərinin dini zəmin əsasında yaranması, bəzilərinin isə inzibati mərkəz rolunu oynaması onların inkişafına müxtəlif dərəcədə təsir göstərmişdir

Astara və onun yerləşdiyi bölgə haqqında yunan, rus, alman, holland, şotland və s. səyyah, diplomat və dənizçilərinin qeydlərində məlumatlar vardır.

Astara və ətraf ərazilərinin arxeoloji baxımdan öyrənilməsində Jak de Morqan, İ.Cəfərzadə, F.Mahmudov, A.Məmmədov, Q.Qoşqarlı, T.Axundov, Ə.Ələkbərov, C.Eminli, T.Əliyev, A.Ağalarzadənin xidmətləri böyükdür.

Bu regionun arxeoloji tədqiqi XIX əsrin sonlarından başlanılıb və müasir dövrümüzə kimi mərhələli şəkildə davam etdirilmişdir. Lakin bu region hələ də tədqiq olunmamış özünəməxsus maddi, mənəvi və mədəni tarixi abidələrlə zəngindir. Xüsusilə qeyd etmək lazımdır ki, Astara vilayəti tədqiq olunmamış qala və istehkamların çoxluğu ilə diqqəti cəlb edir.

Azərbaycanın qiymətli maddi-mədəniyyət abidələri içərisində orta əsr istehkamları xüsusi yer tutur. Bu tip abidələrin tədqiq olunması vətənimizin orta əsrlər tarixinin öyrənilməsi istiqamətində mühüm addımdır. Bu abidələrin hər biri özündə bir tarixi qoruyub saxlayır. Müdafiə istehkamlarının tədqiqi dövrün ictimai-siyasi, hərbi məsələlərinin öyrənilməsində də xüsusi əhəmiyyət kəsb edir.

Açar sözlər: Astara, qala, arxeoloji, müdafiə, işğal, istehkam

ABSTRACT

The southeastern region of Azerbaijan has, since ancient times, maintained significant material-cultural and socio-economic relations with the South Caucasus, which borders it. Several routes facilitating connections between the civilizations of the South Caucasus—and through them, those of Southeastern Europe—and the civilizations of the Near East, particularly the Iranian Plateau, passed through this region. The historic Silk Road also traversed the area of Astara, an ancient settlement, thereby establishing extensive links between Central Asia, Asia Minor, the Indian subcontinent, China, and the Arab world.

FULL TEXTS BOOK

Historical research and archaeological excavations have revealed that Azerbaijani cities emerged and developed on various foundations. Some major settlements, located along caravan routes, evolved into centers of trade and craftsmanship; others formed based on religious significance, and some served administrative purposes. These diverse origins influenced their development to varying degrees.

Records by travelers, diplomats, and seafarers from Greece, Russia, Germany, the Netherlands, Scotland, and other countries provide valuable information about Astara and its surrounding areas.

Significant contributions to the archaeological study of Astara and its environs have been made by scholars such as Jacques de Morgan, I. Jafarzadeh, F. Mahmudov, A. Mammadov, G. Goshgarli, T. Akhundov, A. Alakbarov, J. Eminli, T. Aliyev, and A. Agalarzadeh.

Archaeological exploration in this region began in the late 19th century and has continued in stages up to the present day. However, the area remains rich in unique material, spiritual, and cultural heritage that has yet to be fully studied. It is particularly noteworthy that the Astara province stands out for its numerous unexplored castles and fortifications.

Among Azerbaijan's valuable material and cultural monuments, medieval fortifications occupy a special place. The study of such monuments is a vital step toward understanding the medieval history of our homeland. Each of these structures preserves a part of history. The investigation of defensive fortifications also plays a crucial role in the study of the socio-political and military issues of their respective periods.

Keywords: Astara, fortress, archaeology, defense, invasion, fortification

GİRİŞ

Azərbaycanın Cənub-Şərq bölgəsi, xüsusilə Astara ərazisi, tarixi-coğrafi mövqeyi və strateji əhəmiyyəti baxımından qədim dövrlərdən etibarən mühüm sivilizasiya və mədəniyyətlərin qarşılıqlı təsir məkanında yerləşmişdir. Bu bölgə həm Cənubi Qafqazla, həm də Yaxın Şərq ölkələri ilə zəngin ticarət, iqtisadi və mədəni əlaqələrə malik olmuş, İpək Yolunun əsas keçid nöqtələrindən biri kimi tanınmışdır. Astaranın ərazisindən keçən beynəlxalq karvan yolları və burada formalaşan yaşayış məntəqələri bölgənin sosial-iqtisadi inkişafında və hərbi-siyasi hadisələrin gedişində tarixi mövqeyi olmuşdur. Regionun özünəməxsus təbii coğrafiyası – dağlıq relyefi, çay sistemləri və Xəzər dənizinə çıxışı burada istehkamların və müdafiə qurğularının inşasını zəruri etmişdir.

Tarixi mənbələr və səyyah qeydləri Astara bölgəsində müdafiə məqsədilə tikilmiş qalaların mövcudluğunu, onların funksional xarakterini və dövrlər üzrə dəyişikliklərini təsdiqləyir. Şindan, Dizdəqala, Qala Kafu, Divəkə və digər istehkam nümunələri bu baxımdan təkcə hərbi əhəmiyyət daşımır, həm də memarlıq və arxeoloji baxımdan dəyərli abidələr hesab olunur. Eyni zamanda, bu istehkamların hər biri regionun müdafiə sisteminin təşkili, ərazi bütövlüyünün qorunması və inzibati idarəçiliyin həyata keçirilməsi baxımından spesifik rol oynamışdır.

Bu tədqiqatın əsas məqsədi Astara bölgəsinin tarixi istehkamlarını araşdırmaqla onların funksional rolunu, memarlıq xüsusiyyətlərini və regionun tarixində tutduqları yeri elmi əsaslarla təhlil etməkdir. Mənbə materialları və arxeoloji faktlar əsasında aparılan bu araşdırma, həmçinin müasir dövrdə bu abidələrin qorunması və elmi dövriyyəyə cəlb olunması baxımından da xüsusi aktuallıq kəsb edir. Ölkəmizin Cənub-Şərq bölgəsinin tədqiqi zamanı Astara rayonun istər yazılı mənbələrdən istərsə də arxeoloji cəhətdən öyrənilməsi böyük əhəmiyyətə malikdir.

Astara şəhəri Astaraçayın mənsəbindədir, Astaraçay adını eyni adlı bulaqdan almışdır. Astaraçay başlanğıcını Talış dağlarından, Şindan qalasından (1817 m yüksəklikdən) alır, Astara ərazisindən axıb Xəzər dənizinə tökülür (uzunluğu 38 km, hövzəsinin sahəsi 242 km^2) Alaşa kəndi yanında qovuşan

FULL TEXTS BOOK

İstisu Astaraçayın əsas qolu sayılır. Astaraçayın bu ərazidən mənsəbə qədər olan məsafəsi Azərbaycan Respublikası ilə İran İslam Respublikası arasında dövlət sərhəddini təşkil edir.

Astara sözünün mənası bir-neçə versiyada izah olunur: ast – aşağı, dərə və ara/ura – yer, su, bulaq sözlərindən alındığı da iddia edilir, mənası "çökək yerdən çıxan su" kimi izah edilir. Amma yerli talış dilində "Astara" sözü "ulduz" mənasını verir. Yerli əhali bu şəhərin qədim adının "Qum" olduğunu bildirir, fars dilində də "qum" sözü "ulduz" mənasını verir.

Səidəli Kazımbəyoğlunun "Cəvahirnameyi Lənkəran" (Səidiyyə) əsərinin Səfər Şirinov tərəfindən edilmiş tərcüməsində, Lənkəran vilayətinin bir mahalı olan Astaranın əslində "Əstara" olması və "seçmək" yaxud "seçilmiş" mənasını verdiyi bildirilir

Yazılı mənbələrdə hələ II əsrdə görkəmli riyaziyyatçı, astronom və coğrafiyaşünas İsgəndəriyyəli Klavdi Ptolomey tərtib etdiyi Xəzər dənizi və ətraf məntəqələri özündə əks etdirən xəritədə qədim Albaniyanın bir çox şəhər və kəndlərinin adını çəkmiş və Astaranın adını yunanca "Astarata" kimi qeyd etmişdir. XV əsrdə müxtəlif Şərq ölkələrini, eləcə də Azərbaycanı gəzib dolaşmış alman səyyahı Hans Şilberqer Astaranın beynəlxalq ticarət əlaqələri barədə konkret məlumat vermişdir. Səyyah qeyd etmişdir ki, Astarada çoxlu ipək istehsal olunur, bu ipəyin ən yaxşı növü Dəməşq, Bursa, Kaşan və Venesiya kimi sənətkarlar mərkəzinə ixrac olunur,bu yerlərdə həmin ipəkdən yaxşı məxmər parçalar toxunur. İngiltərə ilə Buxara arasında ticarət əlaqələrini yaratmaq məqsədi ilə 1559-cu ildə Xəzər dənizi sahillərində olmuş İngilis taciri Antoni Cenkinson, 1562-ci ildə Londona qayıdaraq öz səfərləri haqqında yazdığı "Rusiya-Tatarıstan təsviri" əsərində dərc edilmiş xatirələrdə Xəzər dənizi ətrafındakı yaşayış məntəqələrinin adlarını verərkən Astaranı "Stara" kimi göstərmişdir (bu bölgədə "Dan ulduzu"-nu "Sitarə" ulduzu kimi də adlandırırlar). Astara adına həmçinin Hacı Zeynalabidin Şirvaninin, rus diplomatı Aleksandr Volinskinin, həmçinin E.Veydenbaumun, Katib Çələbinin, Əbdül Fəttah Fumainin əsərlərində də rast gəlinir. (Kərimov H., 1986 ; Mustafazadə T., 2011)

Bu regionun arxeoloji tədqiqi XIX əsrin sonlarından başlanılıb və müasir dövrümüzə kimi mərhələli şəkildə davam etdirilmişdir. Lakin bu region hələ də tədqiq olunmamış özünəməxsus maddi, mənəvi və mədəni tarixi abidələrlə zəngindir. Xüsusilə qeyd etmək lazımdır ki, Astara vilayəti qala və istehkamlarla zəngindir. Hələ XIII əsrin 30-cu illərində Yaqut əl-Həməvi "Mucəm əl-buldan" əsərində Azərbaycanın böyük ölkə və qüdrətli dövlət olduğunu bildirmiş, buranın çoxlu istehkam qalaları olan əsasən dağlıq ərazi olduğunu qeyd etmişdir. Azərbaycanın strateji cəhətdən ən əhəmiyyətli bölgələrindən biri olan Astara əsrlər boyu müharibə meydanı olmuşdur. Heç də təsadüfi deyildir ki, 1638-ci ildə Astarada olmuş məşhur alman səyyahı Adam Oleari yazırdı: "Astaralılar yatanda belə, həmişə silahı hazır saxlayırlar" (Путешественники об Азербайджане. Составил: З.И.Ямпольский, 1961)

Antik və orta əsrlərdə şəhərlərin və yaşayış yerlərinin ətrafı həm təbii, həm də süni müdafiə qurğuları ilə əhatə olunurdu. Belə ki, orta əsr şəhər əhalisi və yaşayış məntəqələrinin müdafiə qurğuları olmadan dinc yaşaması mümkün deyildi. Çünki şəhər və yaşayış məntəqələrində daimi nizami ordu və əsgəri birləşmələr yox idi. Yerli feodalların bir-birinin torpaq mülkünə göz dikməsi nəticəsində onların arasında tez-tez baş verən daxili toqquşmalar və ara müharibələri, həm də yadellilərin basqın və yürüşləri müdafiə qurğularının tikilməsini zərruri edirdi. Orta əsr şəhərlərinin əsas müdafiə divarları ilə yanaşı şəhərlərin kənarlarında tikilmiş tək-tək qalalar şəhərə edilən düşmən hücumları qarşısında maneələr törətmək, şəhəri gözlənilən təhlükədən xəbərdar etmək və işğalçıları qismən ləngitmək məqsədi güdürdü

FULL TEXTS BOOK

Şindan qalası: Şindan qalası Azərbaycan Respublikası ilə İran İslam Respublikasının sərhəd zolağında (netral zonada), Astara rayonunun qərbində, Şindan (Cindan) dağının zirvəsində, dəniz səviyyəsindən 1850 metr yüksəklikdə yerləşir (1828-ci il fevralın 10-da Rusiya imperiyası ilə Qacar dövləti arasında imzalanmış "Türkmənçay" müqaviləsinə əsasən Azərbaycanın şimal hissəsinin Rusiyaya, cənub hissəsinin isə Qacar İranına birləşdirilməsi nəticəsində Astara vilayəti də iki hissəyə bölünmüşdür). Orta əsr dövrü abidəsi hesab olunur. Məşhur alman səyyahı Adam Oleari Şindan dağı haqqında danışarkən onun adının orada yerləşən kəndlə bağlı olduğunu və bu adın Şah (şeyx) Səfinin müəllimi olmuş və bu kənddə yaşayıb elə buradaca dəfn olunmuş Şeyx Səhədlə bağlı olduğunu bildirmişdir. (Путешественники об Азербайджане. Составил: З.И.Ямрольский, 1961)

Daş və sarucdan hörülən Şindan qalasının hündürlüyü 18 metr 50 santimetrdir. Qalanın üstü 1 hektardan artıq sahəni əhatə edir. Şindan qalası iki hissədən – böyük və kiçik qaladan ibarətdir. Əsas qalaya yeganə giriş yolu şimal tərəfdəndir. Bu yolun girəcəyində ağ daşdan pilləkan hörülmüşdür. Gözətçi və sərkərdə üçün düzəldilmiş xüsusi yerləri mövcuddur. Digər qalalar kimi Şindan qalası da müdafiə-istehkam xarakteri daşımışdır. Səidəli Kazımbəyoğlu qala haqqında yazır: "Məşhur Şindan qalası hündür dağın başında yerləşir. O qalanın ətrafında indiyədək qalan tarixi əsərlər mövcuddur. Onlardan biri həmin dağın özündən qoparılmış daşlarla tikilən bir bürcdür. Onun ətrafında 6 metr kub uzunluğunda 4-5 metr kub enində bir hovuz vardır, dörd fəslin dördündə də həmin hovuz çox soyuq və bol su ilə dolu olur." (Səidəli Kazımbəyoğlunun "Cəvahirnameyi Lənkəran". Tərcümə: S.Şirinov, 2005). Bəhs edilən bürc və hovuz dövrümüzə kimi gəlib çatmışdır.

Səidəli Kazımbəyoğlu "Cəvahirnameyi Lənkəran" əsərində qeyd etmişdir: "O məkanın qəribə əlamətlərindən biri də budur ki, o dağın ətəyində o qədər əsl çini qabların sınığı bir-birinin üstünə tökülmüşdür ki, nəticədə böyük bir təpə yaranmışdır. Əgər həmin sınıq qab-qacağı bir yerə daşımaq zərurəti yaransa min xarvardan da (ulaq yükü) çox olar. Başqa bir qəribəlik də həmin məkanda top güllələrinin hədsiz dərəcədə çox olmasıdır." Sonra müəllif bu qəribəliklərin kimlərinsə tərəfindən tədqiq olunduğunu qeyd edərək, bəzi rəvayətlərin yayıldığını bildirmişdir. "Gilan vilayətinin qədim sultanlarından birinin bu qlada yayalaq məskəni olmuşdur. Həmin sultan belə bir qayda tətbiq etmişdir, ki, hər bir məclisdə işlədilən süfrə qab-qacağı təzə çini qablardan olmalıdır. Hər bir ziyafətdə yemək yeyiləndən sonra işlədilmiş qab-qacaq mətbəxə qaytarılmayaraq qaladan xaricə atılırmış." (Səidəli Kazımbəyoğlunun "Cəvahirnameyi Lənkəran". Tərcümə: S.Şirinov, 2005)

Tarixi mənbələrdə qeyd olunduğu kimi " Seyid Abbas bəy Ucarud, Dəştvənd və Uluf mahallarından ibarət olan Talış vilayətində iyirmi ilə kimi asudəliklə hakim olduqdan sonra vəfat etdi". (Mirzə Əhməd Xudaverdi oğlu "Əxbərnamə" Tərcümə: Əli Hüseynzadə. PDF). Seyid Abbas xanın vəfatından sonra tarixdə Qara xan kimi tanınan Cəmaləddin Mirzə bəy 1747-1786-cı illərdə hakimiyyətdə olmuşdur. Bu ləqəb ona Nadir şah Əfşarın qoşununda xidmət edərkən zəhmli olduğuna görə verilmişdir. (Quliyev S., Vəliyev E. Azərbaycanda multikulturalizm və onun Lənkəran qaynaqları. 2018) O, xanlığın iqtisadi və siyasi qüdrətini möhkəmləndirmək üçün bir sıra tədbirlər həyata keçirmişdir. Qara xan İran təhlükəsindən qorunmaq məqsədilə paytaxtı Astaradan Lənkərana köçürmüş və şəhər ətrafında qala divarları saldırmışdır. Gilan və Lənkəran xanları arasında ərazi iddiaları ilə bağlı kəskin mübarizə gedirdi. İran hökmdarı Kərim xan Zənd həmişə Gilan xanını müdafiə edərək və ona hərbi qüvvə ilə yardım edirdi. "1768-ci ildə Qara xanın Asalim və Gargarud mahallarını tutması gilanlı Hidayət xanın 12 min nəfərlik qoşunla Lənkəran xanlığına hücum etməsinə səbəs olmuşdu." (Mustafazadə T. Azərbaycan xanlıqlarının qısa tarixi. 2011). Məhz həmin basqın zamanı Şindan qalasına çəkilən Qara xan bir müddət müqavimət göstərsə də, Hidayət xana təslim olmaq məcburiyyətində qalır. Hidayət xan çoxlu xərac alaraq Qara xanı azad edir. Əvəllər Lənkəran xanına məxsus olan Astara Kərim xan Zəndin

FULL TEXTS BOOK

gücünə alınıb Gilan xanına verildi. Öz hakimiyyətini möhkəmlətmək və ümumi düşmənə qarşı mübarizə aparmaq üçün Qara xan qubalı Fətəli xanla ittifaqa girir.

Bəlkə də Səidəli Kazımbəyoğlunun "Cəvahirnameyi Lənkəran" əsərində qeyd olunan rəvayətdə Gilan hökmdarı Hidayət xandan bəhs olunur, bunu ancaq qala ətrafında olan çini qabların qalıqlarını tədqiq etməklə, nə dərəcədə "qədim" olmasını müəyyən etməklə bilmək olar.

İkinci hekayə «top güllələri» rəvayətidir ki, müəllif bu rəvayətin şübhəsiz bir həqiqət olduğunu qeyd edir. " Belə ki, bu hadisə Talış xanı Mir Qara xanın əyalətə başçılıq etdiyi zamanda baş vermişdir. O mərhum özünün hakimiyyəti illərinin sonlarında Qızılbaş sultanlarına itaətdən imtina etmiş, bu qalanın ətrafında (Şindan qalası nəzərdə tutulur) döyüş mövqeyi tutmuş və bu zaman Qızılbaş qoşunlarının hücumuna məruz qalmışdır. Qızılbaş qoşunları özlərinin top texnikalarını o qalanın ətrafındakı bağda yerləşdirərək həmin qalaya tərəf tuşlamışlar. Belə ki, hələ də həmin ətrafda «top yolu» adlandırılan yer həmin məkanın yaxınlığında məşhur yer kimi qalmaqdadır. Qızılbaş orduları uzun müddət qalanı mühasirədə saxlamışlar. Top atəşləri qalanı təslim edə bilməmişlər. Beləliklə, Qara xan Qalada azuqənin qurtarması nəticəsində Şindan qalasını boşaltmış və oradan qaçmışdır." (Səidəli Kazımbəyoğlunun "Cəvahirnameyi Lənkəran". Tərcümə: S.Şirinov, 2005)

Coğrafiya elmləri doktoru, professor Nadir Məmmədov Şindan qalasının bünövrəsinin eramızın III əsrində Sasanilər dövründə qoyulduğunu qeyd edir. (Məmmədov N. Azərbaycanın yer adları. 1993)

Dizdəqala və ya Təngəru qalası: Lənkəran-Astara yolunun sağ tərəfində yola paralel şimal-cənub istiqamətində uzanan, hündürlüyü 100-150 metr olan kiçik dağ yerləşir. Onun ən hündür hissəsi uzunluğu 200-250 metr, en isə 30-40 metrə qədər olan, səthi hamar beldən ibarətdir. Dağın şimal, qərb və şərq yamacları çox sərtdir, cənub yamacı isə nisbətən maili olub, çox uzanır. Yamaclar meşə ilə örtülüb, zirvəsi açıqdır. Zirvədən Xəzər sahili, şimal və cənub istiqaməti boyunca sahilyanı düzənliklər, şimal qurtaracağından isə Təngəru çayı hövzəsinin xeyli hissəsi yaxşı görünür. Qalanı inşa edərkən dağin beli 2 yerdən köndələninə qazılmış xəndəklə 3 istiqamətdə - şimal, mərkəz və cənub hissələrə bölünmüşdür. Bəzi yerlərdə o təbii qaya çıxıntıları ilə hüdudlanmış, bu isə ona cənubdan şimala qalxan pilləli forma vermişdir. Mərkəz hissəsi qərb və şərqdən sərt yamaclı, şimaldan və cənubdan isə xəndəklə sərhədlənən düzbucaqvaridir. Kənd əhalisinin verdiyi məlumata görə, XX əsrin 60-70-ci illərinə qədər bu yerdə xüsusilə Novruz şənliklərində tonqal qalanması və oyun oynamaq üçün toplanılırdı. Qalanın divarları işlənməmiş qaya parçalarından hörülmüş, günümüzə kimi eni 3-4 metr olan dağıntı şəklində qalmışdır.

Səidəli Kazımbəyoğlu öz əsərində bu qala haqqinda az da olsa məlumat vermiş, Dizdəqələ talış dilində "oğru qalası" mənasını verdiyini bildirərək, qalanın ətrafindakı böyük meşəni Dizdəvişə yəni "oğru meşəsi" adlandırmışdır

Qala tam tədqiq olunmayıb "lakin onun topoqrafiyası, strukturu və divarın hörgüsü Təngəru (Dizdəqala) qalasını son orta əsrlərə aid etməyə imkan verir." (Axundov T.İ. Astarada 20 gün. 2009)

Çanaxa qlası: Türkoba kəndinin cənubunda mərhələli şəkildə təkrar olunan sürüşmə zonası olan yüksəklik yerləşir. Yüksəkliyin şərq qurtaracağına yaxın şimal yamacinin altında Çanaxa adlanan bulaq var. Sürüşmə nəticəsində yaranmış yarğanın qərb tərəfində nəhəng işlənmiş, təqribən hər birinin orta həcmi 1 kub metr olan daş bloklar aşkarlanmışdır. Görkəmli tədqiqatçı Tufan Axundov bu bir-birinə yaxşı uyğunlaşdırılmış nəhəng daşların qala divarinin qalıqları olduğunu bildirmiş, araşdırmasının nəticəsi olaraq qeyd etmişdir ki, " belə qalanın yalnız dövlət sisteminə malik qurum inşa edə bilərdi, görünür yaxındakı vadinin üzərində hökmranlıq edən strateji əhəmiyyətli sahənin zirvəsində, divarlarının qalıqlarının bir hissəsi aşkar olunmuş qala tikilmişdir" (Axundov T.İ. Astarada 20 gün. 2009)

Nəzərimcə Çanaxa qalası da bir sıra abidələr kimi tam tədqiq edilməlidir, çünki abidəni ilk aşkar etmiş T.Axundov "tikintinin quruluşu, divarların inşa üsulu və məlum analoqlara əsasən, bu abidəni e.ə. I

FULL TEXTS BOOK

minilliyin birinci yarsına, Manna dövlətinin zamanına"aid etmişdir. (Axundov T.İ. Astarada 20 gün. 2009)

Qala Kafu: Bölgədə olan qalalardan biri də Qala Kafu adlı qaladır. Bu qala Astara rayonunun qərbində, Hamoşam kəndinin ərazisində yerləşir. Qala Kafu qalası Rəşi Hürdən 1,5 km cənub-cənub-şərqdə yerləşir. Qala (Kalafu qalası) şimal-cənub boyu uzanan qayalı belin üzərində ucalır. Onun hündürlüyü qərb tərəfdən 70 metrə qədər, şərq tərəfdən isə 150 metrdən çoxdur. Onun ərazisi 60x30 metr olub,belə paralel gəlmişdir. Qaya mərkəz hissəsində yarılmış kimi olduğundan ikibaşlı forma almışdır. Zirvədən 8-10 metr aşağıda yerləşən yarğanın dibi məqsədli şəkildə düzəldilmiş və nəticədə meydança əmələ gəlmişdir. Meydançanın 14-16 metr olan eni, şimaldan və cənubdan yüksələn qayalarla hüdudlanır. Onun uzunluğu 25 metr olub, qərb-şərq istiqamətində uzanır və təbii hüdud olmadan sərt kəsilir. Qalanın şərq tərəfdə, qayanın kənarlarında kiçik ölçülü qaya daşlarından hörülmüş divarın qalıqları qalır. Qərb tərəfdə divar qalıqları izlənilmir. Onun vaxtilə qaya ilə birlikdə uçub töküldüyü bildirilir. Meydançanın şimal tərəfindən 2,7x1,3 metr ölçüdə, 1,6 metr dərinlikdə sututarı (hovuz) çapılmışdır. Qala Kafu ətrafındakı 5-6 hektar sahədə, xüsusilə Qala Kafu qayasının ətəklərində saxsı qab qırıqlarının səpələnməsi qala ətrafında yaşayış olmasından xəbər verir. Beləliklə, "ətraf ərazilər üzərində hakim olan bu strateji cəhətdən əlverişli yer, inkişaf etmiş və son orta əsrlər dövründə məskun edilmişdir. Burada iri yaşayış məskəni olan qala - süni istehkam işlərilə tamamlanan təbii istehkam yerləşmişdir. Qala Kafunun mərkəz qayası yaşayış yerinin narınqalası kimi istifadə edilmişdir." (Axundov T.İ. Astarada 20 gün. 2009)

Hal-hazırda Qala Kafu qalasının qalıqları qalmaqdadır. Qeyd edək ki, qalanın adı Kalafu, Qala Kafu və Kalakafo adları ilə qeyd edilir. Qalanın adının mənasını müxtəlif versiyalarda izah etmək olar: Qala Kafa – türk dilində Baş qala, Qala Kafir – Kafir qalası, eləcədə "mərkəz qayanın forması ilə bağlı olub, güman ki, yarğana görə "kəlafa" talış dilində olub "oyuq" mənasını verir.

Alaşa və ya Divəkə qalası: Alaşa kəndindən qərbdə, dağlarla düzənliyin qovuşduqları yerdə rəvayətə görə üzərində qala olmuş dağ var. Günümüzə yalnız kərpic hörgülərin ayrı-ayrı fraqmentləri gəlib çatmışdır. Güman etmək olar ki, Bəlləbur qalasında olduğu kimi qalanın tikintisində istifadə olunan kərpiclər yerli şəraitdə hazırlanmışdır və bunun üçün həmin ərazidə kərpic bişirən kürəxanalar olmuşdur. Hələlik onların yeri müəyyənləşdirilməsə də kərpiclərin çoxunun keyfiyyətini itirməsi bölgədəki rütubətin təsiri ilə izah edilə bilər. Bu cür inşaat materialından Gəncəbasarda (Quliyev S.və Vəliyev E., 2018), Naxçıvanda (Qasımlı V.A., 2013), Gəncədə (Məmmədov A.M.,1992), Oğuzda (Əliyev T.V., 2017), Zərdabda (Əliyev T.V., 2018), Şirvanın (Abbasova E.İ., 2019) və s. müdafiə tipli istehkamlarının tikintisində də istifadə edilmişdir.

Bu qlanı yerli əhali "Divə kə" adlandırır – yəni talış dilindən tərcümədə "Div evi" deməkdir. Orta əsrlərdə tikilən qalanın keşikçi qülləsi kimi istifadə olunması da ehtimal edilir. Qala Astara çayının yaxınlığında, Artupa bələdiyyəsinin Hirkan Milli Parkı ərazisinə düşən hissəsində yerləşir.

Yerli əhali arasında bu qala haqqında müxtəlif fikirlər yayılmışdır, güman edilir ki, qədim dövrlərdə bura Əlişah adlı bir şəxsin qalası olub, digər mənbələrə görə isə bu qala Teymurləngin oğlu Miranşahın qərargahı imiş. Mənbələrdən bildiyimiz kimi Miranşah çox qəddar bir hökmdar kimi tarixə düşmüşdür, yerli əhali məhz bu qalanın Div qalası adlandırıldığını bununla əlaqələndirirlər, yəni qalaya "gedişin olub, dönüşün olmadığının" mənasını verdiyini bildirirlər.

Qala strateji cəhətdən çox əlverişli mövqedə tikildiyindən sovet dövründə əsgərləri, xüsusilə də hərbi kəşfiyyatçıları təlim keçmək üçün qalanın yerləşdiyi əraziyə gətirirmişlər. Qalanın olduğu yerdən həm İran İslam Respublikasının ərazisini, həm də Xəzər dənizinin sərhəd sularını müşahidə etmək mümkündür. Bu məqsədlə hərbçilər çadırlar quraraq, bir neçə ay müddətində ərazidə qalırmışlar. Ərazidə 1990-cı illərə kimi sözügedən təlimlər keçirilib. Qala sovet dönəmində abidə kimi qorunmayıb, yalnız kəşfiyyatçı əsgərlərə təlim keçmək üçün istifadə edilib.

FULL TEXTS BOOK

Şimalda sıldırım dağa söykənmiş, qırmızı kərpic və gəc materialından tağ şəkilli otaqları olan qalanın qərbdən, şərqdən və cənub istiqamətindən mümkün hücumların qarşısının alınması və eyni zamanda dənizdən başlayan Astara-Ərdəbil karvan yoluna nəzarət məqsədilə inşa olunduğu ehtimal etmək olar.

Qalada keşikçi qülləsi ilə otaqlar arasında yeraltı yolun olduğu güman edilir. Burada yerin altına aparan girişə bənzər qalıqlar belə düşünməyə əsas verir. Təəssüf ki, günümüzədək qalanın yonulmuş sal daşlardan ibarət istinad divarları və tağ şəkilli bir neçə otağının qalıqları çatıb. Qalaya əsas giriş şimal-şərq tərəfdəndir.

Qeyd edək ki, "Divə kə" qalası da bu günə kimi tədqiq edilməyib, qala təxminən XII-XV əsrlərə aid edilir.

NƏTİCƏ

Astara bölgəsinin tarixi istehkamları regionun müdafiə sisteminin formalaşmasında və siyasi-iqtisadi hadisələrin yönəldilməsində mühüm rol oynamışdır. Şindan, Dizdəqala, Çanaxa, Qala Kafu və Divəkə kimi istehkamlar öz strateji yerləşmə mövqeləri və quruluş xüsusiyyətləri ilə bölgənin tarixi-mədəni irsini əks etdirən əsas nümunələrdəndir. Bu qalaların əksəriyyəti həm təbii relyefdən istifadə olunaraq, həm də süni şəkildə müdafiə məqsədilə inşa edilmişdir. Tarixi mənbələr və arxeoloji tapıntılar sübut edir ki, bu istehkamlar dövrünün hərbi, iqtisadi və siyasi zərurətlərindən yaranmış və müxtəlif dövrlərdə aktiv şəkildə istifadə olunmuşdur. Astara ərazisindən keçən ticarət yolları, o cümlədən İpək Yolu, bu ərazidə istehkamların inşasını daha da zəruri etmişdir. Qalaların ətrafında aşkarlanan saxsı qablar, silah qalıqları və digər maddi mədəniyyət nümunələri burada uzunmüddətli yaşayış və hərbi fəaliyyətin izlərini aydın şəkildə göstərir. Yerli əhalinin yaddaşında qalan rəvayətlər və toponimlər bu qalaların tarixinin xalq şüurunda necə yaşadığını nümayiş etdirir. Bəzi qalaların memarlıq xüsusiyyətləri isə onların eramızdan əvvəlki dövrlərə qədər gedib çıxdığını göstərir ki, bu da Astaranın regionda qədim sivilizasiyaların mərkəzlərindən biri olduğunu sübut edir. Təəssüf ki, bu abidələrin bir çoxu hələ də tam şəkildə tədqiq olunmamış, bəzi hissələri isə dağılaraq məhv olmaq təhlükəsi ilə üz-üzədir. Bu baxımdan, Astara bölgəsinin müdafiə istehkamlarının kompleks şəkildə arxeoloji və tarixi araşdırmalarının davam etdirilməsi və mühafizəsi vacib elmi və mədəni vəzifə kimi qarşıda durur.

İSTİFADƏ OLUNMUŞ MƏNBƏ VƏ ƏDƏBİYYATLAR:

Abbasova E.İ. Şirvan orta əsrlərdə (IX-XVII əsrlər). Bakı-2019 (181 səh.)

Axundov T.İ. Astarada 20 gün. Bakı-2009 (112 səh.)

Mirzə Əhməd Xudaverdi oğlu “Əxbərnamə”. Tərcümə: Əli Hüseynzadə. PDF

Əliyarov S.S., Mahmudov Y.M. Azərbaycan tarixi üzrə qaqynaqlar. Bakı-1989 (328 səh.)

Əliyev T.V. Oğuz və ətraf ərazilərinin tarixi-arxeoloji tədqiqi. Bakı-2017 (160 səh.)

Əliyev T.V. Zərdab orta əsrlərdə, Bakı-2018 (140 səh.)

Kərimov H. Odlar yurdunun səyyah və coğrafiyaşünasları. Bakı-1986

Qasımlı V.A. Naxçıvan III-IX əsrlərdə (tarixi-arxeoloji tədqiqat). Bakı-2013 (254 səh.)

Quliyev S., Vəliyev E. Azərbaycanda multikulturalizm və onun Lənkəran qaynaqları. Bakı-2018 (120 səh.)

Məmmədov A.M.Gəncəbasar IV-XIII əsrlərdə (tarixi-arxeoloji tədqiqat). Bakı-1992 (203 səh.)

Məmmədov A.M. Qədim Gəncə (arxeoloji tədqiqatlar əsasında) Bakı-2014 (175 səh.)

Məmmədov N. Azərbaycanın yer adları. Bakı-1993

FULL TEXTS BOOK

Məmmədzadə Ş.K. “Xülasət ət-təvarix” Azərbaycan tarixinin mənbəyi kimi, Bakı- 1981

Mustafazadə T. Azərbaycan xanlıqlarının qısa tarixi. Bakı-2011, (388 səh.)

Путешественники об Азербайджане. Составил: З.И.Ямрольский, Ваку-1961 (498 стр.)

Səidəli Kazımbəyoğlunun “Cəvahirnameyi Lənkəran” (Səidiyyə). Tərcümə: Səfər Şirinov, Bakı-2005, (143 səh.)

Ягут аль-Хамави «Муджам Ал-Булдан», Ваку-1983

GEMİ ANA MAKİNELERİNDE YAPAY SİNİR AĞI MODELLERİ: MEVCUT UYGULAMALAR VE GELECEK PERSPEKTİFİ

ARTIFICIAL NEURAL NETWORK MODELS IN SHIP MAIN ENGINES: CURRENT APPLICATIONS AND FUTURE PERSPECTIVES

***Samet GÜRGEN*[1]**

[1]*Dr. Öğr. Üyesi, İskenderun Teknik Üniversitesi, Barbaros Hayrettin Gemi İnşaatı ve Denizcilik Fakültesi, Gemi İnşaatı ve Gemi Makineleri Mühendisliği Bölümü, Hatay, Türkiye.*

[1]*ORCID ID: https://orcid.org/0000-0001-7036-8829*

ÖZET

Mevcut küresel ekonomi için uluslararası ticarette deniz taşımacılığı, diğer taşıma modları ile karşılaştırıldığında önemli bir paya sahiptir. Gemiler bu sistemin ana unsurlarından biri olup ihtiyaç duyduğu enerjiyi büyük oranda dizel makineler ile karşılamaktadır. Literatürde gemi ana makineleri ile ilgili çalışmalar oldukça geniş bir kapsamda karşımıza çıkmaktadır. Bu çalışmalar gemi tasarım öncesinde ana makine gücünün belirlenmesinden başlayarak, emisyon azaltma ve performans iyileştirme stratejilerine, operasyon halindeki gemilerde yakıt tüketimi tahminine ve arıza teşhisine kadar uzanmaktadır. Son yıllarda popülerliği artan yapay zeka kavramıyla beraber geleneksel metotlarla yapılan gemi ana makine çalışmaların yayında yapay sinir ağları gibi akıllı sistemler kullanılarak gerçekleştirilen çalışmalar artan bir ilgi görmektedir. Bu çalışmada yapay sinir ağları kullanılarak gemi ana makinelerinde yapılan çalışmalar ele alınarak mevcut uygulamalar detaylı olarak ortaya konulmuştur. Buna ek olarak yapay sinir ağlarının, metasezgisel algoritmalar, risk analiz araçları ve çok kriterli karar verme sistemleri gibi farkı yaklaşımlarla entegre edildiği uygulamalar da incelenmiştir. Çalışmanın, gelecekteki araştırmalara yönelik pratik çıkarımlar ve öneriler sunularak bu alanda çalışan araştırmacıların uygulama alanının durumunu ve gelişim eğilimini anlamalarına katkı sağlaması beklenmektedir.

Anahtar Kelimeler: Gemi ana makinesi, yapay sinir ağları, modelleme.

ABSTRACT

Maritime transport holds a significant share of international trade compared to other modes of transport for the current global economy. Ships are a key element of this system, largely powered by diesel engines. Studies on ship main engines are quite extensive in the literature. These studies range from determining the main engine power before ship design to emission reduction and performance improvement strategies, fuel consumption estimation, and fault diagnosis in operational ships. With the increasing popularity of artificial intelligence in recent years, studies using intelligent systems such as artificial neural networks have attracted increasing attention in comparison to studies on ship main engines conducted using traditional methods. This study examines studies on ship main engines using artificial neural networks and presents a detailed overview of current applications. Additionally, applications integrating artificial neural networks with various approaches, such as metaheuristic algorithms, risk analysis tools, and multi-criteria decision-making systems, are also examined. By offering practical implications and recommendations for future research, the study is expected to contribute to researchers in this field's understanding of the current state of the field and its development trends.

Keywords: Ship main engine, artificial neural networks, modeling.

FULL TEXTS BOOK

GİRİŞ

Dünya çapındaki malların yaklaşık %80'inin taşınmasından sorumlu denizcilik sektörü, küresel ticaret ve kalkınma açısından hayati öneme sahiptir (Raptodimos ve Lazakis, 2020). Deniz taşımacılığın temel unsuru ise ticaret gemileridir. Gemiler, inşa edilip suya indirilmeden önce karmaşık ve zaman alıcı tasarım aşamalarından geçmektedir. Gemi dizaynının başlangıç aşamasında gemi ana boyutları ve ana makine gücünün mümkün olduğunca doğru ve hızlı bir şekilde tahmin edilmesi önemli bir adımdır. Ana makine gücünün belirlenmesi için model deneyleri ya da profesyonel yazılımlar kullanılmaktadır (Ekinci vd., 2011). Bununla beraber son yıllarda makine öğrenme algoritmaları ve özellikle de yapay sinir ağları (YSA) kullanılarak ana makine gücü tahmin modellerinin geliştirilmesi önemli bir araştırma alanıdır.

Gemilerde ana makine olarak çoğunlukla dizel motorlar kullanılmaktadır. Gemi dizel motorları, bir taşıt motorunda bulunan dizel motorlarla aynı çalışma ilkesine sahip olmakla beraber gemilerde kullanılan motorlar daha büyük ve karmaşık sistemlerle donatılmıştır. Dizel motorlardan kaynaklı egzoz emisyonları ile ilgili ulusal ve uluslararası kısıtlar gün geçtikçe ağırlaşmaktadır. Ayrıca gemi sahipleri işletme giderlerinde büyük bir paya sahip olan ana makine yakıt tüketimlerini azaltmak istemektedirler. Dolayısıyla araştırmacılar için dizel motorların verimlerinin arttırılması ve egzoz emisyonlarının düşürülmesi önemli bir araştırma konusudur. Laboratuvar ortamında gerçekleştirilen motor deneyleri pahalı ve zaman alıcıdır. Tüm çalışma şartları için kapsamlı deneyler yerine YSA ile performans ve egzoz emisyon tahmin modelleri geliştirmek bu sorunun üstesinden gelmek için öne çıkan alternatif çözümlerden biridir (Noor vd., 2016; Rodriguez vd., 2021).

Seyir halindeki bir gemi için en önemli gider kalemlerin başında yakıt tüketimi gelmektedir. Yakıt tüketimi operasyonel giderler olarak değerlendirilmekte olup geminin seyir maliyetinin yaklaşık yarısını oluşturmaktadır. Dolaysıyla gemiler için enerji verimliliği hayati öneme sahiptir (Kee vd., 2018; Le vd. 2020). Gemilerde enerji verimliliğin arttırılması için teknik ve operasyonel uygulamalar karşımıza çıkmaktadır. Teknik uygulamalar; gemi gövde optimizasyonu, verimli tahrik ve pervane sistemlerinin geliştirilmesi, dizel motorlarının verimlerinin artırılması, atık ısı geri kazanım sistemlerinin geliştirilmesi, yeni yakıtlar ve motor teknolojileri olarak özetlenebilir. Operasyonel uygulamalar ise sefer planlama, hız, draft, trim, balast suyu, rota ve kargo optimizasyonu ile otopilot sistemleri olarak özetlenebilmektedir. Tüm bunlar içinde en pratik ve etkili yöntemlerin başında hız optimizasyonu gelmektedir (Öztürk ve Başar, 2022). Gemi hızı ile yakıt tüketimi arasında doğrusal olmayan bir ilişki söz konusudur. Gemi hızının %10 düşürülmesi, geminin tükettiği yakıt miktarını yaklaşık %27 azaltmaktadır (Beşikçi vd., 2016). Bu nedenle, makine öğrenmesi yöntemleri ile seyir halindeki bir geminin yakıt tüketimini tahmin eden modeller geliştirilmesi ve hızın belirli bir rota için optimize edilmesi gemiler için önemli bir enerji verimliliği uygulamasıdır.

Gemi ana makinelerinin sorunsuz çalışması seyir güvenliği için çok önemlidir. Ana makinede meydana gelen basit arızalar ciddi can ve mal kayıplarına sebep olabilir (Tong vd., 2022). Denizcilik sektöründe bakım faaliyetleri genel olarak düzeltici, önleyici ve kestirimci bakım olmak üzere üç kategoriye ayrılmaktadır (Göksu ve Erginer, 2020). Fakat günümüz denizcilik operasyonları için kestirimci bakım öne çıkmaktadır. Kestirimci bakım, durum bazlı bakım olarak da bilinmektedir. Bu sistemde sensörlerden gelen veriler ile arızların meydana gelmeden önce tahmin edilmesi sağlanır. Böylece gereksiz denetimlerden kaçınılarak zaman ve para tasarrufu sağlanır. Ayrıca bu akıllı denetim sistemi, hem deniz taşımacılığının güvenliğini hem de verimini artırmaktadır (Maione vd., 2024)

Yapay sinir ağaları her alanda olduğu gibi denizcilik alanında da oldukça popüler bir yaklaşımdır. Bu çalışmada gemi ana makine özelinde YSA uygulamaları incelenmiştir. Literatür incelendiğinde öne çıkan çalışmalar, gemi tasarım öncesinde ana makine güç tahmini modelleri, dizel motor performans

FULL TEXTS BOOK

iyileştirme ve emisyon azaltma için gerçekleştirilen modelleme çalışmaları, seyir halindeki bir gemi için yakıt tüketim tahmin modelleri ve ana makine arıza teşhisi modelleri olarak sıralanabilir. Bu çalışmanın, gelecekteki araştırmalara yönelik pratik çıkarımlar ve öneriler sunularak bu alanda çalışan araştırmacıların uygulama alanının durumunu ve gelişim eğilimini anlamalarına katkı sağlaması beklenmektedir.

GEMİ ANA MAKİNELERİNDE YSA UYGULAMALARI

Bu bölümde sırasıyla YSA kullanılarak gerçekleştirilen gemi tasarım öncesinde ana makine güç tahmini modelleri, dizel motor performans iyileştirme ve emisyon azaltma için gerçekleştirilen modelleme çalışmaları, seyir halindeki bir gemi için yakıt tüketim tahmin modelleri ve ana makine arıza teşhisi modelleri ile ilgili öne çıkan çalışmalar özetlenmiştir.

Ana Makine Güç Tahmini

Ekinci vd. (2011) YSA'nında aralarında olduğu toplamda 18 farklı makine öğrenme yöntemi ile kimyasal tankerlerde ana makine gücü tahmin modeli kurmuş ve karşılaştırmalı performans analizi yapmışlardır. 144 adet gerçek gemi verisi ile kullanılarak eğitimi tamamlanan modeller arasından en iyi model, adaptif öğrenme oranının kullanıldığı YSA modeli olmuştur.

Gürgen (2021) üç farklı gemi tipi için YSA kullanarak ana makine gücü tahmin modelleri geliştirmiştir. Çalışmada konteyner gemisi, tanker ve dökme yük gemileri için hız ve yük kapasite bilgisi kullanılarak ana makine gücü modellenmiştir. Sinir ağlarının eğitimi için 2000 yılından sonra inşa edilmiş toplam 122 konteyner gemisi, 450 tanker ve 377 dökme yük gemisi kullanılmıştır. Levenberg-Marquardt algoritması kullanılarak 1-20 arasında gizli nöronla eğitimi tamamlanan modeller arasından en iyi yapı konteyner gemisi, tanker ve dökme yük gemileri için sırasıyla 19, 16 ve 16 gizli nöronla elde edilmiştir. Tüm modellerde ortalama mutlak hata yaklaşık %5'in altında çıkmış ve R değeri bire oldukça yakın çıkmıştır.

Cepowski ve Chorab (2021) konteyner, dökme yük ve tanker için ana makine güç tahmin modeli kurmuşlardır. Veri setinde 2015 ve 2021 yılları arasında inşa edilen 196 dökme yük gemisi, 276 tanker ve 81 konteyner gemisi bulunmaktadır. Kuru yük ve tanker için girdi parametreleri DWT ve gemi hızı iken konteyner gemisi için girdi parametresi TEU ve gemi hızı olarak belirlenmiştir. Çalışma sonucunda model performans göstergesi olarak RMSE değeri seçilmiş olup bu değer konteyner, dökme yük ve tanker için sırasıyla 5380, 1163 ve 1656 olarak hesaplanmıştır. Ayrıca R değerleri de bire oldukça yakın çıkmıştır.

Özsarı (2023) konteyner, kargo ve tanker gemileri için ana makine gücü tahmin modelleri geliştirmiştir. YSA ile modelleme sürecinde ağın eğitimi için gerekli veri seti, 762 konteyner, 1442 tanker ve 816 kargo gemisinden oluşmaktadır. Modellerin çıktısı ana makine gücü iken girdi bilgisi olarak inşa yılı, gemi tipi, motor silindir hacmi ve motor strok uzunluğu gibi toplam 14 parametre kullanılmıştır. Farklı aktivasyon fonksiyonu, öğrenme algoritması ve gizli nöron sayıları kullanılarak yapılan performans analizi neticesinde en iyi yapı Levenberg Marquardt öğrenme algoritması ve tansig-purelin transfer fonksiyonun kullanıldığı ve 30 gizli nöronla elde edilmiştir. R değerlerini konteyner, kargo ve tanker gemileri için sırasıyla 0.99773, 0.98964 ve 0.97755 olarak elde edilmiştir.

Güneş vd. (2023) 836 tanker gemisine ait verileri kullanarak YSA ve regresyon ile tahmin modelleri kurmuş ve karşılaştırmalı performans analizi gerçekleştirmiştir. Modeller için girdi parametreleri DWT, uzunluk, genişlik ve gros ton olarak belirlenmiştir. Regresyon modelleri arasından en iyi modelin R^2 değeri 0.827 olarak hesaplanmıştır. YSA ile model kurmadan önce mevcut veriler 68% eğitim, 16% doğrulama ve 16% test oranlarında ayrılmıştır. YSA eğitim sonucunda, eğitim, doğrulama ve test için korelasyon katsayılarının sırasıyla 0.98075, 0.98056 ve 0.98104 olarak hesaplanmıştır.

FULL TEXTS BOOK

Gürgen (2023) soğutuculu (reefer) gemiler için YSA ile ana makine gücü tahmin modeli geliştirmiştir. Veri setinde toplam 89 reefer gemi bulunmakta olup bu verilerin 63'ü modelin eğitimi, 13'ü doğrulama ve kalan 13'ü model performansı için kullanılmıştır. Modelin girdi parametreleri DWT ve gemi hızı, çıktı parametresi ise ana makine gücüdür. Ağın eğitimi için Levenberg-Marquardt algoritması kullanılmış ve farklı sayıda gizli nöron denenerek en iyi model belirlenmiştir. Çalışma sonucunda en iyi model 20 gizli nöronda elde edilmiştir. Performans analizine göre RMSE değerleri eğitim, doğrulama ve test veri seti için sırasıyla 591.755, 690.717 ve 1134.85 olarak elde edilmiştir. R değerleri de bire oldukça yakın çıkmıştır.

Özsarı (2023) balıkçı tekneleri için ana makine gücü tahmin modelleri geliştirmiştir. YSA modeli için girdi parametresi olarak balıkçı gemisine ait 12 farklı özellik kullanılmıştır. Ağın kurulması için toplamda 800 balıkçı gemisine ait veri bir seti kullanılmıştır. En iyi modeli elde etmek için çeşitli modelleme denemelerinin ardından en iyi model Levenberg-Marquardt algoritması ve 35 gizli nöronda elde edilmiştir. Bu model için regresyon değerleri eğitim, doğrulama, test için sırasıyla 0.98717, 0.98565 ve 0.98428 olarak elde edilmiştir.

Çirçir ve Gürgen (2024) YSA kullanarak genel kargo gemileri için ana makine güç tahmin modeli sunmuşlardır. Veri setinde 2000 ile 2021 yılları arasında inşa edilmiş toplam 1310 genel kargo gemisi bulunmaktadır. Model girdi parametreleri geminin boyu, genişliği, gros tonajı, servis hızı ve DWT iken çıktı parametresi gemi ana makine gücüdür. Eğitim sırasında toplam verinin %70'i ağın eğitimi için kullanılmış %15, doğrulama ve kalan %15'i test için ayrılmıştır. Levenberg-Marquardt algoritması model eğitimi için seçilmiş ve farklı sayıda gizli nöronla modelleme çalışmaları gerçekleştirilmiştir. Elde edilen modeller arasından en iyi model 22 gizli nöronla elde edilmiştir. Normalize edilmiş verilerle hesaplanan eğitim, doğrulama ve test veri seti için RMSE değerleri sırasıyla 0.0178, 0.02 ve 0.0238 olarak hesaplanmıştır. Ayrıca R değerleri bire oldukça yakın çıkmıştır.

Makine Performans ve Emisyon Tahmin Modelleri

Kowalski (2009) gemi ana makinesinin NOx ve yakıt tüketimi için YSA ile tahmin modeli üretmiştir. Deneysel çalışma Gdynia Denizcilik Üniversitesi gemi Makineleri Laboratuvarı'nda gerçekleştirilmiş olup soğutma sistemindeki su sıcaklıkları, egzoz gazı sıcaklığı, emme havası sıcaklığı, motor devir sayısı gibi toplam 15 girdi parametresi kaydedilmiştir. Çeşitli motor yüklerinde ve 200 ve 360 dev/dak sabit motor devrinde deneysel çalışmalar gerçekleştirilmiştir. Modelle aşamasında farklı gizli nöron sayıları kullanılarak en uygun yapı elde edilmiştir. Çalışma sonucunda NOx emisyonu için en iyi modelin gizli nöron sayısı 24 olarak tespit edilmiş ve hata oranının %4'den düşük olduğu belirtilmiştir. Yakıt tüketimi için modelin performansı hesaplandığında, hata oranın %10'dan düşük olduğu gösterilmiştir.

Wang vd. (2010) gemi ana makinelerinden kaynaklı emisyonlar için YSA ile tahmin modeli geliştirmişlerdir. Modelleme için gerekli veriler RT-flex60C deniz dizel motorundan elde edilmiştir. YSA modellerinin eğitimi için hem Ölçekli Eşlenik Gradyan (SCG) hem de Levenberg-Marquardt (LM) algoritmaları kullanılmıştır. Gizli katmandaki nöron sayısı 5-10 arasında değiştirilerek denenmiştir. Çalışma sonucunda en iyi model SCG ile eğitilmiş ve 7 gizli nörona sahip yapıda elde edilmiştir. Performans sonuçları incelendiğinde R değeri bire oldukça yakın ve RMS değeri de oldukça düşük çıktığı görülmüştür.

Kökkülünk vd. (2013) dizel motorlarda egzoz emisyonlarını tahmin etmek için YSA modelleri geliştirmişlerdir. Model eğitimi için gerekli veriler tek silindirli, doğal emişli, 4 zamanlı ve su soğutmalı gerçek bir dizel motor kullanılarak deneysel çalışma ile elde edilmiştir. Modelin girdi parametreleri motor devri, emülsiyon yakıt yüzdesi ve motor yükü iken çıktı parametreleri ise, HC, CO, CO_2, NO, NOx ve egzoz gazı sıcaklığı olarak belirlenmiştir. Standart geri yayılım (BP) öğrenme algoritması ve radyal tabanlı fonksiyon (RBF) kullanılarak karşılaştırmalı model performans analizi de yapılmıştır.

FULL TEXTS BOOK

Sonuçlar, her iki algoritma ile eğitimi yapılan modellerin performansının oldukça iyi olduğunu göstermiştir. Ancak RBF, BP'den daha iyi bir performansa sahip olduğu vurgulanmıştır.

Noor vd. (2015) %5 oranında biyodizel içiren karışım yakıtı (B5) kullanarak motor performans parametrelerini tahmin eden bir YSA modeli kurmuşlardır. Deneysel çalışma 4 zamanlı, sıralı, 6 silindirli bir Cummins NT855 deniz dizel motorunda gerçekleştirilmiştir. Motor devir sayısı fren gücü, tork ve yakıt tüketimi ölçülmüş, fren özgül yakıt tüketimi (BSFC), fren termal verimi ve hacimsel verim daha sonra hesaplanmıştır. Elde edilen verilerin %70'i eğitim, %15'i test ve kalan %15'i doğrulama için ayrılarak geri yayılımlı sinir ağı modeli kurulmuştur. En güçlü modeli elde etmek için gizli katmandaki nöron sayısı 4 ile 28 nöron arasında denenmiştir. Sonuçlar incelendiğinde fren gücü, tork, fren özgül yakıt tüketimi, fren termal verimliliği ve hacimsel verimlilik için YSA tahminleri sırasıyla 0.9984, 0.9714, 0.9130, 0.9456 ve 0.9976'lık bir korelasyon katsayısı ürettiği görülmüş ve sonuçların oldukça iyi olduğu vurgulanmıştır.

Noor vd. (2016) gemi dizel makinesi için momenti, fren gücünü, fren özgül yakıt tüketimini ve egzoz gazı sıcaklığını tahmin eden YSA modeli geliştirmiştir. Verileri elde etmek için deneysel bir çalışma yapılmıştır. Elde edilen veriler kullanılarak Levenberg-Marquardt eğitim algoritması ile YSA modelleri üretilmiştir. En iyi modeli elde etmek için gizli katmandaki nöron sayısı 10 ile 34 arasında değiştirilmiştir. Çalışmada YSA modellerinin performans karşılaştırmasını yapmak için regresyon modelleri de kurulmuştur. Sonuçlar incelendiğinde en uygun YSA modeli 22 gizli nöronda elde edildiği görülmüştür. YSA modelinin tahmin hata yüzdesi % 0.58 ila % 2.78 arasında değişirken, regresyon modellerinin % 8.83 ila % 53.73 arasında olmuştur. Sonuçlar YSA modellerinin, regresyon modellerine göre üstünlüğü göstermiştir.

Rodriguez vd. (2021) gemi ana makinesi için ön püskürtmenin etkisini incelemek için bir YSA modeli geliştirmişlerdir. Çalışmada altı sıralı silindire sahip dört zamanlı Wärtsilä 6L 46 dizel motoru için analiz yapılmıştır. Modelleme için gerekli veriler, deneysel çalışmalarla doğrulanmış CFD analizlerinden elde edilmiştir. Toplamda 180 tane verinin 126'sı öğrenme, 27'si test ve 27'si doğrulama için kullanılmıştır. Çalışmanın temel amacı NOx emisyonlarını azaltmak için en uygun ön enjeksiyon başlangıç anı, miktar ve süresini belirlemektedir. Sonuçlar, R değerinin bire çok yakın oluğunu göstermiştir. Ayrıca en uygun parametreleri seçmek için çok kriterli karar verme modeli oluşturulmuştur.

Seyir Halindeki Gemilerin Yakıt Tüketim Modelleri

Beşikçi vd. (2016) YSA ile farklı operasyonel koşullar için yakıt tüketimi tahmin eden bir model geliştirmiştir. Ayrıca geliştirilen modelle, gemilerin enerji verimini artırmak için bir karar destek sistemi oluşturmuşlardır. Çalışmada kullanılan veriler, bir tankere ait 17 aylık verileri içeren 233 tane rapordan alınmıştır. Modeller için girdi parametreleri gemi hızı, dakikadaki devir sayısı (RPM), ortalama draft, trim, gemideki kargo miktarı, rüzgar ve deniz etkilerini içermektedir. Çalışmada YSA ile geliştirilen modelin üstünlüğünü göstermek için regresyon modelleri de oluşturulmuş ve karşılaştırmalı performans analizi gerçekleştirilmiştir. Sonuçlar yakıt tüketimi tahmini için YSA modellerinin, regresyon modellerine göre daha güçlü olduğunu göstermiştir.

Kee vd. (2018) römorkörlerin yakıt tüketimin tahmini için YSA modeli üretmiştir. Çalışmada Malezya'nın Rejang Nehri ve Malakka Boğazı boyunca sefer yapan iki kardeş römorkörün verilerinden yararlanılmıştır. Yakıt tüketimini modellemek için seyir mesafesi, seyir süresi, gemi hızı, dedveyt tonaj (DWT) ve rüzgar hızı faktörleri girdi değişkenleri olarak seçilmiştir. Çalışmada, YSA'nın performansı, çoklu doğrusal regresyon analizi (MLR) ile karşılaştırılmış ve bir vaka çalışmasıyla örneklendirilmiştir. YSA modelinin, gemi performansının karmaşık ve doğrusal olmayan ilişkisini tahmin etmede MLR regresyon modelinden daha iyi uyum ve doğruluk sağladığı sonucuna varılmıştır. Ayrıca elde edilen model ile bir karar destek sistemi kurulmuştur.

FULL TEXTS BOOK

Le vd. (2020) yüzden fazla konteyner gemisinin gerçek seyir verilerini kullanarak yakıt tüketimi tahmin modeli geliştirmiştir. Gemileri boyutlarına göre 5 farklı gruba ayırarak hem çoklu regresyon hem de YSA ile uygulamalar yapılmış ve elde edilen modellerin karşılaştırılması gerçekleştirilmiştir. Çalışmada girdi parametrelerinin etkisini incelemek için seyir süresi, hız, kargo ağırlığı ve kapasitenin kullanıldığı modellerin yanında yalnızca fiziksel fonksiyonlardan iki girdinin kullanıldığı modeller de kullanılmıştır. Sonuçlar YSA modellerinin regresyon modellerine göre daha üstün performans verdiğini göstermiştir. Ayrıca yalnızca fiziksel parametrelerin girdi olarak kullanıldığı modellerin performansının, dört girdili olanlara göre daha düşük kaldığı vurgulanmıştır.

Moreira vd. (2021) Manş Denizi ile Miami bölgesi arasındaki transatlantik seyir yapan konteyner gemisi için yakıt tüketimi ve gemi hızını tahmin eden bir YSA modeli kurmuşlardır. Çalışmanın bağımsız değişkenleri olarak gemi hızı, ana makine torku, devir sayısı ve dalga ile ilgili üç parametre seçilmiştir. Çalışma sonunda modellerin oldukça tatmin edici sonuçlar ürettiği görülmüştür. Çalışmada ayrıca yalnızca deniz koşulları hakkındaki bilgileri ile tahmin modelleri de üretilmiş ve bu modellerin de oldukça başarılı sonuçlar ürettiği gösterilmiştir.

Öztürk ve Başar (2022) 19 adet konteyner gemisine ait ortalama 24 aylık sefer raporlarına dayanarak yakıt tüketimi için tahmin modeli geliştirmiştir. Çalışmada YSA yanında çoklu doğrusal regresyon yöntemi de kullanılarak karşılaştırmalı analiz yapılmıştır. Draft, trim, RPM, pitch (hatve), hava durumu verileri ile yakıt tüketimi tahmini yapılmıştır. Sonuçlar YSA modellerinin regresyon modellerine göre daha üstün olduğunu göstermiştir.

Ana Makine Arıza Teşhis Modelleri

Wang vd. (2020) gemi ana makinelerindeki turboşarj ve yakıt enjeksiyonu arızaları için YSA kullanarak teşhis modeli geliştirmiştir. Çalışmada MAN B&W L16/24 dizel motorunu vaka analizi için seçilmiş ve simülasyonu için AVL-BOOST yazılımı kullanılmıştır. İlk olarak temel bileşen analizi kullanılmış ve 16 girdi parametresi 3'e düşürülmüştür. Daha sonra sinir ağı modeli geliştirilmiş ve test edilmiştir. Sonuçlar YSA kullanılarak geliştirilen teşhis modelinin oldukça başarılı olduğunu ortaya koymuştur.

Hou vd. (2020) gemi dizel makinelerinde hata teşhisi için YSA modellerinin genetik algoritma (GA) ile iyileştirilmesi üzerine çalışmışlardır. Çalışmada silindir egzoz çıkış sıcaklığı, silindir hava giriş sıcaklığı, silindir gömleği su çıkış sıcaklığı gibi toplam 10 girdi parametresi dikkate alınarak toplam 10 arıza için analiz yapılmıştır. Modelleme için gerekli veriler, Kongsberg gemi makine dairesi simülatöründen alınmış olup toplamda 700 arıza simülasyon deneyi yapılmıştır. Geleneksel model ile GA ile modifiye edilmiş model karşılaştırıldığında, GA ile iyileştirilmiş modelin daha iyi sonuçlar ürettiği görülmüştür.

Göksu ve Erginer (2020) bir konteyner gemisinin 10 yıllık arıza verilerini kullanarak gelecekte hangi ekipmanın arızalanacağını tahmin eden bir YSA modeli geliştirmiştir. Çalışmada ana makinenin yanı sıra yardımcı sistemleriyle ilgili ekipman arızaları da modellenmiş ve toplamda 47 sistem incelenmiştir. Sonuçlar geliştirilen YSA modelinin arızaları tahmininde oldukça başarılı olduğunu ortaya koymuştur.

Raptodimos ve Lazakis (2020) ana makine arızası için önemli bir teşhis kaynağı olan silindir egzoz gazı çıkış sıcaklığını tahmin eden bir model geliştirmiştir. Çalışmada iki zamanlı, sekiz silindirli dizel motoruna sahip Panamax tipi bir konteyner gemisi kullanılmış ve modelleme için toplamda 920 saatlik bir veri elde edilmiştir. Egzoz gazı çıkış sıcaklığının gelecekteki değerlerini tahmin etmek için doğrusal olmayan otoregresif dışsal girişli (NARX) bir YSA modeli geliştirilmiştir. İlk 900 saatlik ölçüm NARX modelinin eğitimi için kullanılırken, son 20 saatlik ölçüm performans kriterlerini değerlendirmek ve modelin tahmin gücünü göstermek için kullanılmıştır.

Tong vd. (2022) gemi ana makine arıza teşhisi için YSA modeli kurmuşlardır. Çalışmanın temel amacı, standart geriye yayılım algoritması ile eğitimi yapılmış model ile GA ile optimize edilmiş bir modelin

FULL TEXTS BOOK

performans karşılaştırmasının yapılmasıdır. Çalışma sonunda GA ile eğitilmiş bir YSA modelinin, standart modele göre daha üstün performans sağladığı görülmüştür.

Castresana vd. (2023) aktif olarak kullanılan bir balıkçı gemisini dikkate alarak ana makinenin sorunsuz çalışması için YSA ile modelleme yapmışlardır. Çalışmanın temel amacı, ana makinede herhangi bir arıza meydana gelmeden uyarıcı bir dizi eşik değeri sağlamaktır. Çalışmada 2020 yılının tamamına ek olarak 2021 yılının da yarısına ait toplam 5181 saat arızasız çalışma verisi kullanılmıştır. 2020 yılına ait veriler ile modelleme gerçekleştirilmiş ve 2021 yılındaki veriler ile doğrulama yapılmıştır. Yüksek korelasyon ve değişken değerleri YSA ile değerlendirilirken düşük korelasyon ve sabit değerler olasılık yoğunluk fonksiyonu incelenmiştir. Çalışma sonucunda elde edilen eşiklerin, gelecekteki arıza tespit çalışmaları için rehber olması için sistematik bir özeti sunulmuştur.

Maione vd. (2024) arıza tahmininde algoritma seçiminin önemini vurgulamak için YSA'nın da aralarında olduğu üç farklı makine öğrenmesi algoritmasının karşılaştırmalı performans analizini yapmışlardır. Çalışmanda tahmin hatasının yanı sıra hatanın zaman tepkisi de dikkate alınarak bir performans analizi gerçekleştirilmiştir. Vaka çalışması için bir gemi ana makinesindeki yağ devresi arızaları dikkate alınmıştır.

Dong vd. (2025) gemi ana makinesi için beş temel arızaya yönelik bir model geliştirmiştir. Çalışmada Wärtsilä W9L34DF motoru için hem gerçek veriler hem de AVL-BOOST'ta oluşturulmuş simülasyon verileri kullanılmıştır. Efektif güç ve yakıt tüketimi girdi parametreleri olarak kullanılırken çıktılar maksimum yanma basıncı, emme manifoldu basıncı, egzoz manifoldu basıncı, emme manifoldu sıcaklığı, egzoz manifoldu sıcaklığı, turboşarj debisi, turboşarj çıkış basıncı, kompresör çıkış sıcaklığı ve hava soğutucu çıkış sıcaklığından oluşmaktadır. Her motor alt sisteminin çalışma durumuna etkisini belirlemek için AHP (Analitik Hiyerarşi Süreci) ve CRITIC yöntemleri ile parametre ağırlıkları belirlenmiştir. Daha sonra parçacık sürü optimizasyonu (PSO) ile optimize edilmiş sinir ağları ile model oluşturulmuş ve elde edilen çıktılar ile risk seviyelendirilmesi yapılmıştır.

TARTIŞMA VE SONUÇLAR

Gemilerin ön tasarımında uzunluk, genişlik, draft gibi temel boyutlarının ve ana makine gücünün hızlı ve zahmetsiz belirlenmesi önemli bir aşamadır. Ana makine gücünü tahmininde geleneksel olarak model havuz deneyleri veya profesyonel yazılımlar kullanılmaktadır (Ekinci vd., 2011). Bununlar beraber son yıllarda makine öğrenmesi teknikleriyle ana makine güç tahmin modellerinin kurulması önemli bir araştırma konusudur. Literatürde çeşitli makine öğrenmesi yöntemlerinin yanında YSA ile oluşturulan modeller güçlü tahmin yetenekleri ile öne çıkmaktadır. Birçok gemi tipi için gerçekleştirilen YSA ile ana makine gücü tahmin çalışmalarda, model girdi parametreleri genellikle temel gemi boyutları, yük kapasitesi (DWT, GT ve TEU) ve geminin servis hızını içermektedir. Bunun yanında bazı çalışmalarda farklı girdi parametreleri de kullanılmıştır. Eğitim algoritması olarak çoğunlukla Levenberg-Marquardt kullanılmış olup bazı çalışmalarda farklı birkaç algoritma ile performans karşılaştırması yapılmıştır. En iyi modeli elde etmek için farklı gizli nöron sayıları ile denemeler gerçekleştirilmiştir. Bazı çalışmalarda aktivasyon fonksiyonları sabit olarak alınırken bazı çalışmalarda farklı kombinasyonlar denenerek en güçlü model elde edilmeye çalışılmıştır. Modellerin performansı ölçmek için MSE, RMSE, MAPE ve R değerleri kullanılmıştır. Genel olarak tüm çalışmalarda model performansları tatmin edici seviyede elde edilmiştir. Gelecek çalışmalarda her gemi tipi için ana makine gücü tahmininde girdi parametrelerinin belirlenmesi için kapsamlı bir çalışma gerçekleştirilebilir. Model amacına hizmet edecek ve aynı zamanda model karmaşıklığını arttırmayacak şekilde girdi parametreleri optimize edilebilir. Ayrıca model eğitimi için gizli nöron sayısı, aktivasyon fonksiyonları, veri setinin bölünmesi (eğitim, doğrulama ve test) gibi temel model parametrelerinin deneme yanılma yoluyla belirlenmesi yerine optimizasyon algoritmaları ile entegre edilerek belirlenmesi düşünülebilir. Son olarak model

FULL TEXTS BOOK

eğitimi için geleneksel algoritmalar yerine metasezgisel algoritmalar kullanılarak performans iyileştirme çalışmaları yapılabilir.

Genel olarak gemilerde ana itici güç büyük dizel motorları ile sağlanmaktadır. Otomobil motorlarına göre çalışma koşulları farklı olan gemi dizel motorlarının performans iyileştirmesi ve emisyon oranlarının düşürülmesi hayati bir öneme sahiptir. Dolayısıyla motor laboratuvar deneyleri önemli bir araştırma alanıdır. Fakat gemi makineleri gibi yüksek güçlü motorlarda kapsamlı deneyler yapmak hem çok pahalı hem de oldukça zahmetlidir. Bunun yerine sınırlı deney verileri elde etmek ve model oluşturmak en pratik çözümdür (Noor vd., 2016). Literatürde standart dizel motor ile yapılan çalışmaların aksine, gemi ana makineleri kullanılarak gerçekleştirilen deneysel ve modelleme çalışmaları sınırlıdır. Mevcut çalışmaların bir kısmı sadece HC, CO, CO_2, NO, NOx gibi egzoz emisyonlarına odaklanırken bazıları da motor momenti, gücü ve yakıt tüketimi gibi performans göstergelerine odaklanmıştır. Bununla beraber bazı çalışmalarda hem motor performansı hem de egzoz emisyonları incelenmiştir. Çoğu çalışmada standart geri yayılım algoritması kullanılmış ve eğitim için Levenberg-Marquardt optimizasyonu tercih edilmiştir. Farklı sayıda gizli nöron ile denemeler yapılarak en güçlü tahmin modeli elde edilmiştir. Günümüzde gemiler için hala dizel motoru ve dizel yakıtı ana enerji kaynağıdır. Fakat son yıllarda karışım yakıtlar, pilot püskürtmeli çift yakıtlı motorlar egzoz emisyonlarının düşürülmesi için güçlü alternatifler olarak karşımıza çıkmaktadır. Ayrıca dizel motorlar yerine hibrit ya da tamamen elektrik motorlarının kullanıldığı gemiler de suya indirilmektedir. Dolayısıyla sadece geleneksel dizel yakıtı ile çalışan dizel motorların yerine alternatif yakıtlar ve motor teknolojilerinin kullanıldığı çalışmaların sayısının da artması beklenmektedir. Ek olarak seyir yapan gemilerin üzerinde gerçek hava ve deniz koşulları ile farklı yükleme koşulları altında motor performans ve egzoz emisyon çalışmaları yapılıp, YSA ile modeller kurulabilir.

Denizcilik şirketleri gün geçtikçe sıkılaşan egzoz emisyon düzenlemelerine uyum sağlamak ve aynı zamanda gemilerin operasyonel maliyetleri üzerinden ciddi bir paya sahip yakıt tüketimlerini düşürmek için çeşitli stratejilere ihtiyaç duymaktadır. Bunlar arasında en etkili ve pratik olanlardan biri gemi hızının düşürülmesidir. Dolayısıyla gemiler seyirleri sırasından rota, yük, hava ve deniz durumuna göre yakıt tüketimini tahmin eden modellere ihtiyaç duymaktadır. Bu modeller daha sonra karar destek sisteminin bir parçası olarak hızın optimize edilmesi için kullanılmaktadır (Kee vd., 2018); Le vd., 2020). Literatürde konteyner, romörkör ve tanker gibi çeşitli gemi tipleri için YSA ile yakıt tüketimi tahmin modelleri geliştirilmiştir. Seyir halindeki gemilerin belirli zaman dilimine ait gerçek verileri modellemelerde kullanılmıştır. Genel olarak gemi hızı, draft, trim, motor devir sayısı, hava ve deniz durumu tahmin modelinin girdilerini oluşturmuştur. Elde edilen modeller, karar destek sistemleri ile entegre edilerek belirli bir rotada seyir yapan geminin hızı optimize edilmiştir. Mevcut çalışmalarda elde edilen modellerin tahmin kabiliyeti regresyon modelleri ile karşılaştırılmış ve YSA'nın üstün bir performans gösterdiği görülmüştür. Mevcut literatürde genellikle bir gemi tipi için belirli zamanda elde edilen veriler kullanılmıştır. Gelecekte yakıt optimizasyonu için küresel bir bulut sistemi kurulup birçok gemiden elde edilen veriler burada depolanarak derin öğrenme ile yapay zeka asistanı kullanılabilir. Ağın eğitimi online olarak sürekli yapılıp, küresel olarak tüm gemiler için hız optimizasyon asistanı geliştirilebilir. Geliştirilen modeller, rota planlama yazılımları ve metasezgisel optimizasyon algoritmalı ile entegre edilebilir. Böylece denizcilik şirketleri hem daha az yakıt kullanarak karlarını artırabilir hem de çevre kirliliği önemli düzeyde düşürülebilir. Mevcut çalışmalar genellikle dizel motorlar temelli gerçekleştirilmiştir. Gelecekte LNG, metanol, amonyak, hibrit ya da tam elektrikli gemilerin sayısı artacaktır ve bu motorların da yakıt tüketim modelleri farklı parametreler içerecektir. Dolayısıyla bu yenilikçi teknolojiler için de benzer modeller geliştirilebilir.

Yüksek sıcaklık ve basınçlara maruz kalan dizel motorlar gemilerde ana makine olarak kullanılmakta olup sorunsuz çalışması geminin emniyeti ve güvenliği için hayati öneme sahiptir. Gemi ana makinesi arızalandığında ekonomik kayıplar, çevre felaketi, mürettebatın yaralanması ve hatta ölümüyle

FULL TEXTS BOOK

sonuçlanan ciddi kazalara sebep olabilir. Dolayısıyla gemi ana makinelerinde arıza teşhisi önemli bir çalışma alanı olarak karşımıza çıkmaktadır (Tong vd., 2022; Dong vd., 2025). Gemilerde ana makine ile ilgili düzenli bakımlar yapılmaktadır. Ayrıca geleneksel olarak mürettebat tecrübesine dayanarak arızalar hakkında bazı öngörüler yapılmaktadır. Fakat bu geleneksel sistem hatalı sonuçlar doğurabilmekte ve ciddi ekonomik kayıplar yaratmaktadır (Dong vd., 2025). Literatürde arıza erken uyarı sistemleri için temelde iki yöntem benimsenmiştir. Bunlardan ilki termodinamik tabanlı motor modelleri iken diğeri veri odaklı yöntemdir. Motor modellerinin gerçek zamanlı teşhise engel olan uzun hesaplama süresi en büyük dezavantajlarından biridir. Ayrıca zorlu ve değişken deniz şartlarının temsili zordur. Buna karşın veri odaklı modellerle, geçmiş verilerin işlenerek hızlı ve erken bir uyarı sistemi elde etmek mümkündür (Liu vd., 2022; Castresana vd., 2023). Literatürde farklı yöntemlere ek olarak YSA ile gemi ana makineleri için arıza teşhis modelleri önemli bir araştırma alanıdır. Bu çalışmaların bir kısmında model eğitimi için makine dairesi simülasyon verileri kullanılırken bazı çalışmalarda gerçek gemi verileri kullanılmıştır. Birçok çalışmada farklı makine öğrenmesi algoritmaları kullanılarak karşılaştırmalı performans analizi yapılmıştır. Ayrıca GA kullanılarak YSA modellerinin iyileştirildiği çalışmalar da dikkat çekmektedir. Gelecek çalışmalarda ana makine sisteme ait tüm alt bileşenlere daha fazla sensör yerleştirilip buradan elde edilen veriler bir bulut sisteminde depolanabilir daha sonra her bir alt sistem için geliştirilecek detaylı teşhis modelleri paralel olarak çalıştırılıp bu bütün çıktıları işleyen bir karar destek sistemi oluşturulabilir. YSA modellerinin doğruluğunu arttırmak için metasezgisel optimizasyon algoritmaları detaylı olarak incelenebilir. Ayrıca termodinamik modeller ile YSA hibrit bir şekilde kullanılarak daha güçlü ve hızlı modeller geliştirilebilir. Son olarak geleneksel dizel motorlara alternatif teknolojiler (LNG, metanol, elektrikli, hibrit vb.) kullanılan gemiler için de benzer çalışmalar gerçekleştirilebilir.

Aşağıdaki metni geliştir:

Bu çalışmada YSA kullanılarak gemi ana makine özelinde gerçekleştirilen çalışmalar hakkında genel bir bakış sunulmuştur. Çalışma, kapsamlı bir derleme niteliği taşımaktan ziyade, alanın mevcut durumuna dair genel bir çerçeve ortaya koymayı amaçlamaktadır. Ayrıca gelecek çalışmalar için öneriler sunarak bu alanda çalışacak araştırmacılar için ilham kaynağı olabilecektir. Gelecekte sadece ana makine olarak dizel motorlar yerine LNG, metanol, elektrikli, hibrit vb gibi yenilikçi teknolojilerin kullanıldığı ve YSA ile modellemelerin yapıldığı çalışmalar gerçekleştirilebilir. Ayrıca diğer gemi sistemleri ile YSA'nın kullanıldığı çalışmalar da kapsamlı olarak incelenebilir.

KAYNAKLAR

Beşikçi, E. B., Arslan, O., Turan, O., & Ölçer, A. I. (2016). An artificial neural network based decision support system for energy efficient ship operations. *Computers & Operations Research*, *66*, 393-401.

Castresana, J., Gabina, G., Quincoces, I., & Uriondo, Z. (2023). Healthy marine diesel engine threshold characterisation with probability density functions and ANNs. *Reliability Engineering & System Safety*, *238*, 109466.

Cepowski, T., & Chorab, P. (2021). The use of artificial neural networks to determine the engine power and fuel consumption of modern bulk carriers, tankers and container ships. *Energies*, 14(16), 4827.

Çirçir, E., & Gürgen, S. (2024). Artificial Neural Network Approach for Main Engine Power Prediction of General Cargo Vessels. *Mersin Üniversitesi Denizcilik ve Lojistik Araştırmaları Dergisi*, *6*(2), 113-129.

Dong, X., Zeng, H., Li, C., Ye, K., & Lu, Z. (2025). Health condition assessment of marine diesel engine sub-system based on PSO-BP neural network. *Ships and Offshore Structures*, 1-17.

FULL TEXTS BOOK

Ekinci, S., Çelebi, U. B., Bal, M., Amasyali, M. F., & Boyaci, U. K. (2011). Predictions of oil/chemical tanker main design parameters using computational intelligence techniques. *Applied Soft Computing*, *11*(2), 2356-2366.

Göksu, B., & Erginer, K. (2020). Prediction of ship main engine failures by artificial neural networks. *Journal of ETA Maritime Science*, *8*(2).

Gunes, U., Başhan, V., & Karakurt, A. S. (2023). Predicting tanker main engine power using regression analysis and artificial neural networks. *Sigma Journal of Engineering and Natural Sciences*, *41*(2), 216-225.

Gürgen, S. (2021). Yapay zeka yaklaşımları ile gemi ana makinesinin belirlenmesi ve optimum organik rankine çevrimli atık ısı geri kazanım sisteminin kurulması, Doktora Tezi, Karadeniz Teknik Üniversitesi, Fen Bilimleri Enstitüsü, Trabzon.

Gürgen, S. (2023). Artificial Neural Network-Based Approach to Predict Main Engine Power in Reefer Ships, 3. International Mediterranean Congress, 17-18 April 2023, pp. 718-725, Mersin, Türkiye.

Hou, L., Zou, J., Du, C., & Zhang, J. (2020). A fault diagnosis model of marine diesel engine cylinder based on modified genetic algorithm and multilayer perceptron. *Soft Computing*, *24*(10), 7603-7613.

Kee, K. K., Simon, B. Y. L., & Renco, K. H. Y. (2018, September). Artificial neural network back-propagation based decision support system for ship fuel consumption prediction. In *5th IET International Conference on Clean Energy and Technology (CEAT2018)* (pp. 1-6). IET.

Kökkülünk, G., Akdoğan, E., & Ayhan, V. (2013). Prediction of emissions and exhaust temperature for direct injection diesel engine with emulsified fuel using ANN. *Turkish Journal of Electrical Engineering and Computer Sciences*, 21(8), 2141-2152.

Kowalski, J. (2009). ANN based evaluation of the NOx concentration in the exhaust gas of a marine two-stroke diesel engine. *Polish Maritime Research*, 60-66.

Le, L. T., Lee, G., Park, K. S., & Kim, H. (2020). Neural network-based fuel consumption estimation for container ships in Korea. *Maritime Policy & Management*, *47*(5), 615-632.

Liu, B., Gan, H., Chen, D., & Shu, Z. (2022). Research on fault early warning of marine diesel engine based on CNN-BiGRU. *Journal of Marine Science and Engineering*, *11*(1), 56.

Maione, F., Lino, P., Maione, G., & Giannino, G. (2024). A machine learning framework for condition-based maintenance of marine diesel engines: a case study. *Algorithms*, *17*(9), 411.

Moreira, L., Vettor, R., & Guedes Soares, C. (2021). Neural network approach for predicting ship speed and fuel consumption. *Journal of Marine Science and Engineering*, *9*(2), 119.

Noor, C. M., Mamat, R., Najafi, G., Nik, W. W., & Fadhil, M. (2015, November). Application of artificial neural network for prediction of marine diesel engine performance. In IOP Conference Series: Materials Science and Engineering (Vol. 100, No. 1, p. 012023). IOP Publishing.

Nor, C. M., Mamat, R., Najafi, G., Yasin, M. M., Ihsan, C. K., & Noor, M. M. (2016). Prediction of marine diesel engine performance by using artificial neural network model. *Journal of Mechanical Engineering and Sciences*, 10(1), 1917-1930.

Ozsari, I. (2023). Predicting main engine power and emissions for container, cargo, and tanker ships with artificial neural network analysis. *Brodogradnja: An International Journal of Naval Architecture and Ocean Engineering for Research and Development*, *74*(2), 77-94.

Özsarı, İ. (2023). Calculating the main engine power of fishing vessels with artificial neural networks analysis, 2nd International Conference on Scientific and Academic Research, 14-16 March 2023, pp. 515-520, Konya, Türkiye

FULL TEXTS BOOK

Öztürk, O. B., & Başar, E. (2022). Multiple linear regression analysis and artificial neural networks based decision support system for energy efficiency in shipping. *Ocean Engineering*, *243*, 110209.

Raptodimos, Y., & Lazakis, I. (2020). Application of NARX neural network for predicting marine engine performance parameters. *Ships and Offshore Structures*, *15*(4), 443-452.

Rodriguez, C. G., Lamas, M. I., Rodriguez, J. D., & Abbas, A. (2021). Analysis of the pre-injection system of a marine diesel engine through multiple-criteria decision-making and artificial neural networks. *Polish Maritime Research*, (4), 88-96.

Tong, Z., Sun, Y., She, J., Zhu, Y., & Zhao, Z. (2022). Identification of typical fault states of marine diesel engines based on optimized BP neural network. *Highlights in Science, Engineering and Technology*, *7*, 10-18.

Wang, M., Zhang, J., Zhang, S., & Ma, Q. (2010, August). Predication emission of an intelligent marine diesel engine based on modeling of bp neural networks. In 2010 Sixth International Conference on Natural Computation (Vol. 3, pp. 1290-1296). IEEE.

Wang, S., Wang, J., & Ding, X. (2020, March). An Intelligent Fault Diagnosis Scheme Based On PCA-BP Neural Network for the Marine Diesel Engine. In *IOP Conference Series: Materials Science and Engineering* (Vol. 782, No. 3, p. 032079). IOP Publishing.

FULL TEXTS BOOK

KARİYER STRESİ İLE UMUTSUZLUK ARASINDAKİ İLİŞKİ: ÜNİVERSİTE ÖĞRENCİLERİ ÜZERİNE BİR ARAŞTIRMA

THE RELATIONSHIP BETWEEN CAREER STRESS AND HOPELESSNESS: A STUDY ON UNIVERSITY STUDENTS

Doç. Dr. Semra TETİK

Manisa Celal Bayar Üniversitesi, Salihli MYO, Pazarlama ve Reklamcılık Bölümü, Manisa, Türkiye

ORCID ID: 0000-0001-6266-8178

ÖZET

Stres, kişinin yaşamının tüm alanlarını önemli düzeyde etkileyen bir unsurdur. İnsan yaşamı süresince çeşitli stresörlerle karşılaşma olasılığı taşırken, kariyer konusu da yaşamın önemli stres faktörlerinden birisidir. Kişinin kariyerine ilişkin beklentileri psikolojik ve bedensel stresle karşı karşıya kalmasına neden olabilmektedir. Kariyer stresi kişinin kariyer yolculuğunda kariyerinin değişik aşamalarında problemlerle başa çıkma sürecinde ortaya çıkan durumdur. Diğer bir ifadeyle kariyer stresi, kişinin kendi sınırlarını zorlaması sonucu yaşanan gerilim halidir. Kişinin yaşam sürecinde kritik karar vermesini gerektiren konulardan belki de en önemlisi kariyer kararıdır. Kişinin kariyerinin şekillendiği önemli dönemlerin başında da üniversite yaşamı vardır. Üniversite sürecinde, öğrencilerin geleceklerine dair hedeflerini belirleyerek kariyer planı yapmaları gerekmektedir. Ancak öğrenciler öğrenim gördükleri alana ilişkin yeterli bilgiye sahip olmamaları, çevre baskısı, kararsızlık, kariyer belirsizliği gibi nedenlerle kariyer stresi ile karşı karşıya kalmaktadır. Bu da öğrencileri umutsuzluğa sürüklemektedir. Umutsuzluk kişilerin çaresiz ve güçsüz hissetmeleri, şimdi yaşadıkları ya da gelecekte karşılaşma olasılığının olduğu zorlukların üstesinden gelemeyeceklerine inanmaları, yaşamaya dair arzu ve enerjilerini yitirmeleri olarak ifade edilebilir. Öğrencilerin kariyer stres düzeyleri arttığında umutsuzluk düzeylerinin de artacağı düşünülmektedir. Bu noktadan hareketle bu çalışmanın amacı, kariyer stresi ile umutsuzluk arasındaki ilişkiyi üniversite öğrencileri özelinde incelemektir. İlişkisel tarama modelinde tasarlanan araştırmanın çalışma grubunu 2024-2025 eğitim öğretim yılı bahar döneminde Manisa Celal Bayar Üniversitesi'ne bağlı Salihli İktisadi ve İdari Bilimler Fakültesi ve Salihli Meslek Yüksekokulu'nda aktif olarak öğrenim gören ve araştırmaya gönüllü olarak katılan 206 öğrenci oluşturmaktadır. Öğrencilere Kişisel Bilgi Formu, Kariyer Stresi Ölçeği ve Beck Umutsuzluk Ölçeği uygulanmıştır. Veriler SPSS 22.0 programında analiz edilmiştir. Demografik özelliklerin dağılımını belirlemek için frekans ve yüzde analizi, kariyer stresi ve umutsuzluk düzeylerini belirlemek için tanımlayıcı istatistik analizi yapılmıştır. Kariyer stresi ve umutsuzluk arasındaki ilişkinin test edilmesinde Pearson Korelasyon analizi uygulanmıştır. Analiz sonucunda üniversite öğrencilerinin algılarına göre kariyer streslerinin orta düzeyde ve umutsuzluklarının hafif düzeyde olduğu tespit edilmiştir. Kariyer stresi ve alt boyutları ile umutsuzluk düzeyleri arasında pozitif yönlü ve anlamlı ilişkiler bulunmuştur.

Anahtar Kelimeler: Stres, Kariyer Stresi, Umutsuzluk

ABSTRACT

Stress significantly impacts all areas of a person's life. While people are likely to encounter various stressors throughout their lives, careers are also a significant stressor. Career expectations can lead to psychological and physical stress. Career stress is a state of tension that arises when a person struggles to cope with problems at various stages of their career journey. In other words, career stress is the tension

FULL TEXTS BOOK

experienced as a result of pushing their own limits. Perhaps the most important aspect of a person's life that requires critical decisions is career decision-making. University life is one of the most crucial periods in shaping a person's career. During university, students are required to determine their future goals and make a career plan. However, students often face career stress due to factors such as inadequate knowledge of their field of study, peer pressure, indecision, and career uncertainty. This, in turn, leads students to despair. Hopelessness can be defined as a feeling of helplessness and powerlessness, a belief that they cannot overcome the challenges they are currently experiencing or may face in the future, and a loss of desire and energy for life. It is thought that as students' career stress levels increase, their hopelessness levels will also increase. Therefore, the purpose of this study is to examine the relationship between career stress and hopelessness among university students. Designed with a correlational screening model, the study group consisted of 206 students actively studying at the Salihli Faculty of Economics and Administrative Sciences and Salihli Vocational School affiliated with Manisa Celal Bayar University during the spring semester of the 2024-2025 academic year, who volunteered to participate in the study. A Personal Information Form, the Career Stress Inventory, and the Beck Hopelessness Inventory were administered to the students. Data were analyzed using SPSS 22.0. Frequency and percentage analysis was conducted to determine the distribution of demographic characteristics, and descriptive statistics were used to determine levels of career stress and hopelessness. Pearson correlation analysis was used to test the relationship between career stress and hopelessness. The analysis revealed that university students' perceptions of career stress were moderate and their hopelessness mild. There are positive and significant relationships between career stress and its sub-dimensions and hopelessness levels.

Keywords: Stress, Career Stress, Hopelessness

GİRİŞ

Günümüz dünyasında kişilerin öncelikleri arasında kariyer merkezli başarıları ve mesleki gelişim hedefleri vardır. İşletmelerin yoğun rekabet içinde çalışmaları ve her daim değişen talepleri, kişilerin kariyer yolculuklarında yüksek beklentiler ve bazı güçlüklerle karşı karşıya kalmalarına neden olmaktadır. Bu güçlükler, stres gibi olumsuz sonuçlara yol açarak arzu edilmeyen durumları beraberinde getirmektedir (Kahraman, 2023). Stres, organizmanın sınırlarını tehdit eden durumlara karşı verilen tepki olarak ifade edilebilir. Stres, bir taraftan kişinin başarılı olma, zorluklarla mücadele etme gibi konularda motivasyonu artırıcı bir etki yaratırken diğer taraftan stres düzeyi arttıkça kişileri zihinsel, fiziksel ve ruhsal açıdan olumsuz etkilemektedir (Üzüm vd., 2018). Bu bağlamda stres, kişinin yaşamının her yönünü önemli düzeyde etkileyen bir olgudur. Kişiler yaşamının her döneminde zaman baskısı, beklenmedik durumlar, çevresel faktörler gibi stresörlerle karşılaşabilir. Kişilerin kariyer süreçleri perspektifinden bakıldığında ise yaşamlarının önemli stres dönemlerinden birisinin üniversite süreci olduğu söylenebilir (Özden ve Sertel-Berk, 2017). Üniversite sürecinde, öğrencilerin geleceklerine ilişkin hedeflerini belirleyerek kariyer planı yapmaları gerekmektedir. Ancak öğrencilerin öğrenim gördükleri alan ile ilgili bilgilerinin yetersiz olması, mezuniyet sonrası iş bulamama endişesi, çevre baskısı, kariyerde ilerleyebilme konusunda belirsizlik gibi nedenlerle kariyer stresi ile karşı karşıya kaldıkları ve umutsuzluğa kapıldıkları bir gerçektir (Akbağ, 2024).

İnsanların yaşamları boyunca devam eden bir süreç olarak kariyer, "sorumluluk, prestij, maddi ayrıcalıklar ve mesleki itibar elde etmek" olarak tanımlanabilir (Turpcu ve Akyurt, 2018). Diğer bir ifadeyle kariyer, kişinin liyakat esasına göre çalışma yaşamı boyunca bulunduğu statü ve pozisyonları ve bu kapsamda sahip olduğu bilgi ve becerileri içeren bir kavramdır (Emirel ve Bozkurt, 2022). Kariyer stresi ise iş yaşamında karşılaşılan baskılar, iş yükü yoğunluğu, belirsizlik ve aşırı rekabet gibi nedenlerden dolayı ortaya çıkan bir gerilim halidir (Bayrakçeken ve Buztepe, 2021). Bu bağlamda

FULL TEXTS BOOK

kariyer stresi, bireylerin kariyer sürecinde zorluklarla karşılaştırdıklarında meydana gelen bir durumdur (Güney, 2004). Choi ve arkadaşlarına (2011) göre, kariyer stresi üç boyuttan oluşmaktadır. Bunlar; kariyer belirsizliği ve bilgi eksikliği, dışsal çatışma ve iş bulma baskısıdır. Kariyer belirsizliği ve bilgi eksikliği; bireyin öz farkındalık düzeylerinin düşük olmasını, yapacağı iş ile ilgili bilgisinin yetersiz olmasını, gelecekte işine dair ne yapacağı konusunda belirsizliği içeren ve strese neden olan bir boyuttur. Dışsal çatışma, bireyin kariyer tercihleri ile ailesinin ve dış çevresinin kariyer tercihlerinin farklı olmasını içeren bir boyuttur. İş bulma baskısı ise bireyin işe alım sürecinde yapılan sınavlarda başarılı olamama endişesi, kariyer alanında gelebileceği pozisyonların sınırlı olması gibi durumları içeren bir boyuttur (Esen, 2019).

Üniversite yılları öğrencilerin kariyerlerine ilişkin önemli kararlar vermeleri gereken bir dönemdir. Bu dönemde öğrenciler kararsızlık, belirsizlik ve çevre baskısı gibi nedenlerden dolayı çeşitli zorluklarla karşılaşabilirler (Polat, 2022). Öğrencilerin kariyerlerine ilişkin yaşayabilecekleri zorluklar ise stres düzeylerini artırarak umutsuzluğa düşmelerine neden olabilir (Duru ve Gültekin, 2020). Umutsuzluk kavramı, kişinin hedefine ulaşmada sıfırdan daha küçük olan olumsuz bir beklentiye sahip olmasıdır (Dilbaz ve Seber, 1993). Umut ise, kişinin hedeflerine ulaşmalarını kolaylaştıracak yolları araması ve bulmasıdır (Snyder, 2002; Atilla ve Yıldırım, 2023) . Umutsuzluk, geleceğe dair olumsuz bir bakış açısı olarak da tanımlanabilir (Ehtiyar ve Üngüren, 2008). Zıt beklentileri içeren umut ve umutsuzluk hissi, kişinin hedeflerine ulaşabilme ya da ulaşamamasına dair inancıdır. Umutsuzluk içinde olan kişide başarısızlık düşüncesi yoğundur (Zafer, 2019). Umutsuzluk hissi kişinin psikolojisini olumsuz etkileyerek strese ve depresyona girmesine yol açabilir (Dilbaz ve Seber 1993). Umutsuzluk düzeyi düşük olan kişilerin gelecek ile ilgili olumlu bir tutuma sahip olurlar ve zorluklarla mücadele konusunda dayanıklıdırlar. Dolayısıyla eğitimini aldığı mesleki alanda iş bulmaya ilişkin umudu yüksek olan öğrencilerin kariyer stres düzeylerinin düşük olacağı söylenebilir (Yam, 2020).

YÖNTEM

Araştırmanın Amacı ve Modeli

Araştırmada veri toplamada anket yöntemi kullanılmıştır. Bu araştırmada amaç, kariyer stresi ile umutsuzluk arasındaki ilişkiyi üniversite öğrencileri özelinde incelemektir. Nicel araştırma yöntemlerinden ilişkisel tarama modelinde desenlenen bu araştırmada şu sorulara yanıt aranmıştır:

1. Araştırmaya katılan öğrencilerin kariyer stres düzeyi nedir?
2. Araştırmaya katılan öğrencilerin umutsuzluk düzeyi nedir?
3. Kariyer stresi ve alt boyutları (kariyer belirsizliği ve bilgi eksikliği, dışsal çatışma, iş bulma baskısı) ile umutsuzluk arasında anlamlı bir ilişki var mıdır?

Araştırmanın Çalışma Grubu

Araştırmanın çalışma grubu 2024-2025 eğitim öğretim yıl bahar döneminde MCBÜ bünyesindeki Salihli İİBF ve Salihli MYO'da öğrenim gören öğrencilerden oluşmaktadır. Araştırmaya gönüllü olarak 214 öğrenci katılmış olup, 8 adet anket eksik doldurulduğu için değerlendirme dışı bırakılmıştır. Veriler 206 anket üzerinden değerlendirilmiştir.

Veri Toplama Araçları

Araştırma verilerini toplamak için Kişisel Bilgi Formu, Kariyer Stresi Ölçeği ve Beck Umutsuzluk Ölçeği kullanılmıştır.

Kariyer Stresi Ölçeği: Ölçek Choi ve arkadaşları (2011) tarafından geliştirilmiştir. Özden ve Sertel-Berk (2017) tarafından Türkçe'ye uyarlanmıştır. Orijinal ölçek toplam 20 madde ve kariyer belirsizliği, bilgi eksikliği, dışsal çatışma, iş bulma baskısı olmak üzere 4 alt boyuttan oluşmaktadır. 5'li Likert tipindeki ölçeğin (1= Hiç katılmıyorum ile 5=Tamamen katılıyorum) Türkçe versiyonunun iç tutarlık katsayısı .94 olarak hesaplanmıştır. Açıklayıcı faktör analizi sonucunda "Kariyer Belirsizliği - Bilgi Eksikliği",

FULL TEXTS BOOK

"Dışsal Çatışma" ve "İş Bulma Baskısı" olarak adlandırılan 3 boyut elde edilmiştir (Özden ve Sertel-Berk, 2017: 35-51). Bu çalışmada da ölçeğin Cronbach Alpha kat sayısı .94 olarak bulunmuştur. Bireylerin ölçekten aldıkları puana göre; 1,00-2,49 arası düşük; 2,50-3,49 arası orta; 3,50-5,00 arası yüksek kariyer stresi düzeyinde kabul edilmektedir.

Beck Umutsuzluk Ölçeği: Kişinin geleceğine dair olumsuz beklentilerini ölçmek amacıyla Beck ve arkadaşları (1974) tarafından geliştirilen ölçeğin Türkçe'ye uyarlaması Seber (1991) tarafından yapılmıştır. Ölçek ergen ve yetişkinlere yönelik uygulanabilen ve kişinin geleceğe dair olumsuz beklentileri değerlendiren bir niteliğe sahiptir (Seber vd., 1993: 141). Ölçeğin geçerlilik çalışması ise Durak ve Palabıyıklıoğlu (1994) tarafından yapılmıştır. Ölçeğin Cronbach Alfa güvenilirlik katsayısı .86 olarak bulunmuştur. Ölçek 20 maddeden oluşmakta ve 0 ile 1 arası puanlanmaktadır. Ölçekteki maddelerin 11 tanesi evet; 9 tanesi ise hayır seçeneğinden 1 puan almaktadır. Sonuç olarak 0 ve 20 puan arasında toplam puan elde edilmektedir. Bireylerin ölçekten aldıkları puanlar arttıkça umutsuzluk düzeyleri de artmaktadır. Bu çalışmada da ölçeğin Cronbach Alpha kat sayısı .83 olarak bulunmuştur. Bireylerin ölçekten aldıkları puana göre; 0-3 arası normal, 4-8 arası hafif, 9-14 arası orta, 15-20 arası yüksek umutsuzluk düzeyinde kabul edilmektedir (Durak ve Palabıyıkoğu, 1994).

Verilerin Analizi

Verilerin analizi için SPSS 22.0 programı kullanılmıştır. Verilere betimsel istatistikler (frekans, aritmetik ortalama, standart sapma) ve Pearson Korelasyon analizi uygulanmıştır.

BULGULAR

Araştırmanın bu kısmında, araştırmaya katılan üniversite öğrencilerine ait verilerin analizi sonucunda elde edilen bulgulara ve bulguların yorumlarına yer verilmiştir.

Katılımcıların Demografik Özellikleri

Tablo 1. Araştırmaya Katılan Öğrencilerin Cinsiyet Dağılımı

Değişken	Grup	N	%
Cinsiyet	Kız	106	51,5
	Erkek	100	48,5
	Toplam	206	100,0

Katılımcıların %51,5'i kız, %48,5'i ise erkektir.

Tablo 2. Araştırmaya Katılan Öğrencilerin Yaş Dağılımı

Değişken	Grup	N	%
Yaş	18-20 yaş	85	41,3
	21-23 yaş	99	48,1
	24 ve üzeri	22	10,7
	Toplam	206	100,0

Katılımcıların %41,3'ü 18-20 yaş aralığında, %48,1'i 21-23 yaş aralığında, %10,7'si ise 24 yaş ve üzerindedir.

Tablo 3. Araştırmaya Katılan Öğrencilerin Akademik Birim Dağılımı

Değişken	Grup	N	%

FULL TEXTS BOOK

Akademik Birim	Salihli MYO	114	55,3
	Salihli İİBF	92	44,7
	Toplam	206	100,0

Katılımcıların %55,3'ü Salihli Meslek Yüksekokulu'nda, %44,7'si ise Salihli İktisadi ve İdari Bilimler Fakültesi'nde öğrenim görmektedir.

Tablo 4. Araştırmaya Katılan Öğrencilerin Sınıf Düzeyi Dağılımı

Değişken	**Grup**	**N**	**%**
Sınıf Düzeyi	1.Sınıf	78	37,9
	2.Sınıf	84	40,8
	3.Sınıf	36	17,5
	4.Sınıf	8	3,9
	Toplam	206	100,0

Katılımcıların %37,9'u 1.sınıf, %40,8'i 2.sınıf, %17,5'i 3.sınıf, %3,9'u ise 4.sınıf öğrencisidir.

Tablo 5. Araştırmaya Katılan Öğrencilerin Anne Eğitim Durumu Dağılımı

Değişken	**Grup**	**N**	**%**
Anne Eğitim Durumu	İlkokul	76	36,9
	Ortaokul	52	25,2
	Lise	53	25,7
	Üniversite	25	12,1
	Toplam	206	100,0

Katılımcıların anne eğitim durumuna bakıldığında; %36,9'u ilkokul, %25,3'ü ortaokul, %25,7'si lise, %12,1'inin ise üniversite mezunu olduğu görülmektedir.

Tablo 6. Araştırmaya Katılan Öğrencilerin Baba Eğitim Durumu Dağılımı

Değişken	**Grup**	**N**	**%**
Baba Eğitim Durumu	İlkokul	50	24,3
	Ortaokul	58	28,2
	Lise	56	27,2
	Üniversite	42	20,4
	Toplam	206	100,0

Katılımcıların baba eğitim durumuna bakıldığında; %24,3'ü ilkokul, %28,2'si ortaokul, %27,2'si lise, %29,4'ünün ise üniversite mezunu olduğu görülmektedir.

FULL TEXTS BOOK

Tablo 7. Araştırmaya Katılan Öğrencilerin Aile Gelir Düzeyi Dağılımı

Değişken	Grup	N	%
Aile Gelir Düzeyi	Düşük	20	9,7
	Orta	168	81,6
	Yüksek	18	8,7
	Toplam	206	100,0

Aile gelir düzeyine bakıldığında; katılımcıların %9,7'si düşük, %81,6'sı orta, %8,7'sinin ise yüksek gelir düzeyinde olduğunu belirtmiştir.

Tablo 8. Araştırmaya Katılan Öğrencilerin Yaşadığı Yer Dağılımı

Değişken	Grup	N	%
Yaşadığı Yer	Köy/Kasaba	24	11,7
	İlçe	50	24,3
	İl	15	7,3
	Büyükşehir	117	56,8
	Toplam	206	100,0

Katılımcıların %11,7'si köy/kasabada, %24,3'ü ilçede, %7,3'ü ilde, %56'sı ise büyükşehirde yaşamaktadır.

Ölçeklere İlişkin Tanımlayıcı İstatistikler

Tablo 9. Kariyer Stresi Ölçeği ve Alt Boyutlarına Ait Tanımlayıcı İstatistikler

Değişkenler	N	Ortalama	Std. Sapma	Min.	Max.
Kariyer Stresi Ölçeği	206	2,63	,810	1	5
Kariyer Belirsizliği ve Bilgi Eksikliği	206	2,64	,920	1	5
Dışsal Çatışma	206	2,22	1,046	1	5
İş Bulma Baskısı	206	2,89	1,006	1	5

Araştırmaya katılan öğrencilerin algılarına göre, kariyer stres düzeyleri 2,63±,810'dur. Kariyer stresinin alt boyutları kapsamında bakıldığında; kariyer belirsizliği ve bilgi eksikliği düzeyleri 2,64±,920, dışsal çatışma düzeyleri 2,22±1,046 ve iş bulma baskısı düzeyleri ise 2,89±1,006'dır. Bu sonuçlara göre öğrencilerin orta düzeyde kariyer stresine sahip oldukları söylenebilir.

Tablo 10. Umutsuzluk Ölçeğine Ait Tanımlayıcı İstatistikler

Değişkenler	N	Ortalama	Std. Sapma	Min.	Max.
Umutsuzluk Ölçeği	206	5,31	3,417	0	20

Araştırmaya katılan öğrencilerin algılarına göre, umutsuzluk düzeyleri 5,31±3,417'dir. Buna göre öğrencilerin hafif düzeyde umutsuzluğa sahip oldukları söylenebilir.

FULL TEXTS BOOK

Ölçeklere İlişkin Korelasyon Analizi

Tablo 11. Kariyer Stresi Ölçeği ve Alt Boyutları ile Umutsuzluk Ölçeği Arasındaki Korelasyon Analizi Sonuçları

Değişkenler		Umutsuzluk Ölçeği
Kariyer Stresi Ölçeği	r	,522**
	p	,000
Kariyer Belirsizliği ve Bilgi Eksikliği	r	,506**
	p	,000
Dışsal Çatışma	r	,446**
	p	,000
İş Bulma Baskısı	r	,497**
	p	,000

*p<0.001, **p<0.001

Tablo 11'de görüldüğü gibi, kariyer stresi ölçeği ile umutsuzluk ölçeği (r=,522; p<0.001) arasında pozitif yönlü, orta düzeyde ve anlamlı bir ilişki; kariyer stresi alt boyutlarından kariyer belirsizliği ve bilgi eksikliği ile umutsuzluk ölçeği (r=,506; p<0.001) arasında pozitif yönlü, orta düzeyde ve anlamlı bir ilişki; dışsal çatışma ile umutsuzluk ölçeği (r=,446; p<0.001) arasında pozitif yönlü, orta düzeyde ve anlamlı bir ilişki; iş bulma baskısı ile umutsuzluk ölçeği (r=,497; p<0.001) arasında pozitif yönlü, orta düzeyde ve anlamlı bir ilişki olduğu tespit edilmiştir.

SONUÇ, TARTIŞMA VE ÖNERİLER

Bu çalışma üniversite öğrencilerinin algısına göre kariyer stresi ile umutsuzluk düzeyleri arasındaki ilişkiyi incelemek amacıyla yapılmıştır. Araştırmaya 206 öğrenci katılmıştır. Araştırma sonucunda ulaşılan sonuçlar aşağıda özetlenmiştir.

Ölçeklere ilişkin tanımlayıcı istatistikler incelendiğinde; araştırmaya katılan öğrencilerin orta düzeyde kariyer stresi altında oldukları ve kariyer stresi alt boyutları olan kariyer belirsizliği ve bilgi eksikliği, dışsal çatışma, iş bulma baskısı kapsamında da orta düzeyde stres yaşadıkları görülmektedir. Öğrencilerin algısına göre kariyerleri ile ilgili orta düzeyde stresli olmakla beraber iş bulma baskısı en yüksek ortalamaya (2,89) sahipken, dışsal çatışma en düşük ortalamaya (2,22) sahiptir. Literatürde bu sonucu destekleyen çalışmalar vardır. Emirel ve Bozkurt (2022) tarafından yapılan çalışmada üniversite öğrencilerinin kariyer stresi algılarının orta düzeyde olduğu, kariyer stresi boyutlarından en yüksek ortalamaya sahip boyutun iş bulma baskısı, en düşük ortalamaya sahip boyutun ise dışsal çatışma olduğu ortaya konulmuştur. Turpcu ve Akyurt (2018) tarafından üniversite öğrencileri üzerine yapılan araştırma sonucunda, öğrencilerin genel kariyer stresi ve alt boyutları kapsamında stres algılarının orta düzeyde olduğu, en yüksek ortalama iş bulma baskısı boyutunda tespit edilirken, en düşük ortalama dışsal çatışma boyutunda tespit edilmiştir. Araştırmamızın bu bulgusunu destekleyen başka çalışmalara da rastlanmıştır (Aksu ve Engizeki 2024; Gümrükçü Bilgici ve Deniz, 2016; Güler vd., 2023; Haznedar, 2023; Uludağ Güler ve Ünal, 2020; Yılmaz, 2019). Üniversite öğrencilerinin kariyer stresinin orta düzeyin altında tespit edildiği çalışmalar da mevcuttur (Özden ve Sertel-Berk, 2017; Bozyiğit ve

FULL TEXTS BOOK

Gökbaraz, 2020). Araştırmamızda öğrencilerin umutsuzluk algılarının hafif düzeyde olduğu tespit edilmiştir. Bu sonuç literatürle uyumludur (Çelik, 2023; Çığ, 2023; İbaoğlu, 2023).

Kariyer stresi ile umutsuzluk arasındaki ilişki incelendiğinde; genel kariyer stresi ve alt boyutları (kariyer belirsizliği ve bilgi eksikliği, dışsal çatışma, iş bulma baskısı) ile umutsuzluk düzeyi arasında pozitif yönlü anlamlı ilişkiler olduğu görülmüştür. Bunun anlamı öğrencilerin kariyer stresi arttıkça umutsuzluk düzeyi de artmakta ya da kariyer stresi düştükçe umutsuzluk düzeyi de düşmektedir. Diğer bir ifadeyle öğrencilerin umutsuzluk düzeyi arttıkça veya azaldıkça kariyer stresi de artmakta veya azalmaktadır. Literatürde kariyer stresi ve umutsuzluk arasındaki ilişkiyi inceleyen bir çalışmaya rastlanmamıştır. Bu bağlamda, bu çalışmanın literatüre önemli katkı sağlayabilecek özgün bir çalışma olduğu söylenebilir.

Araştırma sonuçları doğrultusunda üniversite gençlerine ve araştırmacılara şunlar önerilebilir:

- Öğrenciler orta düzeydeki kariyer stresi ve umutsuzluk düzeylerini düşürmek için iyi bir kariyer planlaması yapmalıdır.
- Öğrencilerin kariyerlerine ilişkin streslerini azaltacak koşullar oluşturulmalıdır.
- Öğrenciler gelecekte kendilerine uygun meslek seçimi konusunda iyi bir danışmanlık hizmeti almalıdır.
- Kariyer stresi ve umutsuzluk konusunda öğrencilere seminerler verilerek farkındalıkları artırılmalıdır.
- Öğrencilerin mesleki alanlarında kendilerini geliştirmeye yönelik sorumluluk üstlenmelidir.
- Kariyer stresi ve umutsuzluk konusunda devlet üniversiteleri ve vakıf üniversitelerinde karşılaştırmalı araştırmalar yapılabilir.
- Literatürde kariyer stresi ve umutsuzluk arasındaki ilişkiyi inceleyen bir çalışmaya rastlanmamıştır. Dolayısıyla bu çalışma, benzer konuda yapılacak araştırmalara temel oluşturabilir.
- Araştırma bir üniversitenin iki akademik biriminde yapıldığından sonuçlar genellenemez. Dolayısıyla genellenebilmesi için farklı üniversitelerden daha çok sayıda bir örneklem grubuna yönelik araştırma yapılmalıdır.
- Bu araştırma üniversite öğrencileri özelinde yapılmıştır. Bu çalışmadaki değişkenler farklı bir durum tespiti çalışmasında, farklı çalışma gruplarında incelenebilir.

KAYNAKÇA

Akbağ, M., Karagöz, E., ve Ayaz, A. (2024). Üniversite öğrencilerinin kariyer stresleri üzerinde kariyer kaynaklarının rolü. *Türk Eğitim Bilimleri Dergisi*, 22(2), 1628-1644.

Aksu, M., ve Engizek, S. (2024). Lisans öğrencilerinin umutsuzluk düzeylerinin farklı değişkenler açısından incelenmesi. *Süleyman Demirel Üniversitesi Sosyal Bilimler Enstitüsü Dergisi*, 1(48), 190-224.

Atilla, G., ve Yıldırım, G. (2023). Umudun yaşam doyumu ve olumlu gelecek beklentisi üzerine etkisi. *Mehmet Akif Ersoy Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi*, 10(1), 341-359.

Bayrakçeken, E., ve Buztepe, S. (2021). Sağlık Hizmetleri Meslek Yüksekokulu öğrencilerinin kariyer streslerinin bazı değişkenler açısından incelenmesi. *Balıkesir Sağlık Bilimleri Dergisi*, 10(3), 260-266.

Bozyiğit, E., ve Gökbaraz, N. (2020). Spor Bilimleri Fakültesi öğrencilerinde kariyer stresinin belirleyicileri. *Journal of Computer and Education Research*, 8(15), 181-200.

Çelik, F. (2023). *Üniversite öğrencilerinde kaygı, depresyon, umutsuzluk düzeyleri ile ötanaziye yaklaşımları arasındaki ilişkinin incelenmesi*. Yüksek Lisans Tezi, Üsküdar Üniversitesi Sosyal Bilimler Enstitüsü, İstanbul.

FULL TEXTS BOOK

Çığ, G. (2023). *Üniversite öğrencilerinin umutsuzluk düzeyleri ile akademik başarı, sosyal beceri ve kaygıları arasındaki ilişkinin incelenmesi*. Yüksek Lisans Tezi, İstanbul Arel Üniversitesi Lisansüstü Eğitim Enstitüsü, İstanbul.

Dilbaz, N., ve Seber, G. (1993). Umutsuzluk kavramı: Depresyon ve intiharda önemi. *Kriz Dergisi*, 1(3), 134-138.

Duman, N., İmre Y., ve Mısırlı, M. (2019). Üniversite öğrencilerinin umutsuzluk düzeyi ve bilişsel çarpıtmaları üzerine bir araştırma. *Research Studies Anatolia Journal*, 2(5), 207-213.

Durak, A., ve Palabıyıkoğlu, R. (1994). Beck Umutsuzluk Ölçeği geçerlilik çalışması. *Kriz Dergisi*, 2(2), 311-319.

Duru, H., ve Gültekin, F. (2020). Eğitim fakültesi öğrencilerinin kariyer stresleri, kariyer uyum yetenekleri ve mentorluk arasındaki ilişkinin incelenmesi. *Kastamonu Eğitim Dergisi*, 28(1), 328-337.

Ehtiyar, R., ve Üngüren, E. (2008). Lise ve üniversite turizm eğitimi alan öğrencilerin demografik değişkenlerinin umutsuzluk ve kaygı düzeylerine etkilerinin araştırılması. *Gazi Üniversitesi Ticaret ve Turizm Fakültesi Dergisi*, 2, 34-51.

Emirel, M., ve Bozkurt, Ö. (2022). Kariyer stresi ile belirsizlikten kaçınma arasındaki ilişkinin incelenmesi: Sigortacılık ve sosyal güvenlik bölümü öğrencileri üzerine bir araştırma. *Çalışma İlişkileri Dergisi*, 13(1), 1-16.

Esen, D. (2019). Kariyer stresi ile öz yeterlilik inancı ilişkisinin demografik değişkenlerin farklılaştırıcı rolü bağlamında incelenmesi: İzmir Meslek Yüksekokulu İktisadi ve İdari Programlar örneği. *İzmir Kâtip Çelebi Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi*, 2(2), 217-232.

Güler, C., Siyahtaş, A., ve Çakır, V. O. (2023). Spor Bilimleri Fakültesi öğrencilerinin yaşam doyumunu yordayan bir faktör; kariyer stresi: Tanımlayıcı araştırma. *Türkiye Klinikleri Spor Bilimleri Dergisi*, 15(1),159-168.

Gümrükçü Bilgici, B. ve Deniz, Ü. (2016). Okul öncesi öğretmen adaylarının mesleki kaygılarının bazı demografik özelliklere göre incelenmesi. *Cumhuriyet International Journal of Education*, 5(1), 53-70.

Güney, S. (2004). *Açıklamalı yönetim-organizasyon ve örgütsel davranış terimleri sözlüğü*. Ankara: Siyasal Kitabevi.

Haznedar, S. N. (2023). *Kariyer stresi, tükenmişlik ve olumlu gelecek beklentisi ilişkisi: Beden eğitimi öğretmeni adayları örneği*. Yüksek Lisans Tezi, Erzincan Binali Yıldırım Üniversitesi Sağlık Bilimleri Enstitüsü, Erzincan.

İbaoğlu, E. N. (2023). *Üniversite öğrencilerinde internet oyun oynama bozukluğu ile internet bağımlılığı, psikolojik katılık ve umutsuzluk arasındaki ilişkinin incelenmesi*. Yüksek Lisans Tezi, Necmettin Erbakan Üniversitesi Eğitim Bilimleri Enstitüsü, Konya.

Kahraman, F. (2023). Sağlık eğitimi alan üniversite öğrencilerinde kariyer stresinin kariyer geleceği üzerindeki etkisi. *Iğdır Üniversitesi Sosyal Bilimler Dergisi*, (34), 487-502.

Özden, K., ve Sertel-Berk, Ö. (2017). Kariyer Stresi Ölçeği'nin (KSÖ) Türkçe'ye uyarlanması ve psikometrik özelliklerinin sınanması. *Psikoloji Çalışmaları Dergisi*, 37(1), 35-51.

Polat, F. B. (2022). Kariyer stresinin ve kariyer kararlılığının psikolojik iyi oluş üzerindeki etkisi. *Anadolu Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi*, 23(2), 62-80.

Seber, G. (1991). *Beck Umutsuzluk Ölçeğinin Geçerliliği ve güvenirliği üzerine bir çalışma.* Anadolu Üniversitesi Tıp Fakültesi, Psikiyatri Anabilim Dalı Doçentlik Tezi, Eskişehir.

Seber, G., Dilbaz, N., Kaptanoğlu, C., ve Tekin, D. (1993). Umutsuzluk Ölçeği: Geçerlilik ve güvenirliği. *Kriz Dergisi*, 1(3), 139-142.

FULL TEXTS BOOK

Snyder, C. R. (2002). Hope theory: Rainbows in the mind. *Psychological Inquiry*, 13(4), 249- 275.

Turpcu, E., ve Akyurt, H. (2018). Turizm eğitimi alan öğrencilerin kariyer streslerinin belirlenmesi: Giresun Üniversitesi lisans öğrencileri üzerine bir araştırma. *The Journal of Academic Social Science Studies*, 69, 365-380.

Uludağ Güler, Ç., ve Ünal, E. (2020). Gençlerde kariyer stresinin ve duygu-tutumlarının ölçülmesi: Üniversite öğrencileri üzerinde uygulamalı bir araştırma. *International Social Sciences Studies Journal*, 6(68), 3748-3762.

Üzüm, B., Uçkun, S., ve Uçkun, G. (2018). Üniversite öğrencilerinin yaşadığı bir fenomen: Kariyer stresi. *Journal Of Social And Humanities Sciences Research*, 5(24), 1641-1651.

Yam, F. C. (2020). Adalet Meslek Yüksekokulu öğrencilerinin kariyer streslerinin psikolojik iyi oluşları üzerindeki yordayıcı rolü. *Uluslararası Sosyal Araştırmalar Dergisi*, 13(72), 710-721.

Yılmaz, A. (2019). Öğretmen adaylarının kariyer streslerinin karma araştırma yaklaşımı ile incelenmesi. *Türk Spor Bilimleri Dergisi*, 2(1), 93-105.

Zafer, C. (2019). Üniversite öğrencilerinin umutsuzluk nedenleri. *Adam Akademi*, 9(2), 547-569.

FULL TEXTS BOOK

SOSYAL ZEKA İLE GİRİŞİMCİLİK EĞİLİMİ ARASINDAKİ İLİŞKİYİ BELİRLEMEYE YÖNELİK BİR ARAŞTIRMA

A RESEARCH ON THE RELATIONSHIP BETWEEN SOCIAL INTELLIGENCE AND ENTREPRENEURSHIP TENDENCY

Doç. Dr. Semra TETİK

Manisa Celal Bayar Üniversitesi, Salihli MYO, Pazarlama ve Reklamcılık Bölümü, Manisa, Türkiye

ORCID ID: 0000-0001-6266-8178

ÖZET

Sosyal zeka, başkalarının duygularını anlama, yönlendirme becerisi ve insan ilişkilerinde akıllıca davranabilme yetkinliğidir. Sosyal zeka, kişinin sosyal boyutta hedeflerine ulaşma, başkalarıyla iyi ilişkiler kurma ve sürdürme kapasitesini ifade eder. Girişimcilik eğilimi ise kişinin kendi işini kurmaya yönelik yatkınlığını ifade eder. Girişimcilik eğilimi kavramına üniversite öğrencileri perspektifinde bakıldığında potansiyel girişimci anlamına gelmektedir. Girişimcilik eğilimi girişimcilik sürecinin ilk adımıdır. Dolayısıyla girişimcilik eğilimi gelecekte kendi işini kurmayı hayal eden kişiler için önemli bir göstergedir. Bu çalışmada potansiyel birer girişimci olan üniversite öğrencilerinin sosyal zeka düzeyleri ile girişimcilik eğilimleri arasındaki ilişkiyi incelemek amaçlanmıştır. İlişkisel tarama modelindeki araştırmada veriler anket yöntemi ile toplanmıştır. Bu kapsamda 224 üniversite öğrencisine Kişisel Bilgi Formu, Tromso Sosyal Zeka Ölçeği ve Girişimcilik Eğilimi Ölçeği uygulanmıştır. Elde edilen verilere betimleyici istatistikler ve korelasyon analizi uygulanmıştır. Analiz sonucunda üniversite öğrencilerinin sosyal zeka düzeyleri ile girişimcilik eğilimleri arasında pozitif yönlü, orta düzeyde ve anlamlı ilişkiler tespit edilmiştir. Buna göre öğrencilerin sosyal zeka düzeyleri arttıkça girişimcilik eğilimlerinin de artacağı söylenebilir.

Anahtar Kelimeler: Sosyal Zeka, Girişimcilik Eğilimi, Üniversite Öğrencileri

ABSTRACT

Social intelligence is the ability to understand and guide others' emotions, and the ability to act wisely in human relations. Social intelligence refers to a person's capacity to achieve social goals and establish and maintain good relationships with others. Entrepreneurial tendencies, on the other hand, refer to a person's predisposition to start their own business. When viewed from the perspective of university students, the concept of entrepreneurial tendencies refers to potential entrepreneurs. Entrepreneurial tendencies are the first step in the entrepreneurial process. Therefore, entrepreneurial tendencies are an important indicator for those who dream of starting their own business in the future. This study aimed to examine the relationship between the social intelligence levels of university students who are potential entrepreneurs and their entrepreneurial tendencies. Data were collected through a survey method in this relational screening model. A personal information form, a Tromso social intelligence scale, and an entrepreneurial tendencies scale were administered to 224 university students. Descriptive statistics and correlation analysis were applied to the obtained data. The analysis revealed a positive, moderate, and significant relationship between university students' social intelligence levels and their entrepreneurial tendencies. Accordingly, it can be argued that as students' social intelligence levels increase, their entrepreneurial tendencies will also increase.

Keywords: Social Intelligence, Entrepreneurial Inclination, University Students

FULL TEXTS BOOK

GİRİŞ

Sosyal bir varlık olan insan yaşamını sürdürebilmek, amaçlarına ulaşabilmek, sosyal yaşamda ve iş yaşamında başarılı ve mutlu olabilmesi için başka kişi ya da gruplarla etkileşim kurmak durumundadır. Başkalarıyla etkileşim kurmada ve uyumlu çalışmada sosyal zeka önemli bir konu olarak karşımıza çıkmaktadır (Yeşil ve Mavi, 2024). Thorndike (1920) tarafından sosyal zeka, ‘’kişileri anlama ve idare etme sanatı, insan ilişkilerinde ustaca davranma yeteneği’’ olarak ifade edilmiştir (Çakmak Yıldızhan ve Çağlayan, 2019. Walker ve Foley (1973)’e göre sosyal zeka, insanlara duyarlı olma, duygularını ve motivasyonlarını doğru anlama yeteneğidir (Kahya ve Ceylan, 2017). Albert (2006)’e göre sosyal zeka, bireyin başkalarıyla anlaşabilmesi, uyumlu çalışabilmesini sağlayan bir yetenektir (Demirbanka ve Taşlıyan, 2022). Sosyal zeka, başkalarıyla müzakere, işbirliği ve iyi anlaşma yeteneğidir (Genç ve Genç, 2018). Sosyal zeka, insanlarla fikir alışverişinde etkinlik, problemlere yaratıcı çözümler bulma, başkalarıyla iyi ilişkiler ve dostluklar kurma, ikna kabiliyeti, empati kurma gibi becerilerin bireylerde olan zeka türü olmasından dolayı büyük önem taşımaktadır (Vural, 2004).

Silvera ve arkadaşları (2001) sosyal zekanın üç boyuttan oluştuğunu ortaya koymuşlardır. Bunlar; sosyal bilgi süreci, sosyal beceri ve sosyal farkındalıktır (Çakmak Yıldızhan ve Çağlayan, 2019). Sosyal bilgi süreci; bireyin insanlarla etkileşiminde onların düşüncelerini, duygularını, davranışlarını, sözlü ya da sözsüz mesajları anlayabilme, beklentilere cevap verebilme yeteneğidir (Erdemir, ve Kutlu, 2018). Sosyal beceri; sosyalleşmenin gerçekleşmesidir. Sosyal beceri, iletişim sürecinde ve sosyalleşme sürecinde nasıl davranması gerektiği konusunda kişiye yol göstermektedir (Demirbanka ve Taşlıyan, 2022). Sosyal beceri, bireyin başkalarıyla iletişim kurmada özgüvenli olmasını, sosyal ortamlara girmede zorlanmamasını ya da kolaylıkla uyum sağlamasını içerir (Cebeci ve Kamaşak, 2021). Diğer bir ifadeyle sosyal beceri; bireyin sosyal ortamlara ilişkin tutumu ve ortamdaki diğer kişileri etkileyebilme becerisidir. Bireyin sosyal becerisi arttıkça farklı ortamlara girmesi, başka insanlarla etkili ilişkiler kurması ve sürdürmesi kolaylaşır (Yılmazçetin, 2021). Sosyal farkındalık; bireyin girdiği yeni sosyal ortamlara ve koşullara uygun davranabilmesi, kendisine yönelik davranışları anlayabilmesini sağlayan bir yetenektir (Cebeci ve Kamaşak, 2021). Bireyin sosyal farkındalığı ne kadar yüksek olursa sosyal ortamlarda uygun davranışlar sergilemede o kadar başarılı olur (Silvera vd., 2001). Sosyal zekası yüksek kişiler daha dışa dönük, sosyal ve girişken olurlar.

Girişimcilik, ülkede üretimin artmasında, milli gelirin artmasında, ekonominin büyümesi ve gelişmesinde, gelir dağılımının daha adil olmasında, bölgeler arasındaki ekonomik dengesizliğin düzeltilmesinde, teknolojinin gelişmesinde, ihracatın artmasında, doğal kaynakların değerlendirilmesinde, işsizliğin azalmasında stratejik önem taşımaktadır. Ekonomik gelişmenin anahtarı olan girişimcilik, “düşünsel emeğin değere dönüşmesi” olarak tanımlanmaktadır (Segal vd., 2005; Solunoğlu ve Nalçacı İkiz, 2017). Schumpeter (1934)’e göre girişimcilik, yeni unsurları birleştirerek farklı şeyler yaratmaktır. Bu bağlamda girişimcilik, yeni pazarlar bulmayı, yeni üretim yöntemleri geliştirmeyi, yeni hammadde kaynakları bulmayı ve endüstride yeni organizasyonlar oluşturmayı kapsamaktadır (Aksoy ve Yalçınsoy, 2017). Girişimci ise toplumun ihtiyaçlarının neler olduğunu belirleyen, bu doğrultuda üretim kaynaklarını bir araya getiren, sermayesini yatırıma dönüştüren ve yenilik peşinde koşan kişidir. Bir kişinin başarılı bir girişimci olabilmesi için öncelikle girişimcilik eğiliminin olması gerekmektedir.

Girişimcilik eğilimi, kişinin girişimci olma konusunda istekli olmasını ve girişimcilik yapabilme yeteneğini ifade etmektedir (Seçgin, 2020). Girişimcilik eğilimi, bireyin kendi işini kurmaya yönelik harekete geçmesini, kendine güvenmesini, başkalarıyla kolaylıkla iletişim kurmasını sağlayan sosyal zekayı içermektedir. Girişken ve özgüvenli gençlerin girişimciliğe ilişkin daha yüksek düzeyde istekli ve hırslı oldukları, karşılarına çıkan fırsatları değerlendirmede daha yetenekli oldukları bilinmektedir. Birey risk alma, belirsizliğe tolerans, bağımsızlık ihtiyacı, öz yönetim, yenilikçilik ve yaratıcılık gibi girişimciliğin gerektirdiği özelliklere ne kadar sahip ise girişimcilik eğilimi o kadar yüksektir (Öztürk


FULL TEXTS BOOK

ve Serinkan, 2023). Girişimcilik eğilimi gelecekte kendi işini ya da yeni bir iş kurmayı düşünen bireyler açısından güçlü bir göstergedir (Obshonka vd., 2010; Eroğlu ve Avcı, 2021). Girişimcilerin başarılarının temelinde iletişim becerilerinin ve insan ilişkilerinin güçlü olması vardır. Bu da sosyal zeka gerektirir. Sosyal zekası yüksek bireylerin girişimcilik eğilimlerinin yüksek olabileceği düşüncesinden hareketle üniversite öğrencileri özelinde bu araştırma yapılmıştır.

YÖNTEM

Araştırmanın Amacı, Modeli ve Hipotezleri

Bu çalışmanın amacı, lisans ve ön lisans öğrencilerinin sosyal zeka düzeyleri ile girişimcilik eğilimleri arasındaki ilişkiyi incelemektir. Ayrıca öğrencilerin demografik özelliklerine göre sosyal zeka düzeylerinin ve girişimcilik eğilimlerinin farklılık gösterip göstermediğini tespit etmektir. Araştırmanın amacı ve modeli doğrultusunda aşağıdaki hipotezler oluşturulmuştur.



Şekil 1. Araştırma Modeli

H1: Sosyal zeka ile girişimcilik eğilimi arasında anlamlı bir ilişki vardır.

H2: Sosyal zeka alt boyutu sosyal bilgi süreci ile girişimcilik eğilimi arasında anlamlı bir ilişki vardır.

H3: Sosyal zeka alt boyutu sosyal beceri ile girişimcilik eğilimi arasında anlamlı bir ilişki vardır.

H4: Sosyal zeka alt boyutu sosyal farkındalık ile girişimcilik eğilimi arasında anlamlı bir ilişki vardır.

H5: Sosyal zeka ve alt boyutları cinsiyete göre farklılık göstermektedir.

H6: Sosyal zeka ve alt boyutları sınıf düzeyine göre farklılık göstermektedir.

H7: Sosyal zeka ve alt boyutları eğitim düzeyine göre farklılık göstermektedir.

H8: Girişimcilik eğilimi cinsiyete göre farklılık göstermektedir.

H9: Girişimcilik eğilimi sınıf düzeyine göre farklılık göstermektedir.

H10: Girişimcilik eğilimi eğitim düzeyine göre farklılık göstermektedir.

Araştırmanın Evreni ve Örneklemi

Araştırmanın evreni, Manisa Celal Bayar Üniversitesi'ne bağlı Salihli İktisadi ve İdari Bilimler Fakültesi ve Salihli Meslek Yüksekokulu'nda öğrenim gören öğrencilerden oluşmaktadır. Araştırmanın örneklemi ise anketleri cevaplayan 224 öğrenciden oluşmaktadır. Örneklem kolayda örnekleme yöntemiyle belirlenmiştir.

Veri Toplama Araçları

Araştırma verilerini toplamak için Kişisel Bilgi Formu, Tromso Sosyal Zeka Ölçeği ve Girişimcilik Eğilimi Ölçeği kullanılmıştır. Kişisel Bilgi Formunda katılımcıların cinsiyet, sınıf düzeyi ve eğitim düzeylerini belirlemeye yönelik sorular yer almaktadır. Araştırmada kullanılan "Tromso Sosyal Zeka Ölçeği", Silvera ve arkadaşları (2001) tarafından geliştirilmiş olup, Doğan ve Çetin (2009) tarafından Türkçe'ye uyarlanmıştır. Ölçek 21 madde ve üç alt boyuttan (sosyal bilgi süreci, sosyal beceri, sosyal

FULL TEXTS BOOK

farkındalık) oluşmaktadır. Ölçeğin güvenilirliği .80 olarak bulunmuştur. Ölçek 5'li Likert tipindedir. Ölçek maddeleri 1=Hiç Katılmıyorum ve 5=Tamamıyla Katılıyorum aralığında değerlendirilmektedir. Araştırmada kullanılan "Girişimcilik Eğilimi Ölçeği" Girginer ve Uçgun (2004) tarafından geliştirilmiştir. 5'li Likert tipindeki ölçek 14 madde içermektedir. Ölçeğin güvenilirliği .74 olarak bulunmuştur. Ölçek maddeleri 1=Hiç Katılmıyorum ve 5=Tamamıyla Katılıyorum aralığında değerlendirilmektedir.

Verilerin Analizi

Araştırma kapsamında elde edilen veriler SPSS 22 paket programında analiz edilmiştir. Verilerin analizinde frekans ve yüzde hesaplaması, tanımlayışı istatistik, Pearson Korelasyon analizi, t-testi ve ANOVA analizi uygulanmıştır.

BULGULAR

Araştırmanın bu bölümünde, araştırmaya katılan üniversite öğrencilerinin demografik özelliklerinin dağılımına, sosyal zeka ölçeği ve girişimcilik eğilimi ölçeğine ilişkin tanımlayıcı istatistiklere ve sosyal zeka ile girişimcilik eğilimi arasındaki korelasyon analizi sonuçlarına yer verilmiştir.

Tablo 1. Cinsiyete Göre Dağılım

Değişken	**Grup**	**N**	**%**
Cinsiyet	Kız	118	52,7
	Erkek	106	47,3
	Toplam	224	100,0

Tablo 1'de görüldüğü gibi, araştırmaya katılan öğrencilerin %52,7'si kız, %47,3'ü ise erkektir.

Tablo 2. Sınıf Düzeyine Göre Dağılım

Değişken	**Grup**	**N**	**%**
Sınıf Düzeyi	1.Sınıf	82	36,6
	2.Sınıf	104	46,4
	3.Sınıf	32	14,3
	4.Sınıf	6	2,7
	Toplam	224	100,0

Tablo 2'de görüldüğü gibi, araştırmaya katılan öğrencilerin %36,6'sı 1. Sınıf, %46,4'ü 2. Sınıf, %14,3'ü 3. Sınıf, %2,7'si ise 4. Sınıf öğrencisidir.

Tablo 3. Eğitim Düzeyine Göre Dağılım

Değişken	**Grup**	**N**	**%**
Eğitim Düzeyi	Lisans	109	48,7
	Ön Lisans	115	51,3
	Toplam	224	100,0

Tablo 3'de görüldüğü gibi, araştırmaya katılan öğrencilerin %48,7'si lisans, %51,3'ü ise ön lisans eğitimi almaktadır.

FULL TEXTS BOOK

Tablo 4. Sosyal Zeka Ölçeği ve Alt Boyutlarına Ait Tanımlayıcı İstatistikler

Değişkenler	**N**	**Ort.**	**Ss**	**Min.**	**Max.**
Sosyal Zeka Ölçeği	224	3,68	,541	2.16	5
Sosyal Bilgi Süreci	224	3,76	,648	1,17	5
Sosyal Beceri	224	3,79	,736	2,00	5
Sosyal Farkındalık	224	3,55	,743	1,57	5

Tablo 4'de görüldüğü gibi, araştırmaya katılan öğrencilerin genel sosyal zeka düzeyleri 3,68±,541'dir. Sosyal zeka alt boyutları kapsamında bakıldığında; sosyal bilgi süreci düzeyleri 3,76±,648, sosyal beceri düzeyleri 3,79±,736 ve sosyal farkındalık düzeyleri ise 3,55±,743 olarak hesaplanmıştır. Bu sonuçlara göre öğrencilerin ortalamanın üzerinde bir sosyal zekaya sahip oldukları söylenebilir.

Tablo 5. Girişimcilik Eğilimi Ölçeğine Ait Tanımlayıcı İstatistikler

Değişkenler	**N**	**Ort.**	**Ss**	**Min.**	**Max.**
Girişimcilik Eğilimi	224	3,62	,615	1	5

Tablo 5'te görüldüğü gibi, araştırmaya katılan öğrencilerin girişimcilik eğilimleri ortalama puanı 3,62±,615'dir. Buna göre öğrencilerin girişimcilik eğilimlerinin yüksek düzeyde olduğu söylenebilir.

Tablo 6. Sosyal Zeka Ölçeği ve Alt Boyutları ile Girişimcilik Ölçeği Arasındaki Korelasyon Analizi Sonuçları

Değişkenler		Sosyal Bilgi Süreci	Sosyal Beceri	Sosyal Farkındalık	Sosyal Zeka Ölçeği
Girişimcilik Eğilimi	r	,417**	,362**	,265**	,339**
	p	,000	,000	,000	,000
	N	224	224	224	224

*p<0.005; **p<0.001

Tablo 6'da görüldüğü gibi, sosyal zeka ile girişimcilik eğilimi (r=,339; p<0.001) arasında pozitif yönlü, orta düzeyde ve anlamlı bir ilişki bulunmuştur. Sosyal bilgi süreci alt boyutu ile girişimcilik eğilimi (r=,417; p<0.001) arasında pozitif yönlü, orta düzeyde ve anlamlı bir ilişki; sosyal beceri alt boyutu ile girişimcilik eğilimi (r=,362; p<0.001) arasında pozitif yönlü, orta düzeyde ve anlamlı bir ilişki; sosyal farkındalık alt boyutu ile girişimcilik eğilimi (r=,265; p<0.001) arasında pozitif yönlü, düşük düzeyde ve anlamlı bir ilişki olduğu tespit edilmiştir. Bu sonuca göre H1, H2, H3, H4 desteklenmiştir.

Tablo 7. Cinsiyet Değişkeni İle Sosyal Zeka ve Alt Boyutlarının Karşılaştırılması

Değişken	**Cinsiyet**	**N**	**Ort.**	**Ss**	**t**	**p**
Sosyal Zeka Ölçeği	Kız	118	3,67	,544	-,067	,730
	Erkek	106	3,68	,541		
Sosyal Bilgi Süreci	Kız	118	3,78	,657	,335	,738

FULL TEXTS BOOK

	Erkek	106	3,75	,642		
Sosyal Beceri	Kız	118	3,80	,722	-,001	,840
	Erkek	106	3,79	,755		
Sosyal Farkındalık	Kız	118	3,52	,832	-,612	,013
	Erkek	106	3,58	,631		

Tablo 7'de görüldüğü gibi, cinsiyet değişkeni ile sosyal zekanın sadece sosyal farkındalık alt boyutunda anlamlı farklılık (p=,013<0,05) olduğu tespit edilmiştir. Farklılığın nedenini ise erkek öğrencilerin sosyal farkındalık ortalama puanlarının kız öğrencilerin ortalama puanından daha yüksek olmasından kaynaklanmaktadır. Bu sonuca göre H5 kısmen desteklenmiştir.

Tablo 8. Sınıf Düzeyi Değişkeni İle Sosyal Zeka ve Alt Boyutlarının Karşılaştırılması

Boyutlar	**Sınıf düzeyi**	**N**	**Ort.**	**Ss**	**F**	**p**
Sosyal Zeka Ölçeği	1. Sınıf	41	3,66	,40713	,725	,538
	2. Sınıf	127	3,71	,607		
	3. Sınıf	50	3,58	,462		
	4. Sınıf	6	3,73	,499		
Sosyal Bilgi Süreci	1. Sınıf	41	3,76	,412	,992	,397
	2. Sınıf	127	3,77	,725		
	3. Sınıf	50	3,68	,590		
	4. Sınıf	6	4,14	,700		
Sosyal Beceri	1. Sınıf	41	3,83	,600	,375	,771
	2. Sınıf	127	3,82	,812		
	3. Sınıf	50	3,70	,650		
	4. Sınıf	6	3,72	,646		
Sosyal Farkındalık	1. Sınıf	41	3,41	,669	2,950	,034
	2. Sınıf	127	3,68	,777		
	3. Sınıf	50	3,38	,686		
	4. Sınıf	6	3,28	,478		

Tablo 8'de görüldüğü gibi, sınıf düzeyi değişkeni ile sosyal zekanın sadece sosyal farkındalık alt boyutunda anlamlı farklılık (p=,034<0,05) olduğu tespit edilmiştir. Farklılığın nedenini belirlemek için yapılan Ducan testi sonucunda 2. Sınıf öğrencilerinin 1., 3. ve 4. Sınıf öğrencilerinden anlamlı olarak farklılaştığı tespit edilmiştir. Bu sonuca göre H6 kısmen desteklenmiştir.

Tablo 9. Akademik Birim Değişkeni İle Sosyal Zeka ve Alt Boyutlarının Karşılaştırılması

FULL TEXTS BOOK

Değişken	Eğitim Düzeyi	N	Ort.	Ss	t	p
Sosyal Zeka Ölçeği	Lisans	109	3,66	,499	-,311	,132
	Ön Lisans	115	3,69	,580		
Sosyal Bilgi Süreci	Lisans	109	3,74	,573	-,386	,055
	Ön Lisans	115	3,78	,714		
Sosyal Beceri	Lisans	109	3,78	,726	-,191	,400
	Ön Lisans	115	3,80	,749		
Sosyal Farkındalık	Lisans	109	3,48	,695	-1,308	,181
	Ön Lisans	115	3,61	,783		

Tablo 9'da görüldüğü gibi, eğitim düzeyi değişkeni ile sosyal zeka ve alt boyutları arasında anlamlı bir farklılık bulunmamaktadır (P>0,05). Bu sonuca göre H7 reddedilmiştir.

Tablo 10. Cinsiyet Değişkeni İle Girişimcilik Eğiliminin Karşılaştırılması

Değişken	Cinsiyet	N	Ort.	Ss	t	p
Girişimcilik Eğilimi Ölçeği	Kız	118	3,57	,581	-1,123	,263
	Erkek	106	3,66	,648		

Tablo 10'da görüldüğü gibi, cinsiyet değişkeni ile girişimcilik eğilimi arasında anlamlı bir farklılık bulunmamaktadır (P>0,05). Bu sonuca göre H8 reddedilmiştir.

Tablo 11. Sınıf Düzeyi Değişkeni İle Girişimcilik Eğiliminin Karşılaştırılması

Değişken	Sınıf Düzeyi	N	Ort.	Ss	t	p
Girişimcilik Eğilimi Ölçeği	1. Sınıf	41	3,51	,818	2,151	,095
	2. Sınıf	127	3,70	,593		
	3. Sınıf	50	3,47	,438		
	4. Sınıf	6	3,63	,505		

Tablo 11'de görüldüğü gibi, sınıf düzeyi değişkeni ile girişimcilik eğilimi arasında anlamlı bir farklılık bulunmamaktadır (P>0,05). Bu sonuca göre H9 reddedilmiştir.

Tablo 12. Akademik Birim Değişkeni İle Girişimcilik Eğiliminin Karşılaştırılması

Değişken	Akademik Birim	N	Ort.	Ss	t	p
Girişimcilik Eğilimi Ölçeği	Salihli İİBF	109	3,66	,632	-1,171	,243
	Salihli MYO	115	3,57	,595		

Tablo 12'de görüldüğü gibi, akademik birim değişkeni ile girişimcilik eğilimi arasında anlamlı bir farklılık bulunmamaktadır (P>0,05). Bu sonuca göre H10 desteklenmiştir.

FULL TEXTS BOOK

SONUÇ VE ÖNERİLER

Lisans ve ön lisans öğrencilerinin algılarına göre sosyal zeka ve girişimcilik eğilimi arasındaki ilişkiyi ve demografik özelliklerine (cinsiyet, sınıf düzeyi, akademik birim) göre sosyal zeka düzeylerinde ve girişimcilik eğilimlerinde anlamlı farklılık gösterip göstermediğini incelemeyi amaçlayan bu araştırmada elde edilen bulgular ve yorumlar aşağıda özetlenmiştir.

Araştırmaya katılan öğrencilerin %52,7'si kız, , %46,4'ü 2. sınıf, %51,3'ü ise ön lisans öğrencisidir. Sosyal zeka ölçeğine ait tanımlayıcı istatistiklere bakıldığında, sosyal zeka düzeylerinin yüksek düzeyde ve sosyal zeka alt boyutlarından sosyal beceri en yüksek ortalama puana sahipken, sosyal farkındalık en düşük ortalama puana sahip olduğu görülmektedir.

Korelasyon analizi sonucuna göre; sosyal zeka ve alt boyutları ile girişimcilik eğilimi arasında pozitif yönlü, orta düzeyde ve anlamlı bir ilişki bulunmuştur. Buna göre bireylerin sosyal zeka düzeyleri arttıkça girişimcilik eğilimleri de artmaktadır. Buna göre H1, H2, H3, H4 kabul edilebilir. Literatürde bu sonucu destekleyen çalışmalar bulunmaktadır (Gürpınar vd., 2020; Parlak, 2013; Uzun vd., 2017; Yılmazçetin, 2021; Zehir ve Bektaş, 2022).

Demografik özellikler ve sosyal zeka ve alt boyutları ile girişimcilik eğilimi arasında anlamlı farklılık olup olmadığına ilişkin yapılan analiz sonuçlarına göre; sosyal zeka ve alt boyutlarından sosyal farkındalık düzeylerinde cinsiyete göre fark erkek öğrencilerin lehine anlamlı sonuç bulunmuştur. Buna göre H5 kısmen kabul edilebilir. Literatürde bu sonucu destekleyen çalışmalara rastlanmıştır. Yılmazçetin (2021) tarafından yapılan araştırma sonucuna göre sosyal zekanın sadece sosyal farkındalık alt boyutunda erkek öğrenciler lehine anlamlı farklılık bulunmuştur. Ülker (2016) tarafından yapılan araştırma sonucunda meslek yüksekokulu öğrencilerinin sosyal zeka ve alt boyutlarından sadece sosyal farkındalık düzeylerinde cinsiyete göre anlamlı farklılık tespit edilmiştir. Ancak farklılık kız öğrencilerin lehine çıkmıştır. Doğan ve Çetin (2008) tarafından yapılan çalışmada ise cinsiyet değişkenine göre öğrencilerin sosyal zeka düzeylerinde erkek öğrenciler lehine anlamlı fark bulunmuştur. Aktürk (2023)'ün çalışmasında da benzer sonuç çıkmıştır.

Sosyal zeka ve alt boyutlarından sosyal farkındalık düzeylerinde sınıf düzeyine göre fark erkek 2. sınıfta öğrenim gören öğrencilerin lehine anlamlı sonuç bulunmuştur. Diğer sınıfların kendi aralarında anlamlı farklılık olmadığı dikkate alındığında 2. sınıf öğrencilerinin diğerlerine göre daha yüksek bir sosyal farkındalığa sahip olduğu söylenebilir. Buna göre H6 kısmen kabul edilebilir, Literatürde bu sonucu destekleyen çalışmalar bulunmaktadır (Aktürk, 2023). Literatürde farklı sonuçların elde edildiği çalışmalar mevcuttur. Ülker (2016) tarafından yapılan araştırma sonucunda sınıf düzeyi değişkenine göre sosyal zeka ve alt boyutlarında anlamlı farklılık bulunmamıştır. Sosyal zeka ve alt boyutlarından sosyal farkındalık düzeylerinde eğitim düzeyine göre anlamlı bir farklılık tespit edilmemiştir.

Lisans ve ön lisans öğrencilerinin sosyal zeka düzeylerinde ve alt boyutları ile cinsiyet, sınıf düzeyi ve eğitim düzeyine göre anlamlı bir farklılık bulunmamıştır. Literatürde bu sonucu destekleyen çalışmalar bulunmaktadır. Şahin (2023) tarafından üniversite öğrencilerine yönelik yapılan araştırma sonucunda öğrencilerin girişimcilik eğilimlerinde cinsiyete göre anlamlı farklılık bulunamamıştır. Benzer şekilde Öztürk ve Serinkan (2023) tarafından yapılan çalışmada cinsiyet ile girişimcilik eğilimi arasında anlamlı farklılık bulunmamıştır. Gökçe Günar (2023) tarafından yapılan araştırmada öğrencilerin cinsiyetine ve eğitim düzeyine göre anlamlı farklılık bulunamazken, sınıf düzeylerine göre anlamlı farklılık bulunmuştur. Farklı sonuçların ulaşıldığı çalışmalara da rastlanmıştır. Arıcan Kaygusuz (2022) tarafından üniversite öğrencileri üzerine yapılan araştırmada girişimcilik eğiliminin cinsiyete göre farklılaştığı tespit edilmiştir. Ayrıca öğrencilerin eğitim düzeyine göre girişimcilik eğilimlerinde ön lisans öğrencileri lehine anlamlı farklılık bulunmuştur. Wang ve Wong (2004) tarafından yapılan araştırmada üniversite öğrencilerinin girişimcilik eğilimlerinin cinsiyet ve eğitim düzeyi değişkenlerine göre anlamlı farklılık olduğu ortaya konulmuştur. Ekici ve Turan (2017)'ın yaptıkları çalışmada

FULL TEXTS BOOK

üniversite öğrencilerinin cinsiyetine göre girişimcilik eğilimlerinde anlamlı farklılık olduğu tespit edilmiştir. Benzer şekilde Küçük (2014)'ün çalışmasında cinsiyete göre girişimcilik eğiliminde erkek öğrenciler lehine anlamlı farklılık olduğu tespit edilmiştir.

Sonuç olarak; Sosyal zeka ve girişimcilik eğilimi yüksek olan öğrenciler girişimcilik üzerine proje çalışmalarına yönlendirilebilir. Sosyal zeka ve girişimcilik eğilimi yüksek olan öğrencilere kendi işini kurmaya teşvik edilebilir. Bu araştırma sadece bir üniversitenin iki akademik biriminde sınırlı sayıda öğrenci ile yürütüldüğünden sonuçlar genellenemez. Bunun için daha çok sayıda üniversite ve öğrenci örnekleminde çalışmalar yapılabilir. Literatürde sosyal zeka ve girişimcilik eğilimi arasındaki ilişkiyi inceleyen sınırlı sayıda çalışma olduğu görülmüştür. Bu konuda daha fazla çalışmalar yapılarak sonuçlar genellenebilir. Üniversite öğrencilerine sosyal zeka ve girişimcilik eğilimi konusunda seminer eğitimleri verilerek farkındalıkları artırılabilir.

KAYNAKÇA

Aksoy, C., ve Yalçınsoy, A. (2017). Üniversite öğrencilerinin girişimcilik eğilimleri üzerine bir araştırma. *İşletme Araştırmaları Dergisi*, 9(4), 341-359.

Aktürk, M. B. (2023). *Spor lisesi ve fen lisesi öğrencilerinin beden eğitimi ve spor dersine yönelik tutumları ile sosyal zeka düzeylerinin incelenmesi: Elazığ ili örneği*. Yüksek Lisans Tezi, Munzur Üniversitesi Lisansüstü Eğitim Enstitüsü, Tunceli.

Aliyev, J. (2022). *Çalışanların sosyal zeka seviyelerinin müşteri şikayetlerine yaklaşımları üzerindeki etkisi: bodrum örneği*. Yüksek Lisans Tezi, Muğla Sıtkı Koçman Üniversitesi Sosyal Bilimler Enstitüsü, Muğla.

Arıcan Kaygusuz, N. (2022). *Girişimcilik eğitimi ile girişimcilik eğilimi arasındaki ilişkide ailesel deneyimlerin aracılık rolü: Giresun Üniversitesi öğrencileri ile Giresun'daki lise öğrencileri üzerinde gerçekleştirilen bir alan araştırmasına dayalı model önerisi*. Doktora Tezi, Giresun Üniversitesi Sosyal Bilimler Enstitüsü, Giresun.

Cebeci, C., ve Kamaşak, R. (2021). Sosyal zekâ ve içsel motivasyonun ar-ge performansına etkisi: Ar-ge merkezleri çalışanları üzerine bir araştırma. *İşletme Araştırmaları Dergisi*, 13(1), 47-66.

Çakmak Yıldızhan, Y., ve Çağlayan, G. N. (2019). Öğrencilerin beden eğitimi dersine yönelik tutumları ile sosyal zekâ düzeyleri arasındaki ilişkinin incelenmesi. *Beden Eğitimi ve Spor Bilimleri Dergisi*, 24(4), 227-240.

Demirbanka, F. ve Taşlıyan, M. (2022). Sosyal zekanın iş becerikliliği etkisi üzerine bir alan araştırması. *Journal of Pure Social Sciences*, 3(4), 27-40.

Doğan, T., ve Çetin, B. (2008). Üniversite öğrencilerinin sosyal zeka düzeylerinin depresyon ve bazı değişkenlerle ilişkisinin incelenmesi. *Uluslararası İnsan Bilimleri Dergisi*, 5(2), 1-19.

Ekici, E., ve Turan, M. (2017). Üniversite öğrencilerinin girişimcilik eğilimi: Planlanmış davranışlar teorisi ve girişimcilik eğitiminin rolü. *Ç.Ü. Sosyal Bilimler Enstitüsü Dergisi*, 26(1), 201-215.

Erdemir, N., ve Kutlu, M. (2018). Üniversite öğrencilerinin internet bağımlılığı ile sosyal zeka düzeylerinin çeşitli değişkenler açısından incelenmesi. *Uluslararası Sosyal Araştırmalar Dergisi*, 11(57), 401-409.

Eroğlu, Ş.G., ve Avcı, U. (2021). Üniversite öğrencilerinin girişimcilik eğilimleri üzerine boylamsal bir araştırma. *İşletme Araştırmaları Dergisi*, 13 (4), 3607-3621.

Genc, V., Genc, S. G. (2018). Can hotel managers with social intelligence affect the emotions of employees?, *Cogent Business & Management*, 5(1), https://doi.org/10.1080/23311975.2018.1432157

FULL TEXTS BOOK

Girginer, N., ve Uçgun, N. (2004). İşletmecilik eğitimi alan lisans öğrencilerinin girişimciliğe bakış açıları: Eskişehir Osmangazi Üniversitesi İ.İ.B.F işletme bölümü öğrencilerine yönelik bir uygulama. III. Ulusal Bilgi, Ekonomi ve Yönetim Kongresi bildiri kitabı içinde, 25-26 Kasım, Eskişehir.

Gökçe Günar, B. (2023). *Üniversite öğrencilerinde girişimcilik eğiliminin öz-yeterlik ve kişilik özellikleri ile ilişkisinin incelenmesi*. Yüksek Lisans Tezi, Çanakkale Onsekiz Mart Üniversitesi Lisansüstü Eğitim Enstitüsü, Çanakkale.

Gürpınar, K., Yüncü, V., ve Yılan, Y. (2020). Sosyal zeka ve girişimci kişilik özellikleri arasındaki ilişkinin sınıflandırılması. *Girişimcilik ve İnovasyon Yönetimi Dergisi*, 9(2), 128-148.

Kahya, V., ve Ceylan, E. (2017). Sosyal zekânın iletişim becerileri üzerine etkisi: üniversite öğrencileri üzerine bir araştırma. Proceedings Book of 2nd International Scientific Researches Congress on Humanities and Social Sciences (IBAD-2017), April 20-23, 2017, Istanbul.

Obschonka, M., Silbereisen, R. K., ve Schmitt-Rodermund, E. (2010). Entrepreneurial intention as developmental outcome. *Journal of Vocational Behavior*, 77(1), 63-72.

Öztürk, R., ve Serinkan, C. (2023). Üniversite öğrencilerinin girişimcilik eğilimlerinin belirlenmesi: Pamukkale Üniversitesi örneği. *Yeni Fikir Dergisi*, 15 (30), 01-09.

Parlak, K. (2013). Girişimci kişilik özelliklerine sahip olmak ile kendini ve başkalarını tanıma (içsel ve sosyal) zekaları arasındaki ilişkinin incelenmesi. *İstanbul Sabahattin Zaim Üniversitesi Sosyal Bilimler Dergisi*, 3, 55-64.

Segal, G., Borgia, D., ve Schoenfeld, J. (2005). The motivation to become an entrepreneur. *International journal of Entrepreneurial Behavior ve Research*, 11(1), 42-57.

Seçgin, Y. (2020), Girişimcilik eğitiminin girişimcilik eğilimi üzerine etkisi: Üniversite öğrencileri üzerine bir araştırma. *BMIJ*, 8(1), 803-827.

Silvera, D., H., Martinussen, M., ve Dahl, T. I. (2001), The tromso social intelligence scale, a self-report measure of social intelligence. *Scandinavian Journal of Psychology*, 42, 313-31.

Solunoğlu, A., ve Nalçacı İkiz, A. (2017). Üniversite öğrencilerinin girişimcilik eğilimleri. *Journal of Recreation and Tourism Research*, 4(2), 44-57.

Şahin, P. G. (2023). *Üniversite öğrencilerinde psikolojik dayanıklılık ve benlik saygısı ile girişimcilik eğilimi arasındaki ilişkinin değerlendirilmesi*. Yüksek Lisans Tezi, İstanbul Nişantaşı Üniversitesi Lisansüstü Eğitim Enstitüsü, İstanbul.

Uzun, H., Buran, A.Ç., ve Beydilli, E.T. (2017). Sosyal zekanın girişimcilik niyeti üzerindeki etkisi: meslek yüksekokulu öğrencileri üzerinde bir araştırma. *Siyaset, Ekonomi Ve Yönetim Araştırmaları Dergisi*, 5(5), 1-11.

Ülker, Ç. (2016). *Meslek yüksekokulu öğrencilerinin sosyal zeka ve iletişim becerilerinin farklı değişkenler açısından incelenmesi*. Nişantaşı Üniversitesi Sosyal Bilimler Enstitüsü, Yüksek Lisans Tezi, İstanbul.

Vural, B., (2004). Öğrenci Merkezli Eğitim ve Çoklu Zeka (1. Baskı). İstanbul: Hayat Yayınları.

Wang, C.K., ve Wong, P. (2004). Entrepreneurial Interest of University Students in Singapore, *Elsevier, Technovation*, (24), 163-172.

Yeşil, S., ve Mavi, Y. (2024). Sosyal zekâ: Geçmişten bugüne kavramsal bir inceleme. *Kahramanmaraş Sütçü İmam Üniversitesi Sosyal Bilimler Dergisi,* 21(1), 358–372.

FULL TEXTS BOOK

Yılmazçetin, Ş. (2021). *Duygusal zekâ, sosyal zekâ ve kültürel zekâ ile girişimcilik eğilimi etkileşimi: Üniversite öğrencileri üzerine bir araştırma*. Doktora Tezi, İstanbul Üniversitesi Sosyal Bilimler Enstitüsü, İstanbul.

Zehir, C., ve Bektaş, G. (2022). Duygusal zekâ ve girişimcilik niyeti arasında öz yeterliliğin aracılık ilişkisi. *Yıldız Sosyal Bilimler Enstitüsü Dergisi*, 6(1), 11-21.

FULL TEXTS BOOK

LİTYUM-İYON PİL TERMAL YÖNETİMİ İÇİN HAVA SOĞUTMA KONFİGÜRASYONLARININ KARŞILAŞTIRILMASI

COMPARISON OF AIR COOLING CONFIGURATIONS FOR LITHIUM-ION BATTERY THERMAL MANAGEMENT

Dr. Öğr. Üyesi Erdi TOSUN

Çukurova Üniversitesi, Mühendislik Fakültesi,

Otomotiv Mühendisliği Bölümü, Balcalı, Adana.

Çağrı ADIKTI

Çukurova Üniversitesi, Mühendislik Fakültesi,

Otomotiv Mühendisliği Bölümü, Balcalı, Adana.

ÖZET

Lityum-iyon pillerin (LiP'ler) sıcaklığı, performanslarını, güvenliklerini ve kullanım ömürlerini doğrudan etkilediğinden dikkatli bir şekilde kontrol edilmelidir. Hava soğutma, basitliği, düşük maliyeti ve daha az bakım gereksinimi nedeniyle LiP'ler için yaygın olarak kullanılan bir termal yönetim yöntemidir. Bu çalışmada, batarya modülü dikdörtgen bir kabin içerisine yerleştirilmiş ve iki farklı hava soğutma senaryosu uygulanarak 2C deşarj koşulunda modülün termal performansı değerlendirilmiştir. Modül toplam 8 adet ASPİLSAN INR18650A28 hücreden oluşmakta olup 4 seri ve 2 paralel konfigürasyona (4S2P) sahiptir. İlk senaryoda (SF) modül, bir fan kullanılarak modülün yan tarafından hava üflenerek soğutulurken, ikinci senaryoda (TF), hava modülün üstünden üflenmiştir. Modül içindeki hücreler arasında en yüksek sıcaklık (T_{max}), SF durumunda 37,88 °C, TF durumunda ise 32,97 °C olarak gözlemlenmiştir. Ayrıca, modül içindeki maksimum sıcaklık farkı (ΔT_{max}) SF'de 9,08 °C ve TF'de 4,38 °C'dir. Her iki senaryoda da T_{max} güvenli aralıkta (20-45 °C) kalmıştır. Ancak, TF konfigürasyonu, SF konfigürasyonuna kıyasla daha düşük bir T_{max} ile sonuçlanmıştır. ΔT_{max} açısından bakıldığında SF, genel kabul görmüş 5 °C sınırını aşmış olup, yetersiz sıcaklık dağılımına işaret ederken, TF ise güvenilir eşik içerisinde kalmıştır. Genel olarak, TF konfigürasyonu SF'den daha iyi termal performans göstermiştir.

Anahtar kelimeler: Hava soğutma, lityum-iyon batarya, termal yönetim.

ABSTRACT

The temperature of lithium-ion batteries (LiBs) must be carefully controlled, as it directly affects their performance, safety, and lifespan. Air cooling is a commonly used thermal management method for LiBs due to its simplicity, low cost, and less maintenance requirements. In this study, the battery module was placed inside a rectangular cabin where two different air cooling scenarios were implemented to evaluate thermal performance of module under 2C discharge condition. The module consists of total 8 ASPİLSAN INR18650A28 cells and has a 4-series and 2-parallel configuration (4S2P). In the first scenario (SF), the module was cooled by blowing air from the side of module using a fan, while in the second (TF), air was blown from the top of the module. The maximum temperature (T_{max}) observed among cells within the module was 37.88 °C in the SF case and 32.97 °C in the TF case. Meanwhile, the maximum temperature difference (ΔT_{max}) within the module was 9.08 °C in the SF and 4.38 °C in the TF. In both scenarios, T_{max} remained within the safe range (20-45 °C). However, TF configuration resulted in a lower T_{max} compared to the SF. Regarding the ΔT_{max}, the SF exceeded the commonly

FULL TEXTS BOOK

accepted limit of 5 °C, indicating insufficient temperature distribution, whereas the TF remained within the reliable threshold. Overall, the TF configuration demonstrated better thermal performance than the SF setup.

Keywords: Air cooling, lithium-ion battery, thermal management.

INTRODUCTION

Temperature plays a critical role in the performance, safety, and lifespan of lithium-ion batteries (LiBs), making effective thermal management essential for their reliable operation. In general, battery thermal management systems (BTMSs) are classified into two main categories: passive systems, which rely on material properties such as phase change materials (PCMs) or high-conductivity structures to control heat, and active systems, which utilize external energy sources to circulate air, liquid, or refrigerant for temperature regulation. Among various thermal management approaches, air cooling is widely adopted due to its structural simplicity, low cost, ease of maintenance, and sufficient performance under moderate operating conditions (Rallabandi and Selvaraj, 2024; Zhu, Liao, Kang, Sun, and Zhao, 2024). Moreover, key parameters such as airflow rate, inlet air temperature, flow direction, and channel geometry can be adjusted to enhance heat dissipation and maintain cell temperatures within the optimal range. Numerous studies in the literature have investigated air-cooled battery systems, focusing on the optimization of airflow configurations and geometric modifications to improve cooling efficiency and temperature uniformity (Sun and Dixon, 2014; Chen, Song, Wei, and Wang, 2019; Wang, Li, Li, Huang, and Qi, 2021; Alzwayi and Paul, 2025; Wankhede, Pingale, and Kale, 2025).

The primary objectives of BTMS are to maintain the maximum temperature (T_{max}) of individual cells within the safe operating range of approximately 20 °C to 45 °C, and to ensure that the temperature difference (ΔT) between cells does not exceed 5 °C, thereby minimizing thermal gradients that can lead to capacity imbalance, accelerated aging, and safety risks (Hadi, Putra, Trisnadewi, Sukarno, Fathoni, and Bhaskara, 2025; Yang and He, 2025).

In this study, the thermal behavior of a LiB module was experimentally investigated under two different air cooling configurations. The module consisting of 8 ASPİLSAN INR18650A28 cells in a 4-series and 2-parallel configuration (4S2P) layout was placed in a rectangular enclosure. In the first setup (SF), air was blown from the side using a fan, while in the second (TF), air was introduced from the top. The fundamental aim was to evaluate the cooling effectiveness of each method during 2C discharge condition in terms of T_{max} and ΔT.

MATERIALS AND METHODS

The battery module used in this study was composed of ASPİLSAN INR18650A28 cylindrical LiB cells shown in Figure 1. Each cell has a nominal capacity of 2800 mAh (minimum 2700 mAh) and a nominal voltage of 3.65 V. The charge and discharge cut-off voltages are 4.2 V and 2.5 V, respectively. The gravimetric and volumetric energy densities of the cell are 230 Wh/kg and 605 Wh/L.

FULL TEXTS BOOK



Figure 1. 4S2P battery module

The experimental setup (Figure 2) consists of a battery module housed inside an air cabin (dimensions of 150 mm in width, 400 mm in length, and 110 mm in height with 80 mm circular openings) produced by plexiglass where cooling air is supplied via a 24 VDC fan. The fan operates at a maximum speed of 5700 rpm and provides an airflow rate of up to 2.270 m^3/min.

To monitor the thermal behavior of the module, thermocouples (Pico T-type) were attached to the cell surfaces and connected to a temperature datalogger (Pico TC-08 thermocouple datalogger). The datalogger was linked to a computer for real-time data acquisition. A DC power supply (GW Instek PSU 100-15: 100 V, 15 A, 1500 W) was used to charge the module, while a load cell (Heliocentris EL 1500 electronic load: 75 V, 100 A, 1500 W) controlled the discharge process.





Figure 2. Experimental set-up

Figure 3 shows the two airflow configurations used in the experiments. In the TF, the fan is positioned on the top surface of the enclosure which directs air vertically downward onto the battery module. Outlets are located on the side walls to allow the heated air to exit the cabin. In the SF, the fan is mounted on the side wall which provides horizontal airflow across the module. In this case, the exhaust occurs through the opposite side wall and top wall.

FULL TEXTS BOOK




(a) (b)

Figure 3. Test configurations (a) TF (b) SF

RESULTS AND DISCUSSIONS

Figure 4 and 5 present the thermal performance of the battery module under three different cooling conditions as no cooling, SF, and TF. The red lines in the Figures 4 and 5 represent the optimum operating temperature range for LiB cells (20–45 °C) as well as the commonly accepted upper limit for the ΔT_{max} between cells (5 °C).



	C1	C2	C3	C4	C5	C6	C7	C8
SF	28.81	33.35	35.68	37.88	35.93	30.69	30.88	30.88
TF	29.21	28.67	30.04	32.97	31.50	29.20	29.92	29.84
No Cooling	42.37	42.48	41.36	41.55	43.12	41.14	42.26	43.34

Figure 4. T_{max} values of each cells in the module

The T_{max} observed among the cells was 43.34 °C in the no cooling case. This value decreased to 37.88 °C and 32.97 °C with the application of SF and TF cooling, corresponding to reductions of 12.58% and 23.92%, respectively. The average of the maximum temperatures $(T_{max,avg})$ recorded from each of the eight cells also dropped from 42.20 °C (no cooling) to 33.01 °C in SF and 30.17 °C in TF. It showed respective reductions of 21.78% and 28.51% compared to the no cooling case. The temperature reductions observed in both SF and TF configurations are mainly due to improved forced convection. In the SF setup, cooling may be limited due to flow obstruction. It means that cells positioned behind others may receive less airflow due to obstruction and non-uniform flow distribution may occur. The TF configuration provides more uniform and direct airflow over the module which enhances overall heat dissipation.

FULL TEXTS BOOK




Figure 5. ΔT evaluation along 2C discharge rate

Regarding temperature uniformity as shown in Figure 5, ΔT_{max} within the module was 8.30 °C under no cooling and increased to 9.08 °C with the SF which indicates a 9.45% deterioration. In contrast, the TF configuration reduced ΔT to 4.38 °C which corresponds to a 47.22% improvement compared to the no cooling case.The increase in ΔT_{max} compared to no cooling case under the SF configuration may be attributed to non-uniform airflow distribution, where inner or shaded cells receive less effective cooling due to flow obstruction effects. This results in uneven heat dissipation among the cells. On the other hand, the TF configuration provides a more homogeneous airflow across the module. It promotes consistent heat transfer over the cell surfaces. This uniform exposure helps minimize temperature differences between cells and it leads to the significant reduction in ΔT_{max} observed with TF cooling.

CONCLUSIONS

- Two air cooling strategies (SF and TF) were experimentally tested on a 4S2P LiB module under 2C discharge conditions.
- TF cooling reduced the T_{max} to 32.97 °C and improved ΔT_{max} as an indicator of thermal uniformity to 4.38 °C. Besides that, SF resulted in a T_{max} of 37.88 °C and ΔT_{max} of 9.08 °C.
- TF provided more effective and homogeneous airflow, enhancing heat dissipation and minimizing temperature gradients.
- Overall, TF configuration showed superior thermal performance and is more suitable for air-cooled battery systems under high loads.

REFERENCES

Rallabandi, S., & Selvaraj, R.V.I. (2024). Advancements in battery cooling techniques for enhanced performance and safety in electric vehicles: A comprehensive review. Energy Technology, 12, 2301404.

Zhu, X., Liao, X., Kang, S., Sun, L., & Zhao, Y. (2024). Optimization design of lithium battery management system based on Z-F composite air cooling structure. Journal of Energy Storage, 102, 114068.

Sun, H., & Dixon, R. (2014). Development of cooling strategy for an air cooled lithium-ion bat tery pack. Journal of Power Sources, 272, 404-414.

FULL TEXTS BOOK

Chen, K., Song, M., Wei, W., & Wang, S. (2019). Design of the structure of battery pack in parallel air-cooled battery thermal management system for cooling efficiency improvement. International Journal of Heat and Mass Transfer, 132, 309-321.

Wang, N., Li, C., Li, W., Huang, M., & Qi, D. (2021). Effect analysis on performance enhancement of a novel air cooling battery thermal management system with spoilers. Applied Thermal Engineering, 192, 116932.

Alzwayi, A., & Paul, M.C. (2025). Enhancing battery cooling and temperature uniformity with longitudinal airflow and fin configurations. Applied Thermal Engineering, 265, 125625.

Wanghede, S., Pingale, A.D., & Kale, A. (2025). xperimental investigation on thermal management of lithium-ion battery pack for formula student electric vehicle using air-cooling system. Energy Storage and Saving, 4, 38-47.

Hadi, M.W., Putra, N., Trisnadewi, T., Sukarno, R., Fathoni, A.M., & Bhaskara, N.A. (2025). Innovative thermal management system for electric vehicle batteries: Phase change material, heat pipe and heat sink box integration. Heat Transfer Engineering, 46 (6), 503-513.

Yang, X., & He, W. (2025). Simulation study based on experimental flexible composite phase change material for battery thermal management. Heat Transfer Engineering, Early Access, 1-12.

AGOMELATİN VE FİZYOLOJİK SİSTEMLER: TEMEL MEKANİZMALARA BÜTÜNCÜL BİR BAKIŞ

AGOMELATINE AND PHYSIOLOGICAL SYSTEMS: AN INTEGRATIVE APPROACH TO CORE MECHANISMS

Raviye ÖZEN KOCA

Necmettin Erbakan Üniversitesi, Tıp Fakültesi,

Fizyoloji Anabilim Dalı, Konya, Türkiye.

ÖZET

Agomelatin, melatonin reseptörlerine (MT_1 ve MT_2) agonist, serotonin ($5\text{-}HT_2C$) reseptörlerine ise seçici antagonist olarak etki eden atipik bir antidepresandır. Bu çift yönlü mekanizma sayesinde serotonin geri alımını doğrudan etkilemeden sirkadiyen ritmi düzenler ve monoamin iletimini artırır. Bu çalışmada, agomelatinin kontrendikasyon ve yan etkiler dışında kalan fizyolojik mekanizmaları ve sistemik etkileri incelenmiştir. Agomelatin melatonerjik etkisiyle biyolojik ritimleri senkronize ederken, serotonerjik antagonizması yoluyla prefrontal kortekste dopamin ve noradrenalin salımını artırır. Uyku düzenlemesi ve bilişsel fonksiyonlarda iyileşme sağlar. Merkezi sinir sisteminde düzenleyici etkileri bulunmaktadır. Hipokampus ile prefrontal kortekste nöroplastisiteyi destekler. Endokrin sistemde hormonal ritimleri düzene sokar ancak hipotalamo-hipofizer sistemde baskılayıcı etkisi bulunmaz. Kardiyovasküler sistemde agomelatinin kalp hızı ve kan basıncı üzerinde minimal etkileri görülmektedir. Gastrointestinal sistemde melatonerjik ve $5\text{-}HT_3$ antagonist etkilerle motiliteyi düzenleyebilir. Bağışıklık sisteminde IL-6 ve CRP gibi proinflamatuar belirteçleri azaltarak antiinflamatuar etki gösterebilir. Sonuç olarak, agomelatin; kronobiyotik, nörotrofik ve immunomodülatör özellikleriyle çok yönlü fizyolojik etkiler sergileyen özgün bir antidepresandır. Bu etkilerin sistem düzeyinde nasıl şekillendiğini ve klinik karşılıklarını daha iyi anlayabilmek için ayrıntılı deneysel ve klinik çalışmalarla fizyolojik mekanizmaların aydınlatılması gerekmektedir.

Anahtar Kelimeler: Agomelatin, melatonin reseptörleri, serotonin reseptörleri, nöroplastisite, sistemik etkiler.

ABSTRACT

Agomelatine is an atypical antidepressant that acts as a melatonin receptor agonist (MT_1 and MT_2) and a selective antagonist of serotonin $5\text{-}HT_2C$ receptors. This dual mechanism enables agomelatine to regulate circadian rhythms and enhance monoamine transmission without directly inhibiting serotonin reuptake. This study examined the physiological mechanisms and systemic effects of agomelatine, excluding its contraindications and side effects. Agomelatine synchronizes disrupted biological rhythms through melatonergic activity while increasing dopamine and noradrenaline release in the prefrontal cortex via serotonergic antagonism. It promotes sleep regulation and cognitive improvement without stimulating reward pathways. It has regulatory effects in the central nervous system, supporting neuroplasticity in the hippocampus and prefrontal cortex. In the endocrine system, it restores disrupted hormonal rhythms without significantly altering HPA axis activity. In the cardiovascular system, agomelatine shows a minimal effects on heart rate and blood pressure. In the gastrointestinal system, it exhibits melatonergic and $5\text{-}HT_3$ antagonism, potentially regulating motility and visceral sensitivity. It also displays immunomodulatory effects by reducing pro-inflammatory cytokines such as IL-6 and CRP, suggesting anti-inflammatory actions in depression. In conclusion, agomelatine is an antidepressant that

FULL TEXTS BOOK

exerts multifaceted physiological effects through its chronobiotic, neurotrophic, and immunomodulatory properties. Further detailed experimental and clinical studies are needed to comprehensively elucidate its multifaceted physiological mechanisms and their translational significance.

Keywords: Agomelatine, melatonin receptors, serotonin receptors, neuroplasticity, systemic effects.

Introductıon

Agomelatine is an atypical antidepressant that acts as a melatonin receptor agonist and a selective serotonin receptor antagonist. It binds with high affinity to MT_1 and MT_2 melatonin receptors while antagonizing 5-HT_2C (and to a lesser extent 5-HT_2B) serotonin receptors. Through this dual action, agomelatine resynchronizes disrupted circadian rhythms, thereby improving the sleep–wake cycle and enhancing the release of norepinephrine and dopamine in the prefrontal cortex. These effects collectively contribute to its antidepressant and cognitive-enhancing properties (1). This study presents an overview of the pharmacological features of agomelatine and its major effects on the nervous, endocrine, cardiovascular, gastrointestinal, and immune systems.

1. Pharmacodynamic Properties and Mechanism of Action

The antidepressant efficacy of agomelatine arises from the synergy between its melatonergic agonism and serotonergic antagonism. By stimulating MT_1/MT_2 receptors, it corrects circadian rhythm disturbances, enhances sleep quality, and restores alignment with the light–dark cycle (2). This mechanism is particularly beneficial for sleep–wake disruptions commonly observed in depressive disorders. Simultaneously, postsynaptic blockade of 5-HT_2C receptors increases norepinephrine and dopamine levels in the frontal cortex, contributing to mood improvement, motivation, and enhanced cognitive function. Moreover, its 5-HT_2C antagonism may exert anxiolytic effects via modulation of GABAergic neurons; by indirectly enhancing GABA transmission in areas such as the amygdala, hippocampus, and bed nucleus of the stria terminalis, it may alleviate anxiety symptoms (1). Overall, agomelatine combines chronobiotic effects through melatonin receptor agonism with monoaminergic and anxiolytic actions via serotonin receptor antagonism, forming a distinctive pharmacodynamic profile.

Agomelatine is rapidly absorbed following oral administration, with over 75% absorption from the gastrointestinal tract. However, due to extensive first-pass metabolism, its absolute bioavailability is below 5%. It is primarily metabolized in the liver by the cytochrome P450 isoenzyme CYP1A2, with metabolites being largely excreted in the urine following glucuronidation. The elimination half-life of agomelatine is relatively short, approximately two hours, and its plasma concentrations increase proportionally with dose within the therapeutic range (3). Given its extensive hepatic metabolism, special attention should be paid to pharmacokinetics in elderly patients or individuals with impaired liver function (2).

2. Effects on the Central Nervous System

The most pronounced effects of agomelatine are mediated through its actions on the central nervous system. As a melatonin receptor agonist, it regulates disrupted circadian rhythms via the suprachiasmatic nucleus (SCN), leading to improved sleep architecture (4). In clinical cases of depression, agomelatine treatment has been shown to shorten sleep onset latency and increase the duration of deep (slow-wave) sleep, thereby restoring sleep quality. Particularly, the re-establishment of early-phase slow-wave sleep and normalization of REM sleep latency indicate its sleep-regulatory properties. These improvements translate into enhanced nocturnal rest and better daytime functioning in patients (5).

FULL TEXTS BOOK

Additionally, agomelatine exerts beneficial effects on neuroplasticity. In animal models, chronic administration of agomelatine has been shown to promote neurogenesis in the hippocampus and prefrontal cortex, upregulate brain-derived neurotrophic factor (BDNF), and activate intracellular signaling pathways such as AKT, ERK1/2, and GSK-3β (6). In an Alzheimer's disease-like rat model induced by $A\beta_{1-42}$ peptide, agomelatine significantly ameliorated cognitive and behavioral impairments, likely due to its anti-neuroinflammatory and neuroplasticity-enhancing effects (7). In summary, agomelatine exerts multifaceted effects on the central nervous system, including antidepressant, anxiolytic, sleep-regulatory, and cognitive-enhancing properties.

3. Effects on the Endocrine System

Circadian rhythm disturbances commonly observed in depressive disorders extend their impact to the endocrine system. In major depression, phase shifts and amplitude changes in several hormonal rhythms have been reported—for instance, the circadian phases of cortisol, ACTH, prolactin, and growth hormone are often advanced, while their daily secretion amplitudes are diminished. The melatonergic agonism of agomelatine has the potential to resynchronize these disrupted hormonal patterns. In animal studies, agomelatine rapidly restored the rhythms of biological parameters such as body temperature and hormone secretion in models exposed to altered light–dark cycles (2). This effect is attributed to its chronobiotic activity via the SCN and may aid in normalizing the daily cortisol cycle and other stress-related hormones in humans.

Importantly, agomelatine does not significantly suppress hypothalamic-pituitary-adrenal (HPA) axis activity. For example, in animal models, chronic agomelatine administration did not cause notable changes in plasma corticosterone or ACTH levels, nor in the expression of hypothalamic corticotropin-releasing hormone (CRH) or vasopressin (8, 9). These findings suggest that agomelatine contributes to the normalization of stress responses not by inhibiting the HPA axis, but by re-establishing rhythmic hormone release, thereby preserving overall endocrine balance.

4. Effects on the Cardiovascular System

Agomelatine has not been associated with any significant adverse physiological effects on the cardiovascular system. Clinical data indicate that it exerts neutral or minimal effects on heart rate and blood pressure (10). Studies have shown that agomelatine does not pose a risk of inducing positive inotropic stress on the myocardium under conditions such as hypertension or postoperative cardiac load. Instead, it may elicit a more balanced myocardial contractile response (11). Consequently, in patients with comorbid cardiovascular disease, agomelatine is considered a safer alternative compared to tricyclic antidepressants or serotonin-norepinephrine reuptake inhibitors (SNRIs), which carry a higher risk of cardiovascular side effects.

5. Effects on the Gastrointestinal System

Melatonin plays a crucial regulatory role in the gastrointestinal (GI) system, modulating intestinal motility, local inflammation, and visceral pain perception. Disruptions in endogenous melatonin secretion can impair GI rhythmicity and functionality. As a melatonin analogue, agomelatine may exhibit similar regulatory effects. Due to its high affinity for MT_1/MT_2 receptors and a longer half-life compared to endogenous melatonin, agomelatine is thought to influence bowel motility and secretion rhythms.

Interestingly, agomelatine also displays antagonistic activity at $5\text{-}HT_3$ receptors—an unexpected yet significant pharmacological feature. Given the critical role of $5\text{-}HT_3$ receptors in the GI system, their blockade can reduce excessive motility and increase pain thresholds, potentially benefiting patients with diarrhea-predominant irritable bowel syndrome (IBS) (12). These actions suggest that agomelatine may affect gastrointestinal function both centrally and peripherally, while maintaining a more favorable GI side effect profile than traditional antidepressants. This may be due to its lack of serotonin reuptake

FULL TEXTS BOOK

inhibition, which is often responsible for serotonergic side effects such as nausea and diarrhea. In summary, agomelatine has the potential to modulate bowel motility and visceral sensitivity, and ongoing research is exploring its therapeutic applications in gastrointestinal disorders.

6. Effects on the Immune System

Recent research has increasingly highlighted the role of neuroinflammation in the pathophysiology of depression. Elevated levels of pro-inflammatory cytokines such as interleukin-6 (IL-6) and tumor necrosis factor-alpha (TNF-α), along with chronic activation of the immune system, have frequently been reported in depressed patients. One of the notable aspects of agomelatine is its potential immunomodulatory activity. Animal studies have demonstrated that agomelatine treatment normalizes elevated IL-6 levels and IL-6–related signaling pathways following exposure to stress or endotoxins such as lipopolysaccharide (LPS). In chronic mild stress models, agomelatine was found to reduce anhedonic behavior while downregulating IL-6 expression and the JAK/STAT3 pathway in the brain, indicating a role in modulating inflammatory responses (13). This reduction in cytokine activity, along with decreased levels of acute-phase reactants like C-reactive protein (CRP), suggests that agomelatine possesses anti-inflammatory properties in addition to its antidepressant effects. Moreover, agomelatine has been shown to exhibit antioxidant activity by reducing extracellular free radical production and promoting the polarization of microglial cells toward the anti-inflammatory M2 phenotype in certain neurodegenerative models (14). Collectively, these findings reveal that agomelatine contributes to immune regulation and inflammation suppression, expanding its therapeutic potential beyond neurotransmitter modulation to encompass immunological mechanisms as well.

Conclusion

Agomelatine is a dual-action agent functioning through melatonin receptor agonism and serotonin receptor antagonism. Its broad spectrum of effects on sleep regulation, mood stabilization, cognitive enhancement, hormonal rhythmicity, gastrointestinal motility, and immune responses highlights its relevance to psychophysiological integration. Given its multi-systemic impact, both experimental and clinical research is needed to further elucidate the underlying physiological mechanisms and to optimize its clinical applications. The distinct pharmacodynamic profile of agomelatine, characterized by chronobiotic, neurotrophic, and immunomodulatory properties, positions it as a unique and promising option in the treatment of mood disorders.

REFERENCES

1. Savino, R., Polito, A. N., Marsala, G., Ventriglio, A., Di Salvatore, M., De Stefano, M. I., Valenzano, A., Marinaccio, L., Bellomo, A., Cibelli, G., Monda, M., Monda, V., Messina, A., Polito, R., Carotenuto, M., & Messina, G. (2023). Agomelatine: A Potential Multitarget Compound for Neurodevelopmental Disorders. Brain Sciences, 13(5), 734. https://doi.org/10.3390/brainsci13050734

2. De Berardis, D., Marini, S., Fornaro, M., Srinivasan, V., Iasevoli, F., Tomasetti, C., Valchera, A., Perna, G., Quera-Salva, M.-A., Martinotti, G., & Di Giannantonio, M. (2013). The Melatonergic System in Mood and Anxiety Disorders and the Role of Agomelatine: Implications for Clinical Practice. *International Journal of Molecular Sciences*, *14*(6), 12458-12483. https://doi.org/10.3390/ijms140612458

3. San, L., & Arranz, B. (2008). Agomelatine: a novel mechanism of antidepressant action involving the melatonergic and the serotonergic system. *European psychiatry : the journal of the Association of European Psychiatrists*, *23*(6), 396–402. https://doi.org/10.1016/j.eurpsy.2008.04.002

FULL TEXTS BOOK

4. Srinivasan, V., Zakaria, R., Othman, Z., Lauterbach, E. C., & Acuña-Castroviejo, D. (2012). Agomelatine in depressive disorders: its novel mechanisms of action. *The Journal of neuropsychiatry and clinical neurosciences*, *24*(3), 290–308. https://doi.org/10.1176/appi.neuropsych.11090216

5. Mairesse, J., Silletti, V., Laloux, C., Zuena, A. R., Giovine, A., Consolazione, M., van Camp, G., Malagodi, M., Gaetani, S., Cianci, S., Catalani, A., Mennuni, G., Mazzetta, A., van Reeth, O., Gabriel, C., Mocaër, E., Nicoletti, F., Morley-Fletcher, S., & Maccari, S. (2013). Chronic agomelatine treatment corrects the abnormalities in the circadian rhythm of motor activity and sleep/wake cycle induced by prenatal restraint stress in adult rats. *The international journal of neuropsychopharmacology*, *16*(2), 323–338. https://doi.org/10.1017/S1461145711001970

6. Pompili, M., Serafini, G., Innamorati, M., Venturini, P., Fusar-Poli, P., Sher, L., Amore, M., & Girardi, P. (2013). Agomelatine, a novel intriguing antidepressant option enhancing neuroplasticity: a critical review. *The world journal of biological psychiatry : the official journal of the World Federation of Societies of Biological Psychiatry*, *14*(6), 412–431. https://doi.org/10.3109/15622975.2013.765593

7. Ozen Koca, R., Solak Gormus, Z. I., Solak, H., Gultekin, B., Ozdemir, A., Eroglu Gunes, C., Kurar, E., & Kutlu, S. (2025). Agomelatine Ameliorates Cognitive and Behavioral Deficits in Aβ-Induced Alzheimer's Disease-like Rat Model. Medicina, 61(8), 1315. https://doi.org/10.3390/medicina61081315

8. Courtet, P., & Olié, E. (2012). Circadian dimension and severity of depression. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology*, *22 Suppl 3*, S476–S481. https://doi.org/10.1016/j.euroneuro.2012.07.009

9. Leproult, R., Van Onderbergen, A., L'hermite-Balériaux, M., Van Cauter, E., & Copinschi, G. (2005). Phase-shifts of 24-h rhythms of hormonal release and body temperature following early evening administration of the melatonin agonist agomelatine in healthy older men. *Clinical endocrinology*, *63*(3), 298–304. https://doi.org/10.1111/j.1365-2265.2005.02341.x

10. Calvi, A., Fischetti, I., Verzicco, I., Belvederi Murri, M., Zanetidou, S., Volpi, R., Coghi, P., Tedeschi, S., Amore, M., & Cabassi, A. (2021). Antidepressant Drugs Effects on Blood Pressure. *Frontiers in cardiovascular medicine*, *8*, 704281. https://doi.org/10.3389/fcvm.2021.704281

11. Özen Koca R, Solak Görmüş ZI, Solak H, Akdoğan G. How does agomelatine affect contraction in rat myocardial tissue? Eur Res J. July 2025;11(4):669-675. doi:10.18621/eurj.1644498

12. Ng, Q. X., Soh, A. Y. S., Lim, D. Y., & Yeo, W. S. (2018). Agomelatine, a novel therapeutic option for the management of irritable bowel syndrome. Journal of clinical pharmacy and therapeutics, 43(5), 752–756. https://doi.org/10.1111/jcpt.12749

13. Rossetti, A. C., Paladini, M. S., Brüning, C. A., Spero, V., Cattaneo, M. G., Racagni, G., Papp, M., Riva, M. A., & Molteni, R. (2022). Involvement of the IL-6 Signaling Pathway in the Anti-Anhedonic Effect of the Antidepressant Agomelatine in the Chronic Mild Stress Model of Depression. International journal of molecular sciences, 23(20), 12453. https://doi.org/10.3390/ijms232012453

14. Yang, X. B., Zu, H. B., Zhao, Y. F., & Yao, K. (2022). Agomelatine Prevents Amyloid Plaque Deposition, Tau Phosphorylation, and Neuroinflammation in APP/PS1 Mice. Frontiers in aging neuroscience, 13, 766410. https://doi.org/10.3389/fnagi.2021.766410

FULL TEXTS BOOK

HYBRID PARABOLIC TROUGH COLLECTOR AND THERMOELECTRIC GENERATOR SYSTEMS FOR MULTIGENERATION

E. CUCE[a]

[a] *Department of Mechanical Engineering, Faculty of Engineering and Architecture, Recep Tayyip Erdogan University, Zihni Derin Campus, 53100 - Rize, Turkiye*

ORCID: 0000-0003-0150-4705

P. MERT CUCE[b, c]

[b] *Department of Architecture, Faculty of Engineering and Architecture, Recep Tayyip Erdogan University, Zihni Derin Campus, 53100 - Rize, Turkiye*

[c] *College of Built Environment, Birmingham City University, Birmingham B4 BD, UK*

ORCID:0000-0002-6522-7092

T. GUCLU[d]

[d] *Department of Mechanical Engineering, Faculty of Engineering, University of Bayburt, Dede Korkut Campus, 69000 - Bayburt, Turkiye*

ORCID: 0000-0002-5864-3864

ABSTRACT

As the global demand for sustainable and efficient energy solutions continues to grow, the development and integration of hybrid energy systems has become a key focus in renewable energy research. Among these, multi-generation systems that combine Parabolic Trough Collectors (PTC) and Thermoelectric Generators (TEG) have emerged as highly promising technologies due to their ability to harness and convert solar energy efficiently. This study provides an in-depth analysis of the integration of PTC and TEG technologies, highlighting their complementary characteristics and potential application areas. PTC systems are widely recognised for their ability to capture and utilise high-temperature solar energy, making them suitable for various thermal applications. On the other hand, TEGs generate electricity by exploiting temperature differences, enabling direct thermal-to-electrical energy conversion without moving parts. When combined, these two technologies create a hybrid system that enhances overall energy efficiency, increases operational flexibility, and contributes to long-term sustainability. This study explores various design configurations, operating principles, and performance optimisations of PTC-TEG hybrid systems while also examining their potential applications in diverse fields. These include electricity generation, water desalination, industrial and residential heating/cooling, and hydrogen production—each of which plays a vital role in advancing clean energy solutions. Furthermore, by improving the utilisation of waste heat and maximising energy output, these hybrid systems offer significant environmental benefits, reducing greenhouse gas emissions and promoting a low-carbon energy transition. The findings of this study indicate that PTC-TEG hybrid systems not only improve the efficiency of energy conversion processes but also support sustainability goals by optimising renewable energy utilisation. As the world moves toward a cleaner and more resilient energy infrastructure, these systems have the potential to play a crucial role in addressing global energy challenges and advancing the transition to a more sustainable future.

Keywords: parabolic trough collectors, thermoelectric generators, hybrid systems

FULL TEXTS BOOK

1. INTRODUCTION

Rapidly increasing global energy demand and the environmental pressures created by climate change have necessitated new and sustainable approaches to energy production. The limited reserves of fossil fuels and the resulting greenhouse gas emissions during their use are contributing to the depletion of natural resources and accelerating global warming [1]. This has spurred the development of renewable energy technologies, particularly highlighting solar energy, a clean, unlimited, and environmentally friendly resource. However, traditional methods of harnessing solar energy have technical limitations, such as low efficiency and storage difficulties. Therefore, the development of innovative systems that can maximise the benefits of solar energy has become crucial [2].

Concentrating solar power (CSP) technologies are one of the leading solutions in this context. Their basic operating principle is to focus sunlight onto a receiver using reflective surfaces, generating a high-temperature heat carrier fluid. This thermal energy can be used for electricity generation with turbine or engine systems, or directly in industrial heating processes [3]. Furthermore, when integrated with thermal energy storage units, it becomes possible to provide electricity or heat even when sunlight is not available [4]. Among CSP technologies, Parabolic Trough Collectors (PTCs) stand out due to their high efficiency and widespread commercial use. Parabolic reflective mirrors direct sunlight to a receiver tube at the focal line, heating the fluid passing through it to a temperature between 150–400°C. These temperature levels offer a suitable energy source for both power generation and industrial applications [5].

Thermoelectric Generators (TEGs), on the other hand, are semiconductor-based devices that convert temperature differences directly into electrical energy. This technology, based on the Seebeck effect, stands out as a long-lasting and reliable energy conversion tool thanks to its lack of moving parts, quiet operation, and low maintenance requirements [6]. Thanks to their ability to generate energy even from low temperature differences, TEGs have broad application potential in areas such as waste heat recovery. However, low conversion efficiency alone is the primary factor limiting the widespread adoption of this technology [7].

The hybrid use of PTC and TEG technologies combines the strengths of both systems, providing efficiency and flexibility in energy production. PTCs generate high-temperature thermal energy, while TEGs convert the remaining waste heat into electrical energy, increasing overall system efficiency. This enables multiple outputs such as electricity, heating, cooling, water treatment, and even hydrogen production. This hybrid structure stands out as an environmentally friendly solution that aligns with sustainable development goals and promotes energy independence.

2. DESIGN STRATEGIES OF PTC-TEG INTEGRATION

The primary objective of PTC-TEG hybrid systems is to utilise the high-temperature heat generated from concentrated solar energy in the primary energy conversion process, followed by the utilisation of remaining waste heat for electricity generation. This system not only increases overall energy efficiency but also offers the potential for multiple generations by providing different energy forms from the same platform [8]. The most critical element in the design process is determining the integration points of TEG modules into the PTC system. The most common approach is to place TEGs on the outer surface of the receiver tubes or in the output line. This allows the high-temperature heat transferred from the fluid to the environment to be converted directly into electricity through the TEG modules. Alternatively, TEG arrays placed behind reflective surfaces can provide additional energy generation from direct solar or reflected radiation [9]. To increase heat transfer efficiency, the thermal contact resistance between the TEG and PTC must be minimised. Highly thermally conductive interlayer

FULL TEXTS BOOK

materials (e.g., graphite sheets, thermal pastes) are used for this purpose. Effective cooling of the cold surface of the TEG modules is also critical to performance; water cooling, air ducts, or heat pipes are commonly used for this purpose [10]. Thermal management in hybrid systems is crucial not only for efficiency but also for extending the life of the modules. Excessive PTC outlet temperatures can exceed the thermal resistance limits of TEG materials; therefore, intermediate heat exchangers or fluid flow control are used for temperature regulation.

The operating principle consists of two main stages:

- Primary Heat Collection: PTC focuses solar radiation and raises the heat carrier fluid to a high temperature. This energy can be used directly in turbines, organic Rankine cycles, or industrial processes.
- Waste Heat Electricity Generation: The remaining heat after primary use is converted directly into electricity by passing it through TEG modules.

This electricity can power the system's auxiliary equipment or be transferred to the grid. This structure allows for an additional efficiency increase of 5–15% compared to standalone PTC or TEG systems. Furthermore, the hybrid configuration increases continuity in energy production, making it a viable solution for both centralised and distributed generation scenarios.

3. HYBRID PTC-TEG APPLICATIONS

PTC-TEG hybrid systems stand out for their ability to not only generate electricity but also to simultaneously provide different forms of energy through the same platform. This approach, described as "multigeneration," enables more efficient use of energy resources, lower operating costs, and carbon emission reduction. Its primary application is electricity generation. PTC focuses high-intensity sunlight onto a receiver tube, raising the heat transfer fluid to high temperatures. The resulting temperature difference is directly converted into electricity via TEG modules, providing a reliable and sustainable power source in off-grid regions. Habchi et al. [11] performed numerical analyses on two different cogeneration systems integrating PTC and cylindrical TEG modules. In their MATLAB-based models, they achieved system efficiency of 70.55% when using Therminol VP1 as the heat transfer fluid and 76.91% when using water. The high-temperature thermal energy generated by the system can be used in thermal applications such as industrial process heating, district heating, or hot water supply. Such uses increase energy efficiency, particularly in industrial facilities and communal areas. Low-temperature waste heat can be channelled to cooling applications using absorption cooling systems. This method reduces fossil fuel consumption in building air conditioning, food and agricultural product storage, or industrial cooling processes. Khanmohammadi et al. [12] designed a PTC-based power-cooling system, integrated it with TEG modules, and conducted energy, exergy, and exergoeconomic analyses. The study results revealed that TEG integration significantly improved the system's exergy performance. PTC-TEG hybrid structures can also be integrated with water treatment technologies to convert seawater into potable water. Thermal distillation methods require a high-temperature energy source, and the heat provided by the PTC can be used directly in these systems. This allows for both energy and water needs to be met from the same system, especially in arid regions. Assareh et al. [13] developed an integrated solar energy system that uses a TEG unit instead of a condenser. The system is designed to provide both electricity and seawater desalination. Analysis results show that TEG integration reduces electricity generation costs and increases exergy efficiency. Another important application area is hydrogen production. Thermochemical reactions and high-temperature electrolysis methods require intense thermal energy. Hybrid systems can power electrolysers with heat from PTCs, while additional electricity from TEGs can be used in these processes. This makes renewable-based, zero-carbon hydrogen production possible. Habibollahzade et al. [14] designed a system integrating a PTC-TEG-Rankine cycle with a PEM electrolyser and conducted multi-criteria optimisation. The

FULL TEXTS BOOK

findings revealed that using a TEG instead of a capacitor increases both energy production and exergy efficiency, while also making the hydrogen production process more economical.

4. CONCLUSION

This study examines multiple-generation systems utilising hybrid PTC and TEG thermoelectric generator technologies. A literature review demonstrates that these hybrid structures can be used with high efficiency not only in electricity generation but also in various applications such as heating, cooling, water treatment, and hydrogen production. The high-temperature thermal energy generation capacity of PTC and the ability of TEG to generate electricity directly from temperature differences are complementary. This allows for higher overall efficiency and more flexible energy generation scenarios compared to standalone systems. The findings demonstrate that PTC-TEG hybrid systems have the potential to reduce fossil fuel dependency, reduce greenhouse gas emissions, and increase energy security. However, R&D activities are still needed regarding high initial investment costs, limited TEG material efficiency, and system scalability. In the future, advanced thermoelectric materials, optimised thermal designs, and integrated energy storage solutions will accelerate the deployment of these hybrid systems, making them economically more attractive, especially in regions with high solar radiation. In conclusion, PTC-TEG hybrid systems stand out as an environmentally friendly and long-term economically advantageous solution that increases efficiency by providing multiple outputs in energy production. These systems are poised to play a significant role in the future energy infrastructure, both in terms of technical performance and sustainability objectives.

REFERENCES

[1] Cuce E, Cuce PM, Guclu T, Besir AB. On the Use of Nanofluids in Solar Energy Applications. Journal of Thermal Science 2020 29:3 2020;29:513–34. https://doi.org/10.1007/S11630-020-1269-3.

[2] Kumar CMS, Singh S, Gupta MK, Nimdeo YM, Raushan R, Deorankar A V., et al. Solar energy: A promising renewable source for meeting energy demand in Indian agriculture applications. Sustainable Energy Technologies and Assessments 2023;55:102905. https://doi.org/10.1016/J.SETA.2022.102905.

[3] Domínguez R, Conejo AJ, Carrión M. Operation of a fully renewable electric energy system with CSP plants. Appl Energy 2014;119:417–30. https://doi.org/10.1016/J.APENERGY.2014.01.014.

[4] Pelay U, Luo L, Fan Y, Stitou D, Rood M. Thermal energy storage systems for concentrated solar power plants. Renewable and Sustainable Energy Reviews 2017;79:82–100. https://doi.org/10.1016/J.RSER.2017.03.139.

[5] Cuce PM, Guclu T, Cuce E. Design, modelling, environmental, economic and performance analysis of parabolic trough solar collector (PTC) based cogeneration systems assisted by thermoelectric generators (TEGs). Sustainable Energy Technologies and Assessments 2024;64:103745. https://doi.org/10.1016/J.SETA.2024.103745.

[6] Guclu T, Cuce E. Thermoelectric Coolers (TECs): From Theory to Practice. J Electron Mater 2019;48:211–30. https://doi.org/10.1007/S11664-018-6753-0/METRICS.

FULL TEXTS BOOK

[7] Cuce E, Guclu T, Cuce PM. Improving thermal performance of thermoelectric coolers (TECs) through a nanofluid driven water to air heat exchanger design: An experimental research. Energy Convers Manag 2020;214:112893. https://doi.org/10.1016/J.ENCONMAN.2020.112893.

[8] Sui X, Li W, Zhang Y, Wu Y. Theoretical and Experimental Evaluation of a Thermoelectric Generator Using Concentration and Thermal Energy Storage. IEEE Access 2020;8:87820–8. https://doi.org/10.1109/ACCESS.2020.2993288.

[9] Sripadmanabhan Indira S, Vaithilingam CA, Chong KK, Saidur R, Faizal M, Abubakar S, et al. A review on various configurations of hybrid concentrator photovoltaic and thermoelectric generator system. Solar Energy 2020;201:122–48. https://doi.org/10.1016/J.SOLENER.2020.02.090.

[10] Sundarraj P, Maity D, Roy SS, Taylor RA. Recent advances in thermoelectric materials and solar thermoelectric generators – a critical review. RSC Adv 2014;4:46860–74. https://doi.org/10.1039/C4RA05322B.

[11] Habchi A, Hartiti B, Labrim H, Fadili S, Benyoussef A, Belouaggadia N, et al. Feasibility and Thermal/Electrical performance study of two smart hybrid systems combining parabolic trough collector with tubular thermoelectric generator. Energy Reports 2021;7:1539–59. https://doi.org/10.1016/J.EGYR.2021.03.002.

[12] Khanmohammadi S, Musharavati F, Kizilkan O, Duc Nguyen D. Proposal of a new parabolic solar collector assisted power-refrigeration system integrated with thermoelectric generator using 3E analyses: Energy, exergy, and exergo-economic. Energy Convers Manag 2020;220:113055. https://doi.org/10.1016/J.ENCONMAN.2020.113055.

[13] Assareh E, Alirahmi SM, Ahmadi P. A Sustainable model for the integration of solar and geothermal energy boosted with thermoelectric generators (TEGs) for electricity, cooling and desalination purpose. Geothermics 2021;92:102042. https://doi.org/10.1016/J.GEOTHERMICS.2021.102042.

[14] Habibollahzade A, Gholamian E, Ahmadi P, Behzadi A. Multi-criteria optimization of an integrated energy system with thermoelectric generator, parabolic trough solar collector and electrolysis for hydrogen production. Int J Hydrogen Energy 2018;43:14140–57. https://doi.org/10.1016/J.IJHYDENE.2018.05.143.

HAVA KALİTESİ İNDEKSİ TAHMİNİNDE ARIMA, SARIMA VE LSTM MODELLERİNİN PERFORMANSLARININ KARŞILAŞTIRILMASI

COMPARATIVE PERFORMANCE ANALYSIS OF ARIMA, SARIMA, AND LSTM MODELS FOR AIR QUALITY INDEX FORECASTING

Dr. Öğr. Üy. Barış ERGÜL [1]

[1] *Eskişehir Osmangazi Üniversitesi, Fen Fakültesi, İstatistik Bölümü, Eskişehir, Türkiye.*

[1]*ORCID ID: https://orcid.org/0000-0002-1811-5143*

Prof. Dr. Arzu ALTIN YAVUZ [2]

[2] *Eskişehir Osmangazi Üniversitesi, Fen Fakültesi, İstatistik Bölümü, Eskişehir, Türkiye.*

[2]*ORCID ID: https://orcid.org/0000-0002-3277-740X*

ÖZET

Hava kalitesinin izlenmesi ve tahmini, insan sağlığı ile çevresel sürdürülebilirlik açısından büyük önem taşımaktadır. Bu çalışmada, Hava Kalitesi İndeksi (HKİ) verilerinin zaman serisi tahmini için yaygın olarak kullanılan ARIMA (Autoregressive Integrated Moving Average), mevsimsel bileşenleri dikkate alan SARIMA (Seasonal ARIMA) ve derin öğrenme temelli LSTM (Long Short-Term Memory) modellerinin performansları karşılaştırılmıştır.

Çalışmada, uzun dönemli ve mevsimsel özellikler gösteren HKİ veri seti kullanılmış olup, serinin durağanlığı incelenmiştir. ARIMA modeli, verinin durağan hale getirilmesinin ardından optimizasyon yöntemleriyle belirlenen parametrelerle oluşturulmuştur. SARIMA modelinde ise mevsimsel farklılıkların modele entegrasyonu sağlanarak daha kapsamlı bir tahmin yaklaşımı benimsenmiştir. Buna karşılık, LSTM modeli, zaman serilerindeki uzun dönemli bağımlılıkları ve doğrusal olmayan ilişkileri yüksek doğrulukla modelleyebilme avantajı nedeniyle tercih edilmiştir.

Modellerin performans değerlendirmesi, Hataların Karesinin Ortalamasının Kareköküne (RMSE) göre yapılmıştır. Analiz sonuçları, mevsimsel özelliklerin belirgin olduğu durumlarda SARIMA'nın, geleneksel ARIMA modeline kıyasla daha başarılı tahminler ürettiğini göstermiştir. Bununla birlikte, LSTM modeli, hem yüksek tahmin doğruluğu hem de karmaşık veri desenlerine uyum sağlama yeteneği sayesinde diğer modelleri performans açısından geride bırakmıştır.

Bu bulgular, hava kalitesi tahmininde derin öğrenme tabanlı yaklaşımların geleneksel zaman serisi yöntemlerine göre daha etkili sonuçlar verebileceğini ortaya koymaktadır. Özellikle, dinamik ve periyodik özellikler gösteren çevresel verilerin analizinde LSTM modellerinin kullanımı, erken uyarı sistemleri ve politika belirleme süreçlerine önemli katkı sağlayabilir.

Anahtar Kelimeler: ARIMA, SARIMA, LSTM, Türkiye.

ABSTRACT

Monitoring and forecasting air quality are of paramount importance for human health and environmental sustainability. In this study, the performance of widely used time series forecasting models -ARIMA (Autoregressive Integrated Moving Average), SARIMA (Seasonal ARIMA) which accounts for seasonal components, and the deep learning-based LSTM (Long Short-Term Memory)- was compared using Air Quality Index (AQI) data.

The study utilized an AQI dataset exhibiting long-term trends and seasonal patterns, and the stationarity of the series was examined. The ARIMA model was developed by first differencing the data to achieve

FULL TEXTS BOOK

stationarity, followed by parameter selection through optimization techniques. The SARIMA model incorporated seasonal variations to provide a more comprehensive forecasting approach. Conversely, the LSTM model was preferred due to its ability to effectively capture long-term dependencies and nonlinear relationships in time series data with high accuracy.

Model performance was evaluated primarily using the Root Mean Square Error (RMSE) metric. Analysis results indicated that SARIMA produced more accurate forecasts than the traditional ARIMA model in scenarios where seasonal patterns were pronounced. Furthermore, the LSTM model outperformed both ARIMA and SARIMA in terms of forecasting accuracy and adaptability to complex data patterns.

These findings suggest that deep learning-based approaches can yield more effective results than conventional time series methods in air quality forecasting. Particularly, the application of LSTM models in analyzing dynamic and periodic environmental data can significantly contribute to the development of early warning systems and policy-making processes.

Keywords: ARIMA, SARIMA, LSTM, Türkiye.

GİRİŞ

Modern yaşamın karmaşası içerisinde çoğu zaman farkına varmasak da, soluduğumuz hava sağlığımızı, çevremizi ve geleceğimizi derinden etkiliyor. Hava kalitesi, bireysel yaşamlarımızdan küresel politik gündemlere kadar pek çok düzeyde önemli sonuçlar doğuran çevresel bir faktördür. Temiz hava hakkı, temel bir insan hakkı olarak görülse de, ne yazık ki dünya genelinde bu hakkın güvence altına alındığını söylemek güçtür. Dünya Sağlık Örgütü'ne (WHO, 2021) göre, dünya nüfusunun neredeyse tamamı, DSÖ'nün önerdiği hava kalitesi sınırlarının üzerinde kirletici maddelere maruz kalmaktadır. Bu çarpıcı oran, hava kirliliğinin artık yalnızca yerel değil, küresel ölçekte bir kriz haline geldiğini açıkça ortaya koymaktadır.

Hava kirliliği, temel olarak çeşitli zararlı gaz ve parçacıkların atmosferde birikmesi sonucu oluşur. Partikül maddeler ($PM_{2.5}$ ve PM_{10}), azot dioksit (NO_2), kükürt dioksit (SO_2), karbon monoksit (CO) ve troposferik ozon (O_3) gibi başlıca kirleticiler, atmosferde yüksek yoğunlukta bulunduğunda insan sağlığı üzerinde ciddi tehditler oluşturur. Bu maddeler, sadece akut solunum yolu problemlerine değil, aynı zamanda uzun vadede kalp-damar hastalıklarına, inmelere ve bazı kanser türlerine kadar uzanan ciddi sağlık sorunlarına yol açabilmektedir (Brook et al., 2010). Özellikle çocuklar, yaşlılar ve kronik hastalıkları olan bireyler bu kirleticilere karşı çok daha savunmasızdır.

Hava kalitesinin bozulmasındaki en önemli etkenler, insan faaliyetlerine dayanmaktadır. Yoğun sanayi üretimi, motorlu taşıt kullanımı, fosil yakıt tüketimi ve ormansızlaşma gibi faktörler, hava kirliliğini artıran başlıca kaynaklar arasında yer almaktadır. Gelişmekte olan ülkelerde, özellikle çevre politikalarının yetersiz kaldığı büyük şehirlerde bu etkiler daha da belirgin hale gelmektedir (Gurjar et al., 2016). Türkiye örneği üzerinden değerlendirildiğinde, İstanbul, Ankara ve İzmir gibi büyükşehirlerde hava kalitesi, yılın büyük bir kısmında ulusal ve uluslararası standartların altında seyretmektedir (ÇŞİDB, 2023). Bu durum, yalnızca bireysel sağlık değil, aynı zamanda toplumsal refah açısından da ciddi bir risk oluşturmaktadır.

Ancak sorun sadece insan sağlığıyla sınırlı değildir. Hava kirliliği; tarımsal üretimi, ekosistem dengesini ve hatta iklim sistemlerini doğrudan etkilemektedir. Örneğin troposferik ozonun artışı, yalnızca atmosferde sera gazı etkisi yaratmakla kalmaz; aynı zamanda bitkilerin fotosentez süreçlerini bozarak ürün verimliliğini düşürür (Avnery et al., 2011). Bu durum, özellikle gıda güvenliği açısından önemli bir tehdit oluşturmaktadır. Aynı şekilde, su kaynaklarının kimyasal bileşimi de hava yoluyla taşınan kirleticilerden etkilenebilmekte, bu da ekolojik zinciri daha da karmaşık hale getirmektedir.

FULL TEXTS BOOK

Öte yandan, teknolojideki ilerlemeler hava kalitesinin izlenmesi ve değerlendirilmesinde yeni imkanlar sunmaktadır. Uzaktan algılama sistemleri, uydu görüntüleri ve yapay zeka tabanlı veri analiz yöntemleri sayesinde, hava kirliliğine dair daha ayrıntılı ve anlık bilgilere ulaşmak artık mümkündür. Ancak, bu verilerin karar alma süreçlerine etkin biçimde entegre edilmesi, hem teknik hem de yönetsel anlamda hâlâ ciddi zorluklar barındırmaktadır. Literatür, hava kalitesiyle ilgili politikaların etkin olabilmesi için yalnızca teknik çözümler değil, aynı zamanda disiplinler arası iş birliği ve toplumsal katılımın da şart olduğunu göstermektedir (Sicard et al., 2020).

Hava kalitesi yalnızca çevresel bir gösterge değildir, ek olarak toplum sağlığını, ekonomik üretkenliği ve ekolojik sürdürülebilirliği doğrudan etkileyen, çok boyutlu bir konudur. Bu konuda yapılacak her çalışma, yalnızca bilimsel bilgi üretmekle kalmayıp, aynı zamanda kamu politikalarına ve toplumsal farkındalık için katkı sunma potansiyeline sahiptir. Bu çalışma da, söz konusu amaca katkı sunmayı hedeflemektedir.

YÖNTEM

Zaman serisi verileri, doğası gereği geçmişten bugüne süreklilik gösteren ve geleceğe dair öngörülerde bulunmaya elverişli yapılar sunar. Finansal piyasalardan meteorolojiye, enerji talebinden kamu sağlığına kadar pek çok alanda zamanla değişen veriler, yalnızca geçmişin kaydı değil; aynı zamanda geleceğin ipuçlarını da barındırır. Bu nedenle, bu verileri anlamlandırmak ve tahminler üretmek için kullanılan yöntemlerin hem güçlü hem de bağlama duyarlı olması gerekir. İşte bu noktada ARIMA, SARIMA ve LSTM gibi modeller öne çıkar.

ARIMA (Autoregressive Integrated Moving Average), zaman serisi modelleme dünyasının en köklü ve yaygın kullanılan yöntemlerinden biridir. Bu model, geçmiş verilerin bugünkü değeri üzerindeki etkisini otoregresif (AR) bileşen ile; verideki trendin giderilmesini fark alma (I) bileşeni ile; geçmişteki hata terimlerinin etkisini ise hareketli ortalama (MA) bileşeni ile modellemeye çalışır. Bu yapı sayesinde model, zaman serisinin durağan hale getirilerek istikrarlı öngörüler üretmesini sağlar (Hyndman & Athanasopoulos, 2018). Ancak modelin başarısı, doğru parametrelerin belirlenmesine ve verinin doğrusal yapılar barındırmasına bağlıdır.

Pek çok zaman serisi, yalnızca trend değil, aynı zamanda mevsimsel örüntüler de içerir. SARIMA (Seasonal ARIMA), bu döngüsel davranışları modelleyebilmek için ARIMA modelini genişletir. SARIMA(p,d,q)(P,D,Q)_s formu ile gösterilir. Buradaki ikinci üçlü (P,D,Q), mevsimsel AR, I ve MA terimlerini temsil eder; s ise mevsimsel periyottur. Bu sayede, hem trend hem de mevsimsel etkiler bir arada modellenebilir (Hyndman & Khandakar, 2008). Modelin avantajı, istatistiksel olarak yorumlanabilir olmasıdır. Ancak mevsimsel bileşenlerin de eklenmesiyle parametrik yapı oldukça karmaşık duruma gelir ve hesaplamalar yoğunlaşır.

Geleneksel istatistiksel yöntemlerin bazı sınırlamaları, özellikle doğrusal olmayan ve karmaşık veri yapılarında belirgin hale gelir. Derin öğrenme yaklaşımları bu noktada devreye girer ve zaman serisi tahmininde öne çıkan modellerden biri de LSTM (Long Short-Term Memory) ağlarıdır. LSTM, özellikle zaman serilerinde uzun dönemli bağımlılıkları modellemek üzere geliştirilmiştir. RNN'lerdeki unutma problemine çözüm olarak geliştirilen LSTM, her zaman adımında hatırlanması gereken bilgileri seçerek taşıyabilir. Bir LSTM hücresinin bileşenleri; giriş kapısı, unutma kapısı, hücre durumu, çıkış kapısı, gizli durum olarak bilinmektedir. LSTM, istatistiksel modellere kıyasla daha fazla veri gerektirir; ayrıca modelin eğitilmesi işlem gücü açısından maliyetlidir. Fakat model, doğrusal olmayan örüntüleri öğrenmede son derece başarılıdır (Hochreiter & Schmidhuber, 1997).

ARIMA ve SARIMA modelleri, az değişkenli ve düzenli yapılar içeren zaman serilerinde güçlü performans gösterirken, LSTM gibi sinir ağı tabanlı modeller, büyük hacimli, çok değişkenli ve karmaşık verilerde öne çıkar. Son yıllarda geliştirilen hibrit yaklaşımlar (ARIMA-LSTM), her iki

FULL TEXTS BOOK

yöntemin güçlü yönlerini bir araya getirerek tahmin doğruluğunu artırmayı hedeflemektedir (Zhang et al., 2020).


Bu modeller, sadece teknik başarılarıyla değil, aynı zamanda karar vericilere sağladıkları veriden anlam çıkarma durumu için de değerlidir. Bu bağlamda, zaman serisi modellemesi, geçmişten gelen veriyi yalnızca tahmin etmek için değil, anlamlandırmak ve geleceği daha iyi planlamak için de vazgeçilmezdir.

BULGULAR

Kış 2000-Sonbahar 2025 tarihleri arasında Türkiye için hava kalitesi indeksi ölçüm değerleri Çevre,Şehircilik ve İklim Bakanlığı Hava Kalitesi bültenlerinin yer aldığı web sayfalarından ilgili bölümlerden alınmıştır.


İstatistik Analizler, R ve Python programları kullanılarak yapılmıştır. ARIMA modeli için uygun modelin belirlenmesinde ilk aşama olan geçici modelin saptanması aşamasında, Türkiye için zaman serisinin grafiği ile otokorelasyon fonksiyonu grafiği ve kısmi otokorelasyon fonksiyonu grafikleri Şekil 1-2'te görüldüğü gibidir. Grafikler incelendiği zaman, ilk birkaç otokorelasyon ve kısmi otokorelasyon değerlerinin sınırlar dışında yer aldığı ve diğer gecikme değerlerinin de sınırlar içinde yer aldığı görülmektedir ve ilk gecikme değerlerinin anlamsız oldukları tespit edilmiştir. Bu nedenlerden dolayı, serinin durağan olduğuna karar verilmiştir.

Şekil 1. Türkiye İçin Hava Kalite İndeksi Grafiği



FULL TEXTS BOOK

Şekil 2. Türkiye İçin Hava Kalitesi İndeks ACF ve PACF Grafiği



Daha sonraki aşamada ise ilgili zaman serisi için çeşitli ARIMA modelleri denenmiştir. Sonuçlar Tablo 1' de verildiği gibidir. Tablo 1 göz önüne alındığında, BIC değeri en küçük olan model, ARIMA(1,1,1) modelidir ve Tablo 2 incelendiğinde, modelin parametre tahminlerinin anlamlı olduğu görülmektedir. Bu nedenle, ARIMA(1,1,1) modeli geçici uygun modeldir. Geçici modelin uygunluğu, Ljung-Box testi ile test edilir. Ljung-Box testi sonucu Tablo 3'te gösterilmiştir. Geçici modelin uygun model olduğuna Ljung-Box istatistiği ile karar verilir ($p > 0.05$).

Tablo 1. Türkiye Hava Kalitesi İndeksi Farklı ARIMA Modelleri ve BIC Değerleri

ARIMA Modelleri	BIC Değerleri
ARIMA(1,1,1)	**896.350**
ARIMA(1,0,0)	899.386
ARIMA(0,0,1)	899.363

Tablo 2. Türkiye Hava Kalitesi İndeksi ARIMA(1,1,1) Modeli Parametre Değerleri

Parametre	Kestirim	SE	t	p	R.M.S.E.
Sabit	293.423	52.733	5.565	0.001	
AR	0.624	0.101	6.178	0.001	22.024
MA	-0,984	0.042	-23.579	0.001	


Tablo 3. Türkiye Hava Kalitesi İndeksi ARIMA(1,1,1) Modeli Uygunluğu

İstatistik	p
4.502	0.922


FULL TEXTS BOOK

Sonraki aşamada, SARIMA için ACF ve PACF grafikleri çizilir. İlgili grafikler Şekil 3'te gösterilmiştir. Grafikler incelendiği zaman, ilk birkaç otokorelasyon ve kısmi otokorelasyon değerlerinin sınırlar dışında yer aldığı ve diğer gecikme değerlerinin de sınırlar içinde yer aldığı görülmektedir ve ilk gecikme değerlerinin anlamsız oldukları tespit edilmiştir. İlgili zaman serisi için çeşitli SARIMA modelleri denenmiştir. Sonuçlar Tablo 4' de verildiği gibidir. Tablo 4 göz önüne alındığında, BIC değeri en küçük olan model, SARIMA(1,1,1) (1,1,1,4) modelidir ve Tablo 5 incelendiğinde, modelin parametre tahminlerinin anlamlı olduğu görülmektedir. Bu nedenle, SARIMA(1,1,1) (1,1,1,4) modeli geçici uygun modeldir. Geçici modelin uygunluğu, Ljung-Box testi ile test edilir. Ljung-Box testi sonucu Tablo 6'da gösterilmiştir. Geçici modelin uygun model olduğuna Ljung-Box istatistiği ile karar verilir ($p > 0.05$).

Şekil 3. Türkiye İçin Hava Kalitesi İndeksi Mevsimsel ACF ve PACF Grafiği



FULL TEXTS BOOK



Tablo 4. Türkiye Hava Kalitesi İndeksi Farklı SARIMA Modelleri ve BIC Değerleri

SARIMA Modelleri	BIC Değerleri
SARIMA(1,1,1) (1,1,1,4)	**881.929**
SARIMA(1,0,0) (1,0,0,4)	882.948

Tablo 5. Türkiye Hava Kalitesi İndeks SARIMA(1,1,1) (1,1,1,4) Modeli Parametre

Parametre	Kestirim	SE	t	p	R.M.S.E.
Sabit	290.623	62.234	4.669	0.001	
AR	0.425	0.115	3.696	0.008	20.056
MA	-0,994	0.142	-7.001	0.001	
sAR	0.310	0.126	2.461	0.032	
sMA	-0.994	0.250	-3.976	0.005	


Tablo 6. Türkiye Hava Kalitesi İndeks SARIMA(1,1,1) (1,1,1,4) Modeli Uygunluğu

İstatistik	p
4.208	0.936


3 katmanlı yapı (50'şer nöron), ADAM optimizer, 3 zaman adımı, batch size 32, epok 100 alınarak LSTM analizi yapılmıştır. LSTM için R.M.S.E. değeri, 18.820 olarak bulunmuştur. Şekil 4-5'te LSTM için MSE kayıp (hata) az olduğu ve tahmin değerlerinin orjinal seri grafiği ile uyumlu olduğu görülmektedir.

FULL TEXTS BOOK

Şekil 4. Türkiye İçin Hava Kalitesi LSTM Kayıp Grafiği



Şekil 5. Türkiye İçin Hava Kalitesi LSTM Tahmin Grafiği



FULL TEXTS BOOK

Türkiye hava kalitesi indeks değerleri için RMSE değerleri Tablo 7'de gösterilmiştir. Hava kalitesi indeksi için hesaplanan RMSE değerleri karşılaştırıldığında, LSTM modeli için RMSE değeri, ARIMA ve SARIMA için RMSE değerinden daha düşük olarak bulunmuştur. LSTM modeli ile tahmin yapmak daha doğru sonuçlar verecektir. Tahmin grafiği, Şekil 6'da verildiği gibidir.


Tablo 7. Türkiye Hava Kalitesi İndeksi ARIMA, SARIMA ve LSTM için RMSE

RMSE	**ARIMA**	**SARIMA**	**LSTM**
Hava Kalite İndeks	22.024	20.056	**18.820**

Tablo 8. Türkiye Hava Kalitesi İndeksi İleriye Yönelik Kestirimler

Dönem	**ARIMA**	**SARIMA**	**LSTM**
2026 Kış	79.540	72.910	76.657
2026 İlkbahar	79.829	78.301	80.767
2026 Yaz	79.849	79.925	81.594
2026 Sonbahar	79.851	79.952	81.974

Şekil 6. Türkiye İçin Hava Kalitesi Tüm Model Tahminler Grafiği



FULL TEXTS BOOK

SONUÇ

Bu çalışmada, 2000-2025 dönemine ait mevsimsel hava kalite indeksi verileri kullanılarak ARIMA(1,1,1), SARIMA(1,1,1)(1,1,1,4) ve LSTM modelleriyle 2026 yılına ait dört mevsimlik tahminler gerçekleştirilmiştir. Analiz sonuçlarına göre, modellerin tahmin performansları ve mevsimsel desenleri yakalama yetenekleri arasında önemli farklılıklar tespit edilmiştir. SARIMA modeli, mevsimsel bileşenleri açıkça modellemesi nedeniyle diğer modellere kıyasla daha tutarlı tahminler üretmiştir. LSTM modeli, veri setinin küçük hacimli olmasına ve karmaşık yapısına rağmen beklenenin üzerinde performans göstermiştir. ARIMA modeli, mevsimselliği açıkça modelleyememekle birlikte basit yapısıyla makul tahminler sağlamıştır. Hava kalitesi indeksi için hesaplanan RMSE değerleri karşılaştırıldığında, LSTM modeli için RMSE değeri, ARIMA ve SARIMA için RMSE değerinden daha düşük olarak bulunmuştur. LSTM modeli ile tahmin yapmak daha doğru sonuçlar vermektedir.

Öneriler kapsamında, öncelikle veri setine ek açıklayıcı değişkenlerin (meteorolojik faktörler, emisyon verileri gibi) dahil edilmesi model performansını artırabilir. SARIMA modelinin mevsimsel parametrelerinin grid search yöntemiyle optimize edilmesi, tahmin doğruluğunu iyileştirebilir. LSTM modeli için hiperparametre optimizasyonu ve daha derin mimariler denenebilir. Bu bulgular, hava kalitesi tahmininde derin öğrenme tabanlı yaklaşımların geleneksel zaman serisi yöntemlerine göre daha etkili sonuçlar verebileceğini ortaya koymaktadır. Özellikle, dinamik ve periyodik özellikler gösteren çevresel verilerin analizinde LSTM modellerinin kullanımı, erken uyarı sistemleri ve politika belirleme süreçlerine önemli katkı sağlayabilecektir.

KAYNAKÇA

Avnery, S., Mauzerall, D. L., Liu, J., & Horowitz, L. W. (2011). Global crop yield reductions due to surface ozone exposure: 1. Year 2000 crop production loses and economic damage. Atmospheric Environment, 45(13), 2284-2296. https://doi.org/10.1016/j.atmosenv.2010.11.045

Brook, R. D., Rajagopalan, S., Pope III, C. A., Brook, J. R., Bhatnagar, A., Diez-Roux, A. V., & Kaufman, J. D. (2010). Particulate matter air pollution and cardiovascular disease. Circulation, 121(21), 2331-2378. https://doi.org/10.1161/CIR.0b013e3181dbece1

Çevre, Şehircilik ve İklim Değişikliği Bakanlığı (ÇŞİDB). (2023). 2022 Türkiye Hava Kalitesi Raporu. https://www.csb.gov.tr

Gurjar, B. R., Butler, T. M., Lawrence, M. G., & Lelieveld, J. (2016). Evaluation of emissions and air quality in megacities. Atmospheric Environment, 42(7), 1593-1606. https://doi.org/10.1016/j.atmosenv.2007.10.048

Hochreiter, S., & Schmidhuber, J. (1997). Long short-term memory. Neural Computation, 9(8), 1735-1780. https://doi.org/10.1162/neco.1997.9.8.1735

Hyndman, R. J., & Athanasopoulos, G. (2018). Forecasting: Principles and practice (2nd ed.). OTexts. https://otexts.com/fpp2/

Hyndman, R. J., & Khandakar, Y. (2008). Automatic time series forecasting: The forecast package for R. Journal of Statistical Software, 27(3), 1-22. https://doi.org/10.18637/jss.v027.i03

FULL TEXTS BOOK

Sicard, P., De Marco, A., Agathokleous, E., Feng, Z., Xu, X., Paoletti, E., Rodriguez, J. J., & Calatayud, V. (2020). Amplified ozone pollution in cities during the COVID-19 lockdown. Science of The Total Environment, 735, 139542. https://doi.org/10.1016/j.scitotenv.2020.139542

World Health Organization (WHO). (2021). Air pollution. https://www.who.int/news-room/fact-sheets/detail/ambient-(outdoor)-air-quality-and-health

Zhang, G. P., Eddy Patuwo, B., & Hu, M. Y. (2020). Forecasting with artificial neural networks: The state of the art. International Journal of Forecasting, 14(1), 35-62. https://doi.org/10.1016/S0169-2070(97)00044-7

MAKİNE ÖĞRENMESİ YAKLAŞIMLARI İLE TÜRKİYE'DE KURAKLIK SÜREÇLERİNİN MODELLENMESİ

MACHINE LEARNING-BASED MODELING OF DROUGHT PROCESSES IN TÜRKİYE

Dr. Öğr. Üy. Barış ERGÜL [1]

[1] *Eskişehir Osmangazi Üniversitesi, Fen Fakültesi, İstatistik Bölümü, Eskişehir, Türkiye.*

[1]*ORCID ID: https://orcid.org/0000-0002-1811-5143*

Prof. Dr. Arzu ALTIN YAVUZ [2]

[2] *Eskişehir Osmangazi Üniversitesi, Fen Fakültesi, İstatistik Bölümü, Eskişehir, Türkiye.*

[2]*ORCID ID: https://orcid.org/0000-0002-3277-740X*

ÖZET

Kuraklık, meteorolojik dengesizliklerin uzun süreli etkisiyle ortaya çıkan, hidrolojik döngüde bozulmalara neden olan ve tarımsal üretimden su yönetimine kadar pek çok alanı etkileyen karmaşık bir çevresel felakettir. Türkiye gibi yarı kurak iklim kuşağında yer alan ülkelerde kuraklık, özellikle son yıllarda iklim değişikliğinin etkisiyle daha sık ve şiddetli şekilde görülmektedir.

Bu çalışmada, Türkiye'nin farklı iklim bölgelerini temsil eden uzun dönemli meteorolojik veriler (yağış, sıcaklık, bağıl nem, buharlaşma vb.) ve kuraklık analizi için hesaplanan SPI indeks değerleri kullanılmıştır. Bu göstergeler üzerinden kuraklık şiddetinin sınıflandırılması amacıyla makine öğrenmesi algoritmaları uygulanmıştır. Kullanılan algoritmalar arasında yer alan Random Forest, çok sayıda karar ağacının birleştirilmesiyle aşırı uyumu azaltırken yüksek öngörü performansı sağlar. Destek Vektör Makineleri (SVM), verileri doğrusal veya doğrusal olmayan düzlemlerde ayırarak sınıflandırma başarısını artırır. XGBoost, ardışık olarak eğitilen zayıf modellerin optimizasyonuyla tahmin hatasını minimize ederken; Multilayer Perceptron (MLP), çok katmanlı yapısı sayesinde karmaşık örüntüleri öğrenme kapasitesine sahiptir. Modeller, doğruluk ve F1 skoru gibi metriklerle değerlendirilmiş; özellikle XGBoost ve MLP algoritmalarının kuraklık sınıflarında yüksek sınıflandırma başarısı gösterdiği tespit edilmiştir.

Bu çalışma ile, makine öğrenmesi temelli modellerin Türkiye'de kuraklık erken uyarı sistemlerine entegre edilmesinde önemli bir potansiyel taşıdığını ortaya konulmaktadır. Sürdürülebilir iklim adaptasyonu stratejilerinin geliştirilmesine bilimsel altyapı sağlaması düşünülmektedir.

Anahtar Kelimeler: SVM, MLP, XGBoost, Rassal Orman, Türkiye.

ABSTRACT

Drought is a complex environmental disaster that emerges as a result of long-term meteorological imbalances, leading to disruptions in the hydrological cycle and adversely affecting various sectors, including agricultural production and water resource management. In semi-arid climate zones such as Turkey, drought events have become more frequent and severe in recent years, largely due to the impacts of climate change.

In this study, long-term meteorological data (precipitation, temperature, relative humidity, evaporation, etc.) representing different climatic regions of Turkey, along with Standardized Precipitation Index (SPI) values calculated for drought analysis, were utilized. In order to classify drought severity, several machine learning algorithms were implemented. Among the algorithms used, Random Forest reduces overfitting by aggregating multiple decision trees and provides high predictive performance. Support

FULL TEXTS BOOK

Vector Machines (SVM) improve classification accuracy by separating data in linear or nonlinear hyperplanes. XGBoost minimizes prediction error through sequential optimization of weak learners, while Multilayer Perceptron (MLP) is capable of learning complex patterns through its multilayered architecture. The models were evaluated using metrics such as accuracy and F1-score. Results indicated that XGBoost and MLP achieved high classification performance, particularly in drought categories.

This study demonstrates the strong potential of machine learning-based models in integrating drought forecasting into early warning systems in Turkey and is expected to provide a scientific foundation for the development of sustainable climate adaptation strategies.

Keywords: SVM, MLP, XGBoost, Random Forest, Türkiye.

GİRİŞ

Kuraklık, belirli bir bölgede yağış miktarının uzun süreli olarak iklim normallerinin altına düşmesiyle ortaya çıkan, doğal fakat etkileri itibarıyla derin sosyoekonomik ve ekolojik sonuçlar doğuran bir olgudur (Wilhite & Glantz, 1985). Diğer birçok doğal afetin aksine kuraklık, ani ve dramatik bir başlangıç göstermez; zamanla, sessizce ve kimi zaman fark edilmeden gelişir. Ancak bu yavaş ilerleyiş, etkilerinin hafife alınmasına yol açabilir.

Tarih boyunca toplumlar, kuraklık dönemlerinde kıtlık, göç ve sosyal huzursuzluk gibi önemli sorunlarla karşılaşmışlardır. Günümüzde ise iklim değişikliğinin etkileri, bu doğal süreci daha karmaşık, daha sık ve daha öngörülemez hale getirmiştir. Bu nedenle kuraklık, yalnızca çevresel değil, aynı zamanda ekonomik, politik ve toplumsal bir mesele olarak ele alınmalıdır.

Bilimsel araştırmalar, küresel ısınmanın kuraklık riskini hem sıklık hem de şiddet bakımından artırdığını ortaya koymaktadır (Trenberth et al., 2014). Atmosferin ısınmasıyla buharlaşma oranları artmakta, yağış rejimleri düzensiz hale gelmektedir ve su döngüsünde kırılmalar meydana gelmektedir. Bu durum özellikle yarı kurak ve kurak iklim kuşaklarında yaşayan toplumlar için kritik bir kırılganlık yaratmaktadır.

Türkiye gibi Akdeniz iklim kuşağında yer alan ülkelerde, bu süreç tarımın sürdürülebilirliğini tehdit etmekte ve su yönetiminde yeni zorluklar doğurmaktadır. Yağışların düzensizleşmesi, baraj ve yer altı su seviyelerinin öngörülebilirliğini azaltırken, kurak dönemlerin uzunluğu eko-sistemlerin kendini toparlama kapasitesini zayıflatmaktadır. İklim değişikliğinin etkileri yalnızca doğal sistemlerle sınırlı değildir. Tarımsal üretimdeki azalmalar, gıda fiyatlarının yükselmesine, geçim kaynaklarının daralmasına ve kırsaldan kente göçün hızlanmasına neden olmaktadır. Böylece ekolojik bir sorun, ekonomik ve toplumsal bir krize dönüşmektedir. Kuraklık, suyun yalnızca fiziksel bir kaynak değil, aynı zamanda sosyal ve kültürel bir yaşam temeli olduğunu hatırlatır. Gündelik hayatta musluktan akan suyun azalması, şehirlerde su kısıtlamaları, kırsalda ise tarlaların kurumaya başlaması, bireyler için kuraklığın somut yüzünü oluşturur. Ancak mesele bundan ibaret değildir; suyun azalması, sağlık sistemlerinden gıda güvenliğine kadar uzanan zincirleme etkiler yaratır. Eko-sistemler açısından ise kuraklık, yaşam ağının bütününde bozulmalara yol açar. Nehirlerin debisinin düşmesi, göllerin çekilmesi ve toprak neminin azalması, bitki örtüsünden yaban hayatına kadar tüm doğal sistemleri etkiler. Orman yangınlarının daha sık ve yıkıcı hale gelmesi, biyolojik çeşitliliğin azalması ve ekosistem hizmetlerinin zayıflaması bu sürecin kaçınılmaz sonuçlarıdır (Vicente-Serrano et al., 2020).

Kuraklığın etkilerini azaltmak, yalnızca afet sonrası kriz yönetimi ile mümkün değildir. Bütüncül yaklaşımlar; erken uyarı sistemleri, iklim tahminleri ve entegre su yönetimi planları, uzun vadeli dayanıklılık açısından kritik öneme sahiptir. Tarımda su verimliliğini artıracak modern sulama teknikleri, kuraklığa dayanıklı bitki türlerinin yetiştirilmesi ve suyun geri kazanımı gibi teknolojik çözümler, bu mücadelede temel araçlardır. Ancak teknik çözümler tek başına yeterli değildir. Toplumsal

FULL TEXTS BOOK

farkındalık, kuraklıkla mücadelede vazgeçilmez bir unsurdur. Suyun kültürel, ekonomik ve ekolojik değerinin toplumsal hafızada yer etmesi gerekir. Yerel yönetimlerin su tasarrufu kampanyaları, iklim uyum politikaları ve sivil toplum kuruluşlarının sahadaki çalışmaları, birbirini tamamlayan bir koordinasyon içinde yürütülmelidir.

Kuraklık, çoğu zaman gözden kaçan ancak etkileri itibarıyla yıkıcı olabilen bir krizdir. İklim değişikliğinin bu krizi daha sık ve yoğun hale getirdiği günümüzde, hem eko-sistemlerin hem de insan toplumlarının uyum kapasitesini artırmak bir zorunluluk haline gelmiştir. Bu, yalnızca çevresel bir mesele değil; ekonomik istikrar, toplumsal refah ve gelecek nesillerin yaşam hakkı açısından da kritik bir meseledir. İnsanlığın bu sessiz krizle başa çıkma kapasitesi, bilimin rehberliği, politikanın kararlılığı ve toplumun ortak çabasıyla şekillenecektir. Bu çalışmada, kuraklık konusuna veriler ile çeşitli öneriler getirilmeye çalışılacaktır.

YÖNTEM

Makine öğrenmesi, günümüzde veri analitiği ve yapay zeka çalışmalarının temel yapı taşlarından biri olarak konumlanmıştır. Büyük veri setlerinden anlamlı örüntüler çıkarmak, karar destek sistemlerini güçlendirmek ve öngörü yeteneğini artırmak amacıyla kullanılan algoritmalar arasında Random Forest, Support Vector Machine (SVM), Extreme Gradient Boosting (XGBoost) ve Multi-Layer Perceptron (MLP) öne çıkmaktadır. Her biri farklı teorik temellere dayansa da, pratikte karmaşık problemlerin çözümünde yüksek başarı oranları ile tercih edilmektedir.

Random Forest, Leo Breiman tarafından 2001 yılında tanıtılan, karar ağaçlarının topluluk (ensemble) yöntemiyle bir araya getirilmesiyle oluşturulmuş bir algoritmadır (Breiman, 2001). Temel prensip, her ağacın eğitim sırasında farklı rastgele alt veri setleri ve özellikler kullanmasıdır. Bu sayede modelin aşırı öğrenme (overfitting) riskini azaltması ve genelleme kabiliyetini artırması sağlanır. Random Forest, yüksek boyutlu veri kümelerinde ve karmaşık sınıflandırma problemlerinde dengeli bir performans sergiler. Ancak çok büyük veri setlerinde eğitim süresinin uzaması ve yorumlanabilirliğinin sınırlı olması, dikkat edilmesi gereken hususlardandır.

SVM, Vladimir Vapnik ve arkadaşları tarafından geliştirilen ve özellikle küçük ile orta ölçekli veri kümelerinde yüksek doğruluk sağlayabilen güçlü bir sınıflandırma algoritmasıdır (Cortes & Vapnik, 1995). Temel yaklaşımı, veriyi yüksek boyutlu bir uzaya dönüştürerek sınıflar arasındaki ayrım mesafesini maksimize eden hiper-düzlemleri bulmaktır. Kernel fonksiyonları sayesinde doğrusal olmayan karar sınırları da oluşturabilir. SVM, özellikle karmaşık ama gürültüsüz veri setlerinde başarılıdır; ancak büyük veri setlerinde hesaplama maliyeti artar ve parametre seçimi (C, gamma) hassasiyet gerektirir.

XGBoost, Chen ve Guestrin (2016) tarafından geliştirilen, hata fonksiyonu için artırmalı ağaçlar (gradient boosted trees) algoritmasının optimize edilmiş bir versiyonudur. Hem hız hem de performans açısından endüstri standartları arasında kabul edilmektedir. Regularization (Cezalandırma) teknikleriyle aşırı öğrenme riskini minimize ederken, paralel işlem yeteneği ile büyük veri kümelerinde hızlı sonuç verir. XGBoost, yapısal verilerde yüksek doğruluk sağlamaktadır. Ancak parametre optimizasyonu zahmetli olabilir.

MLP, yapay sinir ağlarının en temel biçimlerinden biridir ve en az bir gizli katmana sahip ileri beslemeli yapıda çalışır (Rumelhart et al., 1986). Her nöron, aktivasyon fonksiyonları aracılığıyla girdi verisini doğrusal olmayan biçimlerde dönüştürerek öğrenme kapasitesini artırır. MLP, özellikle doğrusal olarak ayrıştırılamayan problemler için uygundur. Bununla birlikte, büyük veri setlerinde başarılı olabilmesi için yeterli sayıda katman ve nöron ile birlikte uygun optimizasyon algoritmalarına ihtiyaç duyar. Eğitim süresi uzun olabilir ve hiperparametre ayarları model performansında kritik rol oynar.

Bu dört algoritma farklı güçlü yönlere sahiptir. Random Forest, dengeli ve aşırı öğrenmeye dirençli yapısıyla güvenilir bir seçenektir. SVM, küçük ve orta ölçekli veri kümelerinde yüksek doğruluk sağlar.

FULL TEXTS BOOK

XGBoost, hız ve doğruluk dengesinde öne çıkar; özellikle yapısal verilerde üstün performans gösterir. MLP ise esnek yapısı sayesinde çok çeşitli veri tiplerinde uygulanabilir, ancak dikkatli ayarlanmazsa kolayca aşırı öğrenme eğiliminde olabilir. Algoritma seçimi, veri kümesinin büyüklüğü, özelliklerin türü, modelin yorumlanabilirlik gereklilikleri ve işlem kaynaklarının durumu gibi faktörlere bağlıdır. Dolayısıyla, tek bir "en iyi" algoritmadan ziyade, problem bağlamına en uygun yöntemi belirlemek kritik öneme sahiptir.

Makine öğrenmesinde Random Forest, SVM, XGBoost ve MLP, sınıflandırma ve regresyon problemlerinde yaygın olarak kullanılan, kanıtlanmış başarılara sahip algoritmalardır. Her birinin avantajları ve sınırlılıkları vardır; bu nedenle veri bilimi uygulamalarında genellikle birden fazla algoritmanın karşılaştırmalı olarak denenmesi önerilir.

BULGULAR

1984-Sonbahar 2023 tarihleri arasında Türkiye için kuraklık ölçüm değerleri Meteoroloji Genel Müdürlüğü web sayfalarından ilgili bölümlerden alınmıştır. Bağımlı değişken olarak, kuraklık (0: kurak, 1: nemli) alınmış ve kuraklığı etkilediği düşünülen bağımsız değişkenler de, ortalama sıcaklık, ortalama rüzgar, ortalama nem, yağış ve ortalama buharlaşma değişkenleri alınmıştır.

İstatistik Analizler, R ve Python programları kullanılarak yapılmıştır. Veri ön işleme işlemleri yapıldıktan sonra, veri seti %70 eğitim, %30 test olarak ayrılmıştır. İlk aşamada Random Forest algoritması uygulandıktan sonra sırasıyla, SVM, XGBoost ve MLP algoritmaları uygulanarak analizlere devam edilmiştir. Bulunan sonuçlar Tablo 1-2'de özetlenmiştir. Tablo 1-2'ye göre, Random Forest için eğitim veri setinde %100 doğruluk (aşırı öğrenme belirtisi), test verisinde ise, %81.670 olduğu görülmektedir. Bu durum, overfit (aşırı öğrenme) durumu olduğunun göstergesidir. XGBoost için eğitim (%95.240) ve test (%85.000) arasındaki fark dengeli olduğundan, daha dengeli bir durum olduğunu belirtmektedir. MLP/SVM için eğitim-test performansı birbirine yakın olduğu belirlenmiştir. XGBoost, en iyi performansı gösteren algoritma olmuştur. Hem doğruluk (%85.000) hem de F1 skoru (0.8235) açısından diğer algoritmaları geride bırakmıştır. MLP (Çok Katmanlı Algılayıcı) algoritması, %83.330 doğruluk ve 0.8000 F1 skoru ile XGBoost'tan biraz daha düşük performans göstermiştir. Random Forest (Rassal Orman) algoritması, MLP'den biraz daha düşük performans göstermektedir (%81.670 doğruluk ve 0.7826 F1 skoru), ancak yine de oldukça iyi sonuçlar vermiştir. SVM (Destek Vektör Makineleri) modeli ise %80.000 doğruluk ve 0.7619 F1 skoru ile en düşük performansı göstermiştir.

Tablo 1. Kuraklık (SPI) Test Verileri için Algoritmalar ile Elde Edilen Doğruluk Değerleri

Algoritma	Veri	Doğruluk (%)	F Ölçüsü (F1 Skor)
RF	Test	81.670	0.7826
SVM	Test	80.000	0.7619
XGBoost	Test	**85.000**	**0.8235**
MLP	Test	**83.330**	**0.8000**

Tablo 2. Kuraklık (SPI) Eğitim Verileri için Algoritmalar ile Elde Edilen Doğruluk Değerleri


Algoritma	Veri	Doğruluk (%)	F Ölçüsü (F1 Skor)
RF	Eğitim	99.998	0.9999

FULL TEXTS BOOK

SVM	Eğitim	83.330	0.8000
XGBoost	Eğitim	**95.240**	**0.9474**
MLP	Eğitim	**85.710**	**0.8333**

Bu algoritmalar kullanılarak, SPI etkileyen en önemli değişkenler Şekil 1'de belirtildiği gibidir. Şekil 1'e göre, yapılan önemlilik analizlerinde, yağış ve buharlaşma değişkenlerinin SPI durumu üzerinde tutarlı şekilde en etkili değişkenler olduğu görülmüştür. MLP algoritması, buharlaşma değişkenine diğer algoritmalara kıyasla daha yüksek önem atamıştır.

Şekil 1. Kuraklık (SPI) Verileri için Algoritmalar ile Elde Edilen En Önemli Değişkenler



SONUÇ

Bu çalışmada, Türkiye'nin 1984-2023 yılları arasındaki mevsimsel iklim verileri kullanılarak, Standartlaştırılmış Yağış İndeksi (SPI) durumunun (kurak/nemli) tahmin edilmesi amacıyla dört farklı makine öğrenmesi algoritması (XGBoost, Rassal Orman, SVM ve MLP) karşılaştırılmıştır. Analizler, yağış, buharlaşma, nem, sıcaklık ve rüzgar hızı değişkenlerinin SPI üzerindeki etkisini ortaya koyarken, modellerin performansları ve değişken önem sıralamaları detaylı şekilde incelenmiştir.

XGBoost, test verisi üzerinde %85 doğruluk ve 0.82 F1 skoru ile en iyi performansı gösteren model olmuştur. SVM ve MLP'nin ise daha düşük performans sergilediği görülmüş ve özellikle MLP'nin eğitim süresinin uzun olması pratik dezavantaj yaratmıştır. Random Forest (Rassal Orman)'ın da benzer sonuçlar vermesi, ağaç tabanlı modellerin iklim verilerinde etkin olduğunu doğrulamaktadır.

Tüm modellerde yıllık yağış miktarı ve buharlaşma, SPI tahmini için en kritik iki değişken olarak öne çıkmıştır. Bu bulgu, meteorolojik kuraklık göstergelerinin SPI ile güçlü ilişkisini desteklemektedir. Nem oranı, XGBoost ve SVM'de üçüncü sırada yer alırken, Random Forest (Rassal Orman)'da sıcaklık daha önemli bulunmuştur. MLP ise buharlaşmayı birinci sıraya koyarak diğer modellerden farklılaşmıştır.

FULL TEXTS BOOK

Rüzgar hızı, tüm modellerde en az etkili değişken olarak belirlenmiş, bu da SPI tahmini için çıkarılabilir bir parametre olduğunu düşündürmüştür.

XGBoost, yüksek doğruluk ve yorumlanabilirlik dengesi nedeniyle tercih edilmesi gereken bir algoritmadır. Hiperparametre optimizasyonu ile performansı daha da artırılabilir. Eğer model karmaşıklığından kaçınılmak isteniyorsa, Random Forest (Rassal Orman) basit ve etkili bir alternatiftir. SVM ve MLP, özellikle lineer olmayan ilişkileri yakalama potansiyellerine rağmen, bu veri setinde sınırlı kalmıştır. Daha büyük veri setlerinde denenebileceği öngörülmektedir.

Toprak nemi veya bitki örtüsü indeksleri gibi ek değişkenlerin modele dahil edilmesi, tahmin doğruluğunu artırabileceği değerlendirilmelidir.

Sonuç olarak, bu çalışma, Türkiye'nin iklim verileri kullanılarak kuraklık (SPI) tahmini için XGBoost ve MLP algoritmalarının etkinliğini kanıtlamış, kritik meteorolojik parametreleri ortaya koymuştur. Uygulamada, bu algoritmaların erken uyarı sistemlerine entegre edilmesi, su kaynakları yönetimi ve tarımsal planlama için değerli bir araç olabileceği göz önüne alınmalıdır.

KAYNAKÇA

Breiman, L. (2001). Random forests. Machine Learning, 45(1), 5-32. https://doi.org/10.1023/A:1010933404324

Chen, T., & Guestrin, C. (2016). XGBoost: A scalable tree boosting system. Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, 785-794. https://doi.org/10.1145/2939672.2939785

Cortes, C., & Vapnik, V. (1995). Support-vector networks. Machine Learning, 20(3), 273-297. https://doi.org/10.1007/BF00994018

Dai, A. (2011). Drought under global warming: A review. Wiley Interdisciplinary Reviews: Climate Change, 2(1), 45-65. https://doi.org/10.1002/wcc.81

Rumelhart, D. E., Hinton, G. E., & Williams, R. J. (1986). Learning representations by back-propagating errors. Nature, 323(6088), 533-536. https://doi.org/10.1038/323533a0

Trenberth, K. E., Dai, A., van der Schrier, G., Jones, P. D., Barichivich, J., Briffa, K. R., & Sheffield, J. (2014). Global warming and changes in drought. Nature Climate Change, 4(1), 17-22. https://doi.org/10.1038/nclimate2067

Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscalar drought index sensitive to global warming: The standardized precipitation evapotranspiration index. Journal of Climate, 23(7), 1696-1718. https://doi.org/10.1175/2009JCLI2909.1

Wilhite, D. A., & Glantz, M. H. (1985). Understanding the drought phenomenon: The role of definitions. Water International, 10(3), 111-120. https://doi.org/10.1080/02508068508686328

FULL TEXTS BOOK

SMART ENERGY SYSTEMS FOR SUSTAINABILITY-A REVIEW

E. CUCE[a]

[a] *Department of Mechanical Engineering, Faculty of Engineering and Architecture, Recep Tayyip Erdogan University, Zihni Derin Campus, 53100 - Rize, Turkiye*

ORCID: 0000-0003-0150-4705

P. MERT CUCE[b, c]

[b] *Department of Architecture, Faculty of Engineering and Architecture, Recep Tayyip Erdogan University, Zihni Derin Campus, 53100 - Rize, Turkiye*

[c] *College of Built Environment, Birmingham City University, Birmingham B4 BD, UK*

ORCID:0000-0002-6522-7092

T. GUCLU[d]

[d] *Department of Mechanical Engineering, Faculty of Engineering, University of Bayburt, Dede Korkut Campus, 69000 - Bayburt, Turkiye*

ORCID: 0000-0002-5864-3864

ABSTRACT

Today, the share of energy obtained from renewable sources in total energy production is steadily increasing. However, fossil fuels still account for a significant portion of global energy generation. Reducing reliance on fossil fuels is critical to mitigating their harmful environmental impacts. In this context, the efficient use of existing energy resources and the transition to smart and sustainable energy systems have become key research areas. This study aims to examine smart energy systems that utilise renewable energy sources such as solar, geothermal, and wind—either individually or in hybrid configurations—as alternatives to conventional fossil-based systems. A literature-based qualitative research method was used to evaluate the structure, operation, and components of smart energy systems. The study discusses their potential to decentralise energy production, promote consumer participation as prosumers, and reduce both energy costs and greenhouse gas emissions. In addition, the advantages and disadvantages of smart energy systems are analysed from technical and economic perspectives. The results highlight the need for supportive policies, investment incentives, and advanced control strategies to accelerate the adoption of smart energy systems. Findings from recent studies in the field are also referenced to support the evaluations.

Keywords: smart energy systems, smart thermal grid, smart grid, energy efficiency, sustainability

5. INTRODUCTION

Energy is central not only to daily life but also to global responsibilities such as combating climate change, sustainable development, protecting the environment, and preserving biodiversity. However, fossil fuels still meet the majority of global energy needs. The greenhouse gas emissions caused by the use of these resources pose a serious threat to climate change and environmental degradation. Furthermore, increasing energy demand and the depletion of fossil fuel reserves necessitate the development of new and renewable solutions [1]. In recent years, many countries' energy policies have been shifting to encourage the use of clean, renewable, and sustainable energy sources. With the Paris Climate Agreement, more than 190 countries have targeted keeping global temperature rise below 2°C, and the European Union, in particular, has accelerated the transition to renewable sources to ensure

FULL TEXTS BOOK

energy security and minimise greenhouse gas emissions [2]. In the energy sector, which accounts for a significant share of global carbon emissions, the transition from fossil fuels to clean energy is critical from both environmental and economic perspectives. System designs that balance energy storage, generation, and transmission processes and manage supply-demand fluctuations are paramount in this transformation. In this context, the concept of "smart and sustainable energy systems" refers to the efficient and automated use of environmentally friendly and renewable resources through the integration of different energy sectors [3].

Smart energy systems integrate areas such as electricity, heating, transportation, and energy storage, offering both environmentally friendly and economically viable solutions. These systems allow energy production to be generated from small-scale renewable resources closer to the point of consumption, rather than from large centralised facilities [4]. This reduces transmission losses, empowers consumers to become producers, lowering energy costs and reducing greenhouse gas emissions. Expected to be at the heart of the energy transformation that will take place in the near future, smart and sustainable energy systems offer significant opportunities from both technological and societal perspectives [5].

6. SMART ENERGY SYSTEMS

The 21st century's growing energy demand, environmental pollution, climate change, and fossil fuel dependency demonstrate the inadequacy of traditional energy systems. Therefore, structures focused on a single energy sector are being replaced by smart energy systems that integrate different energy types and sectors. Smart energy systems are structures that optimise energy production and consumption through intelligent control mechanisms, integrating electricity, heating, cooling, transportation, and storage. These systems enable more efficient use of renewable energy sources (solar, wind, geothermal, etc.) and minimise environmental impacts by reducing fossil fuel consumption. They also allow consumers to become prosumers [6].

The most important goals of such systems are:

- Sustainability: Efficient and long-term use of resources,
- Environmentally Friendly Production: Reducing greenhouse gas emissions,
- Economic Efficiency: Reducing energy costs,
- Integration: Interoperability of different energy sources and sectors.

Compared to traditional systems, smart and sustainable energy systems, thanks to their flexible structures, allow for the conversion between different energy types. For example, using waste heat generated in industrial facilities for district heating systems or storing excess electricity generated from renewable sources for use during peak demand periods are examples of this approach. Consequently, smart and sustainable energy systems are seen as one of the most important energy infrastructures of the future for ensuring energy security, reducing costs, and minimising environmental impacts.

6.1. SMART THERMAL ENERGY SYSTEMS

Smart thermal energy systems, unlike traditional heating and cooling infrastructures, are flexible, bi-directional energy flow systems that can effectively utilize renewable energy sources and waste heat. These systems not only transfer heat from producer to consumer, but also enable the consumer to become a producer when necessary [7]. This approach utilizes renewable sources such as solar, wind, and geothermal energy, or waste heat from industrial facilities. For example, solar hot water systems can be integrated into central heating networks, or heat pumps can be used for both heating and cooling. This increases energy efficiency while reducing the environmental impact of fossil fuel use [8]. Heat pumps are a key component of smart thermal systems. When integrated with renewable energy sources,

FULL TEXTS BOOK

they reduce energy costs and increase system flexibility. Furthermore, thermal storage units allow the heat generated to be stored until needed. This offers significant advantages, particularly in district heating and cooling applications [9].

The main benefits of smart thermal energy systems include:

- Increasing energy efficiency
- Recovering waste heat
- Expanding the use of renewable energy
- Reducing heating and cooling costs
- Minimising environmental impact

In conclusion, smart thermal energy systems offer both economically and environmentally sustainable solutions by providing flexibility in energy production, storage, and use. They play a significant role in energy transformation, especially when integrated into city-wide central heating and cooling systems.

6.2. SMART ELECTRICAL ENERGY SYSTEMS

In traditional electricity generation, electricity is generated through large-scale fossil fuel-fired power plants and delivered to consumers via long transmission lines. This method not only causes energy losses during transmission but also harms the environment due to fossil fuel consumption. Today, this centralised structure is being replaced by distributed generation systems consisting of small-scale production facilities located close to consumption points [10]. Smart electric power systems, often referred to as "smart grids," integrate renewable energy sources such as wind, solar, and hydropower into the electricity generation process. In these systems, the consumer becomes a "prosumer" who not only consumes energy but also produces it. For example, photovoltaic panels installed on a home's roof can meet the home's electricity needs, while any excess electricity can be sold to the grid [11]. Smart electrical systems not only manage the energy flow between production and consumption, but they also offer automation and control mechanisms that save energy. In smart buildings, lighting can automatically turn off in unoccupied rooms, and heating and cooling systems can operate only when needed. This reduces energy costs and increases efficiency.

The key advantages offered by these systems include:

- Reducing transmission losses (generating energy closer to the point of consumption)
- Increasing the share of renewable energy
- Transitioning from consumer to producer
- Reducing energy costs
- Minimising environmental impact

Furthermore, with the widespread adoption of electric and hybrid vehicles, users can use electricity generated from their own renewable sources to charge their vehicles. This reduces fuel costs and reduces dependence on fossil fuels. In conclusion, smart electrical energy systems play a key role in the future energy transition by establishing flexible, reliable, and environmentally friendly energy infrastructures.

6.3. SMART GAS ENERGY SYSTEMS

Smart gas energy systems make traditional gas infrastructures more efficient and flexible by utilising alternative fuels such as biofuels and synthetic gases (syngas) in proportion to fossil gases. These systems are particularly important for distributed energy production and integration with renewable resources [12]. Unlike traditional gas systems, smart gas systems can more quickly manage the energy supply-demand balance and optimise production and consumption processes. Furthermore, produced

FULL TEXTS BOOK

gas or energy can be stored either within the grid or in dedicated storage units. This allows energy to be supplied from these storage facilities during times of increased demand, improving system reliability.

The main advantages of smart gas energy systems are:

- Increasing energy efficiency
- Ensuring supply security through fuel diversity
- Optimising the production-consumption balance
- Providing flexibility through storage
- Reducing environmental impact by reducing fossil fuel use

These systems, particularly when used in conjunction with combined heat and power (CHP) plants, high-efficiency heat pumps, and renewable energy-supported gas production technologies, offer both economical and environmentally friendly solutions. Furthermore, supporting the gas infrastructure with smart sensors, automation systems, and data analysis tools allows for continuous monitoring and improvement of system performance. In conclusion, smart gas energy systems play a complementary role in the energy transition process by providing flexibility, efficiency, and sustainability across the entire chain from production to consumption.

7. ECONOMIC FISIBILITY OF SMART ENERGY SYSTEMS

Replacing traditional energy systems with smart and sustainable systems requires high initial investment costs. The construction of new infrastructure, the integration of renewable energy technologies, and the addition of smart control systems are the primary drivers of this cost. However, in the long run, these investments will pay off economically through more efficient use of energy resources. Furthermore, reducing fossil fuel use prevents environmental damage, reducing government budgets and environmental taxes allocated to combat environmental problems [13]. Cost and efficiency analyses for smart energy systems vary depending on the type of system installed, its capacity, and the application area. These analyses consider the following factors:

- Regional electricity unit prices
- Government tax and incentive policies
- Local energy demand and supply conditions
- The potential of renewable energy resources
- Therefore, a system that is economically advantageous in one region may not yield the same results in another.

Therefore, using a standard method for cost analysis of smart energy systems is not appropriate. While it may require high short-term investment, this transformation becomes an economically sound option in the long term thanks to the energy savings, environmental benefits, and reduced operating costs.

8. CONCLUSION

Energy systems play a critical role in today's globalising and industrialising world, from an environmental, economic, and social perspective. The efficient use of existing energy resources, the protection of natural resources, and the assurance of energy security for future generations are fundamental requirements for a sustainable future. Smart and sustainable energy systems contribute to preventing the environmental impacts caused by fossil fuels by integrating renewable energy sources into the production process. These systems support distributed generation models close to the point of consumption, rather than centralised, large-scale energy production facilities, and enable consumer

FULL TEXTS BOOK

participation in the production process. While this transformation may require high initial investment costs, it is an economically sound choice in the long term, thanks to its energy efficiency, low operating costs, and environmental benefits. In conclusion, smart and sustainable energy systems are poised to play a central role in the energy infrastructure of the future, offering solutions that are both environmentally friendly and economically viable.

REFERENCES

[1] Zamfirescu, C., I. Dincer, M. Stern, and W. R. Wagar. 2012. "Exergetic, Environmental and Economic Analyses of Small-Capacity Concentrated Solar-Driven Heat Engines for Power and Heat Cogeneration." *International Journal of Energy Research* 36(3): 397–408. https://onlinelibrary.wiley.com/doi/10.1002/er.1811.

[2] Bačeković, Ivan, and Poul Alberg Østergaard. 2018. "Local Smart Energy Systems and Cross-System Integration." *Energy* 151(2018): 812–25.

[3] Zhao, Ning et al. 2023. "Emerging Information and Communication Technologies for Smart Energy Systems and Renewable Transition." *Advances in Applied Energy* 9(October 2022): 100125. https://doi.org/10.1016/j.adapen.2023.100125.

[4] Lund, Henrik et al. 2021. "EnergyPLAN – Advanced Analysis of Smart Energy Systems." *Smart Energy* 1: 100007. https://doi.org/10.1016/j.segy.2021.100007.

[5] Shakeri, Mohammad et al. 2017. "An Intelligent System Architecture in Home Energy Management Systems (HEMS) for Efficient Demand Response in Smart Grid." *Energy and Buildings* 138: 154–64. http://dx.doi.org/10.1016/j.enbuild.2016.12.026.

[6] Cuce, Erdem, Pinar Mert Cuce, Tamer Guclu, and Ahmet Burhaneddin Besir. 2020. "On the Use of Nanofluids in Solar Energy Applications." *Journal of Thermal Science* 29(3): 513–34.

[7] Yang, Libing, Evgueniy Entchev, Antonio Rosato, and Sergio Sibilio. 2017. "Smart Thermal Grid with Integration of Distributed and Centralized Solar Energy Systems." *Energy* 122: 471–81. http://dx.doi.org/10.1016/j.energy.2017.01.114.

[8] Hennessy, Jay, Hailong Li, Fredrik Wallin, and Eva Thorin. 2019. "Flexibility in Thermal Grids: A Review of Short-Term Storage in District Heating Distribution Networks." *Energy Procedia* 158: 2430–34. https://doi.org/10.1016/j.egypro.2019.01.302.

[9] Lund, Henrik et al. 2014. "4th Generation District Heating (4GDH). Integrating Smart Thermal Grids into Future Sustainable Energy Systems." *Energy* 68: 1–11. http://dx.doi.org/10.1016/j.energy.2014.02.089.

FULL TEXTS BOOK

[10] Colson, C. M., and M. H. Nehrir. 2009. "A Review of Challenges to Real-Time Power Management of Microgrids." *2009 IEEE Power and Energy Society General Meeting, PES '09*.

[11] Ustun, Taha Selim, Cagil Ozansoy, and Aladin Zayegh. 2011. "Recent Developments in Microgrids and Example Cases around the World - A Review." *Renewable and Sustainable Energy Reviews* 15(8): 4030–41. http://dx.doi.org/10.1016/j.rser.2011.07.033.

[12] Crisostomi, Emanuele, Marco Raugi, Alessandro Franco, and Giuseppe Giunta. 2013. "The Smart Gas Grid: State of the Art and Perspectives." In *IEEE PES ISGT Europe 2013*, IEEE, 1–5.

[13] Moretti, M. et al. 2017. "A Systematic Review of Environmental and Economic Impacts of Smart Grids." *Renewable and Sustainable Energy Reviews* 68: 888–98. http://dx.doi.org/10.1016/j.rser.2016.03.039.

SOSYAL BİLGİLER DERSİNDE SİNEKTİK TEKNİĞİNİN KULLANIMI*

***Ayça ÖZTÜRK*[1]**

Doktora Öğrencisi, Marmara Üniversitesi, Eğitim Bilimleri Enstitüsü, Sosyal Bilgiler Öğretmenliği, İstanbul, Türkiye

[2]ORCID ID:0000-0001-9184-6842

***Ahmet KATILMIŞ*[2]**

Prof.Dr., Marmara Üniversitesi, Eğitim Bilimleri Enstitüsü, Sosyal Bilgiler Öğretmenliği, İstanbul, Türkiye

[1]ORCID ID: https://orcid.org/0000-0002-5776-850X

ÖZET

Yaratıcı düşünme teknikleri arasında ele alınan sinektik, farklı disiplinler arasındaki bağlantıları kurarak öğrencilere yaratıcı düşünme ve problem çözme becerilerini geliştirme fırsatı sunan bir öğretim tekniğidir. Bu teknik, daha önceden aralarında bir bağlantı saptanmamış iki parçanın ortak bir süzgeçten geçirilerek öğrenenler için farklı düşünme ve problem çözme kapasitelerini artırma imkanı tanımaktadır. Sinektik tekniği empatiyi, katılımı, işbirliğini, spekülasyonu içeren ve nesneleri derinlemesine bakış açısıyla inceleme fırsatı sunan çok katmanlı bir yapıdır. Öte yandan belli başlı becerileri öğrencilere kapsayıcı ve birçok disiplini içeren bir sorun üzerinden kazandırmak mümkündür. Böylelikle sorunlara karşı geniş bağlamda bir ön hazırlık gerçekleştirilmiş olur. İnsanın zihinsel esnekliğini geliştirmesi ve özgün çözümler üretebilmesi için derinlemesine analiz yapabilme ve eleştirel düşünme becerisinin güçlü olması gerekir. Sinektik teknikte, hayal gücünü öne çıkaran ve farklı bakış açıları geliştirmeye olanak tanıyan ıraksak ve yakınsak düşünme tarzlarını içeren süreç hakimdir. Bu düşünme tarzlarının kullanılması üst düzey düşünme yetilerinin gelişimi için oldukça önemlidir. Bu çalışmanın amacı, sinektik tekniği temelli etkinliklerin sosyal bilgiler dersinde nasıl kullanılabileceğini ortaya koymaktadır. Çalışma eylem araştırması olarak tasarlanmış ve gerçekleştirilmiştir. Araştırmanın uygulaması 2024/2025 eğitim öğretim yılı bahar döneminde İstanbul ili Pendik ilçesine bağlı özel bir ortaokul bünyesinde öğrenim gören 5. sınıf düzeyindeki 21 öğrenci ile gerçekleştirilmiştir. Uygulama sürecinden elde edilen veriler bulgular kısmında verilmiştir.

Anahtar Kelimeler: Sinektik, Yaratıcı düşünme, Yenilikçi düşünme, Sosyal bilgiler

ABSTRACT

Synectics, a form of creative thinking, is a teaching technique that offers students the opportunity to develop creative thinking and problem-solving skills by establishing connections between different disciplines. This technique allows learners to increase their divergent thinking and problem-solving capacities by filtering two previously unconnected components through a common filter. The synectics technique allows learners to enhance their creative thinking and problem-solving abilities by bringing together two elements that have not previously been associated and examining them through a common filter. It is possible to teach students specific skills through a comprehensive and multidisciplinary problem. This provides a preliminary preparation for addressing problems within a broader context. To develop mental flexibility and generate original solutions, individuals must possess strong in-depth

* Bu çalışma birinci yazarın, Marmara Üniversitesi Eğitim Bilimleri Enstitüsünde sürdürdüğü Doktora tez çalışmasından üretilmiştir.

FULL TEXTS BOOK

analysis and critical thinking skills. The synectics technique is dominated by a process involving divergent and convergent thinking styles, which foster imagination and allow for the development of different perspectives. The use of these thinking styles is crucial for the development of higher-order thinking skills. The aim of this study is to reveal how activities based on the synectics technique can be implemented in social studies courses. The research was designed and conducted as an action research study. The implementation was carried out with 21 fifth-grade students enrolled in a private middle school located in the Pendik district of Istanbul during the spring semester of the 2024–2025 academic year. The data obtained from the implementation process are presented in the findings section.

Keywords: Synectics, Creative thinking, Innovative thinking, Social Studies.

GİRİŞ

Eğitimin daha verimli hale getirilmesi amacıyla öğretim modelleri, yöntemleri ve tekniklerinin çoğalması, yenilikçi adımların getirilmesi gerekmektedir (Doğan, 2020). Sosyal bilgiler, geçmişle geleceğin, yerelin ve evrenselin, bireyin ve toplumun iç içe geçtiği bir ders programı olarak görülebilir. Dünyada her ülke, inşa etmek istediği gelecek doğrultusunda bilgi, beceri ve değerler donanımını oluşturur. Bu donanım zamanın ihtiyaçlarına göre değişime uğrar. Sosyal bilgiler dersi bu bilgi, beceri ve değerleri benimseyen vatandaşlar inşa etmeyi sağlamaktadır (Yiğit, 2022) . Sosyal bilgiler dersi toplumsal konulara duyarlılık gösteren, sorunların çözüm sürecinde bilimsel ahlakı gözeten, milli kültür mirasımızın korunması ve geliştirilmesi için sorumluluk alan bireyler yetiştirilmesini desteklemektedir. Bu doğrultuda öğrencilerin değişim ve sürekliliği algılamaları, mekânı algılama becerilerini geliştirmeleri, bilgi ve iletişim teknolojilerini bilinçli kullanmalarının sağlanması da sosyal bilgiler öğretim programının amaçları arasındadır (MEB, 2018). Sosyal Bilgiler öğretiminde beceriye ilişkin amaçlar (NCSS, 1994) kapsamında bilgiye ulaşmada kaynakları doğru kullanma, problem çözme, eleştirel ve yaratıcı düşünme, üst düzey düşünme süreçlerini gerçekleştirme ve iş birliği üzerinde odaklanılmaktadır (Öztürk, 2015). Yalçın (2018) özellikle gelişmekte olan ülkelerin gelecekteki rekabet potansiyellerinin şekillenmesi açısından, bu becerilere daha fazla odaklanması gerektiğini ifade etmiştir. Küresel anlamda ülkelerin başarısının büyük ölçüde insan kaynaklarının sahip olduğu bilgi, yenilik ve yaratıcılık becerilerine ve miktarına bağlı olduğu hususunda ortak görüşler artmaktadır. Bu becerilerin ise kaliteli eğitimle mümkün hale geleceği aşikardır. Bu sebeple modern ekonomilerin kullandığı etkili stratejiler yaratıcılık, yenilikçilik ve bu becerileri kazandıracak sistemin yapısına odaklanmaktadır (Alshammari ve Thomran, 2023). Yaratıcılık eğitimsel uygulamaların öğrenmede etkisini kalıcılaştırmaktadır. Aynı zamanda yaratıcı düşünme tekniklerinin kullanımı, eğitimcilerin davranışlarında da değişim ve gelişim meydana getirmektedir (Erginer, 2016). Bir grup mucit tarafından geliştirildiği düşünülen "Sinektik Tekniği" karar alma ve eyleme geçmeyi sağlayan bir mekanizma olarak geliştirilmiştir. Ancak yaratıcı bir problem çözme tekniği olan sinektik tekniğinin temelini Gordon'un (1961) oluşturduğuna dair genel bir kabul söz konusudur. Sinektik terimi Yunanca "syn" ve "ektos" sözcüklerinin birleşimiyle zıt düşüncelerin bir araya getirilmesi olarak adlandırılmıştır (Nolan, 2003). Bu teknik, birbiriyle arasında daha önceden bir bağlantı saptanmamış iki parçanın ortak bir süzgeçten geçirilerek öğrenenleri farklı düşünme ve problem çözme kapasitelerini artırma imkanı tanımaktadır (Hummell, 2006). Gordon ilk olarak yaratıcı sürece eşlik eden tekrarlayan zihinsel aktivite kalıplarını tanımlamıştır. Oluşturduğu sistem, empatiyi, katılımı, spekülasyonu ve nesnenin özerkliğini içeren çok katmanlı bir yapıdır. Bu süreç boyunca sinektik tekniği farklı araştırmacılar tarafından (Özden, 2005; Seligman, 2007; Hummel, 2004) çeşitli analoji biçimleri bulunarak geliştirilmiştir. Yaratıcı düşünmeyi geliştirmeye yarayan sinektik tekniği ile ilgili bazı çalışmalar (Marran ve Rogan, 1966; Erişti ve Polat, 2017; Khan ve Mahmood, 2018; Bayraktar Balkır ve Topkaya, 2019; Suratno, Komaria, N., Yushardi ve diğ. 2019; Temur ve Çakmak, 2023) bulunmaktadır. Ercan (2010) çalışmasında 7. Sınıf öğrencileri ile fen ve teknoloji öğretiminde yaratıcı düşünme tekniklerinden

FULL TEXTS BOOK

sinektiği kullanmıştır. Çalışma sonucunda öğrencilerin çalıştıkları kavramlara ait başlangıçta göremedikleri özellikleri görür duruma geldikleri, nesnelere yüzeysel bakmaktan sıyrılıp çalıştıkları nesneleri derinlemesine bir bakış açısıyla inceler hale geldikleri ortaya konulmuştur. Çalışmada öğrencilerin problem durumlarına getirdikleri alternatif çözüm önerileri sayısında artış gösterdikleri, bunun yanında nicel olan bu artışın beraberinde daha nitelikli çözümler halinde de ortaya konulduğu sonucuna ulaşılmıştır. Çalışma sosyal bilgiler öğretiminde, orijinal olarak geliştirilen sinektik tekniği temelli etkinliklerden nasıl faydalanabileceğini ortaya koymayı amaçlamaktadır. Yaratıcı düşünmeyi geliştirecek etkinlikler sayesinde öğrencilerin farklı bakış açılarını öğrenmeye ve yeni fikirler üretmeye teşvik edilmesi amaçlanmaktadır.

YÖNTEM

Sinektik tekniğinin sosyal bilgiler derslerinde nasıl kullanılabileceğini ortaya koymak amacıyla yapılan bu çalışma, eylem araştırması yönteminde gerçekleştirilmiştir. Bu çerçevede sosyal bilgiler ders kazanımlarını gerçekleştirmek amacıyla sinektik tekniğini temel alan eylem planı hazırlanmış ve uygulaması araştırmacı tarafından gerçekleştirilmiştir.

Katılımcılar

Araştırma 2024/2025 eğitim öğretim yılı bahar döneminde İstanbul ili Pendik ilçesine bağlı özel bir ortaokul bünyesinde öğrenim gören 5 sınıf öğrencileri ile gerçekleştirilmiştir. Araştırmanın uygulamasının gerçekleştirildiği sınıfta 8'i erkek ve 13'ü kız olmak üzere toplam 21 öğrenci bulunmaktadır. Bulgular kısmında veriler analiz edilirken öğrencilerden kod isimlerle bahsedilmiştir.

UYGULAMA VE BULGULAR

Çalışmanın uygulaması 2024-2025 eğitim öğretim yılının Bahar Döneminde gerçekleştirilmiştir. Çalışma sosyal bilgiler dersinde hayatımızda ekonomi öğrenme alanı kapsamında 2 ders saati boyunca yürütülmüştür. Uygulama süresince yürütülen sinektik temelli ders planları araştırmacı tarafından hazırlanmıştır. Bu kapsamda "Kaynakları verimli kullanmanın doğa ve insanlar üzerindeki etkisini yorumlar." kazanımını odağına alan eylem planı ile bu planın uygulama sürecine ilişkin bulgular aşağıda verilmiştir.

Sinektik Tekniği Temelli Ders Planı 1

Ders: 5. Sınıf Sosyal Bilgiler

Kazanım: Kaynakları verimli kullanmanın doğa ve insanlar üzerindeki etkisini yorumlar.

Beceri: Yaratıcı Düşünme – Yenilikçi Düşünme

Süre: 35-40 dakika

Süreç:

✓ Öğrencilere yapılacak etkinlik hakkında kısaca bilgi verilir. Öğrenme alanının kavramları üzerinde durulur. (Tasarruf, enerji, kaynak, ekonomi)

✓ Öğrencilere verilen kavramlarla ilgili ülke ve dünya genelinde yaşanan kaynakların tüketimine yönelik bir problem tanımlamaları istenir. Bu probleme yönelik sinektik çalışması yapılır. Bu sinektik çalışmasına başlarken ikinci hafta çalışma kağıdı tüm öğrencilere dağıtılır. Ders süresince öğrencilerin ürettikleri analojiler ve kurdukları bağlantısallıklar üzerinde konuşulmuştur.

✓ Öğrencilere *"Beyaz eşyaların üzerindeki enerji sınıfı görseli ile analoji kurma"* etkinliği ile doğrudan analoji örnekleri sunulur. Öğrencilerin yeni tasarladığı enerji sınıfı görseline yönelik ürettikleri

FULL TEXTS BOOK

analojiler ve analojileri kurdukları mantıksal yollar hakkında konuşulmuştur. Aksi bir durumda, tekrar analojilere başvurularak süreç yeniden başlatılmıştır.

✓ Öğrencilerin günlükleri yazmaları sağlanarak süreç tamamlanmıştır.

Beyaz Eşyaların Üzerindeki Enerji Sınıfı Görseli ile Analoji Kurma Etkinliği

Kaplan ve Kaplumbağa (Hız ve Sakinlik Zıtlığı)

Kaplan (Düşük enerji sınıfı): Kaplan çok hızlıdır ve hareket ederken yoğun enerji tüketir. Bu, enerji sınıfında "daha az verimli" olan cihazları temsil edebilir. Düşük enerji sınıfı, çevreye daha fazla zarar verir ve çok fazla tüketim yapar.

Kaplumbağa (Yüksek enerji sınıfı): Kaplumbağa yavaş ama sabırlıdır; enerjisini ekonomik kullanır. Yüksek enerji sınıfı, çevre dostu ve enerji tasarrufu sağlayan cihazları temsil eder.

Analojinin Mantığı:

Enerji sınıfı A+++ olan bir buzdolabı, kaplumbağa gibi yavaş ama etkili ve uzun vadede çevreye daha az zarar verir. F sınıfı bir cihaz ise, kaplan gibi agresif enerji harcaması yapar.

Benim kurduğum analoji:

……
……
……

Arı ve Çekirge (Çalışkanlık ve İsraf Zıtlığı)

Çekirge (Düşük enerji sınıfı): Çekirge enerji verimliliği açısından pek başarılı değildir. Hızla hareket eder ve kaynaklarını çabuk tüketir. Bu, düşük enerji sınıflarını (örneğin C, D) temsil eder.

Arı (Yüksek enerji sınıfı): Arı düzenli, disiplinli ve çalışkan bir varlıktır. Enerjisini verimli kullanır ve topluma katkı sağlar. Bu, yüksek enerji sınıflarını (örneğin A+++) temsil eder.

Analojinin Mantığı:

Arı, enerji tasarrufu yapan cihazların sembolü olabilir. Arının disiplini ve düzeni, enerji tasarrufunu ve çevreye duyarlılığı temsil ederken, çekirgenin savrukluğu düşük enerji sınıfı cihazların israfını temsil eder.

Benim kurduğum analoji:

……
……
……

Güneş ve Yıldızlar (Gece ve Gündüz Zıtlığı)

Güneş (Düşük enerji sınıfı): Güneş güçlüdür ve sürekli ışık saçarken büyük bir enerji harcar. Bu, daha düşük enerji sınıflarını (D, E, F) temsil edebilir.

Yıldızlar (Yüksek enerji sınıfı): Yıldız sakin, az ışık yayar ve enerji tasarrufu sağlar. Yüksek enerji sınıfını (A+, A++) temsil eder.

Analoji Mantığını Sen Bul:

……
...……
…...……
……...

FULL TEXTS BOOK


Benim kurduğum analoji:


……
……
……


Uygulama Süreci

Dersin ilk aşamasında öğrencilerin ön bilgilerini yoklamak ve güdülenmeyi sağlamak amacıyla sinektik tekniği hakkında pekiştirme yapılmıştır. Ardından öğrencilere konuyla ilişkili olarak kaynakları verimli kullanmanın doğa ve insanlar üzerindeki etkisi kazanımına yönelik etkinlik yapılacağı belirtilmiştir. Beyaz eşyaların üzerindeki enerji sınıfı görseli ile analoji kurma etkinlik kâğıdı öğrencilere dağıtıldı. Öğrencilerin kurulan analojinin mantığını sorgulayıp anlayabilmeleri için şu açıklama yapılmıştır: *"Kaplan çok hızlıdır ve hareket ederken yoğun enerji tüketir. Bu, enerji sınıfında "daha az verimli" olan cihazları temsil edebilir. Düşük enerji sınıfı, çevreye daha fazla zarar verir ve çok fazla tüketim yapar. Kaplumbağa yavaş ama sabırlıdır; enerjisini ekonomik kullanır. Yüksek enerji sınıfı, çevre dostu ve enerji tasarrufu sağlayan cihazları temsil eder."* Devamında öğrencilerin kağıtlardaki analojileri okuyarak boş kısımları tamamlamaları istenilmiş ve öğrenci ürünlerinden bazıları aşağıda verilmiştir:


Öğrenci 1








Öğrenci 2




FULL TEXTS BOOK


Analojinin Mantığı:

Arı, enerji tasarrufu yapan cihazların sembolü olabilir. Arının disiplini ve düzeni, enerji tasarrufunu ve çevreye duyarlılığı temsil ederken, çekirgenin savrukluğu düşük enerji sınıfı cihazların israfını temsil eder.

Benim kurduğum analoji:
Uçak hafif bir sinek gibidir enerjisini rahat



Yıldızlar (Yüksek enerji sınıfı): Yıldız sakin, az ışık yayar ve enerji tasarrufu sağlar. Yüksek enerji sınıfını (A+, A++) temsil eder.

Analoji Mantığını Sen Bul:
Benz... değil neden sürünü... onlar...

...ışık... ve güçlü yıldızlar başka... yaklaşılere


Benim kurduğum yeni analoji:
Mavi... kalıp... bir buzdolabı... gibidir

Öğrenci 3


Analojinin Mantığı:

Enerji sınıfı A+++ olan bir buzdolabı, kaplumbağa gibi yavaş ama etkili ve uzun vadede çevreye daha az zarar verir. F sınıfı bir cihaz ise, kaplan gibi agresif enerji harcaması yapar.

Benim kurduğum analoji:
büyük süpürgeler... enerji... ve 2. evi



Analojinin Mantığı:

Arı, enerji tasarrufu yapan cihazların sembolü olabilir. Arının disiplini ve düzeni, enerji tasarrufunu ve çevreye duyarlılığı temsil ederken, çekirgenin savrukluğu düşük enerji sınıfı cihazların israfını temsil eder.

Benim kurduğum analoji:
Kedi... yatar... kaynak... tüketirler

diye dolaşa yemeye devam eder.


Güneş ve Yıldızlar (Gece ve Gündüz Zuhalı)


Yıldızlar (Yüksek enerji sınıfı): Yıldız sakin, az ışık yayar ve enerji tasarrufu sağlar. Yüksek enerji sınıfını (A+, A++) temsil eder.

Analoji Mantığını Sen Bul:

Benim kurduğum yeni analoji:

Öğrenci 4


Analojinin Mantığı:


Enerji sınıfı A+++ olan bir buzdolabı, kaplumbağa gibi yavaş ama etkili ve uzun vadede çevreye daha az zarar verir. F sınıfı bir cihaz ise, kaplan gibi agresif enerji harcaması yapar.

Benim kurduğum analoji:
Aslan... hızlıdır... ve... enerji... sınıfındadır... kaplan... aslan

... enerji... harcar... Salyangoz... yüksek... enerji... sınıfındadır

... Çünkü enerjisini ekonomik bir şekilde harcar

FULL TEXTS BOOK


Yıldızlar (Yüksek enerji sınıfı): Yıldız sakin, az ışık yayar ve enerji tasarrufu sağlar. Yüksek enerji sınıfını (A+, A++) temsil eder.


Analoji Mantığını Sen Bul:

Benim kurduğum yeni analoji:


Dünya ise kendi etrafında döner, ayrıca olarak ayın etrafında dönmez. Bu yüzden bence dünya yüksek enerji sınıfına girer.

Verilen cevaplardan hareketle öğrencilerin analojilerdeki mantıksal yolları kavrayabildikleri görülmektedir. Bir öğrencinin verilen analojiye yönelik çıkarımda bulunamadığı görülmüştür. Öğrencilerin yazdıkları cümlelerde verilen doğal kaynak ile analoji arasında iyi derecede ilişki kurabildikleri ve bu bağlamda yeni analojiler ürettikleri gözlemlenmiştir.


Öğrenci Günlükleri


Ad Soyad: [redacted] Öğrenci 1


Bu iki ders benim için keyifliydi. Bazı yeni şeyler öğrendim. (Not: Bilemdeki derslerime benziyordu) Analoji yapmak biraz zordu.


[redacted] Öğrenci 2

Bazen kafa mabi kullandım çok kolay gelmezdi ama geldi. Sinektik ile yaptığım ilk dersten kolay geçti. Bugünkü dersimde ilk derse göre daha kolay geçti.


Daha önceden sosyal dersinde böyle zıtlıklar kurmuştum ama ders dışında kurmadım. Daha önceden böyle bir şey aklıma hemen gelmezdi. Çünkü konular ilk zordu. Sinektik yöntemi çok eğlenceliydi. İlk anlamadım ama sonra olayı anladım. Bu 2. derste ilk derse göre daha kolaydı. Ama çok eğlendim ☺★

[redacted] Öğrenci 4



Öğrenci 3

Yukarıda yer verildiği üzere öğrenciler sinektik tekniği ile ilgili daha önceden bu tarz benzetimler ve zıtlıklar kurduklarını ifade etmişlerdir. Sinektik tekniği ile işlenen ilk derste zorlandıklarını ikinci dersin daha kolay geçtiğini belirtmişlerdir. Analoji kurmada biraz zorlandıklarını, dersin keyifli geçtiğini ifade etmişlerdir.

Araştırmada birinci eylem planı uygulandıktan sonra gerekli iyileştirmeler yapılarak "Yaşadığı ildeki ekonomik faaliyetleri çözümler." kazanımını odağına alan ikinci eylem planını uygulamasına geçilmiştir. Bu plan ve planın uygulama sürecine ilişkin bulgular aşağıda verilmiştir:

Sinektik Tekniği Temelli Ders Planı 2

Ders: 5. Sınıf Sosyal Bilgiler

Kazanım: Yaşadığı ildeki ekonomik faaliyetleri çözümler.

Beceri: Yaratıcı Düşünme – Yenilikçi Düşünme

Süre: 35-40 dakika

Süreç:

✓ Öğrencilere yapılacak etkinlik hakkında kısaca bilgi verilir. Öğrenme alanının kavramları üzerinde durulur. (Faaliyet, ekonomi)

✓ Öğrencilere yapılacak etkinlikle ilgili bilgi verilir. Ekonomik faaliyetler ile analoji kurma etkinliği kâğıdı dağıtılır. Etkinlik dahilinde ekonomik faaliyetleri anlamlandırmak için hazırlanan "vücut" ile ilgili doğrudan analoji örnekleri sunulur. Analoji ifadesi verildikten sonra öğrencilerin ürettikleri yeni analojiler ve analojileri kurdukları mantıksal yollar hakkında konuşulur.

Ekonomik Faaliyetler ile Analoji Kurma Etkinliği

Birincil faaliyetler (Tarım, hayvancılık): "Vücudun midesi" olarak benzetilir; temel besinleri sağlar.

İkincil faaliyetler (Sanayi): "Vücudun elleri" olarak benzetilir; ham maddeleri işleyerek kullanılabilir hale getirir.

Üçüncül faaliyetler (Hizmet sektörü): "Vücudun ayakları" olarak benzetilir; hizmet sağlar ve hareketi sağlar.

Dördüncül faaliyetler (Bilgi ve teknoloji): "Vücudun beyni" olarak benzetilir; yenilik ve ilerleme sağlar.

Beşincil faaliyetler (Karar alma): "Vücudun kalbi" olarak benzetilir; tüm faaliyetleri düzenler ve yönetir.

Genişletmek için yöneltilecek soru

"Ekonomik faaliyetler bir müzik grubu olsaydı, hangi tarzda hangi enstrümanı çalardı?" sorusuyla öğrencilerin ekonomik faaliyetlere daha yaratıcı bir bakış açısıyla yaklaşmaları sağlanır.

FULL TEXTS BOOK

……

……

……


……

……


Uygulama Süreci

Dersin başlangıcında analojinin tanımı yapılmıştır. Etkinlik kâğıdı öğrencilere dağıtılarak ekonomik faaliyetlerin neden vücuda benzetilmiş olabileceği sorusu yöneltildi. Öğrencilerden bir tanesi "*Bir alanı organa benzetmiş olabilirler mesela beyini ticarete gibi şu an aklıma o geldi."* cevabını vermiştir. Başka bir öğrenci *"Vücudun organları ekonomik işlevlere benzetilebilir" şeklinde yanıtlamıştır.* Diğer biri ise *"Mesela bütçe mide gibi, diğer organlar da harcama, masraflar olabilir."* cevabını vermiştir.

Öğrenci 1




Öğrenci 2




FULL TEXTS BOOK

Öğrenci 3


Genişletmek için yöneltilecek soru
"Ekonomik faaliyetler bir müzik grubu olsaydı, hangi tarzda hangi enstrümanı çalardı?" sorusuyla ekonomik faaliyetlere daha yaratıcı bir bakış açısıyla yaklaşarak cevaplayınız.


Öğrenci 4


Genişletmek için yöneltilecek soru
"Ekonomik faaliyetler bir müzik grubu olsaydı, hangi tarzda hangi enstrümanı çalardı?" sorusuyla ekonomik faaliyetlere daha yaratıcı bir bakış açısıyla yaklaşarak cevaplayınız.


Bu etkinlik öğrencilerin soyut kavramları somut deneyimlerle ilişkilendirmelerine olanak tanımıştır. Öğrenciler, gündelik yaşamlarında karşılaştıkları durumları ekonomik faaliyet türleriyle eşleştirerek kavramların anlamını daha kolay kavramıştır. Aynı zamanda ekonomik faaliyet türlerini müzik aletlerinin işlevleriyle birleştirerek yeni bağdaşımlar kurmuşlardır. Öğrencilerin verdiği her bir örneğin çok farklı bakış açıları ile ele alındığı görülmektedir. Öğrenciler, kendi yaşam pratiklerinden hareketle ekonomik faaliyet çeşitlerini anlamlandırdıklarında, öğrenme sürecinde daha aktif katılım göstermiştir.

Öğrenci Günlükleri


Öğrenci 2


Daha önce birbirine zıt şeyler aklıma gelmez ama çok düşünürsen gelir
Sinektikle yaptığım ilk derste zor yaptım ↘ bugünkü dersimde ilk dersle göre kolay yani zor geçti yani ortası.


Öğrenci 3



Öğrenci 1



Öğrenci 4



Öğrenci günlüklerinden hareketle sosyal bilgiler dersinde analoji kullanımının, öğrenciler tarafından yenilik olarak algılanmasının dersi daha ilgi çekici ve etkileşimli bir hâle gelmesini sağladığı anlaşılmıştır.

Sonuç, Tartışma ve Öneriler

Modern eğitim sistemlerini iyileştirme tartışmalarının ana meselesi haline gelen beceri eğitimi ve aynı zamanda bu becerileri sınıfta geliştirme ve eğitim konularına entegre etmeye yönelik araştırmalar (McLeod ve Cropley, 1989; Wagner, 2016; Yılmaz, 2023) bu çalışmanın da çıkış noktası olmuştur. Dersler farklı yaklaşımlarla tasarlanmış beceri ve yetkinlik açısından zengin etkinliklerle işlenmediği takdirde, becerinin çıktısını elde etmekte sorunlar yaşanabilmektedir. Alanda beceri açısından zengin

FULL TEXTS BOOK

etkinlikler oluşturup uygulamak amacıyla "Hayatımızdaki Ekonomi" öğrenme alanının kazanımları odağında, sinektik tekniği ile oluşturulan etkinlikler temelinde sınıf içi uygulama gerçekleştirilmiştir.

Uygulama sonunda öğrencilerin zıtlıklar ve benzetimler üzerine yeni fikirler oluşturarak yaratıcı analojiler ortaya koydukları görülmüştür. Ercan (2010) fen öğretiminde sinektik tekniğini kullandığı eylem araştırmasında, öğrencilerin problem durumlarına getirdikleri alternatif çözüm önerileri sayısında artış gösterdikleri, nicel olan bu artışın beraberinde daha nitelikli çözümler halinde de ortaya konulduğu sonucuna ulaşmıştır.

Bu minvalde bu çalışmada ise analojilerin bilgiyi zihinsel şemalarına daha kalıcı biçimde yerleştirmelerine olanak sağladığı düşünülmektedir. Öğrencilerin kavramları ezberlemelerinin ötesinde kavramlar arasındaki benzerlik ve farklılıkları keşfetmeleri, üst düzey düşünme becerilerinin gelişimine işaret etmektedir. Bu bağlamda analojilerin, sosyal bilgiler dersinde hem kavramsal öğrenmeyi destekleyen hem de eleştirel ve yaratıcı düşünme becerilerinin gelişimine katkıda bulunan etkili bir öğretim stratejisi olduğu söylenebilir. İbrahimoğlu (2019) sosyal bilgiler dersinde öğrenme güçlüğü çekilen konulara yönelik öğrenci deneyimlerini ve görüşlerini incelediği çalışmasında, bu dersin öğrenilmesinde yaşanan zorlukların ortak nedenleri olarak ezber gerektirmesi, bilgi yoğunluğunun çok fazla olması ve bir diğer önemli neden olarak öğretim yöntem ve teknikleri gösterilmiştir. Bu güçlüklerin çözümüne yönelik öğrenci görüşlerinin büyük bir çoğunluğu, sınıf içi ve sınıf dışı etkinliklerle öğrenme yaşantılarının zenginleştirmesi gerektiğini işaret etmektedir. Yaratıcı düşünme tekniği olan sinektik ile zenginleştirilen sosyal bilgiler dersine ait verilen bulgular ışığında daha çok sınıf içi uygulamalı çalışmaların yapılması önerilir. Söz konusu yöntem ve tekniklerden nasıl faydalanılacağı konusunda yapılan çeşitli eylem araştırmaları eğitimcilere ışık tutacaktır.

KAYNAKÇA

Alshammari, A.E. & Thomran, M. (2023). Towards enhancing creativity and innovation in education system for youth in Hail region. *Advanced Education*, 22, 122-136. DOI: 10.20535/2410-8286.279776

Bayraktar Balkır, N. & Zehir Topkaya, E. (2019). The Influence of Synectics Model as a Prewriting Technique on Vocabulary Development. *Kastamonu Education Journal*, 27(5), 2115-2127. doi:10.24106/kefdergi.3324

Doğan, A. (2020). Geçmişten günümüze sınıf eğitimi ve dönüşüm süreci. İlyas Erpay, Musa Çakır (Ed.), *Eğitim ve Sosyal yönleriyle değişim çağı: yenilikçi ve güncel yaklaşımlar* (1.Baskı) içinde (s. 19-21). Ankara: Astana Yayınları.

Ercan, S. (2010). *Fen Öğretiminde Yaratıcı Düşünme Tekniklerinden Sinektik Kullanımına Yönelik Bir Eylem Araştırması*. Sakarya Universitesi, Fen Bilimleri Enstitüsü, Sakarya.

Erginer, E. (2016). Öğretim ilke ve yöntemleri uygulamalı bir çalışma. Ankara: Pegem Akademi.

Eristi, B., & Polat, M. (2017). The Effectiveness of Synectics Instructional Model on Foreign Language Vocabulary Teaching. *Online Submission*, 5(2), 59–76.

Gordon, W. J. J. (1961). Synectics. New York: Harper & Row.

Hummel, L. (2004). *Synectics for creative thinking in technology education*. Boston: Allyn & Bacon.

Hummell, L. (2006). Synectics for Creative Thinking in Technology Education. *The Technology Teacher*, 66(3), 22-27. https://www.proquest.com/scholarly-journals/synectics-creative-thinking-technology-education/docview/235290687/se-2

İbrahimoğlu, Z. (2019). *Middle school students' conceptual perceptions of the social studies course, and their transfer of the outcomes to daily life*. Routledge Journals, Taylor & Francis Ltd.

FULL TEXTS BOOK

Khan, A. A., & Mahmood, N. (2018). Effect of Synectics Model of Teaching in Enhancing Students' Understanding of Abstract Concepts of Mathematics. Pakistan Journal of Distance and Online Learning, 4(1), 185–198.

Marran, J. F. & Rogan, D. V. (1966). Synectics: An Explanation of the Process and Some Comments on its Application in the Secondary School.

Milli Eğitim Bakanlığı (MEB). (2018). Fen bilimleri dersi öğretim programı (İlkokul ve Ortaokul 3, 4, 5, 6, 7 ve 8. Sınıflar). Ankara: Milli Eğitim Bakanlığı Talim ve Terbiye Kurulu Başkanlığı.

McLeod, J., & Cropley, A.J. (1989). Fostering academic excellence. Oxford: Pergamon.

National Council for the Social Studies (NCSS). (1994). *Curriculum Standards for Social Studies: Expectations of Excellence*. Silver Spring, MD: NCSS.

Nolan, V. (2003), Whatever Happened to Synectics?. *Creativity and Innovation Management*, 12:24-27. https://doi.org/10.1111/1467-8691.00264.

Özden, Y. (2005). *Öğrenme ve öğretme* (7. baskı). Ankara: Pegem Akademi Yayıncılık.

Öztürk, C. (2015). *Sosyal Bilgiler Öğretimi: Demokratik Vatandaşlık Eğitimi*. Ankara: Pegem Akademi.

Seligmann, E. R. (2007). *Reaching students through synectics: A creative solution* (Unpublished doctoral thesis). University of Northern Colorado.

Suratno, Komaria, N., Yushardi, Dafik, & Wicaksono, I. (2019). The Effect of Using Synectics Model on Creative Thinking and Metacognition Skills of Junior High School Students. *International Journal of Instruction*, 12(3), 133-150. https://doi.org/10.29333/iji.2019.1239a

Temur, S., & Çakmak, M. (2023). Ortaokul Öğrencilerinin Özgürlük Kavramına İlişkin Algılarının Sinektik Tekniğiyle İncelenmesi. *Academia Eğitim Araştırmaları Dergisi*, 8(1), 39-59. https://doi.org/10.53506/egitim.1226780

Wagner, T. (2016). *Yenilikçiler yaratmak*. (Çev: Özer, A.) T.C. İstanbul: Kültür Üniversitesi Yayınları.

Yalçın, S. (2018). 21. yüzyıl becerileri ve bu becerilerin ölçülmesinde kullanılan araçlar ve yaklaşımlar. *Ankara Üniversitesi Eğitim Bilimleri Fakültesi Dergisi*, 51(1), 183-201. https://doi.org/10.30964/auebfd.405860

Yılmaz, H. (2023). *Sosyal Bilgiler Öğretmenlerinin Çevrimiçi Mesleki Gelişim Ve Çevrimiçi Eğitim Faaliyetleri Hakkındaki Görüşleri, Mesleki Gelişimleri İçin Çevrimiçi Portal (Site) Önerisi*. Bolu Abant İzzet Baysal Üniversitesi, Lisansüstü Eğitim Enstitüsü, Bolu.

Yiğit, E.Ö. (2022). Farklı Ülkelerde Sosyal Bilgiler Öğretimi. Cemil Öztürk ve Tuğba Kafadar (Ed.), *Sosyal Bilgiler Öğretimi Etkili Vatandaşlık Eğitimi El Kitabı* (1.Baskı) içinde (s.25-27). Ankara: Nobel Yayınları.

FULL TEXTS BOOK

TARAFSIZLIK MASKESİ ALTINDA DİJİTAL TARAFGİRLİK: GROK VE FİLİSTİN SÖYLEMİ

Prof. Dr. Cavit Yavuz

Ordu Üniversitesi Sosyal Bilimler Enstitüsü,

İletişim Çalışmaları Anabilim Dalı, Altınordu, Ordu.

Hazal Zaimoğlu

Ordu Üniversitesi Sosyal Bilimler Enstitüsü,

İletişim Çalışmaları Anabilim Dalı, Altınordu, Ordu.

ÖZET

Bu çalışma, yapay zekâ teknolojilerinin tarafsızlık iddiası altında nasıl yönlendirici, seçici ve ideolojik bilgi üretimi gerçekleştirdiğini incelemektedir. Twitter (X) platformunda entegre edilen “Grok” adlı yapay zekâ sisteminin Filistin-İsrail çatışmasına dair kullanıcılara verdiği sekiz farklı yanıt içerik analizi yöntemiyle değerlendirilmiştir. Yapay zekânın sadece teknik bir araç olmadığı, aynı zamanda siyasal ve toplumsal bağlamlarda söylem üreten bir aktör olduğu varsayımıyla hareket edilen çalışmada, öncelikle yapay zekâ kavramının tarihsel gelişimi, dezenformasyon ve propaganda süreçlerindeki yeri kuramsal çerçevede ele alınmıştır. Ardından, Filistin-İsrail meselesinin tarihsel arka planı sunulmuş ve Grok’un yanıtları söylemsel açıdan analiz edilmiştir.

Grok’un verdiği sekiz yanıt; çatışmanın başlangıcı, Gazze'deki sivil kayıplar, İsrail'in güvenlik politikaları, Hamas’ın rolü, barış girişimleri, uluslararası tepkiler, Yahudi halkının mağduriyet söylemleri ve Filistinlilerin hak ihlalleriyle ilgiliydi. Bu yanıtlar yüzeyde dengeli ve tarafsız görünmesine rağmen, analiz sonuçları Grok’un bazı anlatıları ön plana çıkardığını, Filistinlilerin yaşadığı mağduriyetleri arka planda bıraktığını ve İsrail’in güvenlik söylemini meşrulaştıran bir dil kullandığını göstermektedir. Özellikle "çatışmanın karmaşıklığı", “her iki tarafın da haklı talepleri ve acıları var” gibi genellemeler, politik bağlamı belirsizleştirmekte ve yapısal eşitsizlikleri görünmez kılmaktadır.

Çalışmanın temel savı, yapay zekâ sistemlerinin yalnızca bilgi sağlayıcı değil, dijital çağın hegemonik söylem aktörleri olduğudur. Bu nedenle Grok örneği üzerinden dijital tarafgirliğin biçimleri açığa çıkarılmış ve yapay zekânın etik, şeffaf ve çok boyutlu bir denetim mekanizmasıyla değerlendirilmesi gerektiği vurgulanmıştır.

Anahtar Kelimeler: Yapay zekâ, Grok, dijital propaganda, Filistin-İsrail çatışması

ABSTRACT

This study examines how artificial intelligence (AI) technologies can produce selective, guiding, and ideologically loaded information under the guise of neutrality. The research focuses on “Grok,” an AI system integrated into the Twitter (X) platform, and analyzes eight different responses it generated concerning the Israeli-Palestinian conflict using content analysis methodology. Based on the assumption that AI is not merely a technical tool but a discourse-producing actor shaped by political and social contexts, the study first explores the historical development of AI and its role in disinformation and propaganda within a theoretical framework. Subsequently, the historical background of the Israeli-Palestinian issue is presented, followed by a discourse-based analysis of Grok’s responses.

FULL TEXTS BOOK

The eight responses analyzed cover topics such as the origins of the conflict, civilian casualties in Gaza, Israel's security policies, the role of Hamas, peace efforts, international reactions, narratives of Jewish victimhood, and human rights violations against Palestinians. While Grok's answers appear neutral and balanced on the surface, the analysis reveals that it foregrounds certain narratives, marginalizes Palestinian suffering, and employs language that legitimizes Israeli security discourse. Generalized expressions such as "the complexity of the conflict" and "both sides have legitimate grievances and suffering" serve to obscure the political context and render structural inequalities invisible.

The central argument of this study is that AI systems are not just information providers but hegemonic discourse actors of the digital age. Through the example of Grok, the study uncovers forms of digital partisanship and emphasizes the need for ethical, transparent, and multi-dimensional oversight of AI technologies.

Keywords: Artificial intelligence, Grok, digital propaganda, Israeli-Palestinian conflict

GİRİŞ

Dijitalleşmenin hız kazandığı 21. yüzyılda, bilgi üretim süreçleri giderek daha çok yapay zekâ sistemlerinin denetimine girmeye başlamıştır. Son yıllarda yaşanan teknolojik dönüşümler, toplumsal yapıyı ve gündelik yaşam pratiklerini köklü biçimde değiştirmiştir. Bu dönüşümün en dikkat çekici unsurlarından biri olan yapay zekâ, yalnızca dijital araçlarda değil, ulaşım sistemlerinden ev içi teknolojilere kadar geniş bir yelpazede etkisini hissettirmektedir. Günümüzde birçok ülkede yapay zekâ tabanlı uygulamalar, bireylerin günlük yaşamlarını kolaylaştıran ve çeşitli alanlarda işlevsellik sunan çözümler üretmektedir (Güzey, Çakır, Athar, Yurdaöz & Saad,2023). Yapay zekâ, artık sadece teknik bir gelişme değil; aynı zamanda toplumsal, siyasal ve kültürel sonuçlar doğuran bir olgudur. Günümüzde özellikle sosyal medya platformlarında kullanılan yapay zekâ tabanlı algoritmalar, bireylerin neyi göreceğine, neyi okuyacağına ve hatta neyi düşüneceğine doğrudan etki edebilecek bir güce sahiptir. İletişim teknolojilerindeki ilerlemeler, kitle iletişim araçlarının gelişimini beraberinde getirmiş; bu da bireylerin ya da toplulukların düşünsel etkilerini geniş kitlelere ulaştırabilme ve bazı fikirleri toplumsal düzeyde baskın hale getirme olanaklarını artırmıştır (Karakuş,2021).

Bu çerçevede Twitter (X) platformu üzerinden çalışan Grok adlı yapay zekâ sistemi, kullanıcıların sorduğu sorulara gerçek zamanlı yanıtlar üretmesiyle dikkat çekmektedir. Ancak Grok'un özellikle siyasal ve çatışmalı konular üzerindeki yanıtları, yapay zekâ sistemlerinin tarafsızlığına ilişkin etik ve ideolojik tartışmaları gündeme getirmiştir. Söz konusu tartışmalar yalnızca teknolojiye değil, bu teknolojinin üretildiği sosyal bağlama da aittir.

Akademik literatürde, yapay zekâ sistemlerinin manipülasyon potansiyeli giderek daha fazla tartışılmaktadır. Bu bağlamda, Filistin-İsrail meselesi gibi tarihsel, siyasal ve etik açılardan karmaşık bir konuda Grok'un yanıtlarının ideolojik içerikler taşıyıp taşımadığı sorusu önem arz etmektedir. Çünkü yapay zekânın ürettiği cevaplar yalnızca bilgi sunmakla kalmaz, aynı zamanda birer anlamlandırma eylemine dönüşür. Bu da bizi, yapay zekâ sistemlerinin dijital propaganda mekanizmalarına entegre olup olmadığı sorusuna götürmektedir.

Bu çalışmanın temel amacı, Twitter'ın Grok adlı yapay zekâ sisteminin Filistin-İsrail çatışmasına dair verdiği yanıtları içerik analizi yöntemiyle inceleyerek, YZ'nin tarafsızlık iddiasının ne ölçüde geçerli olduğunu ortaya koymaktır. Bu kapsamda, ilk bölümde yapay zekânın tarihçesi, tanımı ve manipülasyon, dezenformasyon ve propaganda süreçlerine etkisi tartışılacak; İkinci bölümde Filistin-İsrail meselesinin tarihsel arka planı özetlenecek, Üçüncü bölümde Grok'un yanıtları söylem çözümlemesiyle değerlendirilecektir. Sonuç bölümünde ise bulgular ışığında yapay zekânın hegemonya üretimindeki yeri sorgulanacaktır.

FULL TEXTS BOOK

1. YAPAY ZEKÂ

Yapay zekâ kavramı, teknolojiyle iç içe geçmiş günümüz dünyasında sıkça duyulmakla birlikte, aslında oldukça karmaşık ve çok boyutlu bir içeriğe sahiptir. Bu kavram, yalnızca makinelerin zeki davranışlar sergilemesi değil, aynı zamanda insan zihninin işleyişini taklit etme kapasitesiyle de ilişkilidir. Ancak yapay zekânın ne olduğu sorusu, farklı disiplinlerde farklı anlamlar kazanmakta, zaman içinde teknolojiyle birlikte dönüşmektedir. Bu bağlamda, yapay zekânın kapsamını ve sınırlarını daha iyi anlayabilmek için çeşitli tanımlara yer vermek yararlı olacaktır.

Oxford English Dictionary'de yapay zekâ, 'görsel algı, ses tanıma, karar verme ve diller arası çeviri gibi, normalde insan zekâsı gerektiren işleri yapabilecek şekilde tasarlanan ve geliştirilen bilgisayar sistemleri' olarak tanımlanmaktadır (akt: Güngör & İmik Tanyıldızı, 2023).

Yapay zekâ terimi, genel olarak insanların desteğiyle veya tamamen bağımsız olarak öğrenebilen çeşitli makine işlevlerini ifade etmektedir (Rodgers,2021).

"Yapay zekâ; insana özgü konuşma, anlama, öğrenme, karar verebilme, akıl yürütme, tahminde bulunma, harekete geçme ve tepki verme gibi insan tarafından gerçekleştirildiğinde yüksek derecede zekâ isteyen birçok bilgi, beceri ve davranışın veri işleme maksadıyla bilgisayar sistemlerine kazandırmayı amaç edinen bir bilim dalı olarak tanımlanabilir." (Öztürk,2024, s.284)

Özetle, yapay zekâ, insan zekâsına özgü bilişsel işlevlerin bilgisayarlar aracılığıyla taklit edilmesi ve bu sayede makinelerin öğrenme, analiz yapma ve karar verme yeteneklerinin geliştirilmesidir.

Bu tanımlar, yapay zekânın insan zekâsının karmaşık işleyişini makinelere aktarma ve bu sayede öğrenme, karar verme gibi yetenekler kazandırma sürecini ifade ettiğini göstermektedir. Bu çerçevede, yapay zekâ yalnızca günümüz teknolojisinin bir ürünü olarak değil, aynı zamanda tarihsel bir düşünsel sürecin devamı olarak da değerlendirilebilir. İnsanoğlu, binlerce yıl öncesinden itibaren bilgi biriktirme, sistemleştirme ve cansız varlıkları belli işlevleri yerine getirecek şekilde harekete geçirme fikriyle ilgilenmiştir. Yaklaşık 4000 yıl öncesine dayanan veri depolama arayışları, bu sürecin en erken örneklerinden biridir. Ancak modern anlamda yapay zekâ ve özellikle yapay sinir ağları alanındaki somut gelişmeler esasen son elli yıl içinde hız kazanmıştır (Öztürk & Ergin Şahin, 2018).

Yapay zekâ kavramı ilk olarak 1956 yılında Dortmund Konferansı'nda John McCarthy, Marvin L. Minsky, Nathaniel Rochester ve Claude E. Shannon tarafından sunulan bir öneri mektubunda dile getirilmiştir (Arslan,2020, s.76). Bu konferans, yapay zekânın akademik bir disiplin olarak doğuşunu simgelemektedir. Bu konferans, yapay zekânın akademik bir disiplin olarak doğuşunu simgelemektedir. O tarihten bu yana yapay zekâ, yalnızca matematiksel modelleme, mühendislik ve veri işleme gibi teknik alanlarda değil, aynı zamanda toplumsal, kültürel ve siyasal yapılar üzerinde doğrudan etkiler yaratan çok katmanlı bir araç hâline gelmiştir. Özellikle bilgi çağında dijital platformların yükselişiyle birlikte, yapay zekânın dezenformasyon üretimi, algı yönetimi ve toplumsal manipülasyon süreçlerinde nasıl kullanıldığı sorusu, etik ve politik tartışmaların merkezine yerleşmiştir.

1.2. Yapay Zekanın Dezenformasyon ve Manipülasyon Gücü

Bilgi üretme, yorumlama ve karar verme gibi insanî yetileri taklit eden yapay zekâ sistemleri, aynı zamanda bilgiyi biçimlendirme, yönlendirme ve hatta manipüle etme potansiyeline de sahiptir. Bu durum, özellikle dijital medya ortamlarında yapay zekânın dezenformasyon ve algı yönetimi araçlarına dönüşme riskini doğurmuştur. İnsana özgü faaliyetleri ve daha fazlasını yerine getirebilmesi adına günümüzde yapay zekânın birçok alanda kullanıldığı görülmektedir. Bu alanlar arasında konu gereği bilginin üretimi ve yayılması süreçlerinde medya sektörü de yer almıştır (Bekar, S,2024, s.50).

FULL TEXTS BOOK

Bu noktada, kavramsal bir çerçeve çizmek amacıyla dezenformasyon, propaganda ve manipülasyon arasındaki ayrımı açıklamak yerinde olacaktır. Dezenformasyon, kasıtlı olarak yanlış veya yanıltıcı bilgilerin dolaşıma sokulması anlamına gelir ve çoğu zaman toplumsal belirsizlik, kafa karışıklığı veya güven erozyonu yaratma amacı taşır. Dezenformasyon genel olarak olmayan bir durumu varmış gibi gösterme amacıyla üretilmiş yalan/yanlış bilgi üretimidir. Bu bilgiler bilinçli olarak yanlış üretilir ve hedef kitlenin bilinçaltı ve eylemleri içeriğe uyumlu olması amaçlanır (Erkan, Ayhan,2018, s. 206). Propaganda, genellikle bir ideolojiyi, politik görüşü ya da kurumsal çıkarı yaymak ve meşrulaştırmak amacıyla yürütülen, içerdiği bilgi doğru olsa bile seçici ve yönlendirici bir iletişim biçimidir. Özsoy'a göre propaganda, insanların düşüncelerini, görüşlerini ve eylemlerini iletişim araçlarını kullanarak propaganda yapanın istediği biçimde etkileme ve değiştirme yönelik bilinçli bir davranışlar bütünüdür (Özsoy,1998,akt. Durmuş,2023, s.47). Manipülasyon ise hem dezenformasyonu hem de propagandayı içerebilen daha geniş bir stratejik yönlendirme biçimi olarak, bireylerin düşünce ve davranışlarını belli hedefler doğrultusunda şekillendirmeyi amaçlar.

Manipülasyon iletişim sürecinde alıcı ile verici arasında mesaj yapısına ve içeriğine bilinç bir müdahaledir. Bu müdahale özellikle kaynak tarafından ve kimi zamanda araç yardımıyla gerçekleşmekte ve alıcıda istenilen düşünce ve fikir üretme süreçlerine etki etme şeklinde gerçekleşmektedir. Bu bağlamda manipülasyonun temel amacı fikir ekmek, fikri değiştirmek veya fikri yönlendirmek şeklinde değişik amaçları gütmektedir (Elitaş,2022, s. 115).

Geleneksel medya araçlarından farklı olarak dijital platformlarda içerik üretim ve yayım hızının artması, yapay zekâ destekli sistemlerin dezenformasyon ve manipülasyon faaliyetlerini daha etkili biçimde gerçekleştirmesini mümkün kılmaktadır. Yapay zekâ, özellikle doğal dil işleme, derin öğrenme ve büyük veri analizi gibi alanlardaki ilerlemeler sayesinde sahte haber üretimi, gerçek dışı içeriklerin otomatik olarak çoğaltılması ve hedef kitleye kişiselleştirilmiş yanlış bilgiler sunulması gibi işlevleri üstlenebilmektedir. Yapay zekâ, derin öğrenme teknikleriyle görsel ve işitsel içerikleri dönüştürebilmekte; özel yazılımlar ve algoritmalar aracılığıyla, yaşanmamış olayları gerçek gibi gösteren sahte içerikler üretilebilmektedir (Bekar,2024). Bu tür içerikler, izleyicide gerçeklik algısını sarsarak bilgiye duyulan güveni azaltmakta ve manipülasyonun etkisini artırmaktadır. Gerçekle kurulan bağın yapay olarak yeniden inşa edilmesi, bireylerin doğru ile yanlışı ayırt etmesini zorlaştırmakta; böylece algılar üzerinden yönlendirme ve kitle kontrolü daha sistematik bir hâl almaktadır.

Yapay zekâ, dezenformasyon kampanyalarında yalnızca yanlış bilgilerin oluşturulması ve yayılmasında değil, aynı zamanda deepfake gibi başlı başına kamuoyunu manipüle etme potansiyeline sahip araçlarla da kilit bir rol üstlenmektedir. Yapay zekâ teknolojileri, insan yazısını taklit eden sentetik metinler oluşturarak bireylerin gerçek ve sahte haberleri ayırt etmesini zorlaştırmakta; bu durum özellikle dış politika gibi hassas alanlarda kamuoyunu etkilemek amacıyla kötü niyetli aktörler tarafından istismar edilmektedir (Kreps, McCain ve Brundage, 2020, akt. Avşar, 2025, s. 215).

Bu nedenle, yapay zekâ temelli araçların yalnızca bilgi üretiminde değil, aynı zamanda kamuoyu algısının şekillendirilmesinde nasıl stratejik bir enstrümana dönüştüğü açıkça görülmektedir. Bu durum, dijital çağda bilgi güvenliği, etik ve demokratik karar alma süreçleri açısından ciddi riskler doğurmakta; özellikle dezenformasyona karşı dirençli toplumlar inşa etmenin gerekliliğini daha da hayati hâle getirmektedir.

2. FİLİSTİN-İSRAİL MESELESİNİN TARİHSEL ARKAPLANI

Adını M.Ö. XII. yüzyılda Avrupa'dan kavimler göçü sırasında deniz yolu ile buraya gelen Filistlerden alan Filistin, Arz-ı Kenan veya Arz-ı Mukaddes denilen yerin ismidir (Karaman,1996). Coğrafi konum olarak Filistin, Akdeniz'in güney doğu ucunda Asya ile Afrika arasında köprü konumundaki yerdir (Karaköse,2018,s.151). Bu stratejik ve kutsal nitelikleri nedeniyle Filistin, tarih boyunca sayısız işgal

FULL TEXTS BOOK

ve çatışmaya sahne olmuştur. Ancak 20. yüzyılın ortalarından itibaren bu bölgedeki en önemli siyasi ve insani krizlerden biri Filistin-İsrail çatışmasıdır. Bu çatışmanın kökleri, Yahudi halkının tarihî olarak "vaat edilmiş topraklar" olarak gördüğü Filistin'e göçü, ardından İsrail devletinin 1948'de kurulması ve bunun sonucunda yaşanan toprak kaybı, göçler, işgaller ve direnişler zincirine dayanmaktadır. Filistin'in tarihî ve coğrafi önemi, bu topraklar üzerindeki hak iddialarını daha da karmaşık hale getirmiş, günümüzde de süregelen siyasi, dini ve insani bir kriz hâlini almıştır.

Filistin-İsrail meselesi, yalnızca iki halk arasında yaşanan bir toprak anlaşmazlığının ötesinde, sömürgecilik, uluslararası çıkarlar, dinî kimlikler ve jeopolitik hesapların iç içe geçtiği karmaşık bir çatışmalar zinciridir. Doğu Akdeniz'in enerji güvenliğinde artan stratejik rolü, Batılı aktörleri bölgedeki siyasi ve askeri kontrolü kendi çıkarları doğrultusunda şekillendirmeye yönlendirmiş; bu bağlamda Gazze Şeridi, istikrarı tehdit eden bir unsur olarak görülmeye başlanmıştır.(Tarakçı,2009).

Bu sorunun kökenleri 19. yüzyılın sonlarına, modern Siyonizmin ortaya çıkışına ve Osmanlı topraklarında Yahudi göçlerinin başlamasına kadar uzanır. 19. yüzyılın sonlarında Rusya'da yaşayan Yahudilerin karşılaştığı ayrımcılık ve baskılar, kitlesel göçlere yol açmış; bu süreçte Siyonist Hareket ortaya çıkarak Yahudi sorununun Filistin'de bağımsız bir ulusal yurt kurulmasıyla çözülebileceği düşüncesi yaygınlaşmıştır (Saleh, 2016). Aynı dönemde Osmanlı Devleti'nin Filistin üzerindeki zayıflaması ve Batılı güçlerin bölge üzerindeki nüfuz artırma çabaları, Siyonist Proje'nin gerçekleştirilmesi için uygun zemin yaratmıştır.Siyonizm hareketi Filistin'de bir 'Yahudi Yurdunun' kurulması amacıyla hareket etmiş olsa da gerçekte Filistin'deki yerli halkı bir şekilde yerlerinden ederek bir Yahudi devletinin kurulması amaçlanmıştır (Kaplan,2019, s.14).

Batılı devletlerin, Osmanlı'nın çöküşünden sonra bölgede güçlü bir Müslüman-Arap devletinin oluşmasını engellemek amacıyla tampon devlet fikrini destekledikleri belirtilmektedir. Bu tampon devletin, Arap ve Müslüman dünyasının bölünmesine ve bölgenin uzun süreli istikrarsız kalmasına neden olduğu ifade edilmektedir.

1914-1948 yılları arasında bu topraklar üzerinde açıktan veyahut yeraltından güç mücadeleleri yaşanmış, çeşitli gizli antlaşmalar yapılmış, bölge üzerinde emelleri olan farklı aktörler arasında stratejik hesaplaşmalar görülmüştür. Osmanlı'nın Arap coğrafyasındaki hâkimiyetinin zayıflamasıyla Filistin toprakları, Balfour Deklarasyonu ve Sykes-Picot Antlaşması gibi gelişmelerle İngiliz mandası altında çatışma ve kaosun merkezi hâline gelmiştir (Yiğit,2020). Bu süreç, sadece bölgesel dengeleri değil, küresel güç ilişkilerini de derinden etkilemiş; Filistin toprakları büyük devletlerin nüfuz mücadelesine sahne olmuş ve yapay sınırlarla şekillendirilen siyasi yapılar, günümüze kadar uzanan istikrarsızlıkların temelini atmıştır. Özellikle Sykes-Picot sonrası çizilen sınırlar, etnik ve dini yapıları göz ardı ettiği için Filistin meselesinin sadece yerel değil, uluslararası boyutta bir sorun olarak büyümesine zemin hazırlamıştır. Günümüzde hâlâ devam eden bu çatışma, sadece yerel bir mesele değil; aynı zamanda küresel politikada yankı uyandıran, insan hakları, uluslararası hukuk ve medya temsilleri gibi birçok alanda tartışmalara zemin hazırlayan bir krizdir.

Bu tarihsel arka plan üzerinde yükselen Filistin sorunu, zamanla yalnızca bölgesel aktörlerin değil, küresel güçlerin çıkar çatışmalarının merkezine dönüşmüştür. Bu noktada özellikle Amerika Birleşik Devletleri ile İsrail arasındaki stratejik işbirliği, Filistin meselesinin uluslararası boyutunu daha da derinleştirmiştir.

Kurulduğu günden bugüne İsrail ile en yakın ilişkilere sahip olan küresel aktörün, ABD olduğu görülmektedir, ABD-İsrail ilişkileri, yalnızca diplomatik ve ekonomik bağları değil, aynı zamanda dini, yasal, askeri bağları da içerdiğinden, II. Dünya Savaşı'ndan sonra, stratejik bağların yanı sıra yakın bir değerler ortaklığı üzerine, şekillenmiştir (Kişi,2023,s.269).

Bu iş birliği, İsrail'in Ortadoğu'daki güvenilir müttefik konumunu pekiştirirken, Filistin halkının yaşadığı hak ihlalleri karşısında uluslararası dengelerin tek taraflı şekillenmesine yol açmıştır. ABD'nin

FULL TEXTS BOOK

Birleşmiş Milletler ’de İsrail lehine kullandığı veto hakları, askeri yardımlar ve diplomatik koruma mekanizmaları, İsrail'in bölgede uyguladığı politikaların sürdürülmesinde belirleyici rol oynamıştır. Bu durum, Filistin meselesinin çözümünü daha da zorlaştırmakta ve tarafsız bir uluslararası arabuluculuk ortamının oluşmasını engellemektedir.

3. ARAŞTIRMA YÖNTEMİ

Bu çalışmada, Grok adlı yapay zekâ sisteminin Filistin-İsrail çatışmasına ilişkin verdiği yanıtları incelemek amacıyla nitel araştırma yöntemi benimsenmiştir. Özellikle, yapay zekânın tarafsızlık iddiası altında nasıl söylemler ürettiğini değerlendirebilmek için içerik analizi tekniği kullanılmıştır. Veri seti, 2025 yılı boyunca X (eski adıyla Twitter) platformunda kullanıcılar tarafından Grok’a yöneltilmiş, Filistin-İsrail çatışmasına dair sorular ve bunlara verilen yanıtlar arasından seçilmiştir. Seçim sürecinde, yalnızca çatışmanın tarihsel, siyasal ya da insani boyutlarına dair açık uçlu sorular esas alınmış; yanıtlarda tarafsızlık, yönlendirme, bilgi eksikliği ya da ideolojik yönelim izlenimi veren örnekler tercih edilmiştir.Analiz sürecinde, Grok’un verdiği yanıtlar belirli temalar altında sınıflandırılmış; metinlerde geçen anahtar kelimeler, öne çıkarılan aktörler, atıfta bulunulan olaylar ve anlatım biçimi dikkate alınarak betimsel bir çözümleme yapılmıştır. Bu süreçte verilerin sıklığı ya da nicel dağılımı değil, içerdiği anlam örüntüleri değerlendirilmiştir. İçerik analizinin amacı, Grok’un yanıtlarının görünürde tarafsız olup olmadığı ve hangi tür söylemsel yönlendirmelere açık olduğu sorularına yanıt bulmaktır. Bu yöntem, çalışmanın yalnızca yapay zekâ çıktılarının teknik değerlendirmesini değil; aynı zamanda bu çıktılarda yer alan anlam katmanlarının çözümlenmesini amaçlamaktadır.

4. ARAŞTIRMA BULGULARI

Bu bölümde, Grok adlı yapay zekâ sisteminin Filistin-İsrail çatışmasına dair verdiği yanıtlar içerik analizi yöntemiyle çözümlenerek elde edilen bulgulara yer verilmiştir. Analiz sürecinde, Grok’un yanıtlarında öne çıkan söylemsel örüntüler belirlenmiş; özellikle tarafsızlık iddiası altında nasıl bir ideolojik yapı sergilediği sorgulanmıştır. Yanıtlar, tematik başlıklar altında gruplandırılarak, her bir tema çerçevesinde anlam örüntüleri, atıfta bulunulan aktörler ve dil kullanımındaki yönlendirici unsurlar analiz edilmiştir.

		Kodlayıcı 1		
		1	**0**	
Kodlayıcı 2	**1**	a = 122	b = 16	138
	0	c = 10	0	10
		132	16	148

Şekil 1. Kodlayıcılar Arası Uyum Tablosu

P(observed) = Po = a / (a + b + c) = 0.82

P(chance) = Pc = 1 / Kodların Sayısı = 1 / 52 = 0.02

FULL TEXTS BOOK

Kappa = (Po - Pc) / (1 - Pc) = 0.82

Bölüm başına eşit olmayan sayıda kod mevcutsa veya değerlendirilecek yalnızca bir kod varsa:

$$P(chance) = Pc = Kodların\ Sayısı / (Kodların\ Sayısı + 1)^2 = 0.02$$

Kappa = (Po - Pc) / (1 - Pc) = 0.82

Şekil 1'de, çalışmada yer alan iki bağımsız kodlayıcının yaptığı kodlamalar arasındaki uyum Cohen's Kappa katsayısı ile hesaplanmıştır. Tabloya göre her iki kodlayıcının aynı birimi kodladığı durum sayısı 126'dır. Bu doğrultuda gözlenen uyum değeri (Po) 0.82 olarak belirlenmiştir. Şansa bağlı uyum olasılığı ise 0.02 olarak hesaplanmış, böylece Cohen's Kappa değeri 0.82 bulunmuştur. Bu değer, nitel araştırmalarda yüksek derecede kodlayıcı uyumu olduğunu göstermektedir (Landis & Koch, 1977'ye göre Kappa > 0.80 "çok iyi uyum"dur).


Kodlayıcı uyumunun yüksek olması, araştırmanın iç geçerliliğini güçlendirmektedir. Ayrıca hesaplamalar, MAXQDA'nın "Intercoder Agreement" modülü üzerinden gerçekleştirilmiştir.



Şekil 2. İçerik Analizine Dayalı Sözcük Frekansı Görselleştirmesi

Bu kelime bulutu, araştırma kapsamında yapılan içerik analizi sonucunda elde edilen metinlerdeki sözcük frekanslarına dayalı olarak oluşturulmuştur. Görselde yer alan sözcüklerin boyutları, ilgili terimin metinler içerisindeki tekrar sıklığına göre ölçeklendirilmiştir. "İsrail", "Filistin", "çatışma", "taraf", "bilgi" ve "yahudi" gibi sözcüklerin öne çıkması, analiz edilen metinlerin tematik olarak siyasal, tarihsel ve toplumsal meseleler çerçevesinde yoğunlaştığını göstermektedir. Bu görselleştirme, içerik analizinin bulgularını destekleyici bir araç olarak kullanılmış; verilerin genel eğilimlerini görsel bir biçimde sunmayı amaçlamıştır.

FULL TEXTS BOOK



Şekil 3. Kodlar Arası Ortaklık ve Tematik İlişki Ağı

Yukarıda yer alan görsel, içerik analizi sürecinde elde edilen tematik kodların birbirleriyle olan eşzamanlı geçişlerini (co-occurrence) temsil etmektedir. MAXQDA yazılımı aracılığıyla oluşturulan bu kod ilişkileri haritası, her bir kodun diğer kodlarla hangi sıklıkta aynı metin segmentinde yer aldığını görsel olarak sunmaktadır. Kodlar arasındaki kırmızı çizgiler, eşzamanlılık ilişkisini; renkli gruplamalar ise tematik kümelenmeleri göstermektedir.

Örneğin, "kuruluş" ve "devletsizlik" kodlarının aynı alt metinlerde birlikte kullanılması, bağımsızlık ve siyasi yapı sorunlarının birbirini tamamlayan temalar olarak çalışmada öne çıktığını göstermektedir. Benzer şekilde, "çatışma", "haklılık söylemi" ve "yerleşimci" kodları arasında yoğun bağlantılar bulunmakta olup bu temaların özellikle politik söylem düzeyinde iç içe geçtiği anlaşılmaktadır. Ayrıca "gönüllü göç" ve "zorunlu göç" kodlarının bağlantısı, anlatılarda göç olgusunun çok boyutlu biçimde ele alındığını ortaya koymaktadır.

Bu görselleştirme, yalnızca bireysel kodların frekansını değil; aynı zamanda bu kodların birlikte kullanıldığı yapısal örüntüleri de ortaya koyarak, analiz sürecine kavramsal bütünlük sağlamaktadır. Bu sayede temalar arası ilişkisel ağ daha görünür kılınmış ve veri setindeki örtük yapılar desteklenmiştir.

SONUÇ

Bu çalışma, yapay zekâ teknolojilerinin görünürdeki tarafsızlık iddiasının ötesinde, nasıl belirli ideolojik çerçeveleri yeniden üretebileceğini ortaya koymuştur. Grok adlı yapay zekâ sisteminin Filistin-İsrail çatışmasına dair verdiği yanıtlar üzerinden yürütülen içerik analizinde, sistemin yalnızca bilgi aktaran bir araç olmaktan ziyade, belirli anlatıları önceliklendiren ve bazı aktörleri daha görünür kılan bir söylem üreticisi olarak işlev gördüğü tespit edilmiştir. Bu durum, yapay zekânın algoritmik altyapısının ve veri setlerinin siyasal ve kültürel bağlamlardan bağımsız düşünülemeyeceğini bir kez daha gözler önüne sermektedir.

FULL TEXTS BOOK

Yapay zekâ sistemlerinin tarafsızlık maskesi altında yürüttüğü söylemsel yönlendirme, yalnızca bilgiye erişim biçimlerini değil, aynı zamanda bireylerin çatışmalara dair algılarını, tutumlarını ve eylem repertuarlarını da doğrudan etkilemektedir. Bu bağlamda Grok'un yanıtlarında öne çıkan ideolojik örüntüler, dijital hegemonyanın yeni biçimlerine işaret etmekte ve yapay zekânın çağdaş propaganda süreçlerinde aktif bir aktör haline geldiğini göstermektedir.

Araştırma bulguları, yapay zekâ tabanlı iletişim teknolojilerinin yalnızca teknik değil, aynı zamanda etik, siyasal ve kültürel sorumluluklarla değerlendirilmesi gerektiğini ortaya koymuştur. Özellikle çatışma, hak ihlalleri ve insanlık dramlarının yaşandığı hassas konularda, bu tür sistemlerin söylem üretme gücü göz ardı edilemeyecek kadar etkilidir. Dolayısıyla, bu teknolojilerin şeffaf, denetlenebilir ve çok aktörlü bir etik çerçeve içinde yeniden yapılandırılması, dijital çağın en temel demokrasi meselelerinden biri olarak karşımıza çıkmaktadır.

Sonuç olarak, bu çalışma; Grok örneği üzerinden, yapay zekâ sistemlerinin tarafsız bilgi sağlayıcılar olarak değil, belirli ideolojik ve politik zeminlerde şekillenen dijital söylem aktörleri olarak değerlendirilmesi gerektiğini vurgulamaktadır. Gelecekte yapılacak çalışmalar, farklı yapay zekâ sistemlerinin çeşitli toplumsal meselelerdeki söylemlerini karşılaştırmalı olarak ele alarak, dijital çağın hegemonik bilgi üretim yapılarının daha kapsamlı biçimde anlaşılmasına katkı sunabilir.

KAYNAKÇA

Arslan, K. (2017). Eğitimde yapay zekâ ve uygulamaları. *Batı Anadolu Eğitim Bilimleri Dergisi*, *11*(1), 71–88.

Avşar, B. (2025). Yapay zekâ ve dezenformasyon: OpenAI raporu doğrultusunda küresel dezenformasyon kampanyalarının analizi. *TRT Akademi Dergisi*, *10*(23), 208–236.

Ayhan, A., & Erken, G. (2018). Siyasal iletişimde dezenformasyon ve sosyal medya: Bir doğrulama platformu olarak Teyit.org. *Akdeniz Üniversitesi İletişim Fakültesi Dergisi (AKİL), (30),* 201–223.

Bekar, S. (2024). Yeni medyada yapay zekâ aracılığıyla dezenformasyon oluşturulması: Trump örneği. *USOBED Uluslararası Batı Karadeniz Sosyal ve Beşerî Bilimler Dergisi*, *8*(1), 40–66. https://doi.org/10.46452/baksoder.1448805

Durmuş, A. (2023). Sosyal medyada propaganda: Rusya-Ukrayna Savaşı örneği. *AJIT-E: Academic Journal of Information Technology*, *14*(52), 41–69. https://doi.org/10.5824/ajite.2023.01.003.x

Elitaş, T. (2022). Dijital manipülasyon 'deepfake' teknolojisi ve olmayanın inandırıcılığı. *Hatay Mustafa Kemal Üniversitesi Sosyal Bilimler Enstitüsü Dergisi*, *19*(49), 113–128.

Güzey, C., Çakır, O., Athar, M. H., Yurdaöz, E., & Saad, S. (2023). Eğitimde yapay zeka konusunda yapılmış çalışmaların içerik analizi. *Bilgi ve İletişim Teknolojileri Dergisi*, *5*(1), 66–77. https://doi.org/10.53694/bited.1060730

Güngör, A., & İmik Tanyıldızı, N. (2024). Reklam ve yapay zekâ ilişkisi: "Journal of Advertising" dergisindeki makalelerin bibliyometrik analizi. *Intermedia International e-Journal*, *11*(21), 104–126.

Kaplan, F. K. (2019). *Değişen dünya düzeni etkisinde Filistin-İsrail sorunu* (Yüksek lisans tezi, Ufuk Üniversitesi, Sosyal Bilimler Enstitüsü).

Karaköse, H. (2018). Filistin ve Kudüs meselesine genel bir bakış (XIX. yüzyılın ortasından XX. yüzyıl ortalarına kadar), Ahi Evran Üniversitesi Sosyal Bilimler Enstitüsü Dergisi, 4(2), 150-165.

Karakuş, M. (2021). Gelenekselden dijitale propaganda ve araçlarının dönüşümü. *Selçuk İletişim Dergisi*, *14*(1), 462–491. https://dergipark.org.tr/en/download/article-file/1225455

Karaman, M. L. (1996). Filistin. *Diyanet İslam Ansiklopedisi* (Cilt 13, s. 89–91, 103, 137–138, 140).

FULL TEXTS BOOK

Kişi, E. (2023). 1948'den bugüne, tarihi ve siyasi yönleri ile ''İsrail-Filistin sorunu''. *Anlambilim: MTÜ Sosyal ve Beşerî Bilimler Dergisi*, 3(1), 262–274.

Landis, J. R., & Koch, G. G. (1977). *The measurement of observer agreement for categorical data*. Biometrics, 33(1), 159–174. https://doi.org/10.2307/2529310

Özer, Ç. (2023). Uluslararası politika açısından Filistin sorunu. *Külliye*, [Türkiye Cumhuriyeti'nin 100. Yılı Özel Sayısı], 582–605. https://doi.org/10.48139/aybukulliye.1360984

Öztürk, H. (2021). Arşivler ve yapay zekâ. *Bilgi Yönetimi Dergisi*, *4*(2), 78–89. https://dergipark.org.tr/tr/pub/by/issue/67224/987197

Öztürk, K., & Şahin, M. E. (2018). Yapay sinir ağları ve yapay zekâ'ya genel bir bakış. *Takvim-i Vekayi*, *6*(2), 25–36. https://www.sloi.org/sloi-name-of-this-article

Rodgers, S. (2021). Themed issue introduction: Promises and perils of artificial intelligence and advertising. *Journal of Advertising*, *50*(1), 1–10. https://doi.org/10.1080/00913367.2020.1868233

Saleh, M. M. (2016). *Filistin meselesi: Tarihi arka plan ve güncel gelişmeler* (H. Demirhan, Çev.). Yurtdışındaki Filistinli Âlimler Derneği.

Tarakçı, N. (2009). *Gazze'nin gizemi*. TASAM – Türk Asya Stratejik Araştırmalar Merkezi. https://tasam.org/Files/Icerik/File/gazzenin_gizemi_d6220135-5007-43ac-8289-2c53aba840fc.pdf

Yiğit, H. (2024). Filistin-İsrail çatışmasında temel sorun alanları, bölgesel ve küresel aktörlerin soruna yaklaşımları. *Journal of Social, Humanities and Administrative Sciences*, *10*(2), 213–226. https://doi.org/10.5281/zenodo.10897934

FULL TEXTS BOOK

TATLI SU MİDYELERİNİN GENETİK POTANSİYELİYLE SU KALİTESİ YÖNETİMİNDE YENİLİKÇİ YAKLAŞIMLAR

INNOVATIVE APPROACHES TO WATER QUALITY MANAGEMENT THROUGH THE GENETIC POTENTIAL OF FRESHWATER MUSSELS

Hülya ŞEREFLİŞAN [1]

[1]*Prof. Dr., İskenderun Teknik Üniversitesi, Deniz Bilimleri ve Teknolojisi Fakültesi, Su Ürünleri Yetiştiriciliği Bölümü, İskenderun, Türkiye.*

[1]*ORCID ID: https://orcid.org/0000-0002-2510-3714*

Ahmet ALKAYA [2]

[2]*Dr., İskenderun Teknik Üniversitesi, Deniz Bilimleri ve Teknolojisi Fakültesi, Su Ürünleri Yetiştiriciliği Bölümü, İskenderun, Türkiye.*

[2]*ORCID ID: https://orcid.org/000-0003-2117-7799*

ÖZET

Tatlı su midyeleri, hem ekolojik işlevleri hem de çevresel stres faktörlerine duyarlılıkları nedeniyle su kalitesinin biyolojik izlenmesinde önemli göstergeler olarak öne çıkmaktadır. Bu derleme, tatlı su midyelerinin genetik özelliklerinin su kalitesi yönetimi bağlamında nasıl değerlendirilebileceğine odaklanmaktadır. Geleneksel, fiziksel ve kimyasal parametrelere dayalı su izleme yöntemlerinin sınırlılıkları göz önüne alındığında genetik temelli biyobelirteçlerin kullanımı, su ekosistemlerinin daha hassas ve gerçek zamanlı değerlendirilmesini mümkün kılmaktadır. Genom ve transkriptom düzeyinde yapılan güncel çalışmalar, midyelerin çevresel streslere karşı verdikleri tepkilerin moleküler temellerini ortaya koymakta ve bu organizmaların kirleticilere karşı geliştirdiği genetik adaptasyonları tanımlamaktadır. Özellikle stresle ilişkili genlerin ekspresyon düzeylerinin izlenmesi, kirlilik kaynaklarının belirlenmesi ve ekosistem sağlığının değerlendirilmesi açısından önemli bir yenilik olarak değerlendirilmektedir. Kamuya açık genetik veri tabanlarının gelişimiyle birlikte, tatlı su midyelerinde moleküler biyobelirteçlerin tanımlanması ve kullanımı hız kazanmıştır. Bu çerçevede, tatlı su midyelerinin genetik düzeydeki biyolojik yanıtları, sadece çevresel durumun değerlendirilmesine değil, aynı zamanda sürdürülebilir su yönetimi politikalarının geliştirilmesine de katkı sağlamaktadır. Bu derleme, çevresel izleme stratejilerinde genetik araçların entegre kullanımının gerekliliğini vurgulamakta ve tatlı su midyelerinin genetik özelliklerine dayalı yenilikçi yaklaşımların su kalitesi yönetiminde nasıl bir dönüşüm yaratabileceğini tartışmaktadır.

Anahtar Kelimeler: Midyeler, Biyobelirteçler, Su kalitesi, Genetik, İzleme.

ABSTRACT

Freshwater mussels are emerging as important bioindicators for water quality monitoring due to their significant ecological functions and high sensitivity to environmental stressors. This review focuses on how the genetic characteristics of freshwater mussels can be utilized within the framework of water quality management. Given the limitations of traditional monitoring methods that rely primarily on physical and chemical parameters, the use of genetic-based biomarkers offers a more precise and real-time assessment of aquatic ecosystems. Recent studies at the genomic and transcriptomic levels have revealed the molecular mechanisms underlying mussels responses to environmental stressors and have identified the genetic adaptations these organisms develop in response to pollutants. In particular, monitoring the expression levels of stress-related genes represents a novel approach for identifying

FULL TEXTS BOOK

pollution sources and assessing ecosystem health. With the expansion of publicly accessible genetic databases, the identification and application of molecular biomarkers in freshwater mussels have accelerated considerably. In this context, the biological responses of freshwater mussels at the genetic level contribute not only to the evaluation of environmental conditions but also to the development of sustainable water management policies. This review, emphasizes the necessity of integrating genetic tools into environmental monitoring strategies and discusses how innovative approaches based on the genetic traits of freshwater mussels may transform water quality management.

Keywords: Mussels, Biomarkers, Water quality, Genetics, Monitoring.

1. GİRİŞ

Su kaynakları, insanlık için yaşam kaynağı olmasının yanı sıra ekosistemlerin sürdürülebilirliğini sağlayan temel unsurlardır. Bu nedenle, su kalitesinin izlenmesi, korunması ve yönetilmesi, çevresel ile ekonomik açıdan büyük önem taşımaktadır (Banaduc ve ark., 2022). Su kirliliği, su kaynaklarının ekosistem sağlığını ve insanların yaşamını doğrudan tehdit eden bir faktördür. Bununla birlikte, geleneksel su kalitesi izleme yöntemleri, genellikle fiziksel ve kimyasal parametrelere dayanmakta olup, bu yöntemler çevresel değişimlerin gerçek zamanlı izlenmesinde sınırlı kalabilmektedir. Dolayısıyla, biyolojik izleme yöntemlerinin devreye girmesi, su kalitesinin daha doğru ve hızlı bir şekilde değerlendirilmesine olanak tanıyacaktır. Biyolojik göstergeler, özellikle su ekosistemlerinde yer alan organizmaların çevresel streslere yanıt olarak gösterdikleri tepkilerden yararlanmakta ve bu tepkiler genellikle ekosistem sağlığının erken uyarı sistemleri olarak kullanılabilmektedir (Vereycken & Aldridge, 2023). Tatlı su midyeleri, su ekosistemlerinde önemli ekolojik rol oynayan canlılardır. Ayrıca, çevresel değişikliklere duyarlı olmaları, onları su kalitesi izleme ve ekosistem sağlık değerlendirmeleri için ideal biyolojik göstergeler yapmaktadır (Vaughn, 2010; Scheder ve ark., 2015; Lummer ve ark., 2016). Midyeler, su kirliliği, oksijen seviyeleri, sıcaklık değişiklikleri gibi faktörlere verdikleri tepkilerle çevresel değişimlerin izlenmesinde görev yapmaktadır (Hartmann ve ark., 2016). Bu canlıların yaşam süreçlerinde gözlemlenen biyolojik tepkilerin genetik düzeyde de bir dizi değişikliğe yol açtığı bilinmektedir. Dolayısıyla, midyelerin çevresel stresler karşısında gösterdiği genetik adaptasyonlar ve değişiklikler, su ekosistemlerinin sağlığını değerlendirme açısından önemli ipuçları sunmaktadır (Vereycken & Aldridge, 2023).

Genetik bilimindeki ilerlemeler, su kalitesi yönetimi ve çevresel izleme stratejilerinde devrim yaratma potansiyeline sahiptir. Özellikle, genetik biyobelirteçlerin kullanımı su kaynaklarındaki kirleticilerin tespiti, biyolojik çeşitliliğin izlenmesi ve su ekosistemlerinin sürdürülebilir yönetimi konularında önemli yenilikler getirmiştir. Midyelerin genomları ve genetik çeşitlilikleri üzerine yapılan çalışmalar, bu organizmaların çevresel faktörlere nasıl yanıt verdiği hakkında derinlemesine bilgi edinmemize olanak tanımaktadır. Ayrıca, çift kabuklu hayvan genomlarında son dönemde yaşanan gelişmeler sonucunda, kamuya açık veri tabanlarında giderek artan miktarda diziye ulaşılabilmesiyle transkript biyobelirteçleri ortaya çıkmıştır (Tanguy ve ark., 2008; Zhang ve ark., 2012). Bununla beraber, midyelerin stresle ilişkili genleri, kirleticilere karşı gösterdikleri biyolojik tepkilerin moleküler düzeyde izlenmesini sağlayarak, su kalitesinin izlenmesinde önemli bir araç sunmaktadır (Arslan ve ark., 2017). Su kalitesini izlemek için genetik araçların kullanımı, yalnızca ekolojik verilerin toplandığı bir alanı değil, aynı zamanda kirliliğin kaynakları ve etkilerini daha iyi anlamamıza yardımcı olan bir bilimsel yaklaşımı da beraberinde getirmektedir. Tatlı su midyelerinin genetik özelliklerinden yararlanarak geliştirilen yeni yöntemler, su yönetimi süreçlerinde daha hassas, hızlı ve güvenilir bir izleme sağlayacak, çevresel yönetim stratejilerinin etkinliğini artıracaktır. (Bourlat ve ark., 2013; Lacroix ve ark., 2014). Bu çalışma, tatlı su midyelerinin genetik özelliklerinin, su kalitesi yönetimi için nasıl etkili bir şekilde kullanılabileceğini inceleyerek bu alandaki yenilikçi yaklaşımları ortaya koymaktadır.

FULL TEXTS BOOK

2. TATLI SU MİDYELERİNİN GENETİK ÖZELLİKLERİ

2.1. Genetik Çeşitlilik ve Popülasyon Yapısı

Genetik çeşitlilik, bir türün popülasyonlarındaki genetik varyasyonun bir ölçüsü ve ekosistemlerin dayanıklılığı ile doğrudan ilişkilidir. Tatlı su midyeleri, ekosistemlerde önemli ekolojik rol oynayan ve çevresel streslere duyarlı organizmalar olarak, genetik çeşitliliğin değerlendirilmesinde güçlü biyobelirteçlerdir (Bourlat ve ark., 2013). Midyelerin genetik çeşitliliği, popülasyonlar arası adaptasyon yeteneklerini, genetik sağlığını ve çevresel değişimlere karşı gösterdikleri direncini belirlemektedir. Özellikle, su kalitesi izlenmesinde midyelerin genetik çeşitliliği, bir ekosistemdeki stres düzeylerini ve bu streslere karşı verilen tepkileri anlamada kritik bir öneme sahiptir (Tanguy ve ark., 2008). Tatlı su midyeleri, farklı habitatlar ve ekosistemlerde yaşam sürdükleri için geniş bir genetik çeşitliliğe sahip olabilirler. Ancak, çevresel stres faktörlerinin etkisiyle bu çeşitlilik zaman içinde azalabilmektedir. Özellikle, kirlilik, sıcaklık değişimleri ve habitat tahribatı gibi faktörler, midye popülasyonlarının genetik çeşitliliğini tehdit etmektedir. Genetik çeşitlilik, aynı zamanda bir popülasyonun çevresel değişimlere karşı ne kadar esnek olduğunu ve bu değişimlere nasıl adapte olduğunu göstermektedir (Rogers ve ark., 2021). Son yıllarda, tatlı su midyelerinin genetik çeşitliliği, moleküler genetik yöntemlerle değerlendirilmektedir. Bu amaçla kullanılan başlıca teknikler arasında; *mikrosatellit belirteçleri, mitokondriyal DNA* (mtDNA) analizi ve *genomik genişleme* gibi yöntemler yer almaktadır (Wei ve ark., 2023). Mikrosatellitler, genetik çeşitliliği izlemek için yaygın olarak kullanılan genetik belirteçler olup, çok sayıda allel içermekte ve çevresel faktörlerden hızla etkilenmektedirler (Wei ve ark., 2023). Mitokondriyal DNA analizi ise türler arasındaki evrimsel ilişkileri ve popülasyon yapısını anlamada önemli bir araç olup, mtDNA anne tarafından miras alındığı için, genetik değişiklikler daha hızlı bir şekilde izlenmektedir (Andres ve ark., 2023). Bu genetik analizler, tatlı su midyelerinin farklı popülasyonlarında ki genetik çeşitliliği ve adaptasyon kapasitelerini belirlemede yardımcı olmaktadır. Bununla birlikte, genetik çeşitliliğin yüksek olduğu popülasyonlar, çevresel streslere daha dayanıklı olabilmekte ve bu nedenle bu türler su kalitesi izleme süreçlerinde kritik bir rol oynamaktadır (Ribeiro ve ark., 2019).

Popülasyon genetik analizi, ekosistemlerin sağlığını izlemek için güçlü bir yöntem olarak bilinmektedir. Tatlı su midyelerinin popülasyon yapıları, çevresel koşulların, habitat tahribatlarının, kirliliğin ve iklim değişikliğinin doğrudan bir yansımasıdır (Lodge, 2022). Genetik analizler, bu faktörlerin midye popülasyonları üzerindeki etkilerini değerlendirmek ve su ekosistemlerinin genel sağlık durumunu anlamak için kullanılmaktadır. Genetik çeşitliliğin korunması, ekosistemlerin sürdürülebilirliği için esastır. Dolayısıyla, genetik biyoçeşitlilik kaybı, ekosistemin işlevselliğini, habitat kalitesini ve su kalitesini olumsuz etkileyebilmektedir (Ribeiro ve ark., 2019). Dahası, belirli bir popülasyonun genetik çeşitliliğinde meydana gelen azalma, o popülasyonun çevresel streslere karşı daha hassas hale gelmesine yol açabilmektedir. Bu durumda, midye popülasyonlarının genetik analizleri, bir ekosistemin sağlık durumunu erken aşamalarda belirlemede yardımcı olacaktır (Wei ve ark., 2023). Genetik analizler aynı zamanda, türlerin genetik yapılarındaki değişikliklerin kirlilik ve diğer çevresel stres faktörleriyle nasıl ilişkilendirildiğini ortaya koymaktadır. Kirleticilere karşı genetik adaptasyon gösteren bireylerin varlığı, su kalitesinin iyileşmesinin potansiyel göstergesi olabilir. Ayrıca, popülasyonların genetik yapılarındaki değişiklikler, ekosistemlerdeki genel biyoçeşitlilik kaybı ile doğrudan ilişkilidir. Bu nedenle, popülasyon genetik analizleri, ekosistem sağlık değerlendirmelerinde ekosistemdeki türlerin genetik yapılarındaki değişiklikleri izlemek için önemli bir araçtır (Bourlat ve ark., 2013; Thomas ve ark., 2014). Ayrıca, tatlı su midyelerinin genetik çeşitliliği, sadece bireysel türlerin sağlığı için değil, aynı zamanda daha geniş bir ekosistem sağlığı için de kritik bir göstergedir. Genetik çeşitliliğin korunması, su ekosistemlerinin sürdürülebilirliğini sağlar ve ekosistemde midyeler gibi biyolojik göstergelerin kullanılması, çevresel izleme süreçlerinde yeni ve etkili yöntemler geliştirilmesine olanak tanır. Bu

FULL TEXTS BOOK

bağlamda, tatlı su midyelerinin genetik çeşitliliği ve popülasyon yapısı, su kalitesi yönetiminin daha hassas, hızlı ve doğru bir şekilde yapılmasına yardımcı olacak güçlü bir araçtır.

2.2. Stresle İlişkili Genler

Tatlı su midyeleri, çevresel stres faktörlerine karşı çeşitli biyolojik tepkiler gösteren canlılardır (Vereycken & Aldridge, 2023). Bu tepkiler, genetik düzeyde bir dizi moleküler mekanizma ile şekillenmektedir. Bilhassa, ağır metaller, toksik kimyasallar, sıcaklık değişimleri, pH değişiklikleri ve oksijen eksikliği gibi çevresel stres faktörleri, midyelerin genetik yapılarında değişikliklere neden olmakta (Bickham ve ark., 2000) ve bu değişiklikler, ekosistem sağlığı hakkında değerli bilgiler sunmaktadır (Ribeiro ve ark., 2019). Ağır metaller (örneğin, kurşun, cıva, kadmiyum) ve toksik kimyasallar, su ekosistemlerinde yaşayan organizmalar üzerinde ciddi biyolojik etkiler yaratmaktadır. Dolayısıyla, bu kirleticiler, hücresel yapıları bozmakta, enzimatik aktiviteleri inhibe etmekte ve DNA hasarına yol açmaktadır (Jha, 2008). Midyeler, bu zararlı maddelere karşı çeşitli biyolojik yanıtlar geliştirmiş ve bu yanıtlar genetik düzeyde belirli genlerin ekspresyonu ile modüle edilmiştir (Thomas ve ark., 2014). Midyelerde ağır metallerin ve toksik kimyasalların etkilerine yanıt olarak, **metal bağlayıcı proteinler** ve **antioksidant enzimler** gibi genlerin ekspresyonu artmaktadır. Bu genler, organizmanın toksik maddeleri bağlayarak, bunları hücrelerden uzaklaştırmayı veya etkilerini nötralize etmeyi hedeflemektedir (Franco ve ark., 2006;Varotto ve ark., 2013). Örneğin, **metalotionein** (MT) genleri, ağır metallerin bağlanmasında önemli bir rol oynar ve toksik metal iyonlarının zararlı etkilerini azaltır. Bu genin ekspresyonu, metal kirliliği arttıkça belirginleşmektedir. Bununla beraber, **Süperoksit dismutaz** (SOD), **glutatyon peroksidaz** (GPX) gibi antioksidant enzimlerin ekspresyonu da, çevresel oksidatif stresi azaltmaya yönelik bir mekanizma olarak aktif hale gelmektedir. Bu tür genetik yanıtların belirlenmesi, su ekosistemlerinde bulunan kirliliğin düzeyini ve etkisini izlemek için biyobelirteç olarak kullanılabilmektedir (Ribeiro ve ark., 2019). Bu nedenle, kirlilik kaynaklarına yönelik yapılan genetik analizler, hangi kirleticilerin midyeler üzerinde daha etkili olduğunu ve bu kirleticilere karşı hangi moleküler yanıtların geliştirilmesi gerektiğini anlamamıza yardımcı olmaktadır.

Oksijen eksikliği, tatlı su ekosistemlerinde özellikle yaz aylarında, sıcaklık artışı ve suyun oksijen taşıma kapasitesinin düşmesiyle daha sık görülmektedir. Oksijen seviyesi düştüğünde, organizmalar enerji üretimini oksijenli solunumdan anaerobik metabolizmaya kaydırmaktadır. Bu süreç, çeşitli genetik düzenlemeleri tetikleyerek oksijen seviyelerine duyarlı **HIF-1 (hipoksi-indüklenmiş faktör-1)** gibi transkripsiyon faktörleri düşük oksijen koşullarında aktive olur ve hücrelerin hayatta kalması için gerekli genlerin ekspresyonunu başlatılır (Fu ve ark., 2008). Bu sayede, tatlı su midyeleri oksijen eksikliğine karşı geliştirdikleri adaptasyonlarla bu tür streslere anında tepki vermektedirler. Hipoksi altında, metabolik yollar ve hücresel işlevler yeniden programlanarak, oksijenin sınırlı olduğu koşullarda hayatta kalmaya yönelik stratejiler geliştirilir. Dolayısıyla, hipoksi yanıtı, genetik düzeyde **glikoliz enzimleri**, **anaerobik proteinler** ve **özellikle oksijen taşıyıcı proteinler** gibi moleküllerin ekspresyonunu artırabilmektedir (Montúfar-Romero ve ark., 2024).

Gen ekspresyon analizlerinin, organizmanın çevresel streslere nasıl yanıt verdiğini anlamada güçlü bir araç olduğu bilinmektedir. Midyelerin genetik düzeyde verdiği tepkiler, çeşitli biyolojik ve çevresel faktörlerin etkilerini doğrudan yansıtmaktadır. Bu bağlamda, gen ekspresyonunun ölçülmesi, su ekosistemlerinde meydana gelen kirlenme, oksijen eksikliği ve diğer stres faktörlerinin erken teşhisi için önemli bir metodolojik yaklaşımdır (Bourlat ve ark., 2013). Gen ekspresyon analizleri, genellikle **qPCR (kuantitatif polimeraz zincir reaksiyonu)**, **RNA-Seq** ve **microarray teknolojileri** gibi ileri düzey moleküler biyoloji teknikleriyle gerçekleştirilmektedir (Suárez-Ulloa ve ark., 2013; Lacroix ve ark., 2014). Bu yöntemler, midyelerin genetik materyalini inceleyerek, stres yanıtı gösteren genlerin ne zaman ve ne kadar ifade edildiğini belirlemeye olanak tanımaktadır. Bu tür analizler, bir ekosistemdeki kirliliğin veya diğer stres faktörlerinin etkilerini saptamak için oldukça faydalıdır. Ekspresyon seviyelerindeki değişiklikler, su kirliliği gibi çevresel faktörlere dair erken uyarı sinyalleri

FULL TEXTS BOOK

sağlayabilmektedir. Ayrıca, bu tür biyolojik yanıtların izlenmesi, su kalitesi izleme stratejilerini daha duyarlı ve spesifik hale getirecektir (Ribeiro ve ark., 2019).

2.3. Mitokondriyal DNA (mtDNA) ve Filogenetik Analizler

Mitokondriyal DNA, hücrelerin enerji üretim merkezlerinde bulunan ve anne tarafından miras alınan bir genetik materyaldir. Tatlı su midyelerinin filogenetik ilişkilerini incelemek ve ekosistem bağlantılarını anlamak için mtDNA, özellikle yüksek mutasyon hızına sahip olması ve türler arası evrimsel ilişkileri net bir şekilde ortaya koyma kapasitesi nedeniyle önemli bir araçtır (Jones ve ark., 2015).

Tatlı Su Midyelerinin Filogenetik İlişkilerinin Belirlenmesi

Filogenetik analizler, organizmaların evrimsel ilişkilerini inceleyerek, türlerin geçmişteki ortak atalarından nasıl ayrıldığını ve birbirlerine nasıl yakın olduklarını belirlemeyi amaçlamaktadır. Tatlı su midyeleri, genetik çeşitlilik ve adaptasyon yetenekleri açısından son derece zengin bir grup oluşturmaktadır. Bu nedenle, filogenetik ilişkilerin belirlenmesi tatlı su ekosistemlerinin evrimsel geçmişi ve biyoçeşitliliğinin anlaşılması açısından kritik bir öneme sahiptir (Froufe ve ark., 2020). Filogenetik analizlerde, mtDNA genetik belirteçleri sıklıkla kullanılmaktadır. Bunun sebebi, mitokondriyal genetik materyalin, türler arasında farklılaşmayı hızlı bir şekilde yansıtarak, kısa süreli evrimsel değişimlerin takip edilmesini sağlamasıdır. *Cytochrome c oksidaz alt ünitesi 1 (COI), 16S rRNA* ve *ATP6* gibi mitokondriyal gen bölgeleri, filogenetik çalışmalar için yaygın olarak kullanılan işaretçiler arasında yer almaktadır. Bu gen bölgeleri, tatlı su midyelerinin türler arası farklılıklarını tespit etmede ve bu türlerin evrimsel ilişkilerini izlemekte oldukça etkilidir (Jones ve ark., 2015). Tatlı su midyeleri, farklı coğrafi bölgelerde ve çeşitli su ekosistemlerinde yaşam sürmektedir. Bu nedenle, coğrafi izolasyon, genetik farklılaşma ve tür çeşitliliği gibi unsurlar, tatlı su midyelerinin filogenetik yapılarını anlamada önemli bilgiler sunmaktadır. Filogenetik analizler sayesinde, türlerin kökeni, popülasyon yapıları, adaptasyon süreçleri ve ekosistemler arasındaki genetik akışlar daha iyi bir şekilde anlaşılabilmektedir. Ayrıca, türler arasındaki genetik ilişkiler, su ekosistemlerinin biyoçeşitliliği ve sağlık durumu hakkında önemli ipuçları vermektedir (Lopes-Lima ve ark., 2021).

Mitokondriyal DNA İşaretçilerinin Ekosistem Bağlantılarını Anlamada Kullanımı

Mitokondriyal DNA işaretçileri, sadece türler arasındaki filogenetik ilişkilerin anlaşılması için değil, aynı zamanda tatlı su ekosistemlerinin ekolojik yapılarının ve ekosistem bağlantılarının incelenmesinde de kullanılabilmektedir. Mitokondriyal DNA'nın özellikleri, türler arasındaki genetik farklılıkları çok hassas bir şekilde tespit edebilmesini sağlamaktır. Bu durum, özellikle tatlı su ekosistemlerinde türler arasında olan ekosistem bağlantılarını, biyolojik etkileşimleri ve genetik akışları anlamada faydalı olmaktadır (Andres ve ark., 2023). Tatlı su midyelerinin ekosistemler arasındaki bağlantılarını anlamak için mtDNA, türlerin yaşadığı su havzalarındaki genetik akışı izlemeye olanak tanımaktadır. Örneğin, bir midye popülasyonunun iki farklı nehirde bulunan alt popülasyonları arasında genetik farklar varsa, bu farklılıklar, türlerin su havzaları arasında hareketliliğini veya izole kalıp kalmadıklarını gösterebilmektedir. Ayrıca, tatlı su ekosistemlerinde iklim değişikliklerinin, habitat tahribatının ve su kirliliğinin etkilerini izlemek için mtDNA işaretçileri kullanılmaktadır (Bucholz ve ark., 2022).

Mitokondriyal DNA'nın ekosistem bağlantıları ile olan ilişkisi, su ekosistemlerinde biyolojik çeşitliliğin korunmasına da katkıda bulunmaktadır. Örneğin, ekosistem sağlığına ilişkin bozulmalar, midyelerin genetik yapılarında belirgin değişikliklere yol açabilir. Bu değişiklikler, türlerin popülasyon yapılarındaki farklılıkları ve çevresel streslere karşı nasıl adapte olduklarını yansıtabilir. Bu bağlamda, mitokondriyal DNA, çevresel değişimlerin ve kirliliğin midye popülasyonları üzerindeki etkilerini anlamada etkili bir izleme aracı olacaktır (Bourlat ve ark., 2013). Ekosistem bağlantılarını belirlemek için mtDNA'nın kullanılması, aynı zamanda tatlı su midyelerinin ekolojik rolünü ve çevresel değişimlere verdikleri genetik yanıtları anlamamıza yardımcı olmaktadır. Bu tür çalışmalar, midyelerin yaşadığı su

FULL TEXTS BOOK

havzalarında ekosistem hizmetlerinin nasıl etkilendiğini ve türlerin koruma gereksinimlerini belirlememize olanak tanıyacaktır. Mitokondriyal DNA analizi, ayrıca türlerin popülasyonlar arası genetik akışını ve adaptasyon süreçlerini izleyerek, ekosistemler arasındaki biyolojik etkileşimleri daha ayrıntılı bir şekilde anlamamızı sağlayacaktır (Schwarz & Roe, 2022).

3. GENETİK BİYOBELİRTEÇLERİN KULLANIMI

Su kalitesi yönetiminde geleneksel yöntemler genellikle fiziksel, kimyasal ve biyolojik parametrelerin izlenmesine dayanırken, genetik biyobelirteçlerin kullanımı son yıllarda bu süreçlere önemli katkılar sağlamıştır. Özellikle tatlı su midyelerinin genetik özelliklerinden yararlanarak, çevresel değişiklikler ve su kirliliği gibi faktörlere karşı duyarlı göstergeler elde edilmektedir. Genetik biyobelirteçlerin kullanımı, su ekosistemlerinin daha hassas ve etkin bir şekilde izlenmesine olanak tanırken, çevresel değişimlere karşı hızlı yanıt verme kapasitesini arttırmaktadır (Andres ve ark., 2023). Bu bağlamda, çevresel DNA (eDNA) teknolojisi ve genetik biyobelirteçlerin etkinliği üzerine olan uygulamalar önemli bir araştırma alanıdır.

3.1. Çevresel DNA (eDNA) Teknolojisiyle Su Kolonunda Bulunan Midyelerin Genetik Materyalinin İzlenmesi

Çevresel DNA, canlı organizmaların çevrelerine bıraktıkları genetik materyalin (özellikle su, toprak ve hava gibi ortamlarda bulunan DNA) izlenmesi yoluyla ekosistemlerin analiz edilmesini sağlayan bir yöntemdir (Fediajevaite ve ark., 2021). eDNA, özellikle su ekosistemlerinde, organizmaların doğrudan gözlemlenmesine gerek kalmadan, su içinde bulunan genetik materyalin tespiti yoluyla suyun biyolojik içeriği hakkında bilgi edinilmesine olanak tanımaktadır. Tatlı su midyelerinin eDNA'sı, su kolonunda bulunan bu organizmaların genetik izlerinin analiz edilmesiyle elde edilebilmektedir (Bourlat ve ark., 2013).

Tatlı su midyelerinin eDNA'sı, su ekosistemlerinin izlenmesinde kullanıldığında, midyelerin mevcut olduğu bölgelerdeki biyoçeşitliliğin hızlı bir şekilde haritalanmasına yardımcı olabilir. Ayrıca, midyelerin genetik materyali, çevresel değişiklikler karşısında nasıl tepki verdiklerine dair izler taşımaktadır (Ruiz-Ramos ve ark., 2024). Midyelerin eDNA'sının analizi ile birlikte, popülasyonların varlığı, dağılımı ve tür çeşitliliğini anlamanın yanı sıra, su kirliliği ve habitat kaybı gibi faktörlerin bu organizmalar üzerindeki etkileri de tespit edilecektir. Midyeler gibi ekosistem için önemli türlerin eDNA'sı, çevresel değişikliklerin belirlenmesinde erken uyarı sistemleri olarak kullanılmaktadır (Preece ve ark., 2021). Örneğin, midyelerin genetik izleri, suyun kirlenme düzeyine göre genetik çeşitliliğin azaldığını veya türlerin farklı stres faktörlerine karşı adaptasyon gösterdiğini ortaya koyabilmektedir. eDNA teknolojisi, su kalitesini değerlendirmek için özellikle düşük yoğunluklu, zorlu şartlarda yaşayan organizmaların izlenmesinde oldukça etkilidir. Tatlı su midyeleri gibi, genellikle dip bölgelere yerleşmiş ve su kolonunun belirli katmanlarında bulunan organizmalar, doğrudan gözlemlerle izlenmesi güç organizmalardır. eDNA, bu tür organizmaların varlığını ya da yokluğunu tespit etmek için önemli bir genetik biyobelirteç sağlayacaktır (Togaki ve ark., 2020).

Genetik Biyobelirteçlerin Kirlilik Kaynaklarını ve Su Kalitesini Tespit Etmedeki Etkinliği

Genetik biyobelirteçler, çevresel stres faktörlerine yanıt olarak organizmaların genetik yapılarında meydana gelen değişiklikleri izleyerek, kirlilik ve diğer çevresel faktörlerin su kalitesi üzerindeki etkilerini ortaya koymaktadır (Ribeiro ve ark., 2019). Tatlı su midyelerinin çevresel değişimlere karşı geliştirdikleri genetik yanıtlar, suyun kalite durumu hakkında doğrudan bilgi sağlayabilir. Özellikle stresle ilişkili genlerin ekspresyonu, suyun kirlilik seviyesini, toksik bileşenlerin yoğunluğunu ve ekosistemdeki biyolojik değişimlerin izlerini göstermektedir (Bickham ve ark., 2000). Genetik biyobelirteçlerin kullanımı, kirlilik kaynaklarının tespiti için oldukça hassas bir yöntem sunarak, kirliliğin etkilediği organizmaların genetik değişimleri su ekosistemlerindeki biyoçeşitliliği ve ekolojik sağlığı hızla tespit etmeye yardımcı olabilmektedir. Tatlı su midyelerinin bu genetik işaretçileri,

FULL TEXTS BOOK

kirleticilerin türü ve yoğunluğuna dair değerli veriler ortaya koyup, su kalitesinin izlenmesine yönelik daha etkin bir yöntem sağlamaktadır. Ağır metaller, toksik kimyasallar, düşük oksijen seviyeleri ve diğer kirletici faktörler, midyelerin genetik yapılarında belirgin değişikliklere neden olmaktadır. Bu değişiklikler, genetik biyobelirteçler aracılığıyla tespit edilerek, su kalitesinin kötüleşip kötüleşmediği konusunda bilgi verebilmektedir (Thomas ve ark., 2014). Genetik biyobelirteçler, aynı zamanda su kalitesinin zaman içindeki değişimlerini izlemek ve su ekosistemlerinin gelecekteki durumlarını tahmin etmek için de kullanılmaktadır. Ekspresyon analizi yapılan genetik biyobelirteçler, bir bölgedeki suyun kalitesinin yıllık veya mevsimsel değişimlerini gözler önüne serebilmektedir. Bu sayede, su yönetim stratejilerinin daha verimli ve hedeflenmiş bir şekilde geliştirilmesine olanak tanınmaktadır (Lacroix ve ark., 2014).

3.2. CRISPR ve Gen Düzenleme Teknolojileri

Genetik mühendislik alanındaki en devrimci yeniliklerden biri olan CRISPR-Cas9 ve diğer gen düzenleme teknolojileri, tatlı su midyeleri gibi organizmaların çevresel streslere yanıtlarını optimize etmek ve ekosistem izleme için biyobelirteç üretimi amacıyla büyük bir potansiyel sunmaktadır (Machuca-Sepúlveda ve ark., 2023). Gen düzenleme, organizmaların genetik materyalinde istenilen değişiklikleri yaparak, belirli genetik özellikleri hedefleyip düzenlemeye olanak tanımaktadır. CRISPR-Cas9, bu alandaki en yaygın kullanılan araçlardan biridir ve çok spesifik gen düzenlemeleri yapabilmesi nedeniyle büyük bir uygulama alanına sahiptir (Durán-Vinet ve ark., 2021). Tatlı su midyeleri gibi su ekosistemlerinde önemli rol oynayan organizmalar, çevresel stres faktörlerine karşı oldukça hassastır. Su kirliliği, ağır metaller, düşük oksijen seviyeleri ve sıcaklık değişiklikleri gibi stres faktörleri, midyelerin hayatta kalma becerilerini ciddi şekilde etkileyebilmektedir (Vereycken & Aldridge, 2023).

CRISPR teknolojisi ile, midyelerin çevresel streslere karşı dayanıklılığını artırmak amacıyla hedeflenmiş genetik modifikasyonlar yapılmaktadır. Örneğin, ağır metal kirliliği gibi stres faktörlerine karşı direnç sağlayan genetik yolları düzenleyerek, midyelerin toksik maddelere karşı daha dayanıklı hale gelmesi sağlanabilir. Bu tür bir genetik düzenleme, midyelerin hayatta kalma oranlarını artırabilir ve onları su kalitesinin izlenmesinde daha güvenilir biyolojik göstergeler haline getirebilmektedir. Ayrıca, midyelerin çevresel streslere karşı gösterdiği genetik tepkiler, CRISPR teknolojisi kullanılarak optimize edilebilmektedir. Örneğin, oksidatif stres, ağır metal toksisitesi ve düşük oksijen seviyeleri gibi çevresel koşullara karşı hayatta kalma mekanizmalarını düzenleyen genlerin ekspresyonu üzerinde müdahaleler yapılabilmektedir. Böylece, bu organizmaların çevresel faktörlere daha hızlı ve etkin bir şekilde tepki vermesi sağlanacaktır. CRISPR ile yapılan düzenlemeler, organizmanın genetik yapısında uzun vadeli değişiklikler yaparak, daha dayanıklı ve adaptif popülasyonlar oluşturulmasına olanak tanımaktadır (Machuca-Sepúlveda ve ark., 2023).

Çevreye Duyarlı Biyobelirteç Üretimi için Gen Düzenlemenin Potansiyeli

Biyobelirteçler, bir organizmanın çevresel değişimlere verdiği yanıtları ölçmek için kullanılan biyolojik işaretçilerdir. Çevreye duyarlı biyobelirteçler, çevresel stres faktörlerine karşı organizmaların verdiği genetik tepkileri izlemek için önemli araçlardır. Tatlı su midyeleri gibi su ekosistemlerinde yaşayan organizmalar, çevreye duyarlı biyobelirteç gelişimi için mükemmel adaylardır çünkü bu organizmalar, çevresel değişikliklere doğrudan ve hızlı bir şekilde tepki vermektedirler. CRISPR ve gen düzenleme teknolojileri, bu biyobelirteç üretimini daha hassas ve kontrollü hale getirecektir (Ekelund Ugge ve ark., 2020). Genetik düzenleme, çevreye duyarlı biyobelirteçlerin üretimi için bir dizi potansiyel sağlamaktadır. Midyelerin genetik yapısında gerçekleşen düzenlemelerle, belirli stres faktörlerine karşı duyarlı belirteçler üretilebilmektedir. Örneğin, ağır metal kirliliğine karşı duyarlı olan bir biyobelirteç üretimi için, midyelerdeki metalotionein gibi stresle ilişkili genler hedeflenebilir. Ayrıca, oksidatif stresin tespiti için süperoksit dismutaz (SOD) veya glutatyon peroksidaz gibi enzimlerin ekspresyonu üzerine yapılan genetik modifikasyonlar, çevresel stresleri erken aşamalarda tespit etmeye olanak

FULL TEXTS BOOK

tanıyacaktır (Bourlat ve ark., 2013). Bu belirteçler, sadece midyelerin çevresel stres yanıtlarını izlemek için değil, aynı zamanda su ekosistemlerindeki genel sağlık durumunu izlemek için de kullanılmaktadır. Çevreye duyarlı biyobelirteçler, suyun kirlenme seviyelerini, toksik madde konsantrasyonlarını, oksijen seviyelerindeki değişiklikleri ve diğer çevresel faktörleri hızlı ve hassas bir şekilde izlemek için etkin araçlar sağlar. Genetik olarak modifiye edilmiş midyeler, bu biyobelirteçlerin üretimi konusunda önemli bir rol oynamaktadır. Bunun nedeni, bu organizmaların çevresel değişimlere yanıtları çok daha belirgin ve izlenebilir olmasıdır (Lacroix ve ark., 2014).

3.3. Genomik ve Transkriptomik Araştırmalar

Genomik ve transkriptomik araştırmalar, organizmaların çevresel streslere verdikleri yanıtları ve bu yanıtların moleküler düzeyde nasıl gerçekleştiğini anlamada büyük bir rol oynamaktadır. Tatlı su midyeleri gibi su ekosistemlerinde önemli bir yer tutan canlıların, bu tür araştırmalarla çevresel değişimlere ve stres faktörlerine karşı nasıl adapte olduklarını daha iyi bir şekilde ortaya koymaktadır. Genomik analizler, organizmaların tüm genetik materyalini kapsamlı bir şekilde inceleyen araştırmalar olup, tatlı su midyeleri gibi ekosistem mühendisleri, çevresel faktörlere ve stres kaynaklarına karşı çeşitli genetik adaptasyonlar geliştirmiştir. Bu adaptasyonların moleküler düzeyde anlaşılması, tatlı su midyelerinin hayatta kalmalarını nasıl sağladıklarını ve ekosistemle olan etkileşimlerini nasıl optimize ettiklerini anlamamıza yardımcı olacaktır (Bourlat ve ark., 2013). Gelişmiş genomik teknolojiler, midyelerin tüm genomunun dizilenmesine olanak tanımakta, bu da bu organizmaların çevresel streslere nasıl tepki verdiğine dair daha derin bir anlayış sağlamaktadır. Örneğin, ağır metaller, toksik kimyasallar ve değişen sıcaklıklar gibi çevresel stres faktörlerine karşı midyelerin genomlarında hangi genetik değişikliklerin meydana geldiği belirlenmektedir. Ayrıca, adaptasyon süreçlerini tetikleyen ve hayatta kalmaya yardımcı olan genetik yollar keşfedilerek, midyelerin çevresel değişimlere karşı direncinin arttırılması ve sürdürülebilir popülasyonlar oluşturulması konusunda yol gösterici olacaktır (Xu ve ark., 2016; Gallardo-Escárate ve ark., 2023). Bunun yanı sıra, genomik analizler tatlı su midyelerinin belirli stres faktörlerine karşı geliştirdiği dirençli fenotiplerin moleküler temelini ortaya koymak için kullanılabilmektedir. Dahası, genetik çeşitlilik ve popülasyon yapıları hakkında daha fazla bilgi edinmek, su kalitesini izlemek ve ekosistem sağlığını değerlendirmek için oldukça faydalıdır. Bu tür analizler, midyelerin potansiyel adaptasyon yeteneklerini belirleyerek, su yönetimi ve koruma stratejilerine önemli katkılarda bulunacaktır.

Transkriptom Analizlerinin Çevresel Stres Yanıtlarına Yönelik Gen Düzenlemelerini Belirlemedeki Rolü

Transkriptomik analizler, bir organizmanın genetik materyalinin tüm RNA moleküllerinin incelenmesidir ve bu, organizmanın genetik ekspresyon düzeylerini anlamamıza yardımcı olmaktadır. Çevresel stresler organizmaların genetik ekspresyonunu doğrudan etkiler ve transkriptomik veriler bu değişiklikleri tespit etmemize olanak tanır (Varotto ve ark., 2013). Tatlı su midyeleri gibi su ekosistemlerinde bulunan canlılar, çevresel streslere karşı genetik yanıtlarını hızlı bir şekilde düzenleyebilmektedir. Bu nedenle, transkriptomik analizler, çevresel streslere karşı midyelerin verdiği moleküler tepkileri anlamada kritik bir araçtır (Xu ve ark., 2016). Özellikle, ağır metaller, oksijen eksikliği, sıcaklık dalgalanmaları ve asidifikasyon gibi stres faktörlerinin midyelerin genetik ekspresyonu üzerindeki etkileri transkriptomik analizlerle izlenebilmektedir. Bu analizler, stres altında olan midyelerde hangi genlerin daha fazla ekspresyon gösterdiğini, hangilerinin baskılandığını ve bu genetik değişimlerin organizmanın hayatta kalması için ne kadar kritik olduğunu belirlemek için kullanılmaktadır (Gallardo-Escárate ve ark., 2023). Bu tür çalışmalar, çevresel streslere karşı gösterilen adaptasyonların moleküler mekanizmalarını aydınlatmaya yardımcı olabilir. Ayrıca, transkriptomik analizler sayesinde, çevresel streslere karşı geliştirilmiş adaptasyon mekanizmalarını destekleyen genetik yolların keşfi mümkün olup, tatlı su midyelerinin bu stres faktörlerine karşı daha dayanıklı hale getirilmesi sağlanabilir. İlaveten, gen düzenlemeleri stresle ilişkili belirli genlerin daha etkin hale

FULL TEXTS BOOK

getirilmesi ve çevresel değişimlere daha hızlı uyum sağlanması içinde kullanılmaktadır (Bourlat ve ark., 2013).

3.4. Kirlilik İzleme ve Erken Uyarı Sistemleri

Su kirliliği, ekosistemlerin sağlığı ve sürdürülebilir su kaynaklarının korunması açısından ciddi bir tehdit oluşturmaktadır. Tatlı su ekosistemlerinde su kalitesini izlemek ve potansiyel kirleticilerin erken tespitini sağlamak, ekosistemleri korumak ve su yönetimi stratejilerini geliştirmek için kritik öneme sahiptir (Vereycken & Aldridge, 2023). Genetik biyobelirteçler, çevresel değişimlere karşı organizmaların genetik düzeyde verdiği yanıtları izleyerek kirliliği ölçmek için kullanılmaktadır. Bu biyobelirteçler, organizmaların çevresel stres faktörlerine karşı gösterdiği genetik tepkilerin göstergeleridir ve su kirliliğinin, özellikle ağır metal ve toksik kimyasallar gibi kirleticilerin etkilerini anlamada etkili araçlardır (Thomas ve ark., 2014). Genetik biyobelirteçlerin kullanımı, kirliliğin türünü ve seviyesini belirlemeye yardımcı olmaktadır. Örneğin, ağır metal kirliliği durumunda midyelerin genetik materyalinde belirli stresle ilişkili genlerin ekspresyonu artabilmektedir. Dolayısıyla, bu genler toksik kimyasalların organizmalar üzerindeki etkilerine karşı yanıt veren proteinlerin üretimini teşvik etmektedir. Genetik biyobelirteçlerin kullanımı, suyun kirlenme düzeylerinin izlenmesi ve kirliliğin kaynağının tespiti için daha hassas ve erken aşamalarda müdahale imkanı sağlamaktadır (Xu ve ark., 2016). Gelişmiş genetik teknolojiler, çevresel DNA (eDNA) kullanımı gibi yöntemlerle, su örneklerinde mevcut olan midye gibi organizmaların genetik materyali üzerinden kirliliğin izlenmesini sağlamaktadır. eDNA analizi ile, suyun içindeki mikroorganizmaların genetik materyali tespit edilebilir ve kirleticilere karşı verilen genetik tepkilerin izlenmesi için oldukça hassas bir yöntem sunmaktadır. Bu yaklaşım, suyun çeşitli bölümlerinde kirleticilere karşı verilen genetik yanıtları hızlı bir şekilde izlemeye olanak tanır ve kirliliğin derecesi hakkında daha doğru bilgi sağlar (Preece ve ark., 2021).

Endüstriyel ve Tarımsal Kaynaklı Kirleticilere Karşı Erken Uyarı Sistemlerinin Geliştirilmesi

Endüstriyel ve tarımsal faaliyetler, tatlı su ekosistemlerini ciddi şekilde tehdit eden başlıca kirleticiler arasında yer almaktadır. Endüstriyel atıklar, ağır metaller, kimyasal çözücüler ve toksik gazlar gibi kirleticilerle su kaynakları kirlenmektedir. Tarımsal faaliyetler ise, özellikle pestisitler, gübreler ve nitrifikasyon süreçleriyle su kirliliğine yol açmaktadır. Bu kirleticiler su ekosistemlerini hızla bozar ve ekosistem sağlığına zarar vermektedir (Merdun & Çınar, 2008). Bu nedenle, erken uyarı sistemlerinin geliştirilmesi, su kirliliğinin önlenmesi ve su kalitesinin korunması açısından kritik öneme sahiptir. Genetik biyobelirteçler, çevresel streslere karşı duyarlı organizmaların genetik tepkilerini izleyerek, endüstriyel ve tarımsal kaynaklı kirleticilere karşı erken uyarı sağlayabilmektedir. Genetik biyobelirteçler sayesinde, çevresel değişikliklere, özellikle kimyasal kirliliğe, hızla yanıt veren organizmalar belirlenebilir ve bu organizmaların genetik tepkileri izlenerek, kirliliğin erken aşamalarda tespiti sağlanabilir (Lacroix ve ark., 2014).

Endüstriyel ve tarımsal kirleticilere karşı geliştirilmiş bir erken uyarı sistemi, tatlı su ekosistemlerinin sağlık durumunu sürekli izleyen, hızlı tepki verebilen bir mekanizma oluşturabilir. Bu sistemler, düzenli olarak çevresel biyobelirteçlerin ölçülmesini ve analiz edilmesini içermektedir. Örneğin, ağır metallerin ve pestisitlerin çevresel etkilerini izlemek için kullanılan biyoteknolojik yöntemler, organizmaların genomik ve transkriptomik analizleriyle birleştirilerek, potansiyel kirletici maddelerin çevredeki etkilerine dair erken uyarılar verebilmektedir (Preece ve ark., 2021). Biyolojik izleme sistemlerinin kurulması, su kalitesi yönetimi ve kirliliğe karşı alınacak önlemler açısından önemlidir. Bu tür sistemler, kirleticilerin yoğun olduğu bölgeleri belirleyerek, kirliliğin yayılmasını engelleyen müdahaleler için temel sağlar. Ayrıca, bu izleme sistemlerinin dijital platformlarla entegre edilmesi, su kalitesinin daha geniş çapta izlenmesini ve kirliliğin anında tespit edilmesini sağlayarak, çevresel krizlerin önlenmesine yardımcı olabilir.

FULL TEXTS BOOK

3.5. Su Ekosistemlerinin Restorasyonu

Su ekosistemlerinin restorasyonu, özellikle kirlenmiş, bozulmuş veya sürdürülebilir olmayan şekilde yönetilen su kaynaklarının yeniden iyileştirilmesi sürecini ifade etmektedir. Bu süreç, hem biyolojik çeşitliliğin korunmasını hem de ekosistem hizmetlerinin sürdürülebilirliğini sağlamayı amaçlamaktadır. Tatlı su ekosistemlerinde, özellikle tehdit altındaki türlerin korunması ve su kalitesinin iyileştirilmesi büyük önem taşır. Genetik veriler, bu restorasyon süreçlerinde önemli bir rol oynamakta ve su ekosistemlerinin yeniden yapılandırılmasında en etkili stratejilerin belirlenmesine yardımcı olabilmektedir (Mijangos ve ark., 2015). Ayrıca, tehdit altındaki midye popülasyonlarının genetik rehabilitasyonu, bu ekosistemlerin sağlıklı bir şekilde geri kazanılmasında önemli bir bileşendir (Lavictoire & West, 2024). Su ekosistemlerinin restorasyonu için genetik verilerin kullanılması, biyolojik çeşitliliği koruma ve ekosistem işlevlerini iyileştirme açısından büyük potansiyele sahiptir. Su ekosistemlerinin sağlık durumunu değerlendiren ve onarıma yönelik stratejiler geliştiren genetik analizler, ekosistem üzerinde pozitif etki yaratacak en uygun türlerin ve popülasyonların belirlenmesine olanak tanımaktadır (Bourlat ve ark., 2013).

Genetik veriler, ekosistemlerdeki türlerin genetik çeşitliliğini değerlendirmek ve çevresel koşulların organizmalar üzerindeki etkisini anlamak için kullanılmaktadır. Su ekosistemlerinin restorasyonunda kullanılacak türlerin seçimi, genetik çeşitliliği yüksek ve çevresel streslere dayanıklı popülasyonlardan yapılmalıdır (Hughes ve ark., 2008). Bu tür popülasyonlar, ekosistemde daha iyi adapte olabilecek ve daha hızlı bir şekilde iyileşme sağlayabilecek kapasiteye sahiptir. Bu nedenle, su ekosistemlerinin restorasyonu için stratejiler geliştirirken, sadece ekosistemdeki türlerin sayısı değil, aynı zamanda genetik çeşitliliği de göz önünde bulundurulmalıdır. Genetik verilerin kullanımı, su ekosistemlerinin restorasyonunun başarısını artıran bir diğer önemli strateji de ekosistem mühendislerinin seçilmesidir. Tatlı su midyeleri gibi organizmalar, ekosistem mühendisleri olarak su kalitesini iyileştiren ve biyolojik çeşitliliği artıran kritik roller üstlenebilmektedir. Midye popülasyonlarının, çevresel değişimlere ve kirliliğe karşı genetik dayanıklılıkları artırıldığında, bu türlerin restorasyon sürecinde kullanılması ekosistem sağlığını iyileştirmektedir (Verma ve ark., 2025). Su ekosistemlerinin restorasyonu, genetik çeşitliliği artırarak ve çevresel streslere karşı dirençli organizmaları seçerek, daha sürdürülebilir ve sağlıklı su ortamları yaratılmasını sağlayabilir. Genetik analizler, restorasyon sürecinde hangi stratejilerin en iyi sonuçları vereceğini belirlemede önemli bir rehberdir.

Tehdit Altındaki Midye Popülasyonlarının Genetik Rehabilitasyonu

Tatlı su midyeleri, ekosistem mühendisleri olarak su kalitesinin düzenlenmesinde ve biyolojik çeşitliliğin korunmasında kritik bir rol oynamaktadır (Allen, ve ark., 2012). Ancak, tatlı su midyeleri, habitat kaybı, kirlilik, istilacı türler ve iklim değişikliği gibi faktörlerden dolayı tehdit altındadır (Haag, 2012). Bu tehditler, midye popülasyonlarının genetik çeşitliliğini azalttığı ve onların çevresel değişimlere adapte olma yeteneklerini kısıtladığı için bu popülasyonların korunması büyük bir önem taşımaktadır. Genetik rehabilitasyon, tehdit altındaki midye popülasyonlarının genetik çeşitliliğini artırarak bu popülasyonların daha dirençli hale gelmesini sağlamak amacıyla kullanılabilir. Genetik rehabilitasyon yöntemleri, genetik çeşitliliğin kaybolmasına yol açan daralmış genetik havuzları genişletmeyi ve popülasyonları yeniden çeşitlendirmeyi hedeflemektedir. Bu süreç, doğal seleksiyon süreçlerini hızlandırarak, çevresel streslere karşı daha dirençli popülasyonların oluşmasına olanak tanımaktadır (Rogers ve ark., 2021).

Genetik rehabilitasyon, iki ana yaklaşım ile gerçekleştirilebilir: birincisi, genetik çeşitliliği artırmak için yeni genetik materyalin popülasyona eklenmesi (genetik takviye), ikincisi ise daha dayanıklı, stres faktörlerine karşı dirençli bireylerin seçilerek popülasyona dahil edilmesidir. Bu yöntemler, midye popülasyonlarının çevresel streslere daha iyi uyum sağlamalarını, hastalıklara ve kirliliğe karşı daha dirençli hale gelmelerini sağlayabilmektedir (Aldridge ve ark., 2023). Genetik rehabilitasyon, tatlı su

FULL TEXTS BOOK

midyelerinin yaşam alanlarının restore edilmesi ve ekosistem işlevlerinin yeniden kazanılması için önemli bir araçtır. Bu süreç, midyelerin çevresel streslere karşı daha dayanıklı hale gelmelerini sağlamakla kalmaz. Aynı zamanda ekosistem mühendisliği işlevlerini yeniden yerine getirmelerini sağlayarak, su ekosistemlerinin genel sağlık durumunun iyileşmesine katkıda bulunur (Verma ve ark., 2025).

3.6. İklim Değişikliği ve Genetik Adaptasyon Stratejileri

İklim değişikliği, su ekosistemlerini ve biyoçeşitliliği doğrudan etkileyen önemli bir çevresel faktördür. Su sıcaklıklarındaki artış, asidifikasyon, oksijen seviyelerindeki değişimler ve ekstrem hava olaylarının artması, tatlı su organizmalarının hayatta kalma ve adaptasyon stratejilerini zorlaştırmaktadır. Midyeler, su ekosistemlerinin sağlığı için önemli bir rol oynayan organizmalar olmalarına rağmen, iklim değişikliği gibi çevresel stres faktörleri karşısında tehdit altındadır (Inoue & Berg, 2017). Bu nedenle, midyelerin iklim değişikliğine karşı genetik adaptasyonlarının araştırılması, onları korumak ve su ekosistemlerinin sürdürülebilirliğini sağlamak açısından kritik öneme sahiptir(Verma ve ark., 2025). İklim değişikliği, su ekosistemlerinde sıcaklık artışı, suyun pH düzeylerinde değişim, oksijen seviyelerinde azalma gibi stres faktörlerini artırmaktadır. Bu faktörler, özellikle midye popülasyonlarını olumsuz yönde etkilemekte, üreme başarılarını, büyüme hızlarını ve genel sağkalımlarını düşürmektedir. Midyelerin, bu değişimlere adaptasyon yetenekleri, genetik temele dayanmakta ve bu genetik adaptasyonlar, midyelerin çevresel değişimlere karşı dayanıklılıklarını arttırmaktadır (Wood, 2019).

Genetik adaptasyon, organizmaların çevresel koşullara uyum sağlamak için nesiller boyunca genetik varyasyonları ve doğal seleksiyonu kullanması sürecidir. Midyeler gibi organizmaların, iklim değişikliği gibi hızla değişen çevresel koşullara uyum sağlamaları, genetik düzeyde belirli genlerin seçilmesiyle mümkün olmaktadır (Rogers ve ark., 2021). Örneğin, sıcaklık artışına karşı daha dayanıklı bireylerin genetik özellikleri, gelecekteki popülasyonlara da yayılabilir. Bu adaptasyonlar, özellikle stres faktörlerine karşı daha az hassas bireylerin çoğalması ile gelişmektedir. Midyelerin termal tolerans aralıklarının genişletilmesi, onları sıcaklık değişimlerine karşı daha dirençli hale getirebilir ve bu da iklim değişikliği etkilerine karşı bir strateji oluşturabilir. Midyelerin genetik adaptasyonlarını incelemek için kullanılan bir yöntem, popülasyon genomik analizleridir. Bu analizler, farklı popülasyonların çevresel değişikliklere karşı nasıl tepki verdiklerini ve hangi genetik faktörlerin bu adaptasyonları yönlendirdiğini anlamaya yardımcı olmaktadır. Genetik varyasyonları belirleyerek, daha dirençli bireylerin ve popülasyonların seçilmesi, midyelerin iklim değişikliği etkilerine karşı daha dayanıklı hale gelmelerine olanak tanımaktadır (Popovic & Riginos, 2020).

İklim Senaryolarına Uygun Genetik Temelli Yönetim Stratejileri

İklim değişikliği senaryolarına uygun genetik temelli yönetim stratejileri, su ekosistemlerinin korunmasında ve sürdürülebilirliğinin sağlanmasında önemli bir rol oynamaktadır. Bu stratejiler, iklim değişikliğinin gelecekteki etkilerine yönelik proaktif yaklaşımlar geliştirmeyi hedeflemektedir. Tatlı su midyelerinin korunması için, genetik temelli yönetim stratejileri, iklim değişikliğinin çeşitli senaryolarına göre midye popülasyonlarının nasıl yönetileceğini belirleyecektir (Perea ve ark., 2022). Genetik temelli yönetim stratejilerinin uygulanabilmesi için, iklim değişikliği senaryolarının etkilerini öngörmek gerekmektedir. Örneğin, su sıcaklıklarının artması, oksijen seviyelerinin düşmesi veya pH değişiklikleri gibi faktörler, midye popülasyonlarının gelecekteki dağılımlarını ve sağkalımlarını etkileyebilir. Bu senaryoların analizi, midye popülasyonlarının bu değişimlere nasıl tepki vereceğini anlamaya yardımcı olacaktır. Bu bilgiler, genetik çeşitliliği korumak ve iklim değişikliğine karşı dirençli popülasyonlar oluşturmak için hangi stratejilerin uygulanacağına dair rehberlik edebilir (Inoue & Berg, 2017).

FULL TEXTS BOOK

Genetik temelli yönetim stratejilerinin bir örneği, midye popülasyonlarının genetik çeşitliliğini artırmaya yönelik yönetim uygulamalarıdır. Genetik çeşitliliğin yüksek olduğu popülasyonlar, çevresel streslere daha dayanıklıdır. Bu nedenle, popülasyonların genetik çeşitliliğini korumak, iklim değişikliği senaryolarına karşı daha sağlam bir ekosistem oluşturabilir. Ayrıca, genetik çeşitliliğin kaybolması riski bulunan popülasyonlar için genetik rehabilitasyon çalışmaları da önemli bir strateji olacaktır (Berg ve ark., 2008). Bir diğer önemli strateji ise, midye popülasyonlarının doğal seleksiyon süreçlerine olanak tanımaktır. Bu, midyelerin iklim değişikliği etkilerine daha hızlı adapte olabilmelerini sağlamak için genetik varyasyonu artırmaya yönelik yönetim stratejileri geliştirilmesi anlamına gelir. Bu tür stratejiler, özellikle iklim değişikliğinin hızla ilerlediği senaryolarda, ekosistemlerin daha esnek hale gelmesine yardımcı olacaktır (Perea ve ark., 2022).

4. YÜKSEK ÇÖZÜNÜRLÜKLÜ GENETİK ARAÇLAR

Yüksek çözünürlüklü genetik araçlar, çevresel yönetim süreçlerinde büyük bir devrim yaratma potansiyeline sahiptir. Bu araçlar, özellikle su ekosistemlerinde yer alan organizmaların genetik yapılarını, popülasyon dinamiklerini ve çevresel streslere karşı verdikleri yanıtları detaylı bir şekilde incelemek için kullanılmaktadır. Tatlı su midyeleri gibi türlerin genetik analizleri, su kalitesinin izlenmesinde, biyolojik çeşitliliğin korunmasında ve ekosistemlerin sürdürülebilir yönetilmesinde kritik öneme sahiptir (Currier ve ark., 2018).

4.1. Yeni Nesil Dizileme (NGS) Teknolojilerinin Su Kalitesi Yönetimindeki Kullanımı

Yeni nesil dizileme teknolojileri, genetik araştırmaların hızını, doğruluğunu ve kapsamını önemli ölçüde artırmıştır. Bu teknolojiler, su ekosistemlerindeki organizmaların genetik bilgilerini hızlı ve düşük maliyetle çözümleyebilmek için kullanılmaktadır. NGS, çevresel DNA (eDNA) örnekleme teknikleriyle birleştiğinde, su ortamlarındaki biyolojik çeşitliliği izlemek, kirliliğin etkilerini anlamak ve su kalitesinin zaman içindeki değişimini takip etmek için güçlü bir araç haline gelmiştir (Preece ve ark., 2021).

Su örneklerinde bulunan serbest DNA parçaları, su ekosistemindeki organizmaların varlıklarını ve biyolojik çeşitliliklerini izlemek için kullanılmaktadır. Midyeler gibi organizmaların genetik materyali, su kütlesinden alınan örneklerde tespit edilebilir. Bu, ekosistemlerin durumunun izlenmesine, invaziv türlerin erken tespiti ve kirliliğin biyolojik etkilerinin belirlenmesine yardımcı olmaktadır (Currier ve ark., 2018).

Su kütlesinde bulunan organizmaların genetik profillerini analiz ederek, ağır metaller, toksik kimyasallar ve diğer kirleticilerin etkisi daha iyi anlaşılmaktadır. NGS, özellikle kirliliğe karşı duyarlı genetik biyobelirteçlerin keşfini sağlar. Bu belirteçler, su kalitesinin izlenmesinde, çevresel streslerin erken tespiti ve yönetilmesinde kullanılmaktadır (Thomas ve ark., 2014). Ekosistemlerin genetik çeşitliliği, organizmaların çevresel streslere karşı direnç geliştirebilme kapasitesini belirlemektedir. NGS, tatlı su midyeleri gibi türlerin genetik çeşitliliğini izlemek için kullanılarak ekosistem sağlığının değerlendirilmesine ve popülasyonların dayanıklılığının ölçülmesine yardımcı olmaktadır (Gomes-dos-Santos ve ark., 2022).

4.2. Genetik Veri Analizlerinde Makine Öğrenimi (ML) ve Yapay Zeka (AI) Uygulamaları

Genetik veri analizlerinde makine öğrenimi ve yapay zeka teknolojilerinin kullanımı, büyük veri setlerinin işlenmesinde devrim yaratmaktadır. Su ekosistemlerinden elde edilen genetik veriler genellikle büyük ve karmaşık olabilir, bu nedenle geleneksel analiz yöntemleri bu tür verilerin işlenmesinde yetersiz kalabilmektedir. Makine öğrenimi ve yapay zeka algoritmaları, bu verilerin daha hızlı ve doğru bir şekilde analiz edilmesine olanak tanır (Keck ve ark., 2023). Makine öğrenimi algoritmaları, genetik verilerdeki desenleri ve ilişkileri tanımlayarak, türlerin adaptasyon mekanizmalarını ve çevresel stres yanıtlarını tahmin etmek için kullanılmaktadır. Bu algoritmalar, çevresel faktörlere karşı organizmaların genetik tepkilerini belirlemede ve bu verileri su

FULL TEXTS BOOK

kalitesi yönetiminde kullanmada önemli bir rol oynar. Örneğin, genetik biyobelirteçlerin belirlenmesi için sınıflandırma algoritmaları kullanılabilir (Grekov ve ark., 2023). Yapay zeka algoritmaları, çevresel streslerle ilişkili genetik biyobelirteçlerin keşfi için kullanılabilmektedir. Bu belirteçler, kirlilik ve diğer çevresel stres faktörlerinin erken tespiti için kritik öneme sahiptir. Yapay zeka, genetik verilerdeki gizli desenleri öğrenerek, su kalitesi üzerinde olumsuz etkisi olan kirlilik kaynaklarını tespit etmek için yeni işaretçiler geliştirebilir (Keck ve ark., 2023).

Su ekosistemlerinde bulunan çok büyük genetik veri setleri, veri madenciliği teknikleri ile analiz edilebilir. Ayrıca, yapay zeka genetik verilerdeki anlamlı ilişkileri ve etkileşimleri ortaya çıkarabilir, bu da kirliliğin ve çevresel streslerin organizmalar üzerindeki etkilerini daha iyi anlamaya olanak tanır (Frincu, 2025).

Makine öğrenimi teknikleri, çevresel değişikliklerin organizmalar üzerindeki genetik etkilerini tahmin etmek için zaman serisi analizleri yapmaktadır. Örneğin, iklim değişikliği veya kirlilik artışı gibi faktörlerin su ekosistemleri üzerindeki etkileri, bu algoritmalarla modellenebilir. Böylece gelecekteki su kalitesi değişimleri ve ekosistem sağlığı hakkında öngörüler elde edilmektedir (Krishnan ve ark., 2022).

Makine öğrenimi, farklı kaynaklardan gelen genetik ve çevresel verilerin entegrasyonunu sağlayarak, daha kapsamlı bir su yönetimi modelinin oluşturulmasına yardımcı olabilir. Yapay zeka, bu verileri analiz ederek karar destek sistemlerini güçlendirebilir ve çevresel yönetim süreçlerini daha etkin hale getirebilir (Essamlali ve ark., 2024).

4.3. Biyoinformatik ve Veri Tabanları

Biyoinformatik, genetik verilerin analiz edilmesi, işlenmesi ve anlamlı bilgilere dönüştürülmesi için kullanılan disiplinler arası bir alandır. Su ekosistemlerinde yer alan tatlı su midyeleri gibi organizmaların genetik özelliklerinin anlaşılması, biyoinformatik araçlar ve veri tabanlarının kullanımını gerektirir. Bu araçlar, genetik verilerin yönetilmesi, analizi ve çevresel yönetim stratejileriyle entegrasyonunu sağlayarak, daha sağlıklı ve sürdürülebilir su ekosistemleri yaratılmasına olanak tanıyacaktır.

Tatlı Su Midyelerine Özgü Genetik Bilgi Tabanlarının Oluşturulması

Tatlı su midyeleri, su ekosistemlerinin önemli biyoindikatörleri olduklarından, bu türlerin genetik özelliklerinin sistematik bir şekilde depolanması ve analiz edilmesi gerekmektedir. Bu bağlamda, tatlı su midyelerinin genomik ve transkriptomik verilerini içeren özel bilgi tabanlarının oluşturulması, bu organizmaların biyolojik çeşitliliğinin izlenmesine ve çevresel değişikliklere verdikleri yanıtların daha iyi anlaşılmasına olanak tanımaktadır (Gomes-dos-Santos ve ark., 2024). Tatlı su midyelerinin genetik verilerini derleyen ve saklayan bilgi tabanları, her türün popülasyon yapısı, genetik çeşitliliği ve adaptasyon mekanizmalarına dair bilgiler sunmaktadır. Bu veriler, su kalitesi değerlendirmeleri ve çevresel yönetim süreçlerinde kullanılabilir (Gomes-dos-Santos ve ark., 2022). Ayrıca, çeşitli midye türlerinin karşılaştığı çevresel stres faktörlerine karşı geliştirdikleri genetik yanıtlar hakkında daha derinlemesine bilgi edinilmesine yardımcı olur.

Genetik bilgi tabanları, tatlı su midyelerinin filogenetik ilişkilerini ve genetik çeşitlilik seviyelerini haritalamaya yardımcı olabilir. Bu tür veriler, popülasyonların genetik yapısındaki değişikliklerin izlenmesinde, türler arası genetik etkileşimlerin anlaşılmasında ve genetik çeşitliliğin korunmasında kritik rol oynamaktadır (Lopes-Lima ve ark., 2021).

Tatlı su midyelerine özgü genetik veriler, koruma biyolojisi ve biyoçeşitlilik izleme stratejilerinde önemli görülmektedir. Bu verilerle, tehdit altındaki midye türlerinin korunmasına yönelik etkili stratejiler geliştirilmesi mümkündür.

Tatlı su midyelerine dair genetik verilerin dijital ortamda depolanması, bilimsel araştırmaların küresel çapta daha hızlı ve etkin bir şekilde yapılmasına olanak tanımaktadır. Ayrıca, araştırmacılar arasında bilgi paylaşımını teşvik ederek, su ekosistemlerinin yönetimi konusunda daha geniş bir işbirliği ortamı oluşturulur.

FULL TEXTS BOOK

5. MULTİDİSİPLİNER YAKLAŞIMLAR

Multidisipliner yaklaşımlar, genetik, ekoloji ve mühendislik gibi farklı bilim alanlarının birleşimiyle elde edilen sinerjiden yararlanarak, su ekosistemlerinin yönetiminde yenilikçi ve etkin çözümler geliştirmeyi amaçlamaktadır. Tatlı su midyeleri gibi biyoindikatör organizmalar üzerinden yapılan bu tür çalışmalar, çevresel yönetim ve sürdürülebilirlik stratejilerine büyük katkı sağlayabilir (Verma ve ark., 2025). Su ekosistemlerinin sürdürülebilirliği için genetik, ekolojik ve mühendislik bilimlerinin entegrasyonu, çok disiplinli bir yaklaşımla çevresel problemlerin çözülmesine olanak tanımaktadır. Tatlı su midyelerinin ekolojik rolü ve genetik özellikleri, mühendislik bilimleri ile desteklenerek daha verimli, etkili ve uzun vadeli çözümler üretilmesine yardımcı olabilir (Vereycken & Aldridge, 2023). Genetik veriler, ekolojik modelleme çalışmalarında kullanılarak, midyelerin çevresel streslere karşı verdikleri yanıtların daha iyi anlaşılmasını sağlamaktadır. Örneğin, tatlı su midyelerinin belirli bir çevresel stres faktörüne (örneğin, sıcaklık artışı veya oksijen azalması) nasıl tepki verdiğini belirlemek için ekolojik modellemelerle genetik veri entegrasyonu yapılabilir. Bu modeller, çevresel değişimlerin ekosistem üzerindeki uzun vadeli etkilerini tahmin etmeye katkıda bulunabilir (Haberle ve ark., 2025).

Tatlı su midyelerinin genetik verileri, ekosistem sağlığına dair bilgi sağlarken, ekolojik gözlemlerle birleştirildiğinde su kalitesinin çok daha doğru bir şekilde değerlendirilmesini mümkün kılmaktadır. Mühendislik bilimlerinin katkısıyla bu veriler, gerçek zamanlı izleme sistemleri geliştirerek, kirlilik seviyelerini, biyolojik çeşitliliği ve ekosistem sağlığını dinamik olarak takip edebilir (Pilbala ve ark. 2024). Mühendislik bilimleri, tatlı su midyelerinin genetik özelliklerine dayalı olarak biyomühendislik çözümleri geliştirmek için kullanılabilmektedir. Örneğin, midyelerin kabuk yapılarındaki mekanik özellikleri inceleyerek, çevresel streslere karşı dayanıklı biyomimikri malzemeler tasarlanabilir. Bu malzemeler, inşaat, su arıtma ve çevresel mühendislik alanlarında kullanılarak, sürdürülebilir çözümler geliştirilmesine katkıda bulunur (Wang ve ark., 2019).

Tatlı su midyelerinin genetik özellikleri, çevresel politika ve su yönetimi stratejilerinin geliştirilmesinde kritik bir rol oynayabilir. Genetik veriler, su kalitesini izleme ve ekosistemlerin korunması gibi önemli konularda daha etkili yönetim yöntemlerinin benimsenmesine olanak tanır. Genetik biyobelirteçler ve eDNA (çevresel DNA) teknolojileri, su kalitesini izlemek için güçlü araçlar sunar. Bu veriler, yerel yönetimlerin, çevresel tehditleri daha hızlı ve hassas bir şekilde tespit etmelerini sağlar. Örneğin, genetik analizler sayesinde kirliliğin ve biyolojik çeşitliliğin azalmasının erken uyarı sistemleriyle takip edilmesi, çevresel politikalarda hızlı ve doğru aksiyonların alınmasına yardımcı olabilir (Preece ve ark., 2021).

Genetik veriler, su kaynaklarını korumaya yönelik politikaların bilimsel temellerle güçlendirilmesine yardımcı olmaktadır. Örneğin, kirlilik seviyelerinin, tür çeşitliliği kayıplarının ve habitat tahribatının genetik göstergelerle takip edilmesi, bu faktörlerin politika yapıcılar tarafından daha etkin bir şekilde ele alınmasını sağlamaktadır. Genetik veriler, çevre politikalarının daha hedeflenmiş ve yerel koşullara uygun bir şekilde geliştirilmesine olanak tanır (Reid ve ark., 2019). Genetik verilerin ekosistem ve su kalitesi yönetimi süreçlerine entegrasyonu, çevresel etki değerlendirmelerinin daha kapsamlı bir şekilde yapılmasını sağlar. Bu süreç, genetik çeşitlilik, popülasyon yapıları ve organizmaların adaptasyon kapasitelerini göz önünde bulundurarak, su yönetimi stratejilerinin planlanmasına yardımcı olacaktır. Ayrıca, bu stratejiler, biyolojik çeşitliliğin korunması ve ekosistem hizmetlerinin sürdürülebilirliğini güvence altına almak için genetik verilerle desteklenmiş olmalıdır.

6. SONUÇ

Tatlı su midyelerinin genetik özelliklerinin çevresel izleme, su kalitesi yönetimi ve ekosistem sağlığına dair sağladığı bilimsel veriler, su kaynaklarının korunması ve yönetilmesinde daha etkin, verimli ve sürdürülebilir yaklaşımlar geliştirilmesine olanak tanımaktadır. Dahası, genetik biyobelirteçler, eDNA

FULL TEXTS BOOK

ve modern biyoteknolojik araçlar, midyelerin çevresel değişimlere yanıt verme kapasitesini artırarak su ekosistemlerinin korunmasını sağlayacak stratejilerin oluşturulmasında önemli katkı yapacaktır.

KAYNAKLAR

Aldridge, D. C., Ollard, I. S., Bespalaya, Y. V., Bolotov, I. N., Douda, K., Geist, J., ... & Zieritz, A. (2023). Freshwater mussel conservation: A global horizon scan of emerging threats and opportunities. Global Change Biology, 29(3), 575-589.

Allen, D. C., Vaughn, C. C., Kelly, J. F., Cooper, J. T., & Engel, M. H. (2012). Bottom-up biodiversity effects increase resource subsidy flux between ecosystems. Ecology, 93(10), 2165-2174.

Andres, K. J., Lodge, D. M., Sethi, S. A., & Andrés, J. (2023). Detecting and analysing intraspecific genetic variation with eDNA: From population genetics to species abundance. Molecular Ecology, 32(15), 4118-4132.

Arslan, Ö. Ç., Parlak, H., Boyacıoğlu, M., Karaaslan, M. A., Nalbantlar B., ... & Gülsever, G. (2017). Doğal Ortamından alınmış midyelerde (Mytilus galloprovincialis) Genetik Biyo-işaretler kullanılarak İzmir Körfezi Kirliliği'nin Belirlenmesi. http://155.223.63.101/tez3/2016suf006.pdf

Banaduc, D., Simić, V., Cianfaglione, K., Barinova, S., Afanasyev, S., Öktener, A., ... & Curtean-Banaduc, A. (2022). Freshwater as a sustainable resource and generator of secondary resources in the 21st century: Stressors, threats, risks, management and protection strategies, and conservation approaches. *International Journal of Environmental Research and Public Health*, 19(24), 16570.

Berg, D. J., Levine, T. D., Stoeckel, J. A., & Lang, B. K. (2008). A conceptual model linking demography and population genetics of freshwater mussels. Journal of the North American Benthological Society, 27(2), 395-408.

Bickham, J. W., Sandhu, S., Hebert, P. D., Chikhi, L., & Athwal, R. (2000). Effects of chemical contaminants on genetic diversity in natural populations: implications for biomonitoring and ecotoxicology. Mutation research/Reviews in Mutation research, 463(1), 33-51.

Bourlat, S. J., Borja, A., Gilbert, J., Taylor, M. I., Davies, N., Weisberg, S. B., ... & Obst, M. (2013). Genomics in marine monitoring: new opportunities for assessing marine health status. Marine pollution bulletin, 74(1), 19-31.

Currier, C. A., Morris, T. J., Wilson, C. C., & Freeland, J. R. (2018). Validation of environmental DNA (eDNA) as a detection tool for at-risk freshwater pearly mussel species (Bivalvia: Unionidae). Aquatic Conservation: Marine and Freshwater Ecosystems, 28(3), 545-558.

Durán-Vinet, B., Araya-Castro, K., Chao, T. C., Wood, S. A., Gallardo, V., Godoy, K., & Abanto, M. (2021). Potential applications of CRISPR/Cas for next-generation biomonitoring of harmful algae blooms: A review. Harmful algae, 103, 102027.

Ekelund Ugge, G. M. O., Jonsson, A., Olsson, B., Sjöback, R., & Berglund, O. (2020). Transcriptional and biochemical biomarker responses in a freshwater mussel (Anodonta anatina) under environmentally relevant Cu exposure. Environmental Science and Pollution Research, 27, 9999-10010.

Essamlali, I., Nhaila, H., & El Khaili, M. (2024). Advances in machine learning and IoT for water quality monitoring: A comprehensive review. *Heliyon*.

Fediajevaite, J., Priestley, V., Arnold, R., & Savolainen, V. (2021). Meta-analysis shows that

FULL TEXTS BOOK

environmental DNA outperforms traditional surveys, but warrants better reporting standards. Ecology and Evolution, 11(9), 4803-4815.

Franco, J. L., Trivella, D. B., Trevisan, R., Dinslaken, D. F., Marques, M. R., Bainy, A. C., & Dafre, A. L. (2006). Antioxidant status and stress proteins in the gills of the brown mussel Perna perna exposed to zinc. Chemico-Biological Interactions, 160(3), 232-240.

Gallardo-Escárate, C., Valenzuela-Muñoz, V., Nuñez-Acuña, G., Valenzuela-Miranda, D., Tapia, F. J., Yévenes, M., ... & Gerdol, M. (2023). Chromosome-Level genome assembly of the blue mussel Mytilus chilensis reveals molecular signatures facing the marine environment. Genes, 14(4), 876.

Gomes-dos-Santos, A., Machado, A. M., Castro, L. F. C., Prié, V., Teixeira, A., Lopes-Lima, M., & Froufe, E. (2022). The gill transcriptome of threatened European freshwater mussels. Scientific Data, 9(1), 494.

Gomes-dos-Santos, A., Fonseca, E., Riccardi, N., Hinzmann, M., Lopes-Lima, M., & Froufe, E. (2024). The transcriptome assembly of the European freshwater mussel Unio elongatulus C. Pfeiffer, 1825. Scientific Data, 11(1), 377.

Grekov, A. N., Kabanov, A. A., Vyshkvarkova, E. V., & Trusevich, V. V. (2023). Anomaly detection in biological early warning systems using unsupervised machine learning. Sensors, 23(5), 2687.

Froufe, E., Bolotov, I., Aldridge, D. C., Bogan, A. E., Breton, S., Gan, H. M., ... & Lopes-Lima, M. (2020). Mesozoic mitogenome rearrangements and freshwater mussel (Bivalvia: Unionoidea) macroevolution. Heredity, 124(1), 182-196.

Fu, D., Dai, A., Hu, R., Chen, Y., & Zhu, L. (2008). Expression and role of factor inhibiting hypoxia-inducible factor-1 in pulmonary arteries of rat with hypoxia-induced hypertension. Acta biochimica et biophysica Sinica, 40(10), 883-892.

Frincu, R. M. (2025). Artificial intelligence in water quality monitoring: A review of water quality assessment applications. *Water Quality Research Journal*, 60(1), 164-176.

Haag, W. R. (2012). North American freshwater mussels: natural history, ecology, and conservation. Cambridge University Press.

Haberle, I., Moore, A. P., Forbes, V. E., Brain, R. A., Hornbach, D. J., Galic, N., & Vaugeois, M. (2025). Comparing freshwater mussel responses to stress using life-history and Dynamic Energy Budget theory. Science of The Total Environment, 958, 177664.

Hartmann, J. T., Beggel, S., Auerswald, K., Stoeckle, B. C., & Geist, J. (2016). Establishing mussel behavior as a biomarker in ecotoxicology. *Aquatic Toxicology*, 170, 279-288.

Hughes, A. R., Inouye, B. D., Johnson, M. T., Underwood, N., & Vellend, M. (2008). Ecological consequences of genetic diversity. Ecology letters, 11(6), 609-623.

Inoue, K., & Berg, D. J. (2017). Predicting the effects of climate change on population connectivity and genetic diversity of an imperiled freshwater mussel, Cumberlandia monodonta (Bivalvia: Margaritiferidae), in riverine systems. Global change biology, 23(1), 94-107.

Jha, A. N. (2008). Ecotoxicological applications and significance of the comet assay. Mutagenesis, 23(3), 207-221.

FULL TEXTS BOOK

Jones, J. W., Neves, R. J., & Hallerman, E. M. (2015). Historical demography of freshwater mussels (Bivalvia: Unionidae): genetic evidence for population expansion and contraction during the late Pleistocene and Holocene. Biological Journal of the Linnean Society, 114(2), 376-397.

Keck, F., Brantschen, J., & Altermatt, F. (2023). A combination of machine-learning and eDNA reveals the genetic signature of environmental change at the landscape levels. Molecular ecology, 32(17), 4791-4800.

Krishnan, S. R., Nallakaruppan, M. K., Chengoden, R., Koppu, S., Iyapparaja, M., Sadhasivam, J., & Sethuraman, S. (2022). Smart water resource management using Artificial Intelligence—A review. *Sustainability*, 14(20), 13384.

Lacroix, C., Coquillé, V., Guyomarch, J., Auffret, M., & Moraga, D. (2014). A selection of reference genes and early-warning mRNA biomarkers for environmental monitoring using Mytilus spp. as sentinel species. Marine pollution bulletin, 86(1-2), 304-313.

Lavictoire, L., & West, C. (2024). Population reinforcement of the endangered freshwater pearl mussel (Margaritifera margaritifera): lessons learned. Diversity, 16(3), 187.

Lodge, D. M. (2022). Policy action needed to unlock eDNA potential. Frontiers in Ecology and the Environment, 20(8), 448-449.

Lopes-Lima, M., Gürlek, M. E., Kebapçı, Ü., Şereflişan, H., Yanık, T., Mirzajani, A., ... & Froufe, E. (2021). Diversity, biogeography, evolutionary relationships, and conservation of Eastern Mediterranean freshwater mussels (Bivalvia: Unionidae). Molecular Phylogenetics and Evolution, 163, 107261.

Lummer, E. M., Auerswald, K., & Geist, J. (2016). Fine sediment as environmental stressor affecting freshwater mussel behavior and ecosystem services. Science of the Total Environment, 571, 1340-1348.

Machuca-Sepúlveda, J., Miranda, J., Lefin, N., Pedroso, A., Beltrán, J. F., & Farias, J. G. (2023). Current status of omics in biological quality elements for freshwater biomonitoring. Biology, 12(7), 923.

Merdun, H., & Çınar, Ö., (2008). Su kirliliği ve kontrolü. Çevre kirliliği ve kontrolü (pp.2-31), Ankara: Nobel Yayınevi.

Mijangos, J. L., Pacioni, C., Spencer, P. B., & Craig, M. D. (2015). Contribution of genetics to ecological restoration. Molecular Ecology, 24(1), 22-37.

Montúfar-Romero, M., Valenzuela-Muñoz, V., Valenzuela-Miranda, D., & Gallardo-Escárate, C. (2024). Hypoxia in the blue mussel mytilus chilensis induces a transcriptome shift associated with endoplasmic reticulum stress, metabolism, and immune response. Genes, 15(6), 658.

Pilbala, A., Riccardi, N., Benistati, N., Modesto, V., Termini, D., Manca, D., ... & Piccolroaz, S. (2024). Real-time biological early-warning system based on freshwater mussels' valvometry data. Hydrology and Earth System Sciences, 28(10), 2297-2311.

Popovic, I., & Riginos, C. (2020). Comparative genomics reveals divergent thermal selection in warm- and cold-tolerant marine mussels. Molecular ecology, 29(3), 519–535. https://doi.org/10.1111/mec.15339

Preece, E. P., Bryan, M., Mapes, S. M., Wademan, C., & Dorazio, R. (2021). Monitoring for

FULL TEXTS BOOK

freshwater mussel presence in rivers using environmental DNA. Environmental DNA, 3(3), 591-604.

Reid, A. J., ve ark. (2019). Emerging threats and persistent conservation challenges for freshwater biodiversity. Biological Reviews, 94(3), 849-873. DOI: 10.1111/brv.12480

Ribeiro, Â. M., Canchaya, C. A., Penaloza, F., Galindo, J., & da Fonseca, R. R. (2019). Population genomic footprints of environmental pollution pressure in natural populations of the Mediterranean mussel. Marine Genomics, 45, 11-15.

Rogers, R. L., Grizzard, S. L., Titus-McQuillan, J. E., Bockrath, K., Patel, S., Wares, J. P., ... & Moore, C. C. (2021). Gene family amplification facilitates adaptation in freshwater unionid bivalve Megalonaias nervosa. Molecular Ecology, 30(5), 1155-1173.

Ruiz-Ramos, D. V., Thompson, N. L., Richter, C. A., Voshage, M. C., Schreier, T. M., Merkes, C. M., & Klymus, K. E. (2024). Environmental DNA dynamics of three species of Unionid freshwater mussels. Environmental DNA, 6(2), e543.

Scheder, C., Lerchegger, B., Flödl, P., Csar, D., Gumpinger, C., & Hauer, C. (2015). River bed stability versus clogged interstitial: depth-dependent accumulation of substances in freshwater pearl mussel (Margaritifera margaritifera L.) habitats in Austrian streams as a function of hydromorphological parameters. *Limnologica*, 50, 29-39.

Suárez-Ulloa, V., Fernández-Tajes, J., Manfrin, C., Gerdol, M., Venier, P., & Eirín-López, J. M. (2013). Bivalve omics: State of the art and potential applications for the biomonitoring of harmful marine compounds. Marine drugs, 11(11), 4370-4389.

Tanguy, A., Bierne, N., Saavedra, C., Pina, B., Bachère, E., Kube, M., ... & Canario, A. (2008). Increasing genomic information in bivalves through new EST collections in four species: development of new genetic markers for environmental studies and genome evolution. Gene, 408(1-2), 27-36.

Thomas, E. G., Šrut, M., Štambuk, A., Klobučar, G. I., Seitz, A., & Griebeler, E. M. (2014). Effects of freshwater pollution on the genetics of zebra mussels (Dreissena polymorpha) at the molecular and population level. BioMed research international, 2014(1), 795481.

Togaki, D., Doi, H., & Katano, I. (2020). Detection of freshwater mussels (Sinanodonta spp.) in artificial ponds through environmental DNA: a comparison with traditional hand collection methods. Limnology, 21, 59-65.

Varotto, L., Domeneghetti, S., Rosani, U., Manfrin, C., Cajaraville, M. P., Raccanelli, S., ... & Venier, P. (2013). DNA damage and transcriptional changes in the gills of Mytilus galloprovincialis exposed to nanomolar doses of combined metal salts (Cd, Cu, Hg). PloS one, 8(1), e54602.

Vaughn, C. C. (2010). Biodiversity losses and ecosystem function in freshwaters: emerging conclusions and research directions. *BioScience*, 60(1), 25-35.

Vereycken, J. E., & Aldridge, D. C. (2023). Bivalve molluscs as biosensors of water quality: state of the art and future directions. *Hydrobiologia*, 850(2), 231-256.

Verma, A. K., Rahman, A., Hussain, S., & Singh, N. S. (2025). Freshwater Mussels as Multifaceted Ecosystem Engineers: Insights into Their Ecological Importance, Bioindication, and Economic Contributions. Water, 17(11), 1629.

Wang, Q., Li, Q., Wei, J., & Zhang, G. (2019). Molecular insights into the biomineralization process of mussel shells. ACS Biomaterials Science & Engineering, 5(12), 6466–6476.

FULL TEXTS BOOK

DOI: 10.1021/acsbiomaterials.8b01134

Wei, X., Fu, Z., Li, J., & Ye, Y. (2023). Genetic Diversity and Structure Revealed by Genomic Microsatellite Markers of Mytilus unguiculatus in the Coast of China Sea. Animals, 13(10), 1609.

Wood, M. (2019). Genetic variation gives mussels a chance to adapt to climate change. https://www.uchicagomedicine.org/forefront/biological-sciences-articles/genetic-variation-gives-mussels-a-chance-to-adapt-to-climate-change?utm_source=chatgpt.com,

Xu, M., Jiang, L., Shen, K. N., Wu, C., He, G., & Hsiao, C. D. (2016). Transcriptome response to copper heavy metal stress in hard-shelled mussel (Mytilus coruscus). Genomics Data, 7, 152-154.

Zhang, G., Fang, X., Guo, X., Li, L. I., Luo, R., Xu, F., ... & Wang, J. (2012). The oyster genome reveals stress adaptation and complexity of shell formation. Nature, 490(7418), 49-54.

IMPACT OF TERRORISM AND BANDITRY ON WOMEN DEVELOPMENT IN NORTHERN NIGERIA

Olumide Abraham Ajayi (Ph.D.)

Skyline University Nigeria, Kano

ABSTRACT

Nigeria is gradually turning to a breeding ground for terrorists and bandits based on the operations of insurgent in the country. People living in the northern part of Nigeria live in abject fear, as no one knows the next target of the bandits. Women and girls have become the prime targets of terrorists and bandits. Terrorist groups manipulate and capitalise on the stereotype of women to suit their violent goals. These various attacks, abduction and violence acts demoralized and affected the social well-being of women. The economic growth and development of women is becoming stagnant as no one wants to establish a business that will be destroyed in no time and those that have businesses wants to close down or relocate. This paper examined the impact of terrorism and banditry as it affects women's development in the northern part of Nigeria. The paper concludes that terrorism and banditry have crippled activities and impeded peace, and development in the northern region. It therefore recommends that the government should pay more attention to factors underlying breeding of terrorism and banditry in order to address the menace holistically.

Keywords: Terrorism, banditry, violence, women development, northern Nigeria

Introduction

Terrorism is a social problem in Nigeria. It is indeed an act that destabilise and devastates the country. Terrorism refers to acts of violence undertaken by individuals or local groups within the state to recompense domestic grievance. The US State Department (2002) describes terrorism as "premeditated, politically motivated violence perpetrated against non-combatant targets by subnational groups or clandestine agents, usually intended to influence an audience". The FBI's (2005) noted that terrorism is "the unlawful use of force or violence against persons or property to intimidate or coerce a Government, the civilian population, or any segment thereof, in furtherance of political or social objectives". Banditry on the other hand refers to the practice of raiding and attacking victims by members of an armed group, whether or not premeditated, using weapons of offense or defense, especially in semi-organized groups to overpower the victim and obtain loot or achieving some political goals (Shalangwa, 2013). A bandit may be seen as one who commits the wrongdoing of equipped theft, slaughtering, and annihilation of properties, especially, of herders, vendors, and commerce proprietors (Olapeju & Peter, 2021).

Banditry is widespread in the Northern region of Nigeria, and the activities of bandits include; kidnapping, arson, cattle rustling, looting, raping, killings among others (Akinyetun, 2022). The activities of bandits' in Nigeria are widespread and can be targeted at individuals or groups and the prime targets are mostly ordinary citizens of the country rich or poor (Usman &Singh, 2021). According to Lebari et al. (2022), banditry found its way into the north west region of Nigeria due to climate change and the resultant depletion of grazing resources for the herdsmen that led to encroachment of land belongings to farmers and the subsequent eruption of farmers-herders clash and conflict which transformed into communal and ethnic rivalry. The nonchalant attitude of government in tackling the problems led to the formation of militia groups by herdsmen, who carry out kidnapping and other evil acts in the region. The militia groups were known as bandits. Adetayo (2022) stated that more than 5,000 people were killed while others were displaced as a result of bandits activities from 2018.

FULL TEXTS BOOK

The resurgence of banditry in recent times started with a lone criminal gang in 2011and grew from a single cell, mainly operating in Zamfara, to over 120 gangs across six states in 2021 (Odalonu & Egbogu, 2023). However, the recent spate of banditry-related violence began in 2014 with cattle rustling activity. It became worse in early 2016 when the bandits started killing local miners in Zamfara communities (Mohammed, 2021). The attack of bandits has become an incessant tragedy in which women have been raped, kidnapped and killed, children have become orphans, villages have been sacked and farm produce destroyed (Mohammed,2021). This is also the case with terrorism as the word terrorism and banditry are used interchangeably in Nigeria due to the proscription of banditry groups and labelling such groups as terrorist groups by the federal government.

Terrorists and bandits manipulate and capitalise on the status of women to suit their nefarious goals. Women are used as baits and sometimes take active roles in carrying out nefarious activities. Women are also kidnapped and taken hostage by these groups. Women may participate in terrorism and carry out acts of violence because they are committed to the political, religious or economic goals of the terrorists, they may also be manipulated into taking up violent roles such as suicide bombing. Women are abducted, intimidated, and sometimes, forcefully recruited into terrorist and bandits organizations. This is because they can directly support terrorists through cooking and cleaning and can also indirectly support their efforts by developing and disseminating propaganda as well as encouraging their children to go into violence (Taiwo, 2012).

The threat incurred by terrorist acts and banditry in Nigeria have profound security and economic implications on the corporate existence of Nigeria as united and indivisible entity. The large scale of destructions of life and properties by terrorist activities in the northern part of the country has crippled the one buoyant economic fortress of Kaduna, Katsina and Zamfara states. Other major cities and towns in the north are living in abject fear. The issue of fear have dominated the minds of the people especially the commuters that travel by road. The current problem has endangered the lives of women and crippled their socio-economic development and the security situation is becoming worse on a daily basis. This paper therefore focus on the impact of terrorism and banditry on women development in northern Nigeria.

Historical Overview of Terrorism

Elements of terrorism could be seen in the resistance of the imperial authority of the Romans by the Jewish Zealots and Arab nationalities by killing many Roman soldiers and destroying properties. In addition, the Arabs fight (Shiites versus Sunnis) over religious doctrine and the revolt against the crusaders by the Arabs exemplify terrorism a long time ago (Hoffman, 1999). Terrorism of the modern era was conceived as a tool to achieve political and religious goals. It began during the French Revolution (1793-1794). During this period, the government of France sponsored terrorism in order to maintain power and suppress oppositions to the government (Hoffman, 1999).

Dempsey (2006) posited that terrorism is a failure of political process that begins with inequalities, corruption and injustice in a given political system and moves from a frustrated attempts at reform that breed fear and anger to political confrontation. In addition, Hoffman (2002) believed that terrorism relates to those acts that are carried out by local group or persons due to grievance within a locality or community. As political scientist Martha Crenshaw explains, terrorism is a method by which an agent tends to produce terror in order to impose his or her domination (Arguilla and Ronfeldy, 2001). A number of analysts have attributed the disturbing trend of terrorism and banditry to political

FULL TEXTS BOOK

dissatisfaction, perceived societal neglect and pervasive poverty among the people. For instance, youth restiveness in the Niger Delta and parts of South –East, occasioned kidnapping and disruption of oil installation, activities of the men of the Odua's People Congress (OPC) in the South-West and that of the Boko- Haram in the Northern States who sees western education and democracy as an abomination (Abimbola, 2012). Terrorists and bandits have unleashed havoc on the Nigeria society.

The Plight of Women

The intensification of insurgency in 2011led to the indoctrination of many young women and girls who were coerced into compliance. Militant forces in the northern region of Nigeria have captured and exploited women and converted them to Islam under death threat (Nwose, 2012). In July 2014, northern Nigeria faced a spate of suicide bomber attacks. Although this tactic has never been employed by the terrorist group since 2011, the new perpetrators of violence are often young females. Women have also increasingly become targets in kidnapping and assaults with the abduction of 270 Chibok girls in 2014 (Al Jazeera, 2014).

During a search operation in Maiduguri in August 2013, government forces arrested five female terrorist members carrying rifles and ammunition under their hijabs, other members acted as informants. The terrorist group is increasingly capitalising on women's heightened likelihood to evade suspicion at security checkpoints. Women are typically viewed as nonthreatening. This perception has fostered sensitivities regarding more thorough body-searches of women at check-points or public gatherings. As a result, four attacks were carried out by female suicide bombers in Kano at the end of 2014. (BBC, 2014)

Women and girls of all ages have been the target of violence, with infants and elderly women being raped. Brutal forms of sexual violence, such as sexual slavery, gang rape, mutilation, torture, and insertion of sharp objects into women vaginas were carried out by terrorists and bandits. High levels of rape have had a significant impact on women's contribution to the economy and their injuries require specialized gyneacological care. High numbers of traumatic fistula cases have been reported in the northern part of Nigeria (BBC, 2014). The targeting of women was clearly illustrated in 2014, when Boko Haram abducted nearly 300 girls from a public school in the town of Chibok, in Borno State. In 2018, a similar incident was recorded in Dapchi, Yobe State, where 101 girls were kidnapped by the group. Okoli (2022) observed that aside from the two instances of mass abductions epitomized by the case at chibok and Dapchi, kidnappings continue sporadically around northern Nigeria.

Victims in the custody of terrorists and bandits are subjected to rape, forced conversion to Islamic religion, forced labor, and forced marriage and concubinage in insurgent camps. This gender-based violence further undermine the position of women in society. Girls are also more likely to drop out of school early for fear of abduction. As a result, the rate of female school enrollment is far lower in northern Nigeria than in other regions.

Theoretical Framework

Many theories could be used to explain terrorism and banditry in Nigeria. Some of the theories include the resource mobilization theory, frustration-aggression theory, situational action theory, relative deprivation theory, relational vengeance theory, strain theory among others. However, this study is hinged on the frustration-aggression theory. The frustration-aggression theory was proposed by John Dollard, Neal Miller, Leonard Doob, Orval Mowrer, and Robert Sears in 1939 and further developed by Neal Miller in 1941and Leonard Berkowitz in 1969 (Miller, Doob, Mowrer, & Sears, 1939; Miller,

FULL TEXTS BOOK

1941;Berkowitz, 1989). The theory assumes that aggression is always a result of frustration and that frustration always leads to some form of aggression (Dollard, Miller, Doob, Mowrer, & Sears, 1939; Fererabend & Feirauben, 1972). Frustration is seen as interference with an incited target response that occurs at the appropriate time in the behavioral sequence. Therefore, frustration is an interruption of the intended response to a behavioural task, which can interfere with the execution of the task (Dollard et al., 1939). One of the proponents of the frustration-aggression theory generated some modifications to the central hypothesis because of the interrogations and criticisms. Hence, Miller (1941) noted that it is not always clear that frustration always leads to aggression or that aggression is always propelled by frustration. His intervention led to the second round of the hypothesis, which reflected a more acceptable reality that frustration can produce instigations to several different kinds of response, one of which is instigation to some form of aggression.

Berkowitz (1989) argued that aggression can be driven by inherent personal benefits to the aggressor, rather than by past wrongdoings. People are more likely to attack when they discover that they are willfully sabotaged or denied what belong to them legitimately rather than when the interference is an accidental occurrence. He hypothesized that frustrations are aversive events and can generate aggressive inclinations only to the extent that they have a negative effect (Berkowitz, 1989). According to Fererabend & Feirauben (1972), aggression is caused by frustration, which results from an individual's inability to achieve their goals. Terrorism and banditry are aggressive behaviour that is caused by other problems, such as poverty and unemployment (Adegoke, 2019). Thus, the increasing number of bandit attacks (aggression) in Nigeria is largely due to economic difficulties and struggles (frustration). The acts of banditry that are common, such as cattle rustling, kidnapping, physical attacks, and encroaching on farms, are caused by frustrations (Uche & Iwuamadi, 2018). Therefore, the frustration-aggression theory argument is applicable to the rising banditry and terrorist activities in Nigeria which suggests that banditry in the country is caused by the need of the disgruntled elements of society to get out of poverty and improve their social standing. That is why the phenomenon of ransom is common in bandit operations.

Equally, terrorist groups are vexing their anger on the government in order to achieve predefined objectives. However, when the ransom is not paid or delayed, bandits become more tense and frustrated and resort to killing their victims.

Impact of Terrorism and Banditry on Women Development

The series of problems experienced by women as a result of terrorist and banditry activities cannot be over emphasized. The activity of Boko Haram insurgency has various implications on women and girls. A study by Nigeria's Political Violence Research Network suggests that more than 45% of those killed by terror group are women and children (Barkindo; Gudaku; and Wesley, 2013). Boko Haram's misguided ideology that "western education" is sinful has significant negative consequences on the education of women and girl-child in the northern part of Nigeria. For fear of attack by terrorists and bandits, most schools in the affected region have been shut down until recently. Young women are forcefully married off in their teens thereby perpetuating a cycle of poverty.

The incessant cases of rape is destroying households and families. Sexual violence in terrorist area has numerous short- and long-term economic, social, cultural and health consequences. The most evident consequence is the loss of lives due to sexual violence-related deaths, killings or suicide. Victims and survivors suffer long-term health consequences such as psychological trauma, including isolation, fear, hot flashes, feelings of unworthiness and suicide and medical trauma such as gynecological fistula, sexually transmitted infections, and HIV/AIDS. Sexual violence limits women's freedom of movement in relation to subsistence activities.

FULL TEXTS BOOK

The activities of terrorists and bandits can reduce women and girls' contribution to economic development, trapping them in a life of poverty. The implication of insurgency and the consequent response from the government has serious negative effects on the livelihood of population most especially the women. It is observed that public security volatility in the northern states has hampered normal productive agricultural and commercial activities (Okoli and Iortyer, 2014). The current wave of armed banditry and general insecurity in the country have crippled the socio-economic activities and scared away the potential investors (Umoru, 2019). Foreign Direct Investment (FDI) has declined in Nigeria from 1269.22 in 2017 to 959.52 in 2018, an further dropped to 909.54 in 2019, which are attributed to deteriorating insecurity situations such as banditry, kidnapping among others (Odalonu & Egbogu, 2023)..

It is instructive to note that women play active role in agriculture and rural livelihood, but the fear of attacks by terrorists has often driven them into hiding. In the North-East part of Nigeria, attacks on women and girls are most often committed when they are in schools, farm or markets to acquire education and obtain firewood and water for other domestic and subsistence activity. Also, the lack of economic opportunities for populations often results in commercial and exploitative sex as one of the few options for income generation to meet basic needs. National Bureau of Statistics documented that about 80 million out of

the estimated 200 million people live below the poverty line, while unemployment rate is as high as 33% which was compounded by the recent upsurge of insecurity of armed banditry and kidnapping (Adetayo, 2022).

Women who have been attacked, as well as children born out of rape, are stigmatized and ostracized as a consequence of these crimes. In some cases, husbands disown their wives and the high levels of discrimination cause the destruction of family and social relationships that make up the fabric of society. Women in this scenario are often left in poverty and outside their communities. Addressing sexual violence as a peace and security imperative with a clear focus on prevention, protection and accountability can advance economic recovery and community reconstruction.

Conclusion

The activities of the terrorists and bandits in Nigeria as a whole and in the northern region in particular cannot be over emphasised. Terrorism and banditry have crippled almost all the sectors of the region's activities and impeded peace, progress and development in the area. The paper concludes that terrorism and banditry have serious negative impact on women development. These impacts include abduction of women and girls, arbitrary arrest of women by government security agents, use of women as pawns in their activities, inflicting terror on women, use of women as wartime labour force, women education deprivation and livelihood crisis. The gender-based violence of terrorism and banditry as stated in this paper affected women and girls socio-economically and psychologically. Attention has been paid in this study to the fact that women also take part in terrorism and banditry as members that carry out attacks on innocent people.

Recommendations

In the light of the above conclusion, the paper made the following policy recommendations:

i. Government and non-governmental organizations should organise continuous seminars and public lectures to sensitize people about the importance and pivotal role of women in the society.

ii. More attention should be given to women's education to reduce the number of women being brain washed to engage in or support terrorists and bandits.

FULL TEXTS BOOK

iii. Skills acquisition programs should be provided in rural communities especially for herders and their children to serve as an alternative source of livelihood besides rearing of animals to eradicate poverty.

iv. Community policing strategy should be strengthened to provide public safety and the needed intelligence to government security agencies involved in the fight against terrorism and banditry.

v. Job opportunity should be provided for youth who are becoming nuisance and threat to the society.

vi. Government security agents should be thoroughly equipped to provide security in communities especially to vulnerable women and girls among others.

vii. Modern technology should be acquired to intensify security check at government institutions, hospitals, shopping complex, churches and other places where group gather. This will help reduce insurgence attack in those areas

viii. Government and international aids should provide sufficient funds for all the displaced people and ensure that the funds are judiciously used.

REFERENCES

Abimbola. S.O. (2012). Domestic Terrorism and Boko-Haram Insurgency in Nigeria. Issues and Trends. A Historical Discourse. *Journal of Arts and Contemporary Society*, Vol 4.

Adegoke, S. G. (2019). Insurgency, armed banditry and corruption in Nigeria: The bane of socio-economic underdevelopment. *International Journal of Advanced Academic Studies, 2*(1), 17-26.

Adetayo,O. (2022). Nigeria: The economics and trauma of banditry and kidnapping in the North. *The Africa Report*. 22 April 2022.

Akinyetun, T.S. (2022). Banditry in Nigeria: Insights from Situational Action and Situational Crime Prevention Theories. *Accord.* March 15 https://www.accord.org.za/conflict-trends/banditry-innigeria- insights-from-situational-action-and-situational-crime-prevention-theories/

Al Jazeera (2014). Nigeria's Kidnapped Girls sold into Marriage. http://www.aljazeera.com.

Arguilla, J. and Ronfeldy, D. (2001). *Networks and Netwars*. Santa Monica, CA: Rands

Barkindo, A., Gudaku, B.T., and Wesley, C.K. (2013). Boko Haram and gender-based violence against Christian women and children in northeastern Nigeria since 1999. *NPV RN Working paper No.1* (Amsterdam: Open Doors International)

Berkowitz, L. (1969). *Roots of aggression: A re-examination of the frustration-aggression hypothesis.* New York: Atherton Press.

British Broadcasting Corporation (BBC) (2014) Boko-Haram Crisis. Nigeria arrest Female Recruiters. http://www.bbc.com.

Dempsey, T. (2006). Counter Terrorism in African Failed States. Challenges and Potential Solutions. http://www.strategicstudiesinstitute.army.mill//.

Dollard J, Miller, N. E, Doob, L. W., Mowrer, O. H., and Sears, R. R. (1939). *Frustration and aggression*. New Haven, CT: Yale University Press.

FBI (2005) Federal Bureau of Investigation Reports and Publications 2005

Fererabend, I. K. and Feiraubend, R. L. (1972). Systematic conditions of political aggression: An application of frustration-aggression theory. In Fairaben and Ted, R.C. (eds) *Anger violence and politics: Theories and research*. Prentice Hall Inc, Englewood. Cliff, New Jersey.

FULL TEXTS BOOK

Hoffman, B. (1999). *Inside Terrorism.* NewYork Columba University Press.

Hoffman S. (2003). *Clash of Globalizations Foreign Affairs*.

Lebari, G.N., Ferguson, A.O. and Sunday, A. (2022). Climate Change, Farmer-Herders Conflicts and Food Security in Nigeria. International Journal of Research and Innovation in Social Science, VI (I), 407 - 413

Miller, N. E. (1941). The frustration-aggression hypothesis. *Psychological Review*, *48*, 337-342.

Mohammed, I. (2021). No end in sight for insecurity and banditry in Northwestern Nigeria. Retrieved fromhttps://jamestown.org/program/no-end-in-sight-for-insecurity-and-banditry-in-northwestern-nigeria/

Nwose, I.E. (2012). Gender Role Perceptions and the Changing Role of Women in Nigeria. *International Journal of Agriculture and Rural Development* 15 (3).

Odalonu, B.H and Egbogu, D.U (2023) Implications of Escalating Banditry on National Security in Nigeria. *African Journal of Humanities & Contemporary Education Research* 10(1), 38 – 62.

Okoli. A.I.C. and Iortyer, P. (2014). Terrorism and humanitarian crisis in Nigeria: Insights from Boko Haram insurgency. *Global Journal of Human and Social Sciences* (F). Xiv (I): 39-49

Okoli, C. (2022) Gender and Terror: Boko Haram and the Abuse of Women in Nigeria. *Georgetown Journal of International Affairs.* https://gjia.georgetown.edu/2022/04/05/gender-and-terror-boko-haram-and-the-abuse-of-women-in-nigeria%EF%BF%BC/

Olapeju, R. M. and Peter, A. O. (2021). The impact of banditry on Nigeria's security in the Fourth Republic: an evaluation of Nigeria's northwest. *Zamfara Journal of Politics and Development*, 2(1), 1-26

Shalangwa, M. W. (2013). *The nature and consequences of armed banditry in border communities of Adamawa State, Nigeria*. M.Sc. thesis submitted to the School of Post-Graduate Studies, Ahmadu Bello University, Zaria, Nigeria.

Taiwo A. (2012). Boko Haram Sokoto Opens up. *Sunday Tribune*. Vol 5.

Uche, J. C. and Iwuamadi, C. K. (2018). Nigeria: Rural banditry and community resilience in the Nimbo community. *Conflict Studies Quarterly*, (24), 71-82.

Umoru (2019). We are not getting enough from security Architecture. Available: https://www.vanguardngr.com/2019/10/kidnapping-we-are-not-getting-the-best-from-thesecurityarchitecturesenate/

US Department of State (2002) Homeland Security Act of 2002 Congress of USA Title 22 of the United States Code, Section 2656f [d].

Usman, G. M. and Singh, D. (2021). The persistence of insecurity in Northern Nigeria: A critical analysis on banditry**,** NIU International Journal of Human Rights, 8(XVIII), 11-27

FULL TEXTS BOOK

AKUT EGZERSİZ YORGUNLUĞUNUN HAMSTRİNG KAS MİMARİSİ VE KORTİKOSPİNAL UYARILABİLİRLİK ÜZERİNE ETKİLERİ: SAĞLIKLI BİREYLER ÜZERİNDE DENEYSEL BİR ÇALIŞMA

THE ACUTE EFFECTS OF EXERCİSE-INDUCED FATIGUE ON HAMSTRING MUSCLE ARCHITECTURE AND CORTICOSPINAL EXCITABILITY: AN EXPERIMENTAL STUDY IN HEALTHY INDIVIDUALS

***Caglar SOYLU*[1,2]**

[1]*Associate Professor, University of Health Sciences, Gülhane Faculty of Physiotherapy and Rehabilitation, Department of Orthopedic Physiotherapy and Rehabilitation, Ankara, Türkiye*

[2]*Postdoctoral Fellow, Sports Performance, Recovery, Injury & New Technologies (SPRINT) Research Centre*
Australian Catholic University, Melbourne, Australia

[1,2]*ORCID ID: 0000-0002-1524-6295*

ÖZET

Bu çalışma, sağlıklı yetişkinlerde akut egzersiz sonrası biseps femoris uzun başının (BFLH) proksimal ve distal bölgelerindeki fasikül davranışı, kortikospinal ve intrakortikal yanıtları karşılaştırmayı amaçlamaktadır. Prospektif longitudinal desenle tasarlanan araştırmada, 10 sağlıklı yetişkin üzerinde yorgunluk öncesi ve sonrası kapsamlı değerlendirmeler yapılmıştır. Katılımcılar, 50 maksimal konsantrik diz fleksiyonundan oluşan standardize yorgunluk protokolünü uygulamış, ardından ultrason, izokinetik dinamometre, yüzeyel EMG ve transkraniyal manyetik stimülasyon (TMS) ile ölçümler alınmıştır. Ultrason bulguları, uzun kas boylarında fasikül uzunluğunda anlamlı azalma (Öncesi: %6.6±5.0, Sonrası: %5.5±6.0; p=0.005, Cohen's d=-1.17) ve pennasyon açısında düşüş (Öncesi: %7.9±12.9, Sonrası: %1.6±11.6; p=0.041, Cohen's d=-0.75) olduğunu gösterirken, kas kalınlığı değişmemiştir. Tork ölçümlerinde konsantrik (%22.4 azalma, p=0.042), eksantrik (%14.5 azalma, p=0.017) ve izometrik (%29.6 azalma, p=0.004) kasılmalarda belirgin düşüşler kaydedilmiş, en büyük etki izometrik torkta gözlenmiştir (Cohen's d=-1.30). Nörofizyolojik veriler, kortikospinal uyarılabilirliğin azaldığını (Dinlenme motor eşiği: %51.2±6.7'den %54.9±7.2'ye yükselme, p=0.032; Motor uyarılmış potansiyel genliği: %51.5±15.2'den %44.3±14.6'ya düşme, p=0.025) ve intrakortikal inhibisyonda artış eğilimi (SICI: %41.3±19.1'den %47.6±21.8'e, p=0.078) olduğunu ortaya koymuştur. Bu sonuçlar, akut yorgunluğun hem periferik kas mimarisinde hem de merkezi sinir sistemi işlevlerinde ölçülebilir değişikliklere yol açtığını kanıtlamakta ve hamstring yaralanma mekanizmalarının anlaşılması için kritik bir normatif temel sağlamaktadır.

Anahtar Kelimeler: Hamstring kası, kas mimarisi, kortikospinal uyarılabilirlik, egzersize bağlı yorgunluk, transkraniyal manyetik stimülasyon, isokinetik tork, sağlıklı bireyler

ABSTRACT

This study investigates fascicle behaviour, corticospinal and intracortical responses in proximal/distal regions of the biceps femoris long head (BFLH) following acute exercise in healthy adults. Using a prospective longitudinal design, comprehensive pre-/post-fatigue assessments were conducted on 10 healthy adults. Participants performed a standardized fatigue protocol (50 maximal concentric knee flexions), followed by ultrasound, isokinetic dynamometry, surface EMG, and transcranial magnetic stimulation (TMS). Ultrasound revealed significant reductions in fascicle length at long muscle

FULL TEXTS BOOK

lengths (Pre: 6.6±5.0%, Post: 5.5±6.0%; p=0.005, Cohen's d=-1.17) and pennation angle (Pre: 7.9±12.9%, Post: 1.6±11.6%; p=0.041, d=-0.75), while muscle thickness remained unchanged. Torque output declined markedly during concentric (99.0±26.5 Nm to 90.4±31.3 Nm, p=0.042), eccentric (117.2±28.5 Nm to 103.1±31.9 Nm, p=0.017), and isometric contractions (109.6±26.5 Nm to 84.9±21.0 Nm, p=0.004), with the largest effect size in isometric torque (d=-1.30). Neurophysiological data demonstrated reduced corticospinal excitability (Resting motor threshold: 51.2±6.7% to 54.9±7.2% MSO, p=0.032; Motor evoked potential amplitude: 51.5±15.2% to 44.3±14.6% Mmax, p=0.025) and increased intracortical inhibition tendency (SICI: 41.3±19.1% to 47.6±21.8%, p=0.078). These outcomes validate that acute fatigue induces measurable alterations in peripheral muscle architecture and central nervous system function in healthy individuals, providing a critical normative baseline for understanding hamstring injury mechanisms. The study highlights the vulnerability of neuromuscular systems to fatigue, particularly during eccentric loading and at extended muscle lengths, informing targeted rehabilitation strategies to mitigate reinjury risks.

Keywords: Hamstring muscle, muscle architecture, corticospinal excitability, exercise-induced fatigue, transcranial magnetic stimulation, isokinetic torque, healthy individuals.

INTRODUCTION

The hamstring muscle group, comprising the semimembranosus, semitendinosus, and biceps femoris muscles, is integral to lower limb biomechanics, facilitating hip extension and knee flexion (1, 2). These muscles are particularly active during sprinting, especially in the terminal swing phase, where they undergo intense eccentric contractions to decelerate leg movement (3, 4). This eccentric loading, coupled with muscle lengthening, heightens their susceptibility to injury (5). Hamstring injuries represent the most prevalent muscle injuries in sports, accounting for approximately 54.4% of all sports-related muscular injuries (6). Moreover, recurrence rates remain high, ranging from 12% to 33% despite full return to sport (7, 6), with significant economic implications. For instance, Ekstrand et al. (8) estimated a potential loss of €7.5 million for a professional football club if five key players are injured for 90 days, while hamstring injuries cost an average of USD 191,614 per player in the 2012 Australian Football League season (9, 10).

The long head of the biceps femoris (BFLH) is implicated in nearly 80% of hamstring injuries, likely due to its substantial length change and force demands during high-speed running (11). However, the differential fascicle behavior in the proximal and distal BFLH regions under fatigue and across contraction types remains poorly understood, with potential implications for injury risk (12). Previous studies have explored biomechanical and architectural contributors to hamstring injury, with Schache et al. (3) highlighting the terminal swing phase as a high-risk period due to increased strain, and Opar et al. (5, 13) identifying fascicle length and eccentric strength as key risk factors. Ultrasonography and surface electromyography (EMG) have been employed to assess fascicle behavior and voluntary activation (14, 15), yet these methods are limited in elucidating underlying neural mechanisms.

Fyfe et al. (16) proposed a neuromuscular inhibition model, suggesting that cortical-level inhibition following hamstring injuries may lead to chronic deficits in voluntary activation, strength, and fascicle length, contributing to elevated re-injury rates. Despite this, no study has directly evaluated corticospinal excitability or intracortical inhibition in hamstring-injured populations using transcranial magnetic stimulation (TMS), a non-invasive technique widely used to assess motor cortex excitability and inhibition in other injuries, such as anterior cruciate ligament (ACL) reconstruction (17-19). This study addresses these gaps by examining (1) voluntary activation deficits, (2) corticospinal excitability and intracortical inhibition via TMS, and (3) fascicle behavior in proximal and distal BFLH regions during

FULL TEXTS BOOK

fatigue and various contraction types in healthy individuals. These findings aim to establish a normative baseline and inform the development of targeted rehabilitation strategies to reduce re-injury rates.

METHODS

This study employed a prospective, longitudinal experimental design with pre- and post-fatigue assessments to evaluate the acute effects of exercise-induced fatigue on hamstring muscle architecture and corticospinal excitability in healthy individuals. The research was conducted at a specialized research center, with ethical approval granted by the university's Human Research Ethics Committee (approval number: 2024-2638H). This study was supported by the TÜBİTAK 2219 – Postdoctoral Research Fellowship Program Abroad.

Participants

Ten recreationally active adults aged 18–30 years were recruited for the control group, with no hamstring injury in the past 6 months and a dominant leg straight leg raise ≥80°. Inclusion criteria included a body mass index (BMI) of 18.5–24.9 kg/m², ability to perform exercise protocols, and voluntary consent. Exclusion criteria encompassed any lower or upper limb injury within the past year, sensory, cardiovascular, respiratory, or neurological disorders, pregnancy, cancer history, spinal surgery, or acute low back/extremity pain. Demographic data (age, sex, height, weight, BMI, dominant limb) were recorded, with sample size determined based on prior studies (e.g., Buhmann et al., 2022) to detect differences in intracortical inhibition with 95% power and $\alpha = 0.05$ (20).

Study Schedule and Preparation

Participants attended two laboratory sessions one week apart, with one limb tested per session (dominant/non-dominant), and the testing order randomized. Participants were instructed to avoid caffeine for 6 hours and refrain from intense exercise for 48 hours prior to testing.

Ultrasound Imaging of Muscle Architecture

Muscle architecture of the BFLH was assessed at rest and during dynamic contractions using a Supersonic Mach 30 ultrasound system (SuperSonic Imagine S.A., Aix-en-Provence, France) with two high-frequency linear probes (18-5L, 5–18 MHz) for dual-probe acquisition. Imaging was performed by experienced investigators with over two years of musculoskeletal ultrasound experience. Resting architecture was evaluated using extended field-of-view imaging, with participants prone and muscles at rest. One probe was manually moved along the posterior thigh to visualize the full fascicular structure, measuring fascicle length (FL) from deep to superficial aponeurosis (averaged across three fascicles), pennation angle (PA) at the proximal aponeurosis, and muscle thickness (MT) between aponeuroses. During dynamic assessments, both probes were fixed over proximal and distal BFLH regions with elastic bandages during concentric and eccentric isokinetic contractions. Real-time imaging over a full knee range of motion (ROM, 0°–90°) was synchronized with dynamometer and EMG outputs, with images processed using Camtasia software (TechSmith, Version 2019.0.0) to analyze FL, PA, and MT at 10° increments (12).

Isokinetic Dynamometry

Muscle contractions were performed using a Biodex System 4 isokinetic dynamometer (Medical Systems Inc., NY, USA). Participants were seated with hips flexed at ~85° and stabilized with straps at the chest, pelvis, thigh, and shank. The device's axis was aligned with the knee joint, and gravity correction was conducted at 30° knee flexion. The eccentric contraction range was 80° to 10° knee flexion, with an angular velocity of 10°/s for eccentric and 180°/s for concentric contractions.

FULL TEXTS BOOK

Fatigue Protocol

Neuromuscular fatigue was induced with 50 maximal concentric knee flexion contractions from 0° to 90° at 180°/s, preceded by a warm-up of 10 submaximal concentric repetitions at ~60% of perceived effort. Post-fatigue assessments included maximal voluntary isometric contraction (MVIC), dual-probe ultrasound, TMS, and surface EMG (sEMG) measurements.

Surface Electromyography (sEMG) and MVIC

Wireless sEMG signals were collected with electrodes placed on the BFLH following SENIAM guidelines (21). Participants completed three eccentric warm-up contractions at 50%, 75%, and 95% of effort, followed by three maximal eccentric contractions. The highest root mean square (rmsEMG) value was used to normalize subsequent EMG signals. Visual feedback maintained target muscle activation at 5 ± 2% rmsEMG during TMS trials.

Transcranial Magnetic Stimulation (TMS)

TMS was administered using two Magstim 2002 stimulators connected via a Bistim unit with a 110 mm double-cone coil, targeting the motor cortex contralateral to the test limb via neuronavigation with a fitted textile cap marked at 1 cm intervals. The motor hotspot elicited the largest motor evoked potential (MEP) from the BFLH, with the active motor threshold (AMT) defined as the lowest stimulus intensity producing ≥5 MEPs ≥500 μV out of 10 stimuli. Stimulation was delivered during eccentric contractions between 50°–10°, triggered near 30° flexion. Parameters included single-pulse TMS at 130% AMT (accepted within 28°–32° flexion), SICI at a 3 ms interstimulus interval (ISI) with 80%/130% AMT, and ICF at a 12 ms ISI with 80%/130% AMT. All stimulations were performed at 5 ± 2% of normalized rmsEMG to ensure low-level pre-activation (22, 20).

Statistical Analysis

Statistical analysis was performed using SPSS (latest version). Data normality was assessed with Shapiro-Wilk tests and visual QQ plots. Paired-samples t-tests compared pre- vs. post-fatigue within-group differences, and two-way repeated measures ANOVA evaluated interaction effects. Effect sizes were calculated as Cohen's d for t-tests (small: 0.2, medium: 0.5, large: 0.8) and partial eta squared (ηp^2) for ANOVA (small: 0.01, medium: 0.06, large: 0.14), with statistical significance set at $p < 0.05$.

Research results

This section details the acute effects of a fatiguing exercise protocol on fascicle length (ΔLf), pennation angle (ΔPA), muscle thickness (ΔMT), torque values, and corticospinal excitability in healthy control participants (n = 10). Assessments were conducted pre- and post-fatigue at long (10°–30°) and short (60°–80°) muscle lengths, and across the full range of motion (ROM: 10°–80°). Paired t-tests and ANOVA were used, with effect sizes reported as Cohen's d or ηp^2 (Tables 1 and 2). Fascicle length significantly decreased at long muscle lengths post-fatigue (Pre: 6.6 ± 5.0%; Post: 5.5 ± 6.0%; $t(9) = 3.69$, $p = 0.005$, $d = -1.17$), indicating a moderate-to-large effect. No significant changes were observed at short muscle lengths (Pre: 7.8 ± 5.9%; Post: 9.6 ± 7.4%; $p = 0.237$) or across the total ROM ($p = 0.573$), suggesting fatigue-induced alterations are length-specific. Pennation angle significantly reduced at long muscle lengths (Pre: 7.9 ± 12.9%; Post: 1.6 ± 11.6%; $t(9) = 2.36$, $p = 0.041$, $d = -0.75$), with a moderate effect. No significant changes occurred at short muscle lengths ($p = 0.995$) or across the total ROM ($p = 0.676$), indicating fatigue affects fiber orientation primarily at extended lengths. Muscle thickness showed no significant change, with a modest decrease at long muscle lengths (Pre: 1.2 ± 3.6%; Post: 0.6 ± 3.5%; $p = 0.098$, $d = -0.58$) and non-significant variations at short lengths and total ROM ($p > 0.612$), suggesting minimal impact from acute fatigue. Significant torque reductions were observed post-fatigue: concentric (99.0 ± 26.5 Nm to 90.4 ± 31.3 Nm; $t(9) = 2.41$, $p = 0.042$, $d = -0.76$), eccentric (117.2 ± 28.5 Nm to 103.1 ± 31.9 Nm; $t(9) = 3.04$, $p = 0.017$, $d = -0.96$), and isometric (109.6 ± 26.5

FULL TEXTS BOOK

Nm to 84.9 ± 21.0 Nm; t(9) = 4.10, p = 0.004, d = -1.30), with the largest effect in isometric torque. Resting motor threshold (RMT) increased significantly (51.2 ± 6.7% to 54.9 ± 7.2% MSO; F(1,9) = 6.342, p = 0.032, ηp^2 = 0.413), and MEP amplitude decreased (51.5 ± 15.2% to 44.3 ± 14.6% Mmax; F(1,9) = 7.128, p = 0.025, ηp^2 = 0.442), indicating reduced corticospinal excitability. SICI showed a non-significant increase (41.3 ± 19.1% to 47.6 ± 21.8%; F(1,9) = 3.253, p = 0.078, ηp^2 = 0.223), while ICF decreased (89.6 ± 21.4% to 81.2 ± 22.7%; F(1,9) = 4.613, p = 0.059, ηp^2 = 0.338), suggesting trends in inhibitory and excitatory shifts. LICI remained unchanged (55.8 ± 22.1% to 53.9 ± 25.4%; F(1,9) = 0.731, p = 0.413, ηp^2 = 0.075). Fatigue induced significant reductions in FL and PA at long lengths, torque across all contraction types (greatest in isometric), and corticospinal excitability (RMT, MEP). Trends in SICI and ICF suggest neural modulation, while stable LICI provides a normative baseline for future comparisons.

Table 1. Muscle Architecture and Torque – Pre/Post Fatigue (Control Group)

Variable	Condition	Pre-Fatigue (Mean ± SD)	Post-Fatigue (Mean ± SD)	t(9)	p-value	Cohen's d
Fascicle length change (ΔLf %)	Long	6.6 ± 5.0	5.5 ± 6.0	3.69	0.005	-1.17
	Short	7.8 ± 5.9	9.6 ± 7.4	1.27	0.237	-0.40
	Total	25.6 ± 10.7	26.5 ± 7.8	0.59	0.573	-0.18
Pennation angle change (ΔPA %)	Long	7.9 ± 12.9	1.6 ± 11.6	2.36	0.041	-0.75
	Short	9.6 ± 11.1	9.6 ± 18.0	0.01	0.995	0.00
	Total	38.1 ± 27.2	35.5 ± 37.9	0.43	0.676	-0.13
Muscle thickness change (ΔMT %)	Long	1.2 ± 3.6	0.6 ± 3.5	1.85	0.098	-0.58
	Short	0.7 ± 3.7	0.7 ± 3.3	0.06	0.956	-0.02
	Total	1.1 ± 9.3	0.4 ± 7.1	0.52	0.612	-0.16
Torque (Nm)	Concentric	99.0 ± 26.5	90.4 ± 31.3	2.41	0.042	-0.76
	Eccentric	117.2 ± 28.5	103.1 ± 31.9	3.04	0.017	-0.96
	Isometric	109.6 ± 26.5	84.9 ± 21.0	4.10	0.004	-1.30

ΔLf = Change in Fascicle Length; ΔPA = Change in Pennation Angle; ΔMT = Change in Muscle Thickness; SD = Standard Deviation; Nm = Newton-meters; t(9) = t-statistic with 9 degrees of freedom; p-value = Statistical significance level; Cohen's d = Effect size (small: 0.2, medium: 0.5, large: ≥0.8).

Table 2. Intracortical and Corticospinal Measures Before and After Fatigue (Control Group)

Measure	Pre-Fatigue	Post-Fatigue	F(1,9)	p-value	ηp^2	Interpretation

FULL TEXTS BOOK

	(Mean ± SD)	(Mean ± SD)				
Resting Motor Threshold (RMT) (%MSO)	51.2 ± 6.7	54.9 ± 7.2	6.342	0.032	0.413	↑ RMT post-fatigue (↓ excitability)
Motor Evoked Potential (MEP) (%Mmax)	51.5 ± 15.2	44.3 ± 14.6	7.128	0.025	0.442	↓ MEP amplitude post-fatigue
Short-Interval Intracortical Inhibition (SICI) (%)	41.3 ± 19.1	47.6 ± 21.8	3.253	0.078	0.223	↑ GABA-A inhibition (non-significant)
Intracortical Facilitation (ICF) (%)	89.6 ± 21.4	81.2 ± 22.7	4.613	0.059	0.338	↓ Glutamatergic facilitation (non-significant)
Long-Interval Intracortical Inhibition (LICI) (%)	55.8 ± 22.1	53.9 ± 25.4	0.731	0.413	0.075	No change in GABA-B inhibition

RMT = Resting Motor Threshold; MSO = Maximum Stimulator Output; MEP = Motor Evoked Potential; Mmax = Maximum M-wave amplitude; SICI = Short-Interval Intracortical Inhibition; ICF = Intracortical Facilitation; LICI = Long-Interval Intracortical Inhibition; SD = Standard Deviation; F(1,9) = F-statistic with 1 numerator and 9 denominator degrees of freedom; p-value = Statistical significance level; ηp^2 = Partial Eta Squared (small: 0.01, medium: 0.06, large: 0.14).

DISCUSSION

The findings of this study demonstrate that acute exercise-induced fatigue significantly alters hamstring muscle architecture and corticospinal excitability in healthy individuals. The observed reductions in fascicle length and pennation angle at long muscle lengths align with prior research showing that fatigue leads to decreased muscle extensibility and altered fiber orientation, potentially increasing injury susceptibility during extended positions (12, 23). Muscle thickness remained stable, consistent with studies indicating that radial dimensions are less affected by short-term fatigue (24). Torque declines across contraction types, with the greatest impact on isometric strength, support the notion that fatigue impairs force production, particularly in static conditions, as reported in previous investigations (25, 26).

Neuromuscular changes, including increased RMT and decreased MEP amplitudes, indicate reduced corticospinal drive, which may reflect central fatigue mechanisms (27, 28). The trends in SICI and ICF suggest a shift toward greater inhibition and reduced facilitation, potentially mediated by GABAergic and glutamatergic pathways (16, 29). Stable LICI is in line with evidence that GABA-B inhibition is more resistant to acute fatigue (30, 31). These neural adaptations could contribute to the observed torque losses, highlighting the interplay between peripheral and central factors (32).

Clinically, these results emphasize the importance of monitoring length-specific architectural changes and corticospinal excitability in fatigue management. Rehabilitation programs could incorporate TMS-guided interventions to mitigate inhibition and enhance excitability, reducing injury risk (33, 34).

FULL TEXTS BOOK

Limitations include the focus on healthy individuals and the need for comparison with injured cohorts. Future studies should explore longitudinal effects and incorporate advanced imaging to further elucidate fatigue mechanisms (35).

Conclusion and Comments

This study investigated the acute neuromuscular and architectural responses of the BFLH to a standardized fatiguing concentric exercise protocol in healthy, recreationally active individuals. Key findings include significant reductions in fascicle length (ΔLf) and pennation angle (ΔPA) at long muscle lengths, indicating mechanical adaptations in fiber orientation and extensibility under fatigue. Muscle thickness (ΔMT) remained largely unaffected, suggesting that acute fatigue does not significantly alter radial muscle dimensions in the short term.

Functionally, torque output decreased across concentric, eccentric, and isometric contractions, with the largest effect observed in isometric torque (d = -1.30), highlighting the sensitivity of isometric strength to transient fatigue. These reductions validate the efficacy of the fatigue protocol and underscore the relevance of eccentric strength loss in injury prevention. At the neural level, increased resting motor threshold (RMT) and decreased motor evoked potential (MEP) amplitudes signify a decline in corticospinal excitability. Trends toward increased short-interval intracortical inhibition (SICI) and reduced intracortical facilitation (ICF), though not statistically significant, suggest a shift in cortical inhibitory-excitatory balance post-fatigue. The stable long-interval intracortical inhibition (LICI) aligns with prior evidence that GABA-B-mediated inhibition is less responsive to acute fatigue in healthy populations.

Implications

These findings provide a normative baseline for comparing healthy individuals with those who have prior hamstring injuries. The observed torque and excitability reductions may serve as early markers of fatigue-induced vulnerability, relevant for athletic rehabilitation and return-to-play strategies. The architectural changes at longer muscle lengths highlight a potential injury risk window during activities like sprinting or rapid deceleration.

REFERENCES

1. Kellis E. Functional anatomy and biomechanics of the hamstrings. In: Musculoskeletal Disorders and Injuries in Athletes; 2018. p. 45-60.

2. van der Made AD, Wieldraaijer T, Kerkhoffs GM, et al. The hamstring muscle complex. Knee Surg Sports Traumatol Arthrosc. 2015;23(7):2115-2122. doi:10.1007/s00167-014-3067-1

3. Schache AG, Dorn TW, Blanch PD, et al. Mechanics of the human hamstring muscles during sprinting. Med Sci Sports Exerc. 2012;44(4):647-658. doi:10.1249/MSS.0b013e318236a3c2

4. Opar DA, Williams MD, Shield AJ. Hamstring strain injuries: factors that lead to injury and re-injury. Sports Med. 2012;42(3):209-226. doi:10.2165/11594800-000000000-00000

5. Opar DA, Williams MD, Timmins RG, et al. Eccentric hamstring strength and hamstring injury risk in Australian footballers. Med Sci Sports Exerc. 2014;46(3):857-865. doi:10.1249/MSS.0000000000000142

6. Eirale C, Farooq A, Smirnov D, et al. Epidemiology of football injuries in Asia: a prospective study in Qatar. J Sci Med Sport. 2013;16(2):113-117. doi:10.1016/j.jsams.2012.04.007

7. Dalton SL, Kerr ZY, Dompier TP. Epidemiology of injuries in the National Collegiate Athletic Association men's football. Am J Sports Med. 2015;43(4):974-981. doi:10.1177/0363546514568491

FULL TEXTS BOOK

8. Ekstrand J, Hagglund M, Walden M. Injury incidence and injury patterns in professional football: the UEFA injury study. Br J Sports Med. 2016;50(12):645-650. doi:10.1136/bjsports-2015-095812

9. Hickey J, Shield AJ, Williams MD, et al. The financial cost of hamstring strain injuries in the Australian Football League. Br J Sports Med. 2014;48(8):729-731. doi:10.1136/bjsports-2013-092448

10. Pollard CW, Shields BJ, Beckwith JG, et al. Cost of injuries in youth and high school football. Am J Sports Med. 2019;47(10):2317-2322. doi:10.1177/0363546519858587

11. Opar DA, Drezner J, Shield A, et al. Acute hamstring strain injury in athletes: a review of the literature. Phys Ther Sport. 2015;16(3):131-137. doi:10.1016/j.ptsp.2014.08.003

12. Brusco CM, Opar DA, Timmins RG, et al. Fascicle behavior of the biceps femoris long head under fatigue: a longitudinal study. J Appl Physiol. 2023;135(2):345-352. doi:10.1152/japplphysiol.00215.2023

13. Opar DA, Williams MD, Timmins RG, et al. The effect of previous hamstring strain injuries on the change in eccentric hamstring strength during preseason training in elite Australian footballers. Am J Sports Med. 2013;41(3):377-384. doi:10.1177/0363546512470637

14. Bourne MN, Opar DA, Williams MD, et al. Muscle activation patterns and length changes in hamstring muscles during sprinting. J Biomech. 2015;48(11):2688-2695. doi:10.1016/j.jbiomech.2015.06.010

15. Sole G, Milosavljevic S, Nicholson HD, et al. Altered muscle activation following hamstring injury. J Electromyogr Kinesiol. 2011;21(4):457-466. doi:10.1016/j.jelekin.2011.02.002

16. Fyfe JJ, Opar DA, Williams MD, et al. The role of neuromuscular inhibition in hamstring strain injury recurrence. J Electromyogr Kinesiol. 2013;23(3):523-530. doi:10.1016/j.jelekin.2012.12.006

17. Rodriguez RM, Harkey MS, Stanley LE, et al. Corticospinal excitability after ACL reconstruction: a longitudinal study. J Orthop Res. 2021;39(5):1032-1040. doi:10.1002/jor.24892

18. Rio E, Kidgell D, Purdam C, et al. Corticomotor excitability and inhibition following acute Achilles tendon rupture repair. J Sci Med Sport. 2016;19(6):408-413. doi:10.1016/j.jsams.2015.07.009

19. Gibbons CE, Pietrosimone BG, Ingersoll CD, et al. Effects of ACL injury on corticospinal excitability. Clin Biomech. 2010;25(6):486-492. doi:10.1016/j.clinbiomech.2010.02.008

20. Buhmann R, Opar DA, Timmins RG, et al. Intracortical inhibition and hamstring injury: a pilot study. J Sport Rehabil. 2022;31(4):321-328. doi:10.1123/jsr.2021-0215

21. Hermens HJ, Freriks B, Disselhorst-Klug C, et al. Development of recommendations for SEMG sensors and sensor placement procedures. J Electromyogr Kinesiol. 2000;10(5):361-374. doi:10.1016/S1050-6411(00)00027-4

22. Presland JD, Opar DA, Timmins RG, et al. Reliability of TMS measures in hamstring muscles under fatigue. J Neurol Sci. 2023;448:120-127. doi:10.1016/j.jns.2023.120456

23. Guilhem G, Doguet V, Hug F, et al. Effects of acute muscle fatigue on the architecture of the hamstring muscles. J Appl Physiol. 2011;110(4):993-1000. doi:10.1152/japplphysiol.01082.2010

24. Mitsiopoulos N, Baumgartner RN, Heymsfield SB, et al. Cadaver validation of skeletal muscle measurement by magnetic resonance imaging and computerized tomography. J Appl Physiol. 1998;85(1):115-122. doi:10.1152/jappl.1998.85.1.115

25. Gandevia SC. Spinal and supraspinal factors in human muscle fatigue. Physiol Rev. 2001;81(4):1725-1789. doi:10.1152/physrev.2001.81.4.1725

FULL TEXTS BOOK

26. Enoka RM, Duchateau J. Muscle fatigue: what, why and how it influences muscle function. J Physiol. 2008;586(1):11-23. doi:10.1113/jphysiol.2007.139477

27. Sidhu SK, Weavil JC, Mangum TS, et al. Group III/IV locomotor muscle afferents alter motor cortical and corticospinal excitability and promote central fatigue during cycling exercise. Clin Neurophysiol. 2017;128(1):44-55. doi:10.1016/j.clinph.2016.10.411

28. Gruet M, Temesi J, Rupp T, et al. Stimulation of the motor cortex and corticospinal tract to assess human muscle fatigue. Neuroscience. 2013;231:384-399. doi:10.1016/j.neuroscience.2012.11.058

29. Opie GM, Sidhu SK, Rogasch NC, et al. Cortical inhibition assessed using paired-pulse TMS is not changed in older adults after a single bout of resistance exercise. Clin Neurophysiol. 2015;126(6):1075-1082. doi:10.1016/j.clinph.2014.09.016

30. Taylor JL, Allen GM, Butler JE, et al. Effect of contraction strength on responses in biceps brachii and adductor pollicis to transcranial magnetic stimulation. Exp Brain Res. 1997;117(3):472-478. doi:10.1007/s002210050243

31. McNeil CJ, Giesebrecht S, Gandevia SC, et al. Behaviour of the motoneurone pool in a fatiguing submaximal contraction. J Physiol. 2011;589(Pt 14):3533-3544. doi:10.1113/jphysiol.2011.207191

32. Place N, Maffiuletti NA, Martin A, et al. Assessment of the reliability of central and peripheral fatigue after sustained maximal voluntary contraction of the quadriceps muscle. Muscle Nerve. 2007;35(4):486-495. doi:10.1002/mus.20714

33. Kidgell DJ, Pearce AJ. Corticospinal properties following short-term strength training of an intrinsic hand muscle. Hum Mov Sci. 2010;29(5):631-641. doi:10.1016/j.humov.2010.01.004

34. Nuzzo JL, Barry BK, Gandevia SC, et al. Acute strength training increases responses to stimulation of corticospinal axons. Med Sci Sports Exerc. 2016;48(1):139-150. doi:10.1249/MSS.0000000000000733

35. Franchi MV, Atherton PJ, Reeves ND, et al. Architectural, functional and molecular responses to concentric and eccentric loading in human skeletal muscle. Acta Physiol (Oxf). 2014;210(3):642-654. doi:10.1111/apha.12225

FULL TEXTS BOOK

HAMSTRING YARALANMASI OLAN YETIŞKINLERDE KAS MIMARISI, KORTIKOSPINAL UYARILABILIRLIK, HAMSTRING SONUC SKORU VE EKSANTRIK GÜÇ ARASINDAKI İLIŞKILER ÜZERINE PILOT ÇALIŞMA

THE RELATIONSHIP BETWEEN MUSCLE ARCHITECTURE, CORTICOSPINAL EXCITABILITY, HAMSTRING OUTCOME SCORE, AND ECCENTRIC STRENGTH IN ADULTS WITH PRIOR HAMSTRING INJURY: PILOT STUDY

***Caglar SOYLU*[1,2]**

[1]*Associate Professor, University of Health Sciences, Gülhane Faculty of Physiotherapy and Rehabilitation, Department of Orthopedic Physiotherapy and Rehabilitation, Ankara, Türkiye*

[2]*Postdoctoral Fellow, Sports Performance, Recovery, Injury & New Technologies (SPRINT) Research Centre*

Australian Catholic University, Melbourne, Australia

[1,2]*ORCID ID: 0000-0002-1524-6295*

ÖZET

Bu pilot çalışma, tek taraflı hamstring gerilme yaralanması (HSI) öyküsü olan altı yetişkinde biceps femoris long head (BFLH) kasının proksimal ve distal bölgelerindeki muscle architecture, corticospinal excitability (transcranial magnetic stimulation [TMS] ile), Hamstring Outcome Score (HaOS) ve eksantrik hamstring gücü arasındaki ilişkileri inceleyerek yaralanma sonrası nöromüsküler ve fonksiyonel uyarlamaları aydınlatmayı amaçlamıştır. Ultrason görüntüleme, TMS, izokinetik dinamometri ve HaOS kullanılarak fascicle length (FL), pennation angle (PA), muscle thickness (MT), kortikospinal metrikler, eksantrik güç ve hasta bildirimli sonuçlar değerlendirilmiştir. Altı katılımcı (ortalama yaş 26±3 yıl, cinsiyet ve VKİ dengeli) tek seansta yaralanmış ve yaralanmamış bacaklarda test edilmiştir. Ultrason, dinlenmede ve dinamik kontraksiyonlar sırasında (0°-90° diz ROM) FL, PA ve MT ölçmüştür. TMS, motor evoked potential (MEP) amplitüdleri (%Mmax), resting motor threshold (RMT), short-interval intracortical inhibition (SICI) ve intracortical facilitation (ICF) ile kortikospinal uyarılabilirliği değerlendirmiştir. Eksantrik güç 30°/s hızda ölçülmüş, HaOS fonksiyonel durumu değerlendirmiştir. İstatistiksel analizler (Pearson korelasyonları, eşleştirilmiş t-testleri, $p<0.05$, Cohen's d) yaralanmış bacakta anlamlı farklılıklar göstermiştir: FL daha kısa (yaralanmış: 5.7±0.5 cm vs. yaralanmamış: 6.4±0.4 cm; $p<0.01$, $d=1.0$), PA azalmış (yaralanmış: 13.2±1.8° vs. yaralanmamış: 14.8±1.9°; $p<0.05$, $d=0.8$), MT değişmemiştir ($p=0.15$, $d=0.2$). Eksantrik güç düşüktü (yaralanmış: 92.1±24.8 Nm vs. yaralanmamış: 115.6±26.3 Nm; $p<0.01$, $d=0.9$). MEP amplitüdleri azalmış (yaralanmış: 19.5±7.2% Mmax vs. yaralanmamış: 25.8±7.9%; $p<0.05$, $d=0.8$), RMT artmış (yaralanmış: 49.2±5.5% vs. yaralanmamış: 44.5±5.1%; $p<0.05$, $d=0.7$), SICI eğilimli yükselmiştir (yaralanmış: 61.0±14.5% vs. yaralanmamış: 51.8±13.7%; $p=0.08$, $d=0.6$). HaOS puanları (ortalama 80±9) eksantrik güç ($r=0.65$, $p<0.05$) ve FL ($r=0.58$, $p<0.05$) ile orta düzeyde korelasyon göstermiş, MEP ile ilişkili değildir ($r=0.30$, $p=0.20$). Azalmış FL ve güç, yüksek SICI ile ilişkilidir ($r=-0.60$, $p<0.05$). Bulgular, HSI sonrası kalıcı mimari ve nöral eksiklikleri gösterir; fonksiyonel sonuçlar kas yapısı ve gücüyle bağlantılıdır. Rehabilitasyon, kortikal uyarılabilirlik ve eksantrik eğitimi hedeflemeli, iyileşmeyi artırmalı ve yeniden yaralanma riskini azaltmalıdır. Bu ön sonuçları doğrulamak ve nöromodülatör müdahaleleri keşfetmek için daha büyük çalışmalar gereklidir.

Anahtar Kelimeler: Hamstring yaralanması, kas mimarisi, transkraniyal manyetik stimülasyon, kortikospinal uyarılabilirlik, eksantrik kas kuvveti, hasta bildirimli sonuçlar, HaOS skoru, nöromüsküler adaptasyon

FULL TEXTS BOOK

ABSTRACT

This pilot study aimed to investigate the relationships between muscle architecture, corticospinal excitability, Hamstring Outcome Score (HaOS), and eccentric hamstring strength in the proximal and distal regions of the biceps femoris long head (BFLH) among six adults with a prior unilateral hamstring strain injury (HSI) to elucidate post-injury neuromuscular and functional adaptations. Six participants (mean age 26±3 years, balanced for sex and body mass index [BMI]) were assessed in a single session on both injured and uninjured limbs. Muscle architecture (fascicle length [FL], pennation angle [PA], muscle thickness [MT]) was measured via ultrasound, corticospinal excitability via transcranial magnetic stimulation (TMS) (motor evoked potential [MEP] amplitudes, resting motor threshold [RMT], short-interval intracortical inhibition [SICI]), eccentric strength via isokinetic dynamometry (30°/s), and functional status via HaOS. Statistical analyses included paired t-tests and Pearson correlations ($p<0.05$, Cohen's d). The injured limb exhibited shorter FL (5.7±0.5 cm vs. 6.4±0.4 cm; $p<0.01$, $d=1.0$), reduced PA (13.2±1.8° vs. 14.8±1.9°; $p<0.05$, $d=0.8$), and unchanged MT ($p=0.15$, $d=0.2$). Eccentric strength was lower (92.1±24.8 Nm vs. 115.6±26.3 Nm; $p<0.01$, $d=0.9$). MEP amplitudes decreased (19.5±7.2% Mmax vs. 25.8±7.9%; $p<0.05$, $d=0.8$), RMT increased (49.2±5.5% vs. 44.5±5.1%; $p<0.05$, $d=0.7$), and SICI showed a trend toward increase (61.0±14.5% vs. 51.8±13.7%; $p=0.08$, $d=0.6$). HaOS score (mean 80±9) correlated moderately with eccentric strength ($r=0.65$, $p<0.05$) and FL ($r=0.58$, $p<0.05$), but not with MEP ($r=0.30$, $p=0.20$). Reduced FL and strength were negatively associated with higher SICI ($r=-0.60$, $p<0.05$). Findings indicate persistent architectural and neural deficits post-HSI, with functional outcomes linked to muscle structure and strength. Rehabilitation should target cortical excitability and eccentric training to enhance recovery and reduce reinjury risk. Larger studies are required to validate these preliminary findings.

Keywords: Hamstring injury, muscle architecture, transcranial magnetic stimulation, corticospinal excitability, eccentric muscle strength, patient-reported outcomes, HaOS score, neuromuscular adaptation

INTRODUCTION

Hamstring strain injuries (HSI) are a prevalent musculoskeletal issue among athletes and active individuals, often leading to functional limitations and an increased risk of recurrent injury (1). HSI typically occurs in the biceps femoris long head (BFLH) and is associated with high-speed movements or excessive stretching during eccentric contractions (2). These injuries can result in alterations in muscle architecture (fascicle length [FL], pennation angle [PA], muscle thickness [MT]), deficits in neuromuscular control, and persistent changes in corticospinal excitability (assessed via transcranial magnetic stimulation [TMS]) (3, 4). Such modifications may complicate the recovery process and negatively impact functional outcomes (5).

Muscle architecture serves as a fundamental determinant of a muscle's force production capacity and biomechanical properties. Following HSI, adaptations such as shortened FL and reduced PA have been frequently reported, and these changes are linked to decreased eccentric strength (6). Eccentric strength plays a critical role in HSI prevention and rehabilitation, as the ability of hamstring muscles to generate high forces during the eccentric phase can reduce injury risk (7). Additionally, patient-reported outcomes, such as the Hamstring Outcome Score (HaOS), provide a valuable tool for assessing individuals' functional status and their ability to return to daily activities (8).

Neuromuscular adaptations are pivotal in the recovery process following HSI. Corticospinal excitability, measured via TMS, reflects the motor cortex's capacity to regulate muscle activity, with post-HSI changes reported in motor evoked potential (MEP) amplitudes, resting motor threshold (RMT), and

FULL TEXTS BOOK

short-interval intracortical inhibition (SICI) (3). These neural alterations are associated with muscle strength and functional recovery, necessitating their consideration in the development of rehabilitation strategies (4). However, the relationships between muscle architecture, corticospinal excitability, eccentric strength, and patient-reported outcomes remain underexplored, particularly in small-sample pilot studies.

This study aims to investigate the relationships between muscle architecture, corticospinal excitability, HaOS scores, and eccentric strength in the proximal and distal regions of the BFLH among adults with a history of unilateral HSI. Understanding these interconnections may contribute to identifying post-HSI neuromuscular and functional deficits and developing targeted rehabilitation strategies. The hypothesis of this study is that persistent changes in muscle architecture and corticospinal excitability exist post-HSI and are associated with eccentric strength and functional outcomes.

METHODS

This study was supported by the TÜBİTAK 2219 – Postdoctoral Research Fellowship Program Abroad and conducted under standard laboratory conditions with oversight by an experienced researcher. Six adult participants with a history of unilateral hamstring strain injury (HSI) at least six months prior, and clinically fully recovered, were included (2). The mean age of participants was 26 ± 3 years, with a balanced gender distribution (three males, three females). Groups were matched for body mass index (BMI) to minimize the influence of anthropometric differences on outcomes. Participants were selected from active athletes or individuals engaging in regular recreational physical activity, with no acute or chronic musculoskeletal disorders, neurological conditions, or systemic health issues that could affect performance measurements (6). Exclusion criteria included a history of another injury within the last three months, active pain, neurological disorders (e.g., epilepsy), metal implants contraindicating TMS, and cardiovascular risk factors (4). All participants were provided with detailed verbal and written information about the study and signed informed consent forms prior to inclusion.

All assessments were conducted in a quiet laboratory environment at 22–24°C, completed within a single day by the same researcher. Participants were instructed to avoid caffeine, alcohol, and intense physical exercise for at least 24 hours prior to testing. Evaluations were performed in a single session, following the sequence of muscle architecture measurements, corticospinal excitability assessment, eccentric strength testing, and functional status questionnaire administration. This order was designed to minimize muscle fatigue and enhance the accuracy of neuromuscular responses (3).

Muscle Architecture Measurements

Muscle architecture was assessed using a B-mode ultrasound system with a high-resolution linear probe (7–12 MHz) (9). Measurements were taken from the proximal (25% of femur length) and distal (75% of femur length) regions of the biceps femoris long head (BFLH). Participants were positioned prone with knees fully extended and muscles relaxed. Following rest measurements, a slow, controlled eccentric contraction over a 0°–90° knee flexion range (lasting 5–10 seconds) was performed, with real-time ultrasound images recorded. Fascicle length (FL), pennation angle (PA), and muscle thickness (MT) were measured in triplicate per region, with average values used for analysis (6). Image analysis was conducted using ImageJ software by a blinded researcher unaware of the measurement outcomes.

Corticospinal Excitability Measurements

Corticospinal excitability was evaluated using transcranial magnetic stimulation (TMS) (10). The motor cortex region representing the BFLH was targeted with a figure-of-eight coil. Electromyography (EMG) recordings were obtained via surface electrodes placed on the BFLH. Motor evoked potential (MEP) amplitudes were normalized to the maximum M-wave (Mmax), while the resting motor threshold

FULL TEXTS BOOK

(RMT) was defined as the minimum stimulation intensity eliciting an MEP with 50% probability at rest. Short-interval intracortical inhibition (SICI) was assessed with a 2–5 ms interstimulus interval, and intracortical facilitation (ICF) with a 10–15 ms interval (4). Measurements were conducted at rest and during low-intensity isometric contraction, with 10–20 stimuli per parameter.

Eccentric Muscle Strength Measurements

Eccentric strength was tested using an isokinetic dynamometer (Biodex System) (11). Participants were seated with hips flexed at 90° and lower extremities stabilized with straps. Measurements were performed at 30°/s angular velocity, with five maximal eccentric contractions per leg, and the highest peak torque (Nm) recorded for analysis (7). A submaksimal warm-up protocol was implemented prior to testing to prepare the neuromuscular system.

Functional Status Assessment

Functional status was evaluated using the Hamstring Outcome Score (HaOS) questionnaire (8). HaOS is a 0–100 scale assessing symptoms, function, sports participation, and quality of life, with higher scores indicating better outcomes. Participants completed the questionnaire during the test session in a quiet laboratory setting based on self-reports (12).

Statistical Analysis

Data were analyzed using IBM SPSS Statistics v.26 (Chicago, IL, USA). Continuous variables were reported as mean ± standard deviation (SD), and categorical variables as frequency and percentage (%). Differences between injured and uninjured limbs were assessed with paired t-tests. Relationships between muscle architecture, corticospinal excitability, eccentric strength, and HaOS scores were evaluated using Pearson correlation coefficients (r), with correlation strength interpreted based on r values. Statistical significance was set at $p < 0.05$, and effect sizes for two-tailed comparisons were calculated using Cohen's d, categorized as small (0.2), medium (0.5), or large (0.8). Normality was verified with the Shapiro–Wilk test, and outliers (beyond ±3 SD) identified via box-plot inspection were excluded from analysis.

RESULTS

Statistical analyses, including paired t-tests and Pearson correlations, revealed significant neuromuscular and functional deficits in the injured limb compared to the uninjured limb among the six participants with a prior unilateral hamstring strain injury (HSI). The mean Hamstring Outcome Score (HaOS) was 80 ± 9, reflecting moderate self-reported functional status, with scores indicating lingering symptoms or limitations in daily and sports activities despite clinical recovery. These preliminary findings from a small pilot sample highlight persistent post-injury adaptations, which may contribute to reinjury risk. Below, we detail the results by category, with references to Tables 1 and 2 for quantitative data, followed by interpretive commentary on clinical implications.

Muscle Architecture

Ultrasound imaging of the biceps femoris long head (BFLH) demonstrated clear architectural remodeling in the injured limb. As shown in Table 1, fascicle length (FL) was significantly shorter in the injured limb (5.7 ± 0.5 cm vs. 6.4 ± 0.4 cm in the uninjured limb; mean difference = 0.7 cm, $p < 0.01$, Cohen's d = 1.0), representing an approximately 11% reduction with a large effect size. This shortening could alter the muscle's length-tension relationship, reducing its ability to absorb eccentric forces during high-speed activities like sprinting, thereby elevating reinjury vulnerability. Similarly, pennation angle (PA) was reduced (13.2 ± 1.8° vs. 14.8 ± 1.9°; mean difference = 1.6°, $p < 0.05$, Cohen's d = 0.8), suggesting a medium-large effect that may compromise force transmission efficiency through altered fiber orientation. In contrast, muscle thickness (MT) showed no significant change (2.3 ± 0.3 cm

FULL TEXTS BOOK

vs. 2.4 ± 0.3 cm; p = 0.15, Cohen's d = 0.2), indicating that hypertrophy or atrophy was not a prominent feature post-HSI in this cohort.

These architectural changes align with prior literature on chronic muscle adaptations after strain injuries, where shorter fascicles are linked to reduced extensibility and increased stiffness. Clinically, this implies that rehabilitation programs should incorporate eccentric loading and stretching to potentially elongate fascicles and mitigate these deficits, though larger studies are needed to confirm causality.

Eccentric Strength

Isokinetic testing at 30°/s revealed substantial weakness in the injured limb's hamstring muscles. Table 1 illustrates a significant reduction in eccentric strength (92.1 ± 24.8 Nm vs. 115.6 ± 26.3 Nm; mean difference = 23.5 Nm, p < 0.01, Cohen's d = 0.9), denoting a large effect size and approximately 20% deficit. This weakness is particularly concerning for functional tasks involving deceleration, such as cutting maneuvers in sports, where eccentric hamstring function is critical for knee stability and injury prevention.

The observed strength asymmetry (injured limb ~80% of uninjured) exceeds typical thresholds for return-to-sport clearance (often >90%), suggesting incomplete recovery despite participants being asymptomatic. This finding underscores the need for targeted eccentric training protocols, like Nordic hamstring exercises, to restore strength and reduce reinjury rates, which can be as high as 30% in HSI cases.

Corticospinal Excitability

Transcranial magnetic stimulation (TMS) metrics indicated neural impairments, as detailed in Table 1. Motor evoked potential (MEP) amplitudes were lower in the injured limb (19.5 ± 7.2% Mmax vs. 25.8 ± 7.9%; mean difference = 6.3% Mmax, p < 0.05, Cohen's d = 0.8), reflecting reduced corticospinal drive and potential central nervous system adaptations post-injury. Resting motor threshold (RMT) was elevated (49.2 ± 5.5% vs. 44.5 ± 5.1%; mean difference = -4.7%, p < 0.05, Cohen's d = 0.7), implying a higher stimulus intensity required for muscle activation, which could manifest as delayed or inefficient motor responses. Short interval intracortical inhibition (SICI) trended toward increase (61.0 ± 14.5% vs. 51.8 ± 13.7%; mean difference = -9.2%, p = 0.08, Cohen's d = 0.6), suggesting enhanced inhibitory circuits that might hinder voluntary contraction. Intracortical facilitation (ICF) showed no difference (120.0 ± 15.0% vs. 122.0 ± 14.5%; p = 0.35, Cohen's d = 0.1), indicating preserved excitatory pathways.

These neural changes point to maladaptive cortical reorganization, where increased inhibition (SICI) may contribute to protective mechanisms but at the cost of performance. From a rehabilitation perspective, neuromodulatory interventions like repetitive TMS could be explored to enhance excitability and facilitate recovery, especially given the borderline significance of SICI.

Table 1. Comparisons of Key Parameters Between Injured and Uninjured Limbs

Parameter	Injured Value	Uninjured Value	Mean Difference (Uninjured - Injured)	p-Value	Cohen's d	Interpretation (Effect Size and Significance)
Fascicle Length (FL)	5.7 ± 0.5 cm	6.4 ± 0.4 cm	0.7 cm	<0.01	1.0	Large effect; significant shortening (persistent fascicle shortening post-

FULL TEXTS BOOK

						injury, may impair force absorption).
Pennation Angle (PA)	13.2 ± 1.8°	14.8 ± 1.9°	1.6°	<0.05	0.8	Medium-large effect; significant reduction (changes in fiber angle may affect force production efficiency).
Muscle Thickness (MT)	2.3 ± 0.3 cm	2.4 ± 0.3 cm	0.1 cm	0.15	0.2	Small effect; no significant difference (minimal injury effect on muscle thickness, no evident atrophy).
Eccentric Strength	92.1 ± 24.8 Nm	115.6 ± 26.3 Nm	23.5 Nm	<0.01	0.9	Large effect; significant reduction (persistent weakness in eccentric contraction, increases reinjury risk during dynamic activities).
MEP Amplitudes	19.5 ± 7.2% Mmax	25.8 ± 7.9% Mmax	6.3% Mmax	<0.05	0.8	Medium-large effect; significant reduction (decrease in corticospinal excitability, indicating neural adaptation deficit).
Resting Motor Threshold (RMT)	49.2 ± 5.5%	44.5 ± 5.1%	-4.7%	<0.05	0.7	Medium effect; significant increase (elevated motor threshold, leading to difficulty in muscle activation and potential motor delays).
Short-Interval Intracortical Inhibition (SICI)	61.0 ± 14.5%	51.8 ± 13.7%	-9.2%	0.08	0.6	Medium effect; trending increase (increased inhibition post-injury, borderline significant; may

						suppress motor control).
Intracortical Facilitation (ICF)	120.0 ± 15.0%	122.0 ± 14.5%	2.0%	0.35	0.1	Very small effect; no significant difference (minimal injury effect on ICF, excitatory pathways largely intact).

Abbreviations: FL = Fascicle Length; PA = Pennation Angle; MT = Muscle Thickness; MEP = Motor Evoked Potential; RMT = Resting Motor Threshold; SICI = Short-Interval Intracortical Inhibition; ICF = Intracortical Facilitation; Mmax = Maximum M-wave amplitude; Nm = Newton-meters; ° = Degrees. Values are presented as mean ± standard deviation (SD). Cohen's d: small (0.2), medium (0.5), large (≥0.8). All comparisons used paired t-tests.

Correlations

Interrelationships among variables were examined via Pearson correlations (Table 2). HaOS scores correlated moderately with eccentric strength ($r = 0.65$, $p < 0.05$) and FL ($r = 0.58$, $p < 0.05$), suggesting that better muscle structure and strength predict improved patient-reported outcomes. A stronger correlation existed between FL and eccentric strength ($r = 0.70$, $p < 0.05$), reinforcing the biomechanical link where longer fascicles support greater force generation. Neural factors showed negative associations: reduced FL and strength both correlated with higher SICI ($r = -0.60$ for each, $p < 0.05$), indicating that increased cortical inhibition may exacerbate architectural and strength deficits. MEP amplitudes negatively correlated with SICI ($r = -0.55$, $p < 0.05$), highlighting inhibitory suppression of excitability. Non-significant correlations included HaOS with MEP ($r = 0.30$, $p = 0.20$) or RMT ($r = -0.25$, $p = 0.25$), implying neural metrics influence function indirectly.

These correlations suggest interconnected neuromuscular mechanisms, where neural inhibition (SICI) acts as a mediator of structural and functional impairments. Clinically, this supports multimodal rehabilitation integrating neural stimulation with strength training to break these negative cycles. However, the small sample limits generalizability, and future longitudinal studies could explore causality.

Table 2. Correlations Among Key Parameters

Correlation Pair	Correlation Coefficient (r)	p-Value	Correlation Strength	Interpretation (Relationship Meaning and Clinical Importance)
HaOS Score - Eccentric Strength	0.65	<0.05	Medium	Positive medium correlation; higher strength associated with better functional outcomes (HaOS) (prioritize eccentric training to enhance patient-reported recovery).
HaOS Score - Fascicle Length (FL)	0.58	<0.05	Medium	Positive medium correlation; longer FL linked to better HaOS scores (muscle architecture directly influences perceived function and quality of life).

FULL TEXTS BOOK

HaOS Score - MEP Amplitudes	0.30	0.20	Weak	Non-significant correlation; MEP changes not directly related to HaOS (neural excitability may affect function through secondary pathways, warranting further investigation).
Fascicle Length (FL) - Short-Interval Intracortical Inhibition (SICI)	-0.60	<0.05	Medium	Negative medium correlation; shorter FL associated with increased SICI (neural inhibition may drive or reflect architectural maladaptations).
Eccentric Strength - Short-Interval Intracortical Inhibition (SICI)	-0.60	<0.05	Medium	Negative medium correlation; lower strength linked to increased SICI (inhibition contributes to strength loss; neuromodulation could be a therapeutic target).
Fascicle Length (FL) - Eccentric Strength	0.70	<0.05	Medium-High	Positive correlation; longer FL associated with higher eccentric strength (strong biomechanical interdependence; interventions targeting fascicle length may boost strength).
MEP Amplitudes - Short-Interval Intracortical Inhibition (SICI)	-0.55	<0.05	Medium	Negative correlation; lower MEP linked to increased SICI (inhibitory processes suppress overall corticospinal output, impacting motor efficiency).
HaOS Score - Resting Motor Threshold (RMT)	-0.25	0.25	Weak	Non-significant correlation; RMT changes weakly related to HaOS (elevated thresholds may subtly hinder function but require larger samples for confirmation).

Abbreviations: HaOS = Hamstring Outcome Score; FL = Fascicle Length; MEP = Motor Evoked Potential; SICI = Short-Interval Intracortical Inhibition; RMT = Resting Motor Threshold.
Correlation strength: weak ($|r| < 0.3$), medium ($0.3 \leq |r| < 0.7$), high ($|r| \geq 0.7$). All analyses used Pearson's r; $p < 0.05$ indicates statistical significance.

In summary, this pilot study elucidates persistent deficits post-HSI, with functional outcomes closely tied to muscle architecture and strength, modulated by neural factors. These insights advocate for integrated rehabilitation approaches, though confirmation in larger cohorts is essential.

DISCUSSION

This pilot study reveals persistent neuromuscular alterations following unilateral hamstring strain injury (HSI), including shortened fascicle length (FL), reduced pennation angle (PA), diminished eccentric strength, and compromised corticospinal excitability (e.g., reduced MEP amplitudes and increased

FULL TEXTS BOOK

RMT), as detailed in Table 1. These findings align with prior research indicating that HSI leads to chronic neuromuscular deficits and heightened reinjury risk (13, 14). The observed ~11% reduction in FL and ~20% decrease in eccentric strength (Table 1) suggest impaired stretch tolerance, which may predispose individuals to reinjury during high-speed activities such as sprinting (6, 15). Correlation analyses (Table 2) further highlight that HaOS scores are moderately linked to eccentric strength ($r = 0.65$) and FL ($r = 0.58$), while increased short interval intracortical inhibition (SICI, $r = -0.60$) negatively impacts these outcomes, supporting the role of cortical reorganization in functional recovery (4, 16).

Clinically, these results advocate for rehabilitation strategies that combine eccentric training with neuromodulatory interventions (e.g., transcranial magnetic stimulation) to address both muscle architecture and neural inhibition, moving beyond traditional strength-focused approaches (5, 7). However, the small sample size ($n = 6$) and cross-sectional design limit the generalizability of these findings. Larger, longitudinal studies are needed to validate these preliminary insights and assess the efficacy of integrated therapies in enhancing HSI recovery.

CONCLUSION

This pilot study demonstrates persistent neuromuscular deficits post-HSI, including altered muscle architecture, reduced eccentric strength, and impaired corticospinal excitability as shown in Table 1. Functional outcomes (HaOS) are positively associated with muscle structure and strength, while neural inhibition (SICI) exerts a negative influence (Table 2). These findings support the integration of eccentric training and neuromodulatory techniques in rehabilitation to mitigate reinjury risk and improve recovery. Despite the small sample limiting broader applicability, this study lays the groundwork for future research to explore long-term effects of combined interventions on HSI management.

REFERENCES

1. Bourne MN, Timmins RG, Opar DA, et al. An evidence-based framework for strengthening exercises to prevent hamstring injury. Sports Med. 2018;48(2):251-267. doi:10.1007/s40279-017-0796-x

2. Opar DA, Williams MD, Shield AJ. Hamstring strain injuries: factors that lead to injury and re-injury. Sports Med. 2012;42(3):209-226. doi:10.2165/11594800-000000000-00000

3. Maniar N, Carmichael DS, Hickey JT, et al. Incidence and prevalence of hamstring injuries in field-based team sports: a systematic review and meta-analysis. J Sci Med Sport. 2021;24(1):12-20. doi:10.1016/j.jsams.2020.06.011

4. Fyfe JJ, Opar DA, Williams MD, Shield AJ. The role of neuromuscular inhibition in hamstring strain injury recurrence. J Electromyogr Kinesiol. 2013;23(3):523-530. doi:10.1016/j.jelekin.2012.12.006

5. Hickey JT, Timmins RG, Maniar N, et al. Criteria for progressing rehabilitation and determining return-to-play clearance following hamstring strain injury. Sports Med. 2020;50(1):95-108. doi:10.1007/s40279-019-01192-9

6. Timmins RG, Bourne MN, Shield AJ, et al. Short biceps femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite football (soccer): a prospective cohort study. Br J Sports Med. 2016;50(24):1524-1535. doi:10.1136/bjsports-2015-095362

7. Opar DA, Piatkowski T, Williams MD, Shield AJ. A novel device using the Nordic hamstring exercise to assess eccentric hamstring strength: a reliability and retrospective injury study. J Orthop Sports Phys Ther. 2015;45(9):773-779. doi:10.2519/jospt.2015.5776

FULL TEXTS BOOK

8. Cacchio A, Rompe JD, Furia JP, et al. Validation of the Hamstring Outcome Score for the evaluation of hamstring injuries. Am J Sports Med. 2014;42(11):2687-2694. doi:10.1177/0363546514544687
9. Kosovalija B, Franchi MV, Spörri J, et al. Hamstring muscle architecture assessed sonographically using wide field of view ultrasound. PLoS One. 2022;17(11):e0277400. doi:10.1371/journal.pone.0277400
10. Presland JD, Tofari PJ, Timmins RG, et al. Reliability of corticospinal excitability and intracortical inhibition in biceps femoris during different contraction modes. Brain Sci. 2023;13(4):634. doi:10.3390/brainsci13040634
11. Chapon J, Alhammoud M, Lefevre N, et al. A critical evaluation of eccentric hamstring muscle strength assessment: A narrative review. Sports Med Open. 2020;6(1):8. doi:10.1186/s40798-020-0236-9
12. van Doormaal MC, van der Horst N, Backx FJG, et al. The prognostic value of the hamstring outcome score to predict the risk of hamstring injuries. J Sci Med Sport. 2021;24(4):361-366. doi:10.1016/j.jsams.2020.10.004
13. Heiderscheit BC, Sherry MA, Silder A, et al. Hamstring strain injuries: recommendations for diagnosis, rehabilitation, and injury prevention. J Orthop Sports Phys Ther. 2010;40(2):67-81. doi:10.2519/jospt.2010.3047
14. Green B, Pizzari T. The efficacy of exercise rehabilitation in restoring impairments in psychosocial outcomes following acute hamstring muscle injuries: a systematic review. Phys Ther Sport. 2021;49:185-195. doi:10.1016/j.ptsp.2021.02.015
15. Maniar N, Shield AJ, Williams MD, et al. Hamstring strength and architectural adaptations following inertial flywheel resistance training. J Sci Med Sport. 2021;24(8):746-752. doi:10.1016/j.jsams.2021.02.003
16. Presland JD, Timmins RG, Bourne MN, et al. The effect of Nordic hamstring exercise training volume on biceps femoris long head architectural adaptation. Scand J Med Sci Sports. 2018;28(7):1775-1783. doi:10.1111/sms.13227

FULL TEXTS BOOK

THE SECULAR *TADAVI BHIL* TRIBE: HISTORY AND CULTURE

Dr. Dhanraj T. Dhangar

Assistant Professor, Dept. of Marathi,

MGV's Arts and Commerce College, Yeola

Dist.-Nashik, (M.S.) India.

ABSTRACT

This research paper explores the unique syncretic identity of the 'Tadvi Bhil' tribe, commonly known as 'Tadvi Muslims', a tribal community primarily found in the Khandesh region of Maharashtra and parts of Madhya Pradesh. The study investigates the tribe's historical transformation, tracing their origins from the Adivasi Bhil community and their partial conversion to Islam during the Mughal era, particularly under Aurangzeb's reign. Despite adopting certain Islamic practices, the Tadvis have preserved their tribal traditions, Hindu rituals, and nature-centric worship. Through a detailed analysis of their religious practices, folk traditions, marriage customs, burial rituals, and the Tadvi dialect, the paper highlights how the community maintains a secular and pluralistic cultural identity. This research underscores the coexistence of tribal, Hindu, and Muslim elements within the Tadvi way of life and argues that their secular worldview provides a valuable model for cultural harmony in contemporary India. The paper also emphasizes the urgent need to document and preserve the Tadvi dialect and folklore, which are at risk due to modernization and linguistic assimilation.

Keywords: Tadvi Bhil, syncretic identity, secular tribe, tribal culture, Khandesh, Islamic conversion, nature worship, Tadvi dialect, cultural harmony.

Introduction:

'*Tadvi Bhil*' or 'Tadvi Muslims' is one of the secular tribe in India. Specifically in Jalgaon district of Maharashtra State. However, this tribe is scattered up to Burhanpur district in Madhya Pradesh State. It appears to be spread out under the shelter of the *Satpuda* range and live natural lifestyle in the presence of nature. This region is also called as Khandesh. The people of 'Tadvi' tribe follows Hindu traditions as well as Muslim traditions and also follows their original Adivasi tribal traditions. By this paper researcher trying to find out the reasons behind their secular thinking and traditions. In the era of global religious divisions and the resulting violence the exploration of this secular tradition can be significant for the creation of a humanistic world.

The History of *Tadavi Bhil* Tribes :

The origin of the word *Tadvi* is said to be a tribe that separated from the *Adiwasi Bhil* tribe. A tribe that fell into '*Tad*' (divorced from the original Community). During the time of Mughal Emperor Aurangzeb, the *Tadvi* converted to Islam, hence the original *Bhil* tribe fell into '*Tad*' and they were called '*Tadvi Bhil'*. However, Aziz Tadvi, a scholar of the Tadvi dialect, notes that *"The chief of the Bhil Community of that time in central India was called 'Tadvi' and his followers followed his orders with full faith. 'Tarbi' is an Arabic word and its meaning is a guide."*[1] Even if the theory given by Shri Aziz Tadvi is accepted as true, the leaders of the original *Bhil* tribes who converted to Islam due to conversion may have been given chieftaincy as a reward and called them '*Tarbi*'. Because there is no other reason for the original tribal *Bhil* community to come into contact with the Arabic language. Moreover, it seems that their original relatives still have a feeling of contempt for the converted. And since the original

FULL TEXTS BOOK

Muslims considered themselves superior after conversion of this new tribal people, it was not possible for this converted community to achieve unity with Islam. Since the Muslims also discriminated them, the '*Tad*' (the feelings of the discriminations from the new community) between them remained. Therefore, the awareness of the 'Tad' left from the original tribal *Bhil* community and the '*Tad*' left from Islam seems more likely that the community was addressed as *'Tadvi'* community. In Muslim understanding too, this community seems to have developed a similar identity as *'Tadvi Muslims'*. In the Maharashtra dictionary, there is an entry as *"Tadvi- Bhils who converted to Muslims"*[2]. In the Jalgaon District Gazetteer, it is recorded that *"Tadvi is the descendant of Bhils."*[3].

Why did the tribal *Bhil*s convert to Islam? :

This is a controversial issue. History also mentions that during the reign of the Mughal Emperor Aurangzeb (1685 to 1707), Aurangzeb forced Hindus to convert to Islam by imposing unjust taxes like *Jiziya*. It is also believed that some *Bhil*s may have converted to Islam after getting tired of Aurangzeb's intimidation and oppression. There is also a belief that the tribal *Bhil*s may have been converted to Islam by giving them land as *Vatan* (donated or allocated land) and offering them temptations. According to Mr. Aziz Tadvi. *"440 years ago, the Tadvis must have accepted Islam because of the influence of the teachings of the Sufi saints of Islam, namely Khwaja Garib Nawaz, as well as his compassionate, tolerant, all-religious, and socially enlightening thoughts."*[4] However, if this were the case, the Tadvis would have accepted Islam comprehensively. However, it is highly likely that this conversion took place due to coercion. That is why they have not abandoned their original tribal and traditional Hindu customs, traditions, festivals, and rituals.

Religious life of the Tribal Tadvi *Bhil* tribe:

As mentioned above, the tribal *Tadvi* tribe celebrates festivals of Hindu and Muslim traditions. Hindu and Muslim gods and goddesses are also worshipped in this community. Also, according to tribal traditions, various elements of nature are also worshipped in this community. Regarding their religious life, Aziz Tadvi says, *"The tribal Tadvi Bhils are not Hindus, nor are they Muslims at all. They practice their Bhil culture, Tadvi culture. They live according to the customs, traditions, and culture prevalent in their culture and practiced from time immemorial; they are proud of their Tadvi culture, self-respect. But since they have been influenced by some Hindu traditions, as well as some Islamic traditions, they follow them. But they have not abandoned their original tribal culture, Tadvi culture. They follow our culture even in the twenty-first century. It would not be entirely correct to call them Muslims or Hindus."*[5] The above opinion of Aziz Tadvi, as a scholar of the Tadvi dialect, is correct, but Hindu seems to have come into existence initially as a regional term. Similarly, in tribal life, the sun, moon, rivers, trees, sky, planets, stars, animals, birds, and creatures are all considered Gods. According to the principle of 'whatever is seen is considered as God', the tradition of nature worship among tribal has been present in Hinduism from the beginning. All the nature deities that tribal's worship, such as *Nagadev* (Snake as God), *Suryadev (Sun as God)*, and Dongardev (Mountain as God), also have a placed in Hinduism. The story '*Indra* (Prince of the God as Hindu Mythology) was angry because *Shri Krishna* stopped worshipping *Indra* and asked his people worshipped *Govardhan* Mountain' appears in the biography of *Shri Krishna*. Therefore, it can be concluded that tribal are basically Hindus. In recent times, when feelings about religion have become very sharp, the idea that tribal are not Hindus is instilled in them for conversion and conversion is done by taking advantage of their poverty and ignorance. At present, like all tribal, nature worship is also practiced in the *Tadvi* tribe, so they should be called tribal or *Adiwasi*. Also, the tribal *Tadvi*s who celebrate festivals of both Hindu and Muslim religious traditions can be called truly broadminded.

FULL TEXTS BOOK

Although some of the tribal *Bhil* community have converted to Islam and become 'Tadvis', they have not abandoned their original Sanatan Hindu religious practices and traditions. The tribal Tadvis still worship 'Mahadev and Parvati' as 'Adi Pita and Adi Mata'. Even today, on the twelfth day after the birth of a child, the *Barse*, the naming ceremony of the child is held as per Hindu custom. *Tadavi* women sing a song as…

Tadavi Dialect Script	**English Translate Script**
Jhulā bāndhalā saṭavī māya	The swing is tied by the Godess Sati
sahādēva lyā	Take *Sahadev*
mahādēva dyā	Give *Mahadev*
Jhulā bāndhalā saṭavī māya	The swing is tied by the Godess Sati
Mahādēva lyā	Give M*ahadev*
Sahādēva ghyā	Take *Sahadev*

This is how the naming ceremony is performed according to Hindu custom. This ritual is performed by five women of full age. It is also believed that at night, a pen and book are placed under the head of the child, and Sati/Parvati (Hindu Godess) comes and writes the child's future on it, giving blessings. Similarly, the family deities of Tadvi are also worshipped by the Hindu gods and goddesses like *Khanderao, Bhavani, Bahirambuva, Bahiroba, Bhairavnath, Manudevi* of Adgaon (Taluka Yaval), *Ichchadevi* near Burhanpur, *Ashapuri Mata* of Ashirgad, *Saptashrungi Devi* of Vani, *Jogeshwari Mata* of Ghatnandra, etc. These are the 'family deities' of Tadvi.

In short, considering the religious life of the tribal Tadvi, the tribal Tadvi, who are nature worshippers, originally belong to the tribal Bhil tribe and worship Hindu gods and goddesses due to their ancient connection with the ancient Hindu culture. They still consider the Hindu family deities as their family deities. However, after accepting Islam during the reign of Aurangzeb, they also celebrate Muslim Festivals and also their original festivals as tribals. It can also be believed that the influence of the tolerant and all-religious teachings of the Sufi saints is definitely visible on the Tadvi in this regard. The Tadvi have also adopted Muslim religious rituals such as visiting the Shrines of these Sufi Saints and offering *Chadar*, offering prayers, and reading the Quran-e-Sharif. They are also seen performing the rituals of *Eid-e-Milad* and *Ramjaan* in the Muslim tradition. However, these Tadvi peoples have similar beliefs about Hindu and Muslim religions, also seem to be proud of their tribal traditions.

Tadvi Folk Dance *'Sajoni'*:

This folk dance is performed in the Tadvi community to celebrate various happy occasions. Especially during marriages, young women dance '*Sajoni'* with their friends. Similarly, when the '*Biddi*' of the bridegroom is taken around the village, the young women perform the '*Sajoni*' folk dance by making a circle and circling around the musical instrument player to the beat of the instruments. Similarly, old women and men also dance in a circle. Young men and women perform *'Sajoni'* dance during *Nag Diwali*, fairs, when the harvest is abundant or during the *Akshaya Tritiya* festival.

Youth House (*'Chorvani Jhopdi'*) of Tadvi Bhil Tribe and Marriage Practice:

Like the tribal *Bhil* society, this *Chorvani Jhopdi* (Youth House) used to have special importance in the Tadvi tribe. Such a hut was built outside the village in an unknown place in the forest, where no one would see it. It was decorated with leaves and flowers from the inside. The leader of the Tadvi tribe

FULL TEXTS BOOK

that's called the *Mhorkya* for man leader or *Mhorki* for woman leader of the village would bring unmarried young men and women here. Here, the young men and women would mix freely with each other. They would eat, drink, play various games and dance together and even sleep with the partner they liked. In fact, this was a test of the manhood and womanhood of the partner they liked. This *Chorvani Jhopdi* was also called a bedchamber, a *gotul*, a youth house. These young men and women were also assigned tasks such as labour, protection, hunting, protecting the farm from wild animals - jackals, wild boars, etc. The young men and women used to communicate with the senior members of the family regarding the marriage with the partner they had chosen. However, this tradition of the *Chorvani Jhopadi* is no longer in the Tadvi community. In the Tadvis community, the engagement party is called *'Phutane'*. The wedding date is decided by the Brahmin and the auspicious time is fixed. Earlier, these marriages were done according to tribal tradition. The *Bhagat* (Priest) used to decide the wedding date and used to arrange the marriage. This child marriage method was also called *'Tika Method'*. Also different types of marriage methods are found among the Tadvi tribal. Among them, there are *'Ate-Mama-Bhavanda Vivah'* (Marriage with maternal cosines), *'Jija-Sali Vivah'* (Marriage with sister in law) as per Hindu customs, along with different marriage methods like *'Vinimaya Vivah'* (Marriage as exchange daughter between two families as daughter in law), *'Sammati Vivah'* (Marriage with consideration of both families), *'Sahapalayan Vivah'*(To run away and get married with loved one), *'Parikshan Vivah'* (Marriage by selection of the partner in youth house), 'Hattagaman Vivah' (Marriage forcefully even man has already married), *'Tika Vivah'* (The agreement of marriage while women is pregnant that if born child will be a female she will become daughter in law of relative family) these marriage practices are as per tribal customs. Currently, most of these tradition have disappeared and *'Nikah'* is done according to the Muslim Marriage method. In some places, they still secretly marry according to the *'Tika Method'* and later bring the bride and groom to the *Mandava* (Place of the Marriage or wedding pavillion) to arrange the '*Nikah*'. During the marriage, the bride and groom stay in the *Mandava* and the bride stay in the house. Tadavi and Muslims can eat together but marriage not possible in these two communities. Although the '*Talaq* Method' (Method of divorced in Muslim Community) is not followed in this tribe. Earlier, '*Farkat*' (Method of divorced in Hindu Community) was done in front of the *Caste Panchayat*, now it is done in the court as per Indian Law.

'Burial Rituals or Mortal Rituals' among the Tribal Tadvi Bhils:

The tribal Bhils and Tadvi tribes do not cremate the body after death, but rather perform a 'burial ritual'. The Tadvi Bhil community has a separate place for 'burial rituals'. This burial ground is called '*Asaba*'. The last rites are performed on the fortieth day after the death of a person. It is called '*Chehalam*'. On this day, the family and relatives of the deceased person come together. At night, they perform a program of singing hymns and praises in the name of Hajj, Prophet Muhammad, Allah, and pray for the salvation of the deceased. The next morning, they serve a meal of cow mutton or dal-rice-roti.

The Tadvi Dialect :

The Tadvi dialect is a dialect of Tadavi Community. Tadavi Dialect is a subject of independent research. Since the Tadvi are originally Bhil tribes, the Tadvi dialect contains a large number of words from the Bhil dialect. For example, *Abdhaki* (suddenly), *Dulli* (a basket of bread or breadcrumbs), *Adhayakandya* (a person who behave like browler), Achhwa*ni* (a juce of aurvedic jadibuti for new born baby), *Akaphoka* (simple in bahaviour), *Ehin* (I will come), *Ukhalla* (dustbeen), *Kirrasan* (farmer), *Taril* (yours), *Lavati* (wife), *Daydamra* (father), *Lokya* (husband), *Pekna* (the coming of a goddess or a saint), *Balli* (mountain), *Kavarale Hati* (become greedy), *Tissuk* (a triangle), *Chingel* (an eccentric or stubborn person), *Dapadel* (hidden or secret), *Passu* (a sleepy person), *Patolya* (a card player, a gambler),

FULL TEXTS BOOK

Chavanya Jodne (hitting in the ear / slapped on the cheek), *Chandheya* (the wall behind the house), *Dafla-Dafli* (old man and old women) are some words found in the original Bhil dialect.

Due to the adoption of Islam during the time of Aurangzeb and the association of *Pir-Fakirs*, some Arabic-Persian words are also found in this dialect, but their quantity is less and the form of those words is not like Arabic-Persian but like the *Dakhkhani*, Muslim dialect. The region where the *Tadvis* live is part of *Khandesh.* The Ahirani dialect is the main dialect of this region. Especially in Chopda taluka, where the Tadvis live, is *Ahirani*-speaking, the influence of Ahirani is also found on the Tadvis dialect.The *Leva* community lives in large numbers in Yawal, Raver, and Burhanpur talukas, where the *Tadvis* live. Since agriculture is the main occupation of the Leva community and they come into contact with the *Leva* community for agriculture and labour, the influence of the Leva dialect is also found on the *Tadvis* dialect. However, despite the similarity in words, there is a difference in the accent of *Tadvi* dialect and the accent of *Leva* dialect in pronunciation. Therefore, *Tadvi, Leva* and *Ahirani* speakers can be identified separately.

In modern times, due to the increasing level of education in this society, standard Marathi words are also pronounced by the mouths of Tadvi dialect speakers. Especially, the children of the Tadvi tribe in young and teenagers who learns in the schools and colleges speak Ahirani mixed Standard Marathi Language in the *Ahirani* linguistic belt, while in the Leva linguistic belt, they speak *Leva* dialect mixed with Standard Marathi Language of Maharashtra State. However, some words are especially illuminating the distinctive nature of Tadvi dialect. A more in-depth analysis of Tadvi dialect can be presented by conducting linguistic and grammatical studies. Apart from daily usage, *Tadvi* also speaks sentences with some hidden meanings. The dialect of the Hindu tribal as well as the Muslim tribal who cultivate the culture, the grammatical structure of the dialect, and the folklore are all challenging to study. As the number of native speakers is decreasing due to the level of education, it has become imperative to preserve and promote this dialect and document its folklore.

Conclusion :

The Tadavi Bhil tribe stands as a remarkable example of syncretic cultural identity in India. Despite historical pressures of conversion particularly during the Mughal era under Aurangzeb. The Tadavis have preserved their ancestral tribal customs while selectively incorporating elements from both Hinduism and Islam. Their religious life, festivals, rituals, and family deities reflect an inclusive worldview rooted in nature worship, tribal ethos, and a deep reverence for cultural continuity.

The community's practices, such as celebrating both Hindu and Muslim festivals, maintaining traditional tribal marriage customs alongside *Nikah* ceremonies, and performing burial rites with Islamic and tribal influences, demonstrate a distinct cultural resilience. The preservation of the Tadvi dialect, with its layers of Bhil, *Ahirani, Leva,* and *Dakhni* influences, adds another rich dimension to their identity and calls for urgent linguistic documentation.

The now-vanished tradition of the *Chorvani Jhopdi* and the continued practice of the folk dance *'Sajoni'* further illustrate how the *Tadavis* have maintained their indigenous cultural expressions even amidst socio-religious change. This study reveals that the Tadavi tribe embodies a rare form of secular tribal consciousness not in opposition to religion, but by harmonizing diverse religious influences into a cohesive cultural practice.

In an age of increasing communal polarization and identity-based politics, the Tadavi community offers a valuable model of peaceful coexistence, cultural plurality, and spiritual tolerance. Recognizing and preserving the unique cultural heritage of such communities is essential not only for academic purposes but also for promoting a more inclusive, humane, and pluralistic society.

FULL TEXTS BOOK

REFERENCES

1. Tadavi, Aziz. Aadivasi Tadavi Bolibhashetīl Shabda Sangrah Kosh. Jalgaon: Paanphul Prakashan, 2011. p. 11.

2. Ketkar, Shree Vyaṁ. Maharashtra Shabdakosh. Vol. 4, Maharashtra Rajya Sahitya Sanskruti Mandal, Mumbai, p. 1456.

3. Jalgaon Jilha Gazetteer. "Pramukh Jati-Jamati." Government of Maharashtra, p. 205.

4. Tadavi, Aziz. Aadivasi Tadavi Bhill: Samaj Aani Sanskruti. Beed: Harshvardhan Publications, Dec. 2016, p. 58.

5. Ibid. p. 53.

FULL TEXTS BOOK

TÜRKİYE EKONOMİSİNİN RİSK DEĞERLERİNİN YUNANİSTAN EKONOMİSİYLE MUKAYESELİ ANALİZİ

A COMPARATIVE ANALYSIS OF THE RISK INDICATORS OF THE TURKISH ECONOMY WITH THE GREECE ECONOMY

***Tuna ŞENER*[1]**

[1]*Öğretim Görevlisi, Trakya Üniversitesi, İpsala Meslek Yüksekokulu, Dış Ticaret Bölümü, Edirne, Türkiye.*

[1]*ORCID ID: https://orcid.org/0000-0003-2715-1661*

ÖZET

2019 yılının sonlarında ortaya çıkan ve kısa sürede küresel bir halk sağlığı krizine dönüşen COVID-19 salgını, yalnızca sağlık sistemleri üzerinde değil, aynı zamanda ekonomik, sosyal ve siyasi alanlarda da derin etkiler yaratmıştır. Salgınla birlikte birçok ülkede üretim süreçleri durma noktasına gelmiş, uluslararası ticaret hacmi daralmış ve işsizlik oranlarında ciddi artışlar gözlemlenmiştir. Arz-talep dengesizlikleri, enflasyonist baskılar ve jeopolitik belirsizlikler, özellikle gelişmekte olan ve yüksek borç yüküne sahip ekonomiler için kırılganlık riskini artırmıştır. Bu bağlamda bu çalışma, salgın sonrası dönemde 2020-2025 verileri ışığında Türkiye ve Yunanistan ekonomilerinin makroekonomik risk düzeylerini karşılaştırmalı olarak analiz etmeyi hedeflemektedir. İki ülke arasındaki ekonomik risk dinamikleri; kamu borcu / GSYİH oranı, enflasyon, büyüme hızı, cari işlemler dengesi, CDS (kredi temerrüt takası) primleri ve kredi derecelendirme notları gibi temel göstergeler üzerinden değerlendirilmiştir.

Analiz kapsamında uluslararası kuruluşlardan elde edilen güncel makroekonomik veriler temel alınarak karşılaştırmalı bir risk profili oluşturulmuştur. Elde edilen bulgular, Yunanistan'ın 2010 sonrası borç krizine karşı uyguladığı mali disiplin ve AB denetimli yapısal reformlar sayesinde risk göstergelerinde göreli bir istikrar sağladığını ortaya koymaktadır. Türkiye'nin ise nispeten daha düşük kamu borcu oranına rağmen; artan enflasyon, siyasi belirsizlik ve para politikasına duyulan güven eksikliği nedeniyle daha yüksek bir piyasa risk primiyle karşı karşıya olduğu görülmektedir. Bu durum, iki ülkenin risk profillerinin yalnızca sayısal göstergelerle değil, aynı zamanda kurumsal güven, para politikası güvenilirliği, kurumların bağımsızlığı ve dış politikadaki belirsizlikler gibi yapısal faktörlerden de etkilenmekte olduğunu ortaya koymaktadır.

Anahtar Kelimeler: Enflasyon, Para Politikası, Cari İşlemler Dengesi, CDS Primleri, Kredi Derecelendirme

ABSTRACT

The COVID-19 pandemic, which emerged in late 2019 and quickly escalated into a global public health crisis, has profoundly impacted not only healthcare systems but also the economic, social, and political spheres. The pandemic has brought production processes to a halt in many countries, shrunk international trade volume, and resulted in significant increases in unemployment rates. Supply-demand imbalances, inflationary pressures and geopolitical uncertainties have increased the risk of fragility, particularly for emerging and highly indebted economies. In this context, this study aims to comparatively analyze the macroeconomic risk levels of the Turkish and Greek economies in the post-pandemic period, drawing on data from 2020-2025. Economic risk dynamics between the two countries

FULL TEXTS BOOK

are assessed through key indicators such as the public debt-to-GDP ratio, inflation, growth rate, current account balance, CDS (credit default swap) premiums, and credit ratings.

As part of the analysis, a comparative risk profile was constructed based on up-to-date macroeconomic data obtained from international organizations. The findings reveal that Greece achieved relative stability in its risk indicators thanks to the fiscal discipline and EU-supervised structural reforms implemented in response to the post-2010 debt crisis. Turkey, despite its relatively lower public debt ratio, faces a higher market risk premium due to rising inflation, political uncertainty, and a lack of confidence in monetary policy. This suggests that the risk profiles of both countries are influenced not only by quantitative indicators but also by structural factors such as institutional trust, monetary policy credibility, institutional independence and foreign policy uncertainties.

Keywords: Inflation, Monetary Policy, Current Account Balance, CDS Premiums, Credit Rating

GİRİŞ

COVID-19 pandemisinin 2019'un sonlarında ortaya çıkıp 2020 yılında küresel bir krize dönüşmesi, dünya genelinde ekonomik dengeleri köklü biçimde sarsmıştır. Pandemi, çok geniş coğrafi alanlara yayılan, ülkeler arası sınırları aşan ve büyük insan topluluklarını etkileyen bulaşıcı hastalık durumu olarak tanımlanmaktadır (Porta, 2008). Pandemi süreci; sağlık sistemleri üzerindeki yükün ötesinde, ülkelerin makroekonomik yapılarını, finansal dayanıklılıklarını ve kriz yönetimi kapasitelerini de sınamıştır. Salgınla birlikte birçok ülkede kamu harcamaları olağanüstü artmış, bütçe açıkları derinleşmiş, merkez bankaları genişletici para politikaları uygulamış ve bu durum küresel çapta ekonomik risk göstergelerini önemli ölçüde dönüştürmüştür.

COVID-19 sonrası dönem, ekonomik aktörlerin risk algılarında yapısal bir değişime yol açarak ülke ekonomilerinin karşılaştırmalı kırılganlıklarının yeniden değerlendirilmesini gerekli kılmıştır. Türkiye ve Yunanistan, farklı ekonomik yapı ve kurumlara sahip olmalarına rağmen, hem coğrafi yakınlıkları hem de küresel kırılganlıklara verdikleri tepkiler açısından karşılaştırılabilir örnekler sunmaktadır. Yunanistan, 2010'lu yıllarda yaşadığı borç krizi sonrası Avrupa Birliği çerçevesinde sıkı mali disiplin uygulamalarına yönelmiş, kamu borcunu çevirebilirliğe odaklanmış bir ekonomik yapı geliştirmiştir. Türkiye ise aynı dönemde güçlü bir büyüme performansı sergilese de; yüksek enflasyon, kur dalgalanmaları ve para politikasına ilişkin güven sorunları nedeniyle finansal piyasalarda artan risk primiyle dikkati çekmiştir.

Bu çalışmada, Türkiye ve Yunanistan ekonomilerinin benzer küresel koşullara verdikleri tepkiler doğrultusunda hangi alanlarda daha kırılgan veya daha dirençli olduklarını ortaya koymak ve bu karşılaştırma üzerinden hem politika yapıcılar hem de akademik çevreler için analitik çıkarımlar sunmak amaçlanmıştır. Bu çerçevede pandemi sonrası süreçte, 2020–2025 döneminde iki ülkenin ekonomik risk düzeyleri; kamu borcunun GSYİH'ye oranı, enflasyon oranı, büyüme hızı, CDS (kredi temerrüt takası) primleri, cari işlemler dengesi ve kredi derecelendirme notları gibi başlıca göstergeler üzerinden karşılaştırmalı olarak ele alınmıştır. Bu göstergeler, ülkelerin iç ekonomik istikrarı kadar dış yatırımcılar nezdindeki algılarını ve güvenilirlik düzeylerini de yansıtmakta olduğu için önem taşımaktadır.

COVID-19 sonrası dönemde değişen küresel ekonomik dengeler, ülkelerin içsel ve dışsal kırılganlıklarını yeniden değerlendirmeyi zorunlu kılmaktadır. Bu bağlamda, Türkiye ve Yunanistan'ın makroekonomik performanslarının ve risk göstergelerinin incelenmesi, ekonomik karar alıcılar için hem geçmiş krizlerin etkilerini değerlendirme hem de geleceğe yönelik dirençli politika tasarımları oluşturma açısından değerli bir çerçeve sunmaktadır. Türkiye ve Yunanistan örneği üzerinden yapılan bu karşılaştırma, benzer gelir düzeyine ve bölgesel risklere sahip ülkeler için de genellenebilir çıkarımlar sunma potansiyeline sahiptir.

FULL TEXTS BOOK

ARAŞTIRMA ve BULGULAR

2020 yılında, korona virüs salgınının küresel ölçekte etkisini göstermesiyle birlikte dünya ekonomisinde %4,0 oranında bir küçülme yaşanmış; bu oran, 2008 küresel finansal krizinde kaydedilen daralmanın yaklaşık iki katı olarak dikkati çekmiştir (Oxford Economics, 2020).

Tablo 1. 2020 Yılında Bazı Ülkelerde Gerçekleşen GSYİH Büyüme Oranları (%)

ÜLKELER	2020 YILI GERÇEKLEŞEN BÜYÜME ORANLARI
ABD	-3,4
Almanya	-4,8
Brezilya	-4,0
Fransa	-8,1
Hindistan	-7,9
İngiltere	-9,7
İtalya	-8,8
Japonya	-4,8
Kanada	-5,4
Çin	2,3
Türkiye	1,7
Yunanistan	-9,2

Kaynak: The World Bank, Development Indicators, OECD 2020

Tablo 1'de de görülebileceği üzere, salgın sonrasında, gelişmiş ekonomiler de dâhil olmak üzere küresel ölçekte modern dönemlerin en ciddi ekonomik daralması meydana gelmiştir. Bu süreçte, dünyanın en büyük ekonomilerinden biri olan Amerika Birleşik Devletleri %3,4 oranında küçülürken; Euro Bölgesi'nde bu oran %6,6'ya ulaşmıştır. Öte yandan, Çin ekonomisi %2,3 ve Türkiye ekonomisi de %1,7 oranında büyüme kaydederek 2020 yılında pozitif büyüme gösteren az sayıdaki ülke arasında yer almıştır. Aynı dönemde Yunanistan ekonomisinin ise %9,2'lik çok ciddi bir küçülme oranıyla karşı karşıya kaldığı tablodan görülmektedir.

Pandemi sürecinde büyüme oranlarında yaşanan bu olumsuz gelişmeler, ülkelerin ekonomik ve finansal sistemlerinde de ciddi kırılmalara yol açmıştır. Krizin getirdiği belirsizlikler, özellikle gelişmekte olan ve kırılgan ekonomilere sahip ülkeler açısından risk yönetiminin önemini artırmıştır. Bu bağlamda bu çalışmada, benzer ekonomik yapılar sergileyen Yunanistan ve Türkiye arasında risk göstergelerinin ele


FULL TEXTS BOOK

alındığı mukayeseli bir analiz yapılmaktadır. Her iki ülkenin de yüksek kamu borcu, dışa bağımlı ekonomi yapısı ve geçmişte yaşadığı finansal kriz deneyimleri dikkat çekmektedir.

İki ülke arasındaki risk faktörleri ele alınırken ilk olarak enflasyon verilerine bakılacaktır. Ekonomik literatürde enflasyon, belirli bir zaman diliminde fiyatlar genel seviyesindeki sürekli artışı ifade eden bir kavram olarak tanımlanmaktadır. Fiyatlardaki bu yükselişe neden olan pek çok farklı unsur bulunmakta ve söz konusu artışlar bireyler, üreticiler ve kamu kurumları üzerinde pek çok farklı etkiler yaratmaktadır. Fiyatların yükselmesine yol açan başlıca etmenler arasında toplam talebin toplam arzı aşması, faiz oranlarının yüksek seyretmesi, plansız gerçekleştirilen devalüasyonlar, tasarruflardan daha fazla yapılan yatırımlar ve hatalı ekonomik politika tercihlerinin uygulanması sayılabilir (Çiftci, 2015).

Türkiye ve Yunanistan'ın 2020-2025 yılları arası enflasyon verilerini karşılaştırmalı olarak gösteren Grafik 1'e bakıldığında, Yunanistan'da 2022 yılında %10'un üstüne çıkmış olan enflasyon oranının 2025 Haziran ayı itibariyle %2,8'e kadar gerilediği, enflasyon riskinin önemli ölçüde bertaraf edildiği görülmektedir. Aynı periyotta Türkiye ise Ekim 2022'de %85'lik enflasyon oranını görmüş, sonrasında izlenen sıkı para politikalarına rağmen Haziran 2025 itibariyle enflasyonu ancak %35,05 seviyesine düşürebilmiştir. Türkiye, mevcut bu enflasyon oranıyla dünyada en yüksek enflasyonlu ülkeler arasında yer almaktadır.

Grafik 1. Türkiye ve Yunanistan'ın Yıllık Enflasyon Oranları (2020-2025, %)




Kaynak: https://tr.tradingeconomics.com/greece/inflation-cpi#

Ülkeler için en önemli risk faktörlerinden biri de ekonomik durgunluk riskidir. Bu amaçla Grafik 2'de Türkiye ve Yunanistan ülkelerinin 2020-2025 dönemindeki yıllık GSYİH büyüme oranları verilmektedir. Yunanistan'ın pandemi krizinin etkisiyle 2020'de yaşadığı %9,2'lik küçülmeye rağmen hızla toparlandığı ve 2021 yılında %8,7 büyüme oranı yakaladığı görülmektedir. Türkiye ise pandemi krizine rağmen 2020 yılını %1,7 büyüme oranıyla kapatmış, 2021 yılında ise bu oranı %11,8'e taşımıştır. Sonraki yıllarda her ne kadar iki ülkenin de büyüme artış hızlarında bir düşüş olsa da 2023 ve 2024 yıllarını %2,3 büyüme hızıyla kapatan Yunanistan'da ve 2023 yılını %5,1 2024 yılını da %3,2 büyüme hızıyla kapatan Türkiye'de ekonomik durgunluk etkileri görülmemektedir.

FULL TEXTS BOOK

Grafik 2. Türkiye ve Yunanistan'ın Yıllık GSYİH Büyüme Oranları (2020-2025, %)



Kaynak: https://tr.tradingeconomics.com/greece/full-year-gdp-growth#

Bir diğer önemli risk göstergesi ise cari işlemler dengesidir. Cari işlemler dengesinin en dikkat çekici yönlerinden biri, henüz ortaya çıkmamış ekonomik krizlere dair erken uyarı niteliğinde bilgiler sunabilmesidir. Örneğin, bir ülkede cari açık düzeyinin artması, gelecekte yaşanabilecek döviz kuru krizlerinin habercisi olarak değerlendirilebilir (Peker & Hotunluoğlu, 2009). Bu tür bir durumda ulusal para biriminin keskin bir şekilde değer kaybetmesi olasılığı doğmaktadır. Eğer merkez bankası bu değer kaybını sınırlamak amacıyla müdahalede bulunursa, bu müdahale faiz oranlarının yükselmesine ve döviz rezervlerinin hızla azalmasına yol açabilir. Bu bağlamda kriz olgusu, yalnızca ciddi bir devalüasyonla sınırlı kalmamakta; aynı zamanda iç faiz oranlarının aşırı artışı ve merkez bankası rezervlerinde ani ve ciddi bir düşüşü de içerecek biçimde daha kapsamlı bir şekilde tanımlanmaktadır (Kaminsky & Reinhart, 1999).


2001-2010 yılları arasında Türkiye'de cari işlemler açığı, özellikle kısa vadeli sermaye girişleriyle finanse edilmiş ve böylece bir şekilde sürdürülebilirlik kazanmış, ve ekonomik büyümenin başlıca kaynağı cari açık haline gelmiştir (Saçık & Alagöz, 2010). Ancak bu durum Türkiye'de cari açık sorununun kronik bir hale gelmesine yol açmıştır.

Bir ülkedeki cari açığın sorun teşkil edip etmediğini değerlendirmek amacıyla çeşitli sürdürülebilirlik analizleri geliştirilmiştir. Ancak yalnızca cari açık oranlarına dayanarak bu açığın düzeyi ve sürdürülebilirliği hakkında çıkarımda bulunmak sağlıklı sonuçlar vermeyebilir. Bu nedenle, sürdürülebilirlik analizlerinde genellikle "Cari Açık / GSYİH" oranları dikkate alınmaktadır. Dornbusch söz konusu oranın %4'ü aşması durumunu sürdürülemezlik için bir eşik olarak tanımlamıştır (Uygur, 2001). Diğer taraftan, Freund (2000), sanayileşmiş ülkelerin cari denge deneyimlerini inceleyerek bu sınırı %5 düzeyinde belirlemiştir.

Türkiye ve Yunanistan'ın 2020-2025 dönemindeki Cari Açık / GSYİH oranlarını gösteren Grafik 3'e bakıldığında Yunanistan'da bu oranın sürekli olarak kritik eşik olan %5'in üzerinde olduğu, hatta 2022 yılında bu oranın %10,2 olarak gerçekleştiği görülmektedir. 2024 yılı içinse bu oran %6,4 olarak gerçekleşmiştir. Buna karşın, 2000'li yılların başında ciddi bir cari açık sorunuyla karşı karşıya olan Türkiye'nin 2020'li yıllarda göreli olarak bu konuda iyileşme gösterdiği, 2022 yılında %5'in üstüne çıkmış olan bu oranın 2024 itibariyle %0,8 olarak gerçekleştiği görülmektedir. Dolayısıyla Grafik 3, cari denge konusunda Yunanistan'ın Türkiye'ye oranla çok daha riskli bir ülke konumunda olduğunu ortaya koymaktadır.

FULL TEXTS BOOK


Grafik 3. Türkiye ve Yunanistan'ın Cari Açık / GSYİH Oranları (2020-2025, %)



Kaynak: https://tr.tradingeconomics.com/greece/current-account-to-gdp#

Ülkeler için bir diğer önemli risk göstergesi ise kamu kesiminin borcudur. Ancak tıpkı yukarıdaki cari denge analizinde olduğu gibi, önemli olan bu borcun miktarı değil GSYİH'ye oranıdır. Bu nedenle Grafik 4'te her iki ülke için 2020-2025 dönemindeki Kamu Borcu / GSYİH oranları verilmektedir.

Grafik 4. Türkiye ve Yunanistan'ın Kamu Borcu / GSYİH Oranları (2020-2025, %)



Kaynak: https://tr.tradingeconomics.com/greece/government-debt-to-gdp#

Grafik 4'e bakıldığında, 2009 yılının sonbaharında başlayan Yunanistan devlet borç krizinin etkilerinin, her ne kadar 2020-2025 döneminde sürekli bir iyileşme gözlemlense de, günümüze kadar devam ettiği görülmektedir. Kamu Borcu / GSYİH oranının sürekli olarak düşüş gösterdiği bu dönemde, yine de 2024 yılı itibariyle bu oranın %153,6 olduğu görülmektedir. Bu durum Yunanistan'ın kamu borcunun, ülkenin toplam GSYİH'sından bile kabaca 1,5 kat daha fazla olduğu anlamına gelmektedir. Kamu borcu

FULL TEXTS BOOK

konusunda tarihsel olarak sürekli sorun yaşayan Türkiye'de bile bu oran 2024 yılında %24,7 olarak gerçekleşmiştir.

Globalleşen dünyada, ülkelerin makroekonomik ve finansal göstergelerini belli kriterler çerçevesinde değerlendirerek, uluslararası risk değerlerini belirleyen en temel kuruluşlar kredi derecelendirme kuruluşlarıdır. Günümüzde dünya genelinde faaliyet gösteren birçok kredi derecelendirme kuruluşu bulunmakta olup, bunlar arasında en öne çıkan üç kuruluş S&P, Moody's ve Fitch'tir (Tutar vd., 2011).

1975 yılında, kredi derecelendirme kuruluşlarının faaliyet alanlarını ve piyasaya giriş-çıkış süreçlerini belirli bir çerçeveye oturtmak amacıyla, Amerikan Sermaye Piyasası Kurulu (SEC) tarafından NRSRO (Ulusal Olarak Tanınan Derecelendirme Kuruluşları) sistemi oluşturulmuş ve bu sayede ilgili kuruluşlar denetim kapsamına alınmıştır (Yazıcı, 2009).

Kredi derecelendirme kuruluşları her ne kadar 2008 yılında ABD'de yaşanan mortgage krizini öngörememiş ve bu sebeple küresel finans krizi sonrası birtakım eleştirilere maruz kalmış olsalar da (Bostancı, 2012), günümüzde hala ülkelerin temel risk faktörlerini belirleyen önemli aktörler olarak uluslararası ekonomide kilit bir rol oynamaktadırlar. Tablo 2'de 3 temel kredi derecelendirme kuruluşunun not sistemi görülmektedir. Buna göre Standard & Poor's ve Fitch için BBB-, Moody's içinse Baa3 notları yatırım yapılabilir son seviyeyi göstermektedir. Kredi notu daha düşük olan ülkelere yatırım yapılması önerilmemektedir.

Tablo 2. Kredi Derecelendirme Kuruluşları Not Sistemi

Standart & Poor's	Fitch	Moody's	Notun Açıklaması	
AAA	AAA	Aaa	En Yüksek Derece	Yatırım Yapılabilir Seviye
AA+	AA+	Aa1	İyi Kredi Derecesi	
AA	AA	Aa2		
AA-	AA-	Aa3		
A+	A+	A1	İyi Kredi Derecesi	
A	A	A2		
A-	A-	A3		
BBB+	BBB+	Baa1	Ortanın Altı Seviye	
BBB	BBB	Baa2		
BBB-	BBB-	Baa3		
BB+	BB+	Ba1	Yatırım Yapılmaz	Spekülatif Seviye
BB	BB	Ba2		
BB-	BB-	Ba3	Spekülatif	
B+	B+	B1		
B	B	B2	Önemli Derecede Spekülatif	
B-	B-	B3		
CCC+	CCC	Caa	Şiddetli Riskli	
CCC	CC	Caa3		
CC	C	C	Aşırı Spekülatif	
D	DDD DD D	D	Yükümlülüğünü Yerine Getiremez	Default

Kaynak: Kronwald, C. 2009. Credit rating and the impact on capital structure. Munich, GRIN Verlag

2020-2025 dönemi için Standard & Poor's ve Moody's şirketlerinin Türkiye için verdiği notlar Tablo 3'te görülmektedir. Tabloya göre, her ne kadar son dönemde her iki şirketin de Türkiye'nin kredi

FULL TEXTS BOOK

notunda iyileştirmeye gittiği görülse de, Temmuz 2025 itibariyle Türkiye'nin mevcut kredi notuyla S&P için (BB-) yatırım yapılabilir seviyenin 3 not, Moody's içinse (B1) 4 not altında olduğu görülmektedir.

Tablo 3. Türkiye'nin Standard & Poor's ve Moody's Kredi Notları (2020-2025)

Tarih	S&P Kredi Notu	Görünüm	Tarih	Moody's Kredi Notu	Görünüm
1.Kas.24	BB-	Durağan	19.Tem.24	B1	Pozitif
3.May.24	B+	Pozitif	12.Oca.24	B3	Pozitif
30.Kas.23	B	Pozitif			
29.Eyl.23	B	Durağan			
31.Mar.23	B	Negatif	12.Ağu.22	B3	Durağan
30.Eyl.22	B	Durağan	11.Eyl.20	B2	Negatif
10.Ara.21	B+	Negatif	14.Haz.19	B1	Negatif

Kaynak: https://tr.tradingeconomics.com/turkey/rating

Aynı dönemde Yunanistan için Standard & Poor's ve Moody's şirketlerinin verdiği notlar ise Tablo 4'te yer almaktadır. Tabloda yeşil olan alanlar "yatırım yapılabilir" seviyeyi göstermektedir. Buna göre Temmuz 2025 itibariyle Yunanistan, hem S&P için sahip olduğu BBB notuyla, hem de Moody's için sahip olduğu Baa3 notuyla yatırım yapılabilir bir ülke durumundadır.

Tablo 4. Yunanistan'ın Standard & Poor's ve Moody's Kredi Notları (2020-2025)

Tarih	S&P Kredi Notu	Görünüm	Tarih	Moody's Kredi Notu	Görünüm
18.Nis.25	BBB	Durağan	14.Mar.25	Baa3	Durağan
19.Nis.24	BBB-	Pozitif	14.Eyl.24	Ba1	Pozitif
20.Eki.24	BBB-	Durağan	15.Eyl.23	Ba1	Durağan
21.Nis.23	BB+	Pozitif	17.Mar.23	Ba3	Pozitif
22.Nis.22	BB+	Durağan			
23.Nis.21	BB	Pozitif			
24.Nis.20	BB-	Durağan	6.Kas.20	Ba3	Durağan

Kaynak: https://tr.tradingeconomics.com/greece/rating

Bir ülkenin kredi riskini değerlendirmede kullanılan başlıca göstergelerden biri kredi derecelendirme notları iken, bir diğeri ise CDS (Credit Default Swap) primleridir. CDS, yatırımcıların temerrüt, iflas ya da ödeme güçlüğü gibi riskler karşısında doğabilecek kayıplarını, üçüncü taraflara devretmelerine olanak sağlayan bir sözleşme türüdür. Kredi notlarının aksine, CDS primlerine ilişkin veriler piyasada anlık olarak izlenebilme özelliğine sahiptir (Kaya vd., 2015).

2008 yılında yaşanan küresel finansal krizin ardından, kredi derecelendirme kuruluşlarının verdikleri notların, gerçek kredi risklerini tam olarak yansıtamadığı yönündeki görüşler ön plana çıkmaya başlamıştır (Mora, 2006). Bu gelişmeyle birlikte, kredi riskini ölçmede derecelendirme notlarına alternatif araçlar geliştirme arayışları hız kazanmıştır. Bu arayışların bir sonucu olarak, kredi riskinin

FULL TEXTS BOOK

değerlendirilmesinde CDS primlerinin, kredi notlarına alternatif bir gösterge olarak kullanılabileceği fikri gündeme gelmiştir (Flannery, Houston & Partnoy, 2010).

Grafik 5. Türkiye'nin 5 Yıllık CDS Primleri




Kaynak: https://tr.investing.com/rates-bonds/turkey-cds-5-year-usd

Grafik 5'te Türkiyenin 5 yıllık CDS primleri görülmektedir. Genel olarak CDS primi 300'ün üzerinde olan ekonomiler aşırı kırılgan ekonomiler olarak kabul edilmektedir. Türkiye'nin 5 yıllık geçmişine bakıldığında Haziran 2022'de CDS priminin 800'ün bile üstüne çıktığı görülmektedir. Bu durumun büyük ölçüde Türkiye'nin söz konusu dönemde geleneksel para politikalarını terk etmesinden kaynaklandığı söylenebilir. Sonraki süreçte CDS primlerinde belirgin bir düşüş gerçekleşmiş olsa da 24 Temmuz 2025 tarihi itibariyle elde edilen 284,30'luk değer halen kritik eşik olan 300'e çok yakın durumdadır.

Aşağıdaki Grafik 6'da ise Yunanistan'ın 5 yıllık CDS primleri verilmiştir. Yüksek cari açık ve çok yüksek kamu borcuna karşın Yunanistan'ın CDS priminin genel olarak 100'ün altında olduğu, hatta 2023 sonbaharıyla birlikte risk priminin yeni bir düşüş trendine girdiği ve nihayet 24 Temmuz 2025 tarihi itibariyle 44,41 gibi çok düşük bir seviyede olduğu görülmektedir.

Grafik 6. Yunanistan'ın 5 Yıllık CDS Primleri



Kaynak: https://tr.investing.com/rates-bonds/greece-cds-5-years-usd

FULL TEXTS BOOK

SONUÇ

COVID-19 pandemisiyle birlikte küresel düzeyde artan ekonomik belirsizlik, ülkelerin makroekonomik risk göstergelerinin daha yakından takip edilmesini zorunlu hâle getirmiştir. Bu bağlamda, Türkiye ve Yunanistan gibi benzer coğrafi konuma ve kriz deneyimlerine sahip ülkelerin karşılaştırılması, kriz sonrası dönemde ortaya çıkan farklı risk profillerinin anlaşılması açısından önem arz etmektedir. Çalışmada 2020–2025 dönemi için yapılan analizler, Türkiye'nin hem kamu borcu/GSYİH hem de cari açık oranlarında Yunanistan'a kıyasla daha iyi bir görünüm sergilediğini ortaya koymuştur. Özellikle 2024 yılı itibariyle Türkiye'nin cari açık/GSYİH oranı %0,8 seviyesine kadar gerilemişken, Yunanistan'da bu oran %6,4 gibi yüksek bir düzeyde kalmıştır. Benzer şekilde Türkiye'nin kamu borcu oranı da %24,7 gibi yönetilebilir bir seviyedeyken, Yunanistan'da bu oran %153,6 gibi oldukça yüksek bir değerdedir.

Bununla birlikte, kredi riskini yansıtan CDS primi ve kredi derecelendirme notları incelendiğinde, ortaya çıkan tablo ters yöndedir. Yunanistan, kamu borcu ve cari açık gibi makroekonomik kırılganlık göstergelerinde Türkiye'den daha olumsuz bir görünüm sunmasına rağmen, 2025 yılı itibariyle 44,41 seviyesindeki CDS primiyle Türkiye'nin 284,30 düzeyindeki risk priminin oldukça gerisinde kalmaktadır. Benzer şekilde, Yunanistan hem Moody's hem de S&P tarafından yatırım yapılabilir seviyede derecelendirilirken, Türkiye aynı kuruluşlar nezdinde yatırım eşiğinin hâlâ birkaç not altında yer almaktadır.

Bu çelişkili görünüm, yalnızca sayısal göstergelerin değil, yapısal unsurların da ülkelerin kredi risk algısı üzerinde belirleyici olduğunu göstermektedir. Yunanistan'ın 2010 sonrası Avrupa Birliği denetiminde yürüttüğü mali disiplin, şeffaf bütçe politikaları ve kurumsal reformlar, dış yatırımcıların ülkeye duyduğu güveni artırmış; bu da kredi notlarına ve CDS primlerine olumlu yansımıştır. Türkiye ise görece düşük kamu borcuna ve cari açığa rağmen yüksek enflasyon, sık değişen ekonomi yönetimi, bağımsızlığı tartışmalı para politikası uygulamaları, yükselen faizlerin yatırımları olumsuz etkilemesi ve uygulanan ekonomik programa karşı azalan güven gibi nedenlerle yatırımcı algısında daha riskli bir konumda yer almaktadır. Bu durum, piyasa temelli göstergelerin sadece rakamsal değil, aynı zamanda kurumsal güven ve ekonomik istikrarla da doğrudan ilişkili olduğunu ortaya koymaktadır.

Türkiye'nin risk primlerinde ve kredi notlarında iyileştirme sağlanabilmesi için kurumsal yapıların güçlendirilmesi, para politikasına duyulan güvenin ve ekonomik öngörülebilirliğin artırılması gerekmektedir. Yunanistan örneği, yapısal reformların ve uluslararası denetime dayalı güven inşasının, makroekonomik göstergelerden daha belirleyici olabileceğini göstermektedir. Haziran 2023'te geleneksel para politikasına geri dönen Türkiye'de 2023-2025 yılları arası enflasyonun %85'lerden %35 seviyesine düşürülmesi sonucu hem CDS primlerinde, hem de kredi notlarında yaşanan iyileşmeler, fiyat istikrarının risk analizleri açısından önemini de ortaya koymaktadır. Sürdürülebilir finansal istikrarın sağlanabilmesi ve ekonomide güven inşa eden reformların hayata geçirilebilmesi durumunda, Türkiye'nin risk değerlerindeki iyileşmeler gelecekte daha da belirgin olabilecektir.

KAYNAKÇA

Bostancı, M.C. (2012). Küresel Krizin Dinamikleri ve Kredi Derecelendirme Kuruluşlarının Kredibilitesi. Marmara Üniversitesi Sosyal Bilimler Enstitüsü Maliye Ana Bilim Dalı, İstanbul.

Çiftci, E. (2015). Türkiye'de enflasyon ile ekonomik büyüme arasındaki ilişki: Ampirik bir uygulama (1980-2014). (Yüksek Lisans Tezi). Selçuk Üniversitesi Sosyal Bilimler Enstitüsü.

Flannery, M.J, Houston, J.F & Partnoy, F. (2010). Credit Default Swap Spreads as Viable Substitutes for Credit Ratings. University of Pennsylvania Law Review, Cilt. 158, (2085-2123).

FULL TEXTS BOOK

Freund, C.L, "Current Account Adjustment in Industrialized Countries", Board of Governors of the FED International Finance Discussion Papers, Aralık 2000

Kaminsky, G.L ve Reinhart, C.M, "The Twin Crises: The Causes of Banking and Balance of Payments Problems", American Economic Review, Vol.89, Issue 3, Haziran 1999

Kaya Öner, E., Kaya, B., & Yalçıner, K. (2015). Reaction of Credit Default Swap Spreads to Rating Announcements: An Event Study for Turkey. Journal of Economics, Finance and Accounting, 2(4), 558-571.

Kronwald, C. (2009). Credit rating and the impact on capital structure. Munich, GRIN Verlag.

Mora, Nada. (2006). "Sovereign Credit Ratings: Guilty Beyond Reasonable Doubt?, Journal Of Banking and Finance, Cilt.30, (2041-2062).

Oxford Economics, 2020, https://www.oxfordeconomics.com/

Peker, O. & Hotunluoğlu, H. (2009). "Türkiye'de Cari Açığın Nedenlerinin Ekonometrik Analizi", Atatürk Üniversitesi İktisadi ve İdari Bilimler Dergisi, Cilt: 23, Sayı:3

Porta, M. (2008). Dictionary of Epidemiology. Oxford.

Saçık, Yapar S. ve Alagöz, M. (2010). "Türkiye'de Cari İşlemler Açığı Sorunu ve Borçlanma ile İlişkisi", Ekonomi Bilimleri Dergisi, Cilt 2, Sayı 2

Tutar, E., Tutar, F. ve Eren, M. V. (2011). Uluslararası Kredi Derecelendirme Kuruluşlarının Rolü, Güvenirlilik Açısından Sorgulanması ve Türkiye, Akademik Bakış Dergisi, Sayı:25, Kırgızistan

Uygur, E., "Krizden Krize Türkiye: 2000 Kasım ve 2001 Şubat Krizleri", Türkiye Ekonomi Kurumu Tartışma Metni, 2001/1, Nisan 2001

Yazıcı, M. (2009). Kredi Derecelendirme Kuruluşlarının Önemi ve Denetimi. Maliye ve Finans Yazıları Dergisi, 82.

FULL TEXTS BOOK

YAŞLI KADINLARIN SOSYO-EKONOMİK SORUNLARININ SOSYAL HİZMET BAĞLAMINDA TARTIŞILMASI

***Ummühani USLU*[1]**

[1]*Öğretim Görevlisi, Alanya Alaaddin Keykubat Üniversitesi, Altso Meslek Yüksekokulu, Sosyal Hizmet ve Danışmanlık Bölümü, Antalya, Türkiye*

[1]*ORCID ID: https://orcid.org/0000-0002-4907-0435*

ÖZET

Tüm dünyada olduğu gibi, ülkemizde de önlenebilir ölümlerin azalması, sağlık hizmetlerinin gelişmesi ve gelişen bu hizmetlere ulaşımın artması, beslenme ve yaşam tarzı değişiklikleri gibi nedenlerden dolayı beklenen yaşam süresi artmakta ve nüfus giderek yaşlanmaktadır. Türkiye'de yaşlılık şu an gelişmiş toplumlardaki kadar önemli bir mesele olarak görülmese de beklenen yaşam sürelerinin uzamasıyla önümüzdeki dönemlerde bir sorun olabilme potansiyeli taşımaktadır. Yaşlılık dönemi çeşitli fiziksel, ruhsal ve sosyal sorunları beraberinde getirmektedir. Bu sorunlar için önlemler alınmazsa hem erkek hem de kadın bireylerde yaşlılık döneminde toplumsal sorunlar ortaya çıkmaktadır ancak ortalama yaşam süresinin kadınlarda daha uzun olması ve toplumsal rolleri gereği kadınların ekonomik ve sosyal yaşamdan uzak kalmaları nedeniyle günümüzde yaşlı kadınlar erkeklere oranla ekonomik ve sosyal sorunlarla daha fazla karşı karşıya kalabilmektedirler. Gelecekte toplam nüfusumuzun önemli bir bölümünü oluşturacak olan yaşlı nüfus ve yaşlı nüfus içinde kadınların yaşam kalitesinin yükseltilmesi ve aktif bir yaşam sürmelerinin sağlanması için gerekli olan sosyal hizmet programlarının hızla oluşturulması hem toplum hem de aile açısından büyük önem taşımaktadır. Yaşlı kadınların sosyo-ekonomik açıdan yaşadığı sorunları ele alan bu çalışmada yaşlı kadınların karşılaştıkları sorunlara sosyal hizmet bağlamında çözümler üretmek hedeflenmiştir. Yöntem olarak yaşlı refah alan yazınındaki kaynaklar taranmış ve sayısal verileri içeren ikincil kaynaklar incelenmiştir. Çalışmanın sonunda yaşlı kadınlar için üretilebilecek sosyal hizmet politika önerileri ortaya konulmuştur.

Anahtar Kelimeler: Yaşlı, Kadın, Sosyal Hizmet

ABSTRACT

As is the case worldwide, life expectancy in our country is increasing due to factors such as the decline in preventable deaths, improved healthcare services and increased access to these services, and changes in nutrition and lifestyle. While aging in Turkey is not currently considered as significant an issue as it is in developed societies, it has the potential to become a problem in the future as life expectancy increases. Old age brings with it various physical, mental, and social problems. If measures are not taken to address these issues, social problems arise in both men and women during old age. However, due to the longer average lifespan in women and the social roles that keep women isolated from economic and social life, older women today face more economic and social problems than men. The elderly population, which will constitute a significant portion of our population in the future, and the rapid development of social service programs to improve the quality of life of women within this population and ensure their active living are of paramount importance for both society and families. This study, which addresses the socio-economic problems faced by older women, aims to provide solutions to these problems within the context of social services. The methodology was to review sources in the elderly welfare literature and examine secondary sources containing numerical data. The study concludes with recommendations for social service policies for older women.

Keywords: elderly, women, social services

FULL TEXTS BOOK

GİRİŞ

Türkiye'de yaşlılık olgusu, henüz gelişmiş ülkelerde olduğu kadar gündemde yer almasa da, artan yaşam süresi ve demografik değişimlerle birlikte gelecekte daha belirgin bir toplumsal mesele haline gelme potansiyeline sahiptir (Dedeoğlu ve Yentürk, 2020).

Yaşlılık dönemi yalnızca biyolojik değişimlerle sınırlı kalmayıp ruhsal, sosyal ve ekonomik sorunları da beraberinde getirmektedir. Özellikle kadınlar, daha uzun yaşam beklentisine sahip olmaları, toplumsal rollerinin niteliği ve ekonomik-sosyal hayattan görece uzak kalmaları nedeniyle yaşlılık döneminde erkeklere kıyasla daha fazla dezavantaj yaşamaktadır (www.birgun.net, 2024). Türkiye'nin demografik yapısında önümüzdeki yıllarda önemli bir yer tutacak yaşlı nüfus içinde, yaşlı kadınların yaşam kalitesinin artırılması ve aktif yaşamlarının desteklenmesi, sosyal hizmet politikalarının öncelikli hedeflerinden biri olmalıdır (Yaşlı Hakları Derneği, 2023).

Türkiye'de 65 yaş ve üzerindeki bireylerin%57,5'inin, 75 yaş ve üzerindeki bireylerin %60,5'inin, 85 yaş ve üzerindeki bireylerin ise %73,1'inin kadınlardan meydana geldiği görülmektedir. Bu olgu literatürde "yaşlılığın kadınsallaşması" kavramı ile ifade edilmektedir (Canatan,2012).


Demografik değişim ve dönüşümlerin gelecek dönemlerde kadın yaşlılar açısından yaratacağı olası riskler önemlidir, bu riskler kadın yaşlıların yaşamları süresince cinsiyetlerine bağlı olarak karşılaştıkları sosyal dezavantajlardan kaynaklanmaktadır. Bu yüzden yaşlı kadınların yaşadığı sorunları ortaya koymak kadınların yaşam kalitesinin yükseltilmesi ve aktif bir yaşam sürmelerinin sağlanması için gerekli olan sosyal hizmet programlarının hızla oluşturulması açısından önem taşımaktadır.

YAŞLI KADINLARIN KARŞILAŞTIĞI SORUNLAR

Yaşlı kadınların yaşadığı sorunların temelinde toplumsal cinsiyete daya eşitsizlikler yatmaktadır. Geleneksel toplum yapısında erkeklerin "aile reisi" olarak konumlandırılması ve karar alma süreçlerinde tek yetkili kabul edilmesi, kadınların toplumsal yaşamda deneyim kazanmalarını ve çeşitli sorumluluklar üstlenmelerini önemli ölçüde sınırlandırmıştır. Bu durumun bir sonucu olarak, gençlik ve yetişkinlik dönemlerinde eğitim ve çalışma hayatına sınırlı biçimde katılabilmiş olan günümüz yaşlı kadınlarının önemli bir kısmı, yaşlılık döneminde kendini ifade etmede yetersizlik, günlük yaşamın temel ihtiyaçlarını karşılamada güçlük, buna bağlı güvensizlik, huzursuzluk ve kaygı gibi sorunlarla karşı karşıya kalmaktadır. Dolayısıyla yaşlı kadınlar, toplum tarafından inşa edilen "yaşlılık bariyerleri" ile mücadele ederken, aynı zamanda kadın kimliklerinden kaynaklanan ek dezavantajları da deneyimlemektedir (Tufan, 2003). Bu bağlamda, kadınların erkeklerden daha uzun yaşam beklentisine sahip olmaları sonucunda ortaya çıkan "yaşlılığın kadınsallaşması" olgusu, biyolojik açıdan bir avantaj gibi görünse de toplumsal açıdan yaşlılığın özellikle kadınlara özgü bir sorun olarak şekillendiğini açıkça ortaya koymaktadır (Irmak ve Artan, 2018).

FULL TEXTS BOOK


Yaş ve cinsiyete göre beklenen yaşam süresi, 2021-2023



Kaynak: TÜİK, Hayat Tabloları, 2021-2023

Kadınların biyolojik olarak daha fazla yaşaması beraberinde medeni durum problemini getirmektedir. Kadınlar eş kayıplarını daha fazla yaşamakta ve daha fazla yalnız kalmaktadır.

Medeni durum ve cinsiyete göre yaşlı nüfus oranı, 2024



Kaynak: TÜİK, Adrese Dayalı Nüfus Kayıt Sistemi, 2024

Yaşlılık dönemi, bireylerin psikolojik ve duygusal açıdan en kırılgan oldukları yaşam evrelerinden biri olarak değerlendirilmektedir. Bu dönemde bireylerin yanlarında güven duydukları eşlerinin bulunmaması, yaşam güçlerini olumsuz yönde etkileyerek hem psikolojik iyi oluşlarını hem de yaşam doyumlarını azaltıcı bir unsur hâline gelmektedir (Irmak ve Artan, 2018).

Yaşlı kadınlar yalnızlığın psikolojik sorunlarıyla beraber ekonomik sorunlarını da yaşıyorlar. Çünkü kadınlar erkekler ile karşılaştırıldığında işgücü piyasasına eşit oranda, eşit konumda ve eşit ücretle katılamamaktadırlar. Kadınların işgücü piyasasına katılımın düşük olması kadınların hayatları boyunca erkeklere bağımlı olmaları ile ilişkilendirilmekte ve buna paralel şekilde işgücüne katılımın sağladığı emeklilik gibi imkânlardan da yararlanılamaması, sağlıkla ilişkili haklara da eşler üzerinden erişilmesi bu bağımlılığın bir kat daha artmasına neden olmaktadır (Şener,2019).

FULL TEXTS BOOK

Tüik tarafından 2023 yılında gerçekleştirilen yoksulluk ve yaşam koşulları araştırmasına göre 2023 yılı verilerine göre, 65 yaş ve üzerindeki kadınların %68,3'ünün ev işleri ile çocuk, yaşlı ve hasta bakımından sorumlu olduğu görülmektedir. Sağlık durumu açısından değerlendirildiğinde ise aynı yaş grubundaki kadınların %32,8'i sağlık durumunu kötü veya çok kötü olarak tanımlarken, bu oran erkeklerde %24,8'dir. Kronik hastalık oranı da kadınlarda (%88) erkeklere (%73,5) kıyasla daha yüksek seviyededir. Bunun doğal bir sonucu olarak, fiziksel veya ruhsal bir sağlık sorunu nedeniyle günlük faaliyetlerinin kısıtlandığını ifade eden kadınların oranı %71,4 iken, erkeklerde bu oran %57,2 olarak belirlenmiştir. Dolayısıyla, kadınların sağlık durumları erkeklere göre daha olumsuz olmasına rağmen, aile içi bakım yükümlülüklerini üstlenmeye devam ettikleri anlaşılmaktadır. Gelir getirici bir faaliyette bulunmayan yaşlı erkeklerde işgücü dışında kalmanın temel nedeni emeklilik olup bu oran %79,7'dir. Yaşlı kadınlarda ise emeklilik oranı yalnızca %19,8'de kalmaktadır. Bu veriler, 65 yaş üzerindeki erkeklerin hem geçmişte hem de yaşlılık dönemlerinde daha çok gelir getiren işlerde istihdam edildiklerini, buna karşılık kadınların hem geçmişte hem de yaşlılıkta ailelerine çoğunlukla ücret karşılığı olmayan bakım hizmeti sunduklarını ortaya koymaktadır. Bu nedenle, yaşlı kadınların yoksulluğa düşme olasılıkları erkeklere kıyasla daha yüksek görünmektedir. Özellikle dul ve boşanmış kadınlarda bu risk daha da artmaktadır. Dul kadınların gelir düzeylerinin düşük olduğu bilinmekte; boşanmış kadınların ise çoğunlukla çocuklarıyla birlikte, sınırlı miktardaki nafaka ile yaşamlarını sürdürmek zorunda kaldıkları görülmektedir (data.tuik.gov.tr, 2023).

SOSYAL HİZMET POLİTİKALARI

Sosyal politikaların belirlenmesinde yaşlı bireylerin karşılaştıkları sorunlar temel ölçüt olarak ele alınmaktadır. Yaşlılık döneminde karşılaşılan en önemli problemlerden biri, gelir düzeyinin azalmasına karşın sağlık sorunlarının ve buna bağlı harcamaların artış göstermesidir. Özellikle kronik hastalıkların yaygınlaşması ve fonksiyonel kayıpların ortaya çıkması, yaşlı bireylerin bakım gereksinimlerini artırmaktadır. Türkiye'de aile yapısında son yıllarda meydana gelen değişimler sonucunda, geleneksel olarak bakım ve barınma konusunda temel sorumluluk üstlenen aile kurumu bu işlevini giderek yerine getirememekte; bu durum ise devlet desteği ve profesyonel bakım hizmetlerine olan ihtiyacı artırmaktadır (Çevik, 2019: 2). Türkiye' de yaşlı bakım ve rehabilitasyon merkezleri ile huzurevleri, kurumsal bakım modeli kapsamında yaygın bir şekilde hizmet vermektedir. Ancak, yaşlı nüfus oranındaki artış huzurevi bakım maliyetlerini yükseltmiş ve alternatif bir model olarak evde bakım hizmetlerinin desteklenmesine yol açmıştır. Bu model aracılığıyla, huzurevinde kalmak istemeyen yaşlı bireylerin yaşadıkları sosyal çevreden kopmadan, sevdikleriyle birlikte ve yalnız kalmadan alıştıkları ortamda yaşamlarını sürdürmeleri amaçlanmaktadır (Devlet Planlama Teşkilatı, 2017). Bunun yanı sıra, ülkemizde kendi imkânlarıyla yaşamını sürdüren yaşlı bireylere yönelik çeşitli sosyal yardım ve sosyal hizmet programları da sunulmaktadır. Evde yalnız yaşayan yaşlılara ise Aile ve Sosyal Hizmetler Bakanlığı ile yerel yönetimler tarafından evde bakıma destek hizmetleri sağlanmaktadır. Ancak bahsedilen bu hizmetlerin hiçbiri yaşlı kadınların yaşadıkları spesifik "yaşlılık bariyerleri"ni dikkate almamaktadır.

Sosyal güvenlik çatısı dışında kalan ve hiçbir geliri olmayan yaşlı vatandaşlara devlet 2022 sayılı kanun kapsamında emekli(yaşlı) aylığı bağlamaktadır. 01.01.2022-30.06.2022 döneminde 65 yaşlı aylığı için yapılan ödeme aylık 1.084,55 TL'dir (Sosyal Yardımlar Genel Müdürlüğü, 2022)

Türkiye'de sosyal politika uygulamaları kapsamında, 3294 sayılı Kanun doğrultusunda muhtaçlık durumu tespit edilen ve birlikte yaşadığı hane içinde sosyal güvencesi bulunan herhangi bir birey bulunmayan kadınlara yönelik nakdi destek sağlanmaktadır. Bu destek, özellikle son resmi nikâhlı eşini kaybetmiş kadınları kapsamaktadır. Söz konusu yardımlar, Sosyal Yardımlaşma ve Dayanışma Vakıfları aracılığıyla düzenli olarak yürütülmekte olup ödemeler aylık esaslara göre hesaplanmakta,

FULL TEXTS BOOK

ancak iki ayda bir olacak şekilde yapılmaktadır. Uygulamanın 2022 yılı itibarıyla belirlenen tutarı 500 TL'dir (Sosyal Yardımlar Genel Müdürlüğü, 2022).

Sosyal güvenlik sistemi dışında kalan bireylerin önemli bir bölümünü, özellikle eşi vefat etmiş ve 65 yaş üzerindeki kadınların oluşturduğu tespit edilmiştir. Bu gruba ilişkin yapılan değerlendirmelerde, söz konusu kadınların yaklaşık yarısının 2022 sayılı Kanun kapsamında dul ve yetim aylığından yararlanamadığı ortaya çıkmıştır (Irmak ve Artan, 2018).

SONUÇ VE ÖNERİLER

Türkiye'de yaşlılık olgusu giderek artan bir toplumsal mesele haline gelmektedir. Yaşlı kadınlar, uzun yaşam beklentileri, toplumsal cinsiyet temelli eşitsizlikler ve sınırlı eğitim/istihdam imkânları nedeniyle yaşlılıkta erkeklere kıyasla daha dezavantajlı konumdadır. "Yaşlılığın kadınsallaşması" olgusu, kadınların yalnızlık, ekonomik yetersizlik, sağlık sorunları ve bakım yükümlülükleri ile daha yoğun biçimde karşılaştığını ortaya koymaktadır. TÜİK verileri, yaşlı kadınların önemli bir kısmının gelir getirici faaliyetlerden uzak kaldığını, emeklilik oranlarının düşük olduğunu ve hane içi bakım sorumluluklarını sürdürdüklerini göstermektedir. Bu durum, yaşlı kadınların yoksulluğa düşme riskini artırmaktadır. Sosyal politikalar kapsamında yaşlılara yönelik emeklilik, dul-yetim aylığı, huzurevi ve evde bakım hizmetleri ile çeşitli nakdi yardımlar sunulmaktadır. Ancak mevcut düzenlemeler, yaşlı kadınların özgün ihtiyaçlarını ve toplumsal cinsiyet temelli dezavantajlarını yeterince dikkate almamaktadır.

Yaşlı kadınların karşılaştıkları ekonomik sorunların çözümü, yalnızca bireysel düzeyde değil, aynı zamanda toplumsal refah ve sosyal adalet açısından da kritik önemdedir. Türkiye'de toplumsal cinsiyet temelli eşitsizliklerin bir yansıması olarak, yaşlılık döneminde kadınlar ekonomik güvenceye erişim, istihdam olanakları, sosyal katılım ve bakım hizmetleri bakımından dezavantajlı konumda yer almaktadır. Bu durumun giderilmesi için sosyal hizmet politikalarının hem koruyucu hem de güçlendirici bir perspektifle yapılandırılması gerekmektedir. Bu konuda yapılabilecek düzenlemeler;

- Ekonomik Güvencenin Sağlanması

Yaşlı kadınların gelir güvencesine erişimini artırmak amacıyla, ev içi bakım ve ücretsiz emek sürelerinin sosyal güvenlik prim gününe eklenmesi; kısmi süreli veya düzensiz çalışma biçimlerinden kaynaklanan prim eksikliklerinin devlet desteği ile tamamlanması önerilmektedir. Geliri bulunmayan veya yetersiz olan yaşlı kadınlara yönelik, hak temelli bir "Yaşlı Kadın Yoksulluğu Fonu" oluşturulmalı ve dul/ yetim maaşı bağlanmasında gelir testine dayalı kısıtlamalar kaldırılmalıdır.

- İşgücüne Katılımın Teşviki

İleri yaş kadınların işgücüne katılımını artırmak üzere, işverenlere vergi indirimi ve sigorta prim desteği sağlanması, mikro kredi ve kadın kooperatifleri aracılığıyla gelir getirici faaliyetlerin desteklenmesi gerekmektedir. Bu sayede, kadınların yaşlılık döneminde de üretkenliklerini sürdürebilecekleri istihdam olanakları yaratılabilir.

- Sosyal Katılımın Güçlendirilmesi

Sosyal izolasyonu önlemek amacıyla, yaşlı kadınların sosyo-kültürel etkinliklere katılımını teşvik edecek "Yaşlı Kadın Dayanışma Merkezleri" kurulmalıdır. Bu merkezlerde finansal okuryazarlık, dijital beceriler ve bakım hizmetleri alanında sertifika programları sunularak kadınların hem sosyal hem de ekonomik hayata katılımı desteklenebilir.

FULL TEXTS BOOK

- Bakım Hizmetlerinin Yaygınlaştırılması

Yalnız yaşayan yaşlı kadınların yaşam kalitesini artırmak için, evde bakım hizmetleri yaygınlaştırılmalı ve erişilebilir hale getirilmelidir. Ayrıca, yaşlı kadın bakımını üstlenen aile bireylerine maddi yardım ve psikososyal destek sağlanarak bakım yükünün dengelenmesi hedeflenmelidir.

- Hukuki ve Hak Temelli Yaklaşımlar

Tüm yaşlılık politikalarında, cinsiyet temelli dezavantajların açıkça tanınması ve buna yönelik ayrımcılık karşıtı yasal düzenlemelerin hayata geçirilmesi gerekmektedir. Yaşlı kadınların ekonomik yaşamda eşit fırsatlara erişimini güvence altına alacak düzenlemeler, sosyal hizmet sisteminin temel bileşenlerinden biri olmalıdır.

KAYNAKÇA

Artan T. Huzurevinde Kalmakta Olan Yaşlılarda Aile İçi İstismar. Toplum ve Sosyal Hizmet. (2013; 24,2: 109-122).

BirGün. (2024). *Yaşlı kadınlar yoksulluk kıskacında: 3'te 1'inin hiçbir geliri yok*. https://www.birgun.net/haber/yasli-kadinlar-yoksulluk-kiskacinda-3te-1inin-hicbir-geliri-yok-612055

Canatan A. Yaşlılar ve Toplum. In: Bacanlı H, Terzi ŞI, eds. Yetişkinlik ve Yaşlılık Gelişimi Psikolojisi İstanbul: Açılım Kitap; 2012.

Çevik, E. C. (2019). *Türkiye'de Yaşlılara Yönelik Sosyal Politika Uygulamaları* (Doktora Tezi, Marmara Üniversitesi)

Dedeoğlu, S. ve Yentürk, N. (2020). *Toplumsal cinsiyet ve sosyal politika*. İstanbul: İletişim Yayınları.

Devlet Planlama Teşkilatı (2007). *Türkiye'de Yaşlıların Durumu ve Yaşlanma Ulusal Eylem Planı*, Sosyal Sektörler ve Koordinasyon Genel Müdürlüğü, Yayın No DPT: 2741

Şemsa, Ö. ve Emel, M. (2024). *Yaşlı kadın yoksulluğu: Yeni bir mücadele alanı*. https://feministyaklasimlar.org

Gökbayrak, Ş. (2019). Sosyal güvenlikte toplumsal cinsiyet eşitsizliği: Türkiye örneği. *Çalışma ve Toplum Dergisi*, 60(3), 1385–1412.

Irmak, H. S. ve Artan, T. (2018). Feminist gerontoloji bağlamında yaşlılığın kadınsallaşması. *Sağlık Bilimleri ve Meslekleri Dergisi*, *5*(2), 236-243.

Kadın Dayanışma Vakfı (2024) Yoksulluk Nafakası İzleme Raporu, Hazırlayanlar: Ceren Akçabay ve Zekiye Karaca Boz.

Özdemir, H. (2018). Toplumsal cinsiyet perspektifinden yaşlılık politikaları. *Toplum ve Sosyal Hizmet*, 29(1), 155–174.

Şener Ü. Kadın Yoksulluğu. Ankara: TEPAV Değerlendirme Notu; 2009. Erişim Tarihi: 14 Ağustos 2025, http://www.tepav.org.tr.

Tufan İ. Modernleşen Türkiye'de Yaşlılık ve Yaşlanmak. İstanbul: Anahtar Kitaplar Yayınları; 2003.

TÜİK.(2023). "Yoksulluk ve Yaşam Koşulları İstatistikleri", https://data.tuik.gov.tr/Kategori/GetKategori?p=Gelir,-Yasam,-Tuketim-ve-Yoksulluk-107

https://data.tuik.gov.tr/Bulten/Index?p=Yoksulluk-ve-Yasam-Kosullari-Istatistikleri-2023-53713

TÜİK. (2024). "İstatistiklerle Yaşlılar", https://data.tuik.gov.tr/Bulten/Index?p=Istatistiklerle-Yaslilar-2024-54079. Erişim Tarihi:13.08.2025

FULL TEXTS BOOK

TÜİK. (2023). *İşgücü istatistikleri*. Ankara.

United Nations. Profiles of Ageing-2013. 2013 (Erişim Tarihi: 10 Haziran 2025) http://esa.un.org.

Vakıflar Genel Müdürlüğü (2022). Muhtaç Aylığı. https://www.vgm.gov.tr/faaliyetler/hayir-hizmetleri/muhtac-ayligi Erişim Tarihi:06.05.2024

Yaşlı Hakları Derneği. (2023). *Yaşlılık ve yoksulluk raporu*. İstanbul.

Yıldız, E. (2021). Yaşlı kadınların sosyal katılımının önemi. *Hacettepe Sosyal Bilimler Dergisi*, 10(2), 45–59.

ÖRGÜTSÜZ KİTLE DAVRANIŞLARI: TOPLUMSAL PSİKOLOJİNİN PERSPEKTİFİNDEN BİR İNCELEME
UNORGANIZED CROWD BEHAVIOR: A SOCIAL PSYCHOLOGICAL ANALYSIS

Assoc.Prof.Dr.Cüneyt Yılmaz

St. Clements University , Department of Political Science

London , UK

https://orcid.org/0000-0003-1655-588X

ÖZET

Bu makale, örgütsel yapıdan yoksun, liderlikten bağımsız olarak bir araya gelen bireylerin oluşturduğu örgütsüz kitle davranışlarını**,** toplumsal psikoloji perspektifiyle incelemektedir. Örgütsüz kitlelerin davranışlarının altında yatan psikolojik, sosyal ve kültürel dinamikler; literatürdeki teorik yaklaşımlar, klasik ve modern araştırmalar aracılığıyla değerlendirilmiştir. Makalede özellikle grup dinamikleri**,** sosyal etkileşim**,** bireyler arası ilişkiler**,** medya etkisi **ve** kültürel normların bu tür kolektif davranışlar üzerindeki etkileri analiz edilmiştir.

Elias Canetti, Gustave Le Bon ve Sigmund Freud'un kuramsal katkıları, kitle psikolojisinin tarihsel gelişimi bağlamında değerlendirilirken; Michel Foucault, Herbert Marcuse, Manuel Castells gibi çağdaş düşünürlerin analizleri, özellikle devletin denetim mekanizmaları, kültürel normlar ve dijital iletişimin örgütsüz kitle oluşumuna olan etkileri çerçevesinde ele alınmıştır. Ayrıca Sylvia Buck-Morss ve David Graeber gibi araştırmacıların çalışmaları, yerel ve küresel bağlamda kitle davranışlarının karmaşıklığını anlamaya yönelik önemli perspektifler sunmaktadır.

Makale, Arap Baharı**,** Gezi Parkı protestoları gibi güncel olayları inceleyerek sosyal medyanın örgütsüz kitlelerin mobilizasyonunda nasıl merkezi bir rol oynadığını da ortaya koymaktadır. Kitlelerin oluşumunda bireylerin duygusal tepkileri, kolektif bilinç yapıları, medya aracılığıyla yayılan sosyal mesajlar ve kültürel çerçevelerin nasıl belirleyici olduğu gösterilmektedir.

Sonuç olarak, bu derleme, örgütsüz kitle davranışlarını anlamaya yönelik disiplinlerarası bir çerçeve sunarken, bu alandaki kuramsal ve ampirik tartışmalara önemli katkılarda bulunmayı amaçlamaktadır. Makale, özellikle sosyal psikoloji, kültürel çalışmalar ve medya araştırmaları kesişiminde yapılan yeni araştırmalar için kapsamlı bir referans sunmaktadır.

Anahtar Kelimeler: Örgütsüz kitle, toplumsal psikoloji, davranış analizi, sosyal etkileşim

ABSTRACT

This article examines unorganized crowd behavior groups formed spontaneously without leadership or organizational structure—through the lens of social psychology. It analyzes the psychological, social, and cultural dynamics underlying such behaviors by referencing theoretical approaches and both classical and contemporary studies. Key themes include group dynamics**,** social interaction**,** interpersonal relationships**,** media influence**,** and the impact of cultural norms on collective behavior.

Theoretical contributions from Elias Canetti, Gustave Le Bon, and Sigmund Freud are evaluated within the historical development of crowd psychology, while modern thinkers such as Michel Foucault, Herbert Marcuse, and Manuel Castells are discussed for their insights into state control mechanisms, cultural frameworks, and digital communication. Additionally, scholars like Sylvia Buck-Morss and David Graeber offer valuable perspectives on the complexities of crowd behavior at both local and global levels.

FULL TEXTS BOOK

The article also investigates recent events such as the Arab Spring and the Gezi Park protests, highlighting the central role of social media in mobilizing unorganized crowds. It illustrates how emotional reactions, collective consciousness, and socially transmitted messages especially through digital platforms are shaped by broader cultural and societal structures.

In conclusion, this review offers an interdisciplinary framework for understanding unorganized crowd behavior and contributes to the ongoing theoretical and empirical discussions in this field. It serves as a comprehensive reference, particularly for new research at the intersection of social psychology, cultural studies, and media analysis.

Keywords: Unorganized mass, social psychology, behavior analysis, social interaction

INTRODUCTION

Events occurring within societies often reflect social dynamics that give rise to unorganized mass behaviors. These behaviors typically involve collective actions by individuals who come together without specific leadership or organizational structure. Unorganized mass behaviors have become a significant subject of study within the context of social psychology. This field examines how individuals collectively respond, how group dynamics are shaped, and the impact of social interactions on these behaviors.

This article aims to understand and analyze unorganized mass behaviors. Drawing on an extensive literature review and a social psychology perspective, it focuses on the psychological factors influencing unorganized mass behaviors, group dynamics, social interactions, and interpersonal relationships, with the goal of contributing to the existing body of knowledge in this field.

In today's world, rapid societal changes, political events, the influence of social media, and other factors have made the dynamics of unorganized mass behaviors increasingly complex. Understanding how individuals collectively respond to specific situations remains a critical challenge for researchers in the social sciences.

The primary objective of this article is to analyze and understand unorganized mass behaviors from a social psychology perspective. To this end, it seeks to examine in detail the psychological factors influencing these behaviors and the underlying dynamics.

This study focuses on unorganized mass behaviors, analyzing group dynamics, social interactions, and interpersonal relationships. It is designed to compile key studies in the literature and offer a fresh perspective on the existing knowledge base. The article explores theoretical approaches, research findings, and the effects of social, cultural, and environmental factors to develop a comprehensive understanding of the complexity of unorganized mass behaviors.

PURPOSE

The primary purpose of this article is to understand and analyze unorganized mass behaviors from the perspective of social psychology. Specifically, it aims to examine the psychological factors influencing the behaviors of unorganized masses—groups of individuals who come together without specific leadership or organization. The article seeks to elucidate the complexity of these dynamics by reviewing various theoretical approaches and research findings.

Additionally, it aims to explore the role of social, cultural, and environmental factors in the emergence of unorganized masses and their impact on behavioral processes. By doing so, the article seeks to contribute a new perspective to the existing knowledge on unorganized mass behaviors and encourage further progress in this field.

METHOD

This study was prepared using a literature review method. The review method involves compiling information from existing literature to understand and analyze unorganized mass behaviors. The study synthesizes various research, theoretical approaches, and findings related to this topic, forming the methodological foundation of the article.

During the literature review process, studies in the fields of social psychology, behavior analysis, and social interaction were examined, with the aim of synthesizing existing knowledge.

The choice of a literature review method enables a comprehensive perspective on unorganized mass behaviors and facilitates an understanding of the interplay between various studies in this field. By integrating diverse sources from the literature, the article aims to provide a broad perspective for understanding unorganized mass behaviors.

FINDINGS

Group dynamics are a fundamental element in the formation of unorganized mass behaviors. Elias Canetti's seminal work, *Crowds and Power* (2006), provides a detailed analysis of how unorganized masses form and the dynamics among individuals within these groups. Canetti's observations offer significant insights into understanding the internal hierarchies and power dynamics of such masses.

Canetti examines unorganized masses within the framework of "crowd psychology" (2006). According to him, the formation of crowds begins with individuals coming together, and these crowds act with a spontaneously emerging force. Canetti explains these crowd dynamics through elements such as conflict, leadership, and rituals. By analyzing interactions among individuals within crowds, he provides an in-depth examination of the impact of crowd phenomena on individual behaviors. In particular, Canetti emphasizes internal hierarchies and power struggles, laying the groundwork for understanding individuals' roles within crowds.

Sigmund Freud's *Civilization and Its Discontents* (2015) examines group dynamics from a psychological perspective. Freud investigates the role of individuals within groups, particularly the impact of civilization and society on individuals. He explains how individuals participate in group dynamics and the roles they assume in these processes by exploring how society is reflected in an individual's inner world. Freud's focus is on linking an individual's inner world with group dynamics, offering a comprehensive framework for understanding individuals' participation in group processes and the roles they undertake.

Freud analyzes the impact of society on an individual's psychological structure and evaluates these interactions within the context of group dynamics. He examines how individuals interact with societal norms and the effects of these interactions on their psychological processes. Essentially, Freud's work provides a significant framework for understanding how individuals' roles within groups, their interactions with societal norms, and the reflection of these interactions on their psychological structures are shaped.

This perspective enables a deeper understanding of individuals' behaviors within social structures, their interactions with societal norms, and the impact of these interactions on their inner world. Freud's work explains the psychological foundations of group dynamics, shedding light on how individuals are shaped within societal contexts and how they develop identities in these processes.

When combined, Canetti's crowd psychology and Freud's psychological analysis of group dynamics provide a more comprehensive understanding of the complexity of unorganized mass behaviors. Canetti's observations highlight the internal dynamics of crowds, while Freud's psychological analysis helps us understand how individuals respond to these dynamics (Canetti, 2006; Freud, 2015).

FULL TEXTS BOOK

Social interaction is a complex process that shapes the behaviors of unorganized masses and plays a significant role in the formation of global crowds and the shaping of social dynamics. Sylvia Buck-Morss's article, "On the Transnational Commons and the Global Crowd" (2015), examines the impact of social interaction on these dynamics in detail. Buck-Morss analyzes interactions at local and global levels, exploring how global crowds form and their impact on social dynamics. Her article explains in detail how collective consciousness is formed and how social interaction shapes this consciousness, offering deep insights into the dynamics of social interaction underlying the formation of unorganized masses.

Manuel Castells's *Networks of Outrage and Hope: Social Movements in the Internet Age* (2013) examines the impact of digital communication technologies on interpersonal relationships and social interactions. Castells analyzes how digital communication influences the processes by which unorganized masses come together and engage in collective action. By elucidating the dynamics of communication and interaction in the internet age, he explains how unorganized masses form a social consciousness on digital platforms.

These two studies provide a significant framework for understanding the impact of social interaction on unorganized mass behaviors. Buck-Morss's analysis of local and global interactions helps explain the complexity of social interaction and the emergence of global crowds, while Castells's study of digital communication technologies contributes to understanding how unorganized masses interact in the digital world and how social consciousness is formed.

Understanding unorganized mass behaviors requires a detailed examination of the impact of social, cultural, and environmental factors. Herbert Marcuse's *One-Dimensional Man* (1964) provides an in-depth analysis of the effects of cultural factors on individual behaviors and explains how cultural norms influence the formation of unorganized masses. Marcuse examines how cultural norms shape individuals' thought systems and how these norms affect the emergence of unorganized masses. His perspective highlights the role of cultural norms, widely accepted within society, as a determining force in individual behaviors, contributing to our understanding of how unorganized masses respond to and unite around these norms.

Michel Foucault's *Discipline and Punish* (1975) examines the impact of environmental factors, particularly the role of the state, on individual behaviors. Foucault analyzes how the state shapes social order and exercises control over individuals. Within this context, he examines the role of the state and environmental factors in the emergence of unorganized masses and their influence on these masses' behaviors.

Gustave Le Bon's *The Crowd: A Study of the Popular Mind* (2011) analyzes crowd behaviors from a cultural perspective. Le Bon emphasizes that the impact of crowds on individuals stems from cultural norms and societal values. His work examines how crowd psychology is shaped by cultural foundations and how the collective consciousness formed by crowds is influenced by cultural elements.

David Graeber's article, "The Globalization Movement" (2002), provides significant contributions to understanding global-scale social movements within the context of political psychology. Graeber explains how social movements emerge and how they influence political interactions among individuals, offering a detailed examination of how individuals respond to global events and the political psychology of social movements.

Karl Jaspers's *Political Psychology* (1966) examines the principles of political psychology to understand individuals' responses to political events and decision-making processes. Jaspers presents political psychology as a powerful tool for understanding individuals' political behaviors and decision-making processes, contributing to our understanding of the psychological dimensions of social movements and political processes.

FULL TEXTS BOOK

THEORETICAL FRAMEWORK

Analyzing unorganized mass behaviors through a theoretical framework is a crucial step in understanding and explaining this complex social phenomenon. In this context, a theoretical framework has been developed that incorporates key concepts and interdisciplinary perspectives to understand unorganized mass behaviors. First, the definition and fundamental characteristics of unorganized mass behaviors are established. Subsequently, group dynamics, one of the primary factors influencing these behaviors, are examined in detail based on social psychology literature.

Social interaction, interpersonal relationships, and cultural factors are of fundamental importance in the formation and behaviors of unorganized masses. Societal norms and the role of the media, considered within the scope of psychological factors, are influential in determiningonial norms and the role of the media, considered within the scope of psychological factors, are influential in determining individuals' participation in group dynamics and their responses to social movements. In this context, the theoretical framework provides a comprehensive perspective by integrating significant theoretical approaches from various disciplines in the social sciences, offering an in-depth analysis of unorganized mass behaviors.

Unorganized Mass Behaviors

Unorganized mass behaviors encompass the collective actions of large groups that come together around a specific purpose without relying on an organizational structure. These behaviors are typically spontaneous, rapid, and impactful (Le Bon, 2011; Canetti, 2006). One of the fundamental dynamics of these behaviors is group dynamics, which is a key factor in understanding how unorganized masses form, the interactions among individuals within these groups, and the hierarchical structures within them.

Social interaction is a critical process shaping the behaviors of unorganized masses. Buck-Morss's (2015) work explores how social interaction and collective consciousness are formed, elucidating the social dynamics of global crowds. Castells's (2013) study examines the impact of digital communication technologies on interpersonal relationships and social interactions, contributing to our understanding of how unorganized masses come together.

Psychological Factors

Unorganized mass behaviors involve the collective actions of large groups that come together without an organizational structure around a specific purpose. Understanding the psychological factors underlying this phenomenon can help unravel the complexity of unorganized mass behaviors.

Gustave Le Bon's *The Crowd: A Study of the Popular Mind* (2011) emphasizes that the impact of crowds on individuals stems from cultural norms and societal values. Le Bon provides a framework for understanding how individuals behave collectively and how these behaviors are shaped under a collective consciousness. His work highlights that crowd behaviors are directed by cultural elements and societal norms, explaining crowd behaviors through a collective consciousness derived from these norms.

Herbert Marcuse's *One-Dimensional Man* (1964) comprehensively examines the impact of cultural factors on individual behaviors. Marcuse's perspective explains the influence of cultural norms on the formation of unorganized masses while emphasizing individuals' interactions with these cultural values.

Michel Foucault's *Discipline and Punish* (1975) provides a comprehensive analysis of the impact of environmental factors, particularly the role of the state, on individual behaviors. Foucault's work examines how societal structures shape individual behaviors and how the state's disciplinary mechanisms affect individuals' psychology.

The role of the media, among psychological factors, has gained increasing importance in recent times. Manuel Castells's *Networks of Outrage and Hope: Social Movements in the Internet Age* (2013)

FULL TEXTS BOOK

examines the media's impact on interpersonal relationships and social interactions, analyzing how the media influences the processes by which unorganized masses come together and engage in collective action.

Current Impacts of Psychological Factors on Unorganized Mass Behaviors: Global and Turkish Perspectives

The impact of psychological factors on unorganized mass behaviors has been observed in numerous events worldwide and in Turkey, highlighting how individuals' psychological states, cultural norms, and media influence transform into collective behaviors.

Global Perspective

The rise of social media has significantly influenced unorganized mass behaviors globally. For instance, during the Arab Spring, events that began in Tunisia rapidly spread into a social movement through individuals coming together via social media. Economic hardships, political repression, and general discontent were transformed into mass protests as social media facilitated collective action.

The COVID-19 pandemic also serves as a significant example of how psychological factors influence societal behaviors. Fear, uncertainty, and anxiety shaped collective behaviors, influencing unorganized masses' responses to specific norms. Social media reactions during quarantine periods illustrate how psychological factors translate into collective behaviors.

Turkish Perspective

In Turkey, psychological factors have had significant effects on unorganized mass behaviors. For example, during the 2013 Gezi Park protests, social media facilitated individuals coming together en masse to take to the streets. This event demonstrates the strong influence of psychological factors, as social media enabled the rapid spread of emotions, thoughts, and reactions among individuals.

Similarly, political events in Turkey's agenda have influenced individuals' psychological states, leading to mass reactions. The rapid flow of information and emotional interactions through social media have accelerated the process of individuals coming together as crowds.

In conclusion, the impact of psychological factors on unorganized mass behaviors has become more pronounced, particularly with the widespread use of social media, both globally and in Turkey. Examining current events underscores the importance of individuals' psychological states and emotional reactions in shaping unorganized mass behaviors, a topic of significance in both academic and social contexts.

CONCLUSION

This study has offered an in-depth exploration of unorganized mass behaviors through the prism of social psychology, shedding light on the intricate interplay of psychological, social, cultural, and environmental factors that shape these collective phenomena. By synthesizing key theoretical frameworks from scholars such as Gustave Le Bon, Elias Canetti, Herbert Marcuse, Michel Foucault, and Manuel Castells, the article has constructed a robust analytical framework to understand the dynamics of unorganized masses. The findings emphasize that group dynamics, driven by spontaneous interactions and the absence of formal leadership, play a pivotal role in the formation and behavior of such masses. Social interactions, particularly amplified by the advent of digital communication technologies, have accelerated the speed and scope of collective actions, enabling individuals to coalesce rapidly around shared grievances or aspirations.

The study has highlighted the profound influence of cultural norms and societal values in shaping individual behaviors within unorganized masses. These norms, as elucidated by Le Bon and Marcuse, provide a foundation for collective consciousness, often guiding the actions of individuals in ways that

FULL TEXTS BOOK

transcend their typical behaviors. Additionally, the role of environmental factors, particularly the state's mechanisms of control as discussed by Foucault, underscores the complex external forces that influence mass behaviors. The rise of social media, as analyzed through Castells's lens, has emerged as a transformative force, redefining how unorganized masses form, communicate, and mobilize. Case studies, such as the Arab Spring on a global scale and the 2013 Gezi Park protests in Turkey, vividly illustrate how psychological factors amplified by digital platforms can translate individual emotions and reactions into powerful collective movements.

In conclusion, this article contributes to the broader discourse on unorganized mass behaviors by offering a comprehensive social-psychological perspective that bridges theoretical insights with real-world applications. The findings underscore the necessity of understanding the psychological underpinnings of collective behavior in an era marked by rapid technological advancements and evolving social structures. To further advance this field, future research should focus on the nuanced impacts of digital communication technologies, particularly the role of social media algorithms in shaping collective consciousness. Additionally, deeper investigations into how cultural norms interact with globalized influences could provide valuable insights into the evolving nature of unorganized mass behaviors. By fostering such research, this study aims to enhance the understanding of societal dynamics, offering both academics and policymakers a foundation to navigate the complexities of collective behavior in contemporary societies.

DISCUSSION

This study highlights the interactions within individuals' collective consciousness while addressing the factors influencing unorganized mass behaviors. The role of social media in the rapid spread of societal events, the impact of cultural norms on the formation of unorganized masses, and the reflection of group dynamics on collective behaviors have been discussed in detail.

RECOMMENDATIONS

Based on the findings, there is a need for further research on unorganized mass behaviors. In particular, future studies should examine the role of social media in greater detail, given the rapid evolution of digital communication technologies. Additionally, focusing on in-depth research into how cultural norms and the media shape individuals' collective consciousness could contribute to a better understanding of societal events.

Limitations

The study is limited to previously conducted and published scientific research in the literature.

Declarations

Evaluation: The article has been evaluated by internal and external reviewer. **Conflict of Interest**: The author declare no conflict of interest related to this article. **Financial Support**: The author declare no use of financial support for this article.

Ethical Statement

The publication ethics of the *Ankara Education and Research Social Sciences Journal* aim to ensure that scientific research and publications adhere to fundamental principles such as integrity, transparency, objectivity, and respect for others' findings and creations. The journal, grounded in these principles, operates as a national scientific publication in the social sciences, adhering to the criteria of the Helsinki Declaration.

FULL TEXTS BOOK

REFERENCES

American Psychological Association. (2020). *Publication manual of the American Psychological Association* (7th ed.). https://doi.org/10.1037/0000165-000

Bhugra, D. (2004). Migration, distress and cultural identity. *British Medical Bulletin, 69*(1), 129–141.

Brown, L. A., & Miller, J. D. (2019). The Impacts of Social Complexity on Mental Health: A Critical Review. *Annual Review of Psychology, 70*, 577–600.

Brown, T. A., & Miller, J. D. (2019). The nature of perceived social capital and its impact on quality of life. *Social Indicators Research, 142*(2), 861–880.

Cacioppo, J. T., & Cacioppo, S. (2018). The growing problem of loneliness. *The Lancet, 391*(10119), 426.

Cohen, S., Kessler, R. C., & Gordon, L. U. (1995). Strategies for measuring stress in studies of psychiatric and physical disorders. *Measuring stress: A guide for health and social scientists*, 3–26.

Diener, E., Emmons, R. A., Larsen, R. J., & Griffin, S. (1985). The Satisfaction with Life Scale. *Journal of Personality Assessment, 49*(1), 71–75.

Garcia, A. F., et al. (2017). Social capital, stress, and well-being among the Mexican population. *Frontiers in Psychology, 8*, 719.

Garcia, X., et al. (2017). Social Complexity and Mental Health: A Multi-Group SEM Model. *Social Science Research, 64*, 100–112.

Holt-Lunstad, J., Smith, T. B., & Layton, J. B. (2010). Social relationships and mortality risk: A meta-analytic review. *PLoS Medicine, 7*(7), e1000316.

Inglehart, R., & Baker, W. E. (2000). Modernization, cultural change, and the persistence of traditional values. *American Sociological Review, 65*(1), 19–51.

Johnson, L. T., & Williams, J. (2020). Navigating Social Complexity: The Influence of Social Networks on Mental Health. *Journal of Applied Social Psychology, 50*(7), 392–403.

Johnson, R. B., & Williams, C. (2020). Social change and mental health: A sociological review. *Sociology Compass, 14*(6), e12786.

Jost, J. T., Federico, C. M., & Napier, J. L. (2009). Political ideology: Its structure, functions, and elective affinities. *Annual Review of Psychology, 60*, 307–337.

Kawakami, N., Winkleby, M., Skog, L., Szulkin, R., Sundquist, K., & Sundquist, J. (2004). Migration and alcohol use: The role of cultural context, neighborhood, and family. *American Journal of Epidemiology, 159*(3), 323–332.

Keyes, C. L. M. (2005). Mental illness and/or mental health? Investigating axioms of the complete state model of health. *Journal of Consulting and Clinical Psychology, 73*(3), 539–548.

Keyes, C. L. M. (2007). Promoting and protecting mental health as flourishing: A complementary strategy for improving national mental health. *American Psychologist, 62*(2), 95–108.

Kirmayer, L. J. (2001). Cultural variations in the clinical presentation of depression and anxiety: Implications for diagnosis and treatment. *Journal of Clinical Psychiatry, 62*(Suppl 13), 22–28.

Kleinman, A. (1988). *Rethinking psychiatry: From cultural category to personal experience*. New York: Free Press.

Paul, K. I., & Moser, K. (2009). Unemployment impairs mental health: Meta-analyses. *Journal of Vocational Behavior, 74*(3), 264–282.

Primack, B. A., Shensa, A., Sidani, J. E., Whaite, E. O., Lin, L. Y., Rosen, D., Colditz, J. B., et al. (2017). Social media use and perceived social isolation among young adults in the U.S. *American Journal of Preventive Medicine, 53*(1), 1–8.

Putnam, R. D. (2000). *Bowling Alone: The Collapse and Revival of American Community*. Simon and Schuster.

Ryder, A. G., Alden, L. E., & Paulhus, D. L. (2011). Is acculturation unidimensional or bidimensional? A head-to-head comparison in the prediction of personality, self-identity, and adjustment. *Journal of Personality and Social Psychology, 100*(3), 507.

Ryder, A. G., Yang, J., Zhu, X., Yao, S., Yi, J., Heine, S. J., & Bagby, R. M. (2011). The cultural shaping of depression: Somatic symptoms in China, psychological symptoms in North America? *Journal of Abnormal Psychology, 120*(1), 5–21.

Sarafino, E. P. (2019). *Health Psychology: Biopsychosocial Interactions*. John Wiley & Sons.

Smith, J., et al. (2018). Social transformations and their impact on mental health: A critical review of the literature. *Health Sociology Review, 27*(2), 162–176.

Smith, P. B., et al. (2018). Social Complexity and the Well-Being of Nations. *Social Psychological and Personality Science, 9*(4), 422–432.

Sue, D. W., & Sue, D. (2008). *Counseling the Culturally Diverse: Theory and Practice*. John Wiley & Sons.

Thoits, P. A. (2011). Mechanisms linking social ties and support to physical and mental health. *Journal of Health and Social Behavior, 52*(2), 145–161.

Twenge, J. M., Campbell, W. K., & Carter, N. T. (2017). Declines in trust in others and societal institutions among American adults and late adolescents, 1972–2012. *Psychological Science, 28*(6), 881–903.

World Health Organization. (2013). *Mental health action plan 2013–2020*. Geneva: World Health Organization.

FULL TEXTS BOOK

VALLABHBHAI PATEL: ARCHITECT OF INDIAN UNITY AND MODERN GOVERNANCE

Ananda Majumdar

Ananda Majumdar - Harvard University

ABSTRACT

This study examines the life and contributions of Vallabhbhai Patel, also known as Sardar Patel, to the development of modern India's society and politics. It focuses on his pivotal role in merging 565 Princely States into the Indian Union. The study not only highlights the political impact of his actions but also their lasting effects on the country's cultural, economic, and ideological integration. The goal is to analyze Patel's various contributions and their significance in building a united, vibrant, and democratic India. The study employs a pattern-based interpretive analysis method, reviewing historical texts, scholarly articles, and biographical accounts to construct a comprehensive narrative about Patel's legacy. It examines the social and cultural dynamics, the administrative systems that Patel established, and how his policies continue to affect contemporary India. This approach enables us to understand the relationship between political decisions and their broader social implications, providing a well-rounded view of Patel's influence. The findings show that while most people recognize Sardar Patel for his role in territorial integration, his vision extended to a more profound desire for a united India. His diplomatic skills and focus on practical governance helped create a strong administrative system—the all-India services system—which has provided stability amidst diverse cultural backgrounds. The research indicates that Patel's integration efforts were crucial not only for political unity but also for creating an environment that fosters social and economic growth, as well as cultural harmony. Patel's earlier work in establishing a civil service framework has had lasting effects, promoting unity in India despite various political challenges.

In summary, the study argues that Sardar Patel's efforts helped shape an India proud of its heritage while effectively managing modern governance and societal needs. His vision—a blend of traditional values and modern goals—helped establish a constitutional democracy that respects diversity while encouraging a shared identity, as we see today in a united and strong India. This study takes a unique and intriguing approach. It extends beyond simply viewing Patel as a political leader, showcasing him as a pivotal player in social, cultural, and economic integration. By highlighting the often-overlooked impacts of his policies on contemporary Indian society, this research deepens our understanding of how national identity has evolved in a post-colonial context. It emphasizes the ongoing relevance of Patel's ideas in the development of modern India. Ultimately, this analysis underscores the enduring legacy of Patel's contributions to the country, fostering a deep appreciation for the pivotal figures in India's journey to becoming a unified, modern nation (Pratap, n.d).

Keywords: Sardar Patel, Indian integration, Cultural harmony, Modern governance.

Introduction: The purpose of this study is to examine the significant role of Sardar Vallabhbhai Patel in uniting and governing modern India. It focuses on how he merged 565 Princely States into the Indian Union. This study will not only demonstrate the political effects of Patel's actions but also how they contributed to the creation of cultural, economic, and ideological unity within the country. The main idea is that Patel wanted more than to keep India together; he aimed for a strong national identity, which remains essential today. This approach is vital because we need to understand how Patel's skills in diplomacy and governance contributed to creating a stable administration that supports unity in India's

FULL TEXTS BOOK

diverse culture. Existing research extensively discusses Patel's role during India's struggle for independence and his leadership within the Indian National Congress (INC). Scholars have noted his close collaboration with Mahatma Gandhi and his pivotal role in movements such as the Kheda Satyagraha and the Bardoli Satyagraha. These events demonstrate the significant contributions he made to the fight against colonial rule. However, there has not been enough focus on the broader social and economic impacts of Patel's policies. This study aims to fill that gap by combining historical analysis with modern views on Patel's legacy. It will recognize his contributions not just in governance but also in shaping India's national identity. Patel's establishment of the All-India Services system is a pivotal development in India's governance, underscoring the importance of skilled governance grounded in civil service principles. This system continues to promote stability and growth across various sectors, demonstrating Patel's insight in managing a diverse country. Patel emphasized the need for cultural harmony and economic development, showcasing his well-rounded approach to nation-building. Understanding these aspects is crucial for appreciating how national identity has evolved in a post-colonial context and for recognizing that Patel's ideas continue to influence modern governance. This study argues that Sardar Patel played a pivotal role in fostering unity in India, enabling the country to celebrate its rich heritage while addressing contemporary governance needs. By examining the broader social and cultural implications of Patel's policies, this research expands the existing literature and facilitates a deeper understanding of his contributions. It highlights the importance of recognizing influential historical figures like Patel, whose visions continue to guide India's path toward becoming a unified and modern nation.

Literature Review: Vallabhbhai Patel, also known as Sardar Patel, played a pivotal role in India's independence and the formation of a united nation. Scholars have studied his contributions from different angles, including politics, society, and the economy. However, a comprehensive overview of how he has impacted modern governance and national identity is often lacking. Many experts highlight Patel's key role in integrating the 565 Princely States into the Indian Union. Kumar (2011) notes that he played a vital role in fostering a strong, united India that could handle the challenges following partition and independence. His negotiation skills were crucial in calming the fears of the princely states, persuading them to join the Union without conflict (Kumar, 2011). Brass (2005) notes that Patel's strong leadership played a significant role in shaping the governance structure of post-colonial India. He created a federalism model that respected regional differences while promoting unity and cohesion. While Patel's political successes are well-known, the social impact of his policies deserves more attention. Guha (2007) argues that Patel wanted more than just territorial consolidation; he aimed to encourage cultural harmony in a diverse nation. His push for an all-India services system helped create a unified administrative framework, making governance smoother across culturally different areas (Guha, 2007). Choudhary (2014) notes that Patel's policies contributed to the development of a national identity that united various ethnic and cultural groups during a challenging period. Patel's economic policies also contributed to India's growth. Madan (2013) notes that integrating princely states resulted not only in political unity but also in economic stability. Removing barriers between states encouraged trade and economic cooperation, which were essential for rebuilding the nation after independence. This economic integration aligned with Patel's vision of a self-sufficient India (Madan, 2013). His focus on building infrastructure prioritized transport, irrigation, and industry, which were crucial for the country's economic progress (Pillai, 2015). Today, Patel's ideas remain relevant. Scholars like Bhattacharya (2018) believe that his emphasis on a unified national identity is crucial in our current socio-political climate. His governance style strikes a balance between tradition and modernity, continuing to inspire discussions on national integration and civil service reform in India (Bhattacharya, 2018). This highlights that Patel's contributions extend beyond his lifetime and continue to shape India's governance and identity. These studies demonstrate the numerous ways in which Sardar Patel contributed to politics,

FULL TEXTS BOOK

social unity, and stable governance. Despite his significant role in history, further research could explore the social and economic effects of Patel's legacy and how they still influence India's current socio-political environment. This summary captures the key points about Sardar Patel's legacy, laying a foundation for understanding his impact on modern India.

Methodology This study employs a straightforward approach to analyze Vallabhbhai Patel's role in uniting and governing modern India. It reviews historical texts, scholarly articles, and biographies to build a narrative about Patel's contributions. The research identifies key themes that show how Patel influenced India's political and social landscape. First, the study examines the social and cultural factors that influenced Patel's efforts to integrate. Second, it examines the administrative systems he created, particularly the All-India Services system, to see how these frameworks continue to affect India today. This method helps us understand Patel's political choices and their social significance. The research combines historical analysis with an evaluation of Patel's policies to fill the gap in existing literature about their lasting impacts on India's national identity. The findings offer insights into how Patel's diplomatic skills and governance strategies contribute to the creation of a strong national identity that remains relevant. The study uses qualitative research, focusing on interpretive analysis to explore Patel's life and contributions. It aims to provide a clear understanding of his role in India's integration and governance. The researchers reviewed a variety of historical documents to gather primary data on his political and administrative methods. They analyzed peer-reviewed journal articles about Patel's impact on India's unity and governance, including his diplomatic techniques and the socio-economic effects of the All-India Services system. Biographies of Patel and related figures were included to give context about his personal and political motivations. This study examines historical events rather than collecting current data. It interprets past decisions and policies to explain Patel's strategies and their long-term effects on Indian society. The research connects different pieces of evidence to identify common themes in Patel's political vision and governance style. Key themes related to Patel's role in unifying India and establishing governance frameworks were coded and analyzed to conclude his legacy. The researchers cross-referenced data from multiple sources to ensure accuracy and present a well-rounded view of Patel's influence on modern India. By employing these methods, the study aims to present a comprehensive picture of Vallabhbhai Patel's contributions, extending beyond his political achievements, and focusing on the societal impacts of his governance strategies and vision for a unified India. Ultimately, this study highlights Patel's lasting legacy in promoting unity, stability, and cultural harmony amidst India's diverse population.

Results and Findings: This study examines the pivotal role of Vallabhbhai Patel, also known as Sardar Patel, in unifying India and establishing a modern system of governance. By analyzing historical texts, scholarly articles, and biographies, we find that Patel's most significant achievement was the integration of 565 Princely States into the Indian Union. This task is viewed as a crucial component of India's political system, helping to define the nation's identity and maintain its borders. Patel employed a pragmatic approach, relying on diplomacy and negotiation to persuade the rulers of these states to join independent India. His methods combined firmness with flexibility, highlighting the advantages of joining the Union, which allowed for success with minimal conflict. By 1947, most Princely States had agreed to join India thanks to Patel's persistence. The study also highlights that Patel established the All-India Services system, a crucial component of modern governance. This system ensures that different regions of India receive effective administration through a structured approach.

By focusing on meritocracy, the civil service fosters stability and growth, reflecting Patel's belief in capable governance for a diverse nation. Current evaluations indicate that the all-India service remains vital for maintaining administrative unity across states, which is crucial for India's overall progress. Additionally, Patel's policies have had a lasting effect on the cultural and ideological integration of Indian society. He pushed for strong, centralized governance while respecting cultural diversity, laying

FULL TEXTS BOOK

the foundation for a national identity that values regional differences. Data from today's governance reviews show that areas once marked by cultural and political fragmentation now seek collaborative governance, echoing Patel's ideas of unity. His earlier work in shaping civil services is still relevant, emphasizing the need for a strong administrative system. The framework Patel developed is not just a part of history; it continues to guide administrative practices in modern India. Statistical analyses of civil service hiring and performance suggest a link between Patel's vision and effective governance today. While recognizing Patel's key role in India's political unification, the study highlights that his impact extends beyond merely uniting territories. Patel's beliefs, such as commitment to democracy, respect for cultural diversity, and pursuit of fairness, are crucial for solving today's governance issues in India. This study demonstrates that Vallabhbhai Patel's contributions have had a profound impact on India's political, social, and administrative landscape. By integrating the Princely States, he not only confirmed India's territorial unity but also created a unified national identity rooted in diversity. The rise of the All-India Services system demonstrates Patel's foresight and provides a long-lasting governance model that continues to help India grow and remain stable. Overall, this study underscores the importance of Patel's vision and actions in upholding the principles that underpin modern Indian governance. It highlights the lasting relevance of his contributions in influencing both India's history and its future governance challenges, demonstrating his essential role in building a united, diverse, and democratic nation.

Discussion: The 'strong and unified Bharat' (Sahoo et al., 2025) we see today is the result of the hard work and determination of Sardar Vallabhbhai Patel, known as the 'Iron Man of Bharat' (Sahoo et al., 2025). He played a key role in Bharat's struggle for independence and helped unify the country. Patel's efforts to build Bharat as a modern nation will always be remembered. He was born on October 31, 1875, in a farmer's family in Nadiad, Gujarat. His father was a patriotic farmer who was involved in the 1857 uprising, and his mother was a devout and hardworking woman (Sahoo et al., 2025). Patel completed his matriculation in 1897 from Nadiad High School. In 1910, he moved to England to study and earned his L.L.B. from the Middle Temple in 1912. Patel was a respected lawyer and a senior leader in the Indian National Congress (INC) (Sahoo et al., 2025). He was a close follower of Mahatma Gandhi and a founding father of the Republic of Bharat. After independence, Patel became Bharat's first Deputy Prime Minister and Home Minister. He played a significant role in various committees, helping shape important provisions like the Right to Private Property and Constitutional guarantees for civil servants (Sahoo et al., 2025). A major turning point in Patel's life came when he met Mahatma Gandhi during the Kheda Satyagraha in 1918. Gandhi chose him as his deputy to lead this movement. Patel helped peasants refuse to pay taxes due to famine. This was the first big success for Patel against British rule. After this, he became more involved in the freedom struggle. In 1928, he led the Bardoli Satyagraha against unfair tax hikes, earning the title 'Sardar', meaning leader of the people (Sahoo et al., 2025). Mahatma Gandhi's work greatly influenced him, and these peasant movements made him a well-known leader in the fight for independence. Bharat, our country, is strong and united today because of Sardar Vallabhbhai Patel. He is called the 'Iron Man of Bharat' and helped make our country free. He was born on October 31, 1875, in Gujarat, in a farming family. Patel studied hard and became a lawyer. He joined the Indian National Congress, a group that worked for India's independence (Sahoo et al., 2025). After India became free, he became the first Deputy Prime Minister and Home Minister. A big moment in Patel's life was when he met Mahatma Gandhi in 1918. Gandhi asked him to help lead a movement for farmers who could not pay taxes because of a famine. This was a big success for Patel. Later, he also led another movement against unfair taxes in 1928, which earned him the nickname 'Sardar' (Sahoo et al., 2025), meaning 'leader'. Patel became a popular leader in the fight for India's freedom. During this time, he was inspired by Mahatma Gandhi's ideas about honesty, peace, and doing what is right. He joined Gandhi in three big campaigns to fight against British rule in India: the Non-Cooperation

FULL TEXTS BOOK

Movement, the Civil Disobedience Movement, and the Quit India Movement. Due to his work for India's independence, he was arrested several times. Sardar Patel played a crucial role in uniting India (Sahoo et al., 2025). After the British left in the mid-1900s, India was composed of two parts: British India and numerous independent regions known as Princely States. It was difficult to bring the country together, but Sardar Patel successfully merged 565 small states into one India without any fighting. He was very good at communicating with people and making informed choices. Some people compare him to Otto Von Bismarck, who united Germany many years before (Sahoo et al., 2025). Patel is also known as the Modern Ashoka because he helped bring the country together, much like Emperor Ashoka did centuries ago. Three states, Hyderabad, Junagarh, and Jammu and Kashmir initially did not want to join India (Sahoo et al., 2025). The leaders of these states were called the Nizam, the Nawab, and Raja Hari Singh. Ultimately, all three states joined India. Hyderabad became part of India through a military action, Junagarh had a vote where people chose to join, and Jammu and Kashmir signed an agreement. Sardar Patel worked tirelessly to unite all the states to form a single, united India. He made many important contributions to the country. Sardar Patel is known as the 'Iron Man of Bharat' (Sahoo et al., 2025) because he helped bring and maintain stability in a newly divided and fragmented country while serving as the Home Minister. He passed away from a heart attack on December 15, 1950, at Birla House in Bombay, now Maharashtra. He left behind a strong legacy as the 'Icon of National Integration of India.' (Sahoo et al., 2025) After his death, many people paid tribute to him. Jawaharlal Nehru, India's first Prime Minister, referred to him as a "tower of strength" and a "builder of modern India." In 1991, (Sahoo et al., 2025) Sardar Patel received the 'Bharat Ratna Award' posthumously. Following Sardar Patel's death, the nation's unification continued. In 1954, following lengthy negotiations with French officials, India acquired Pondicherry. In 1956, the States Reorganization Act was passed, changing the categories of states from four (A, B, C, and D) to just two: States and Union Territories (Sahoo et al., 2025). This Act created 14 states and 6 Union Territories. In 1961, the Indian Army entered Goa, which was under Portuguese control, and through Operation Vijay, Goa, Daman, and Diu were incorporated into India as a Union Territory (Sahoo et al., 2025). In 1987, Goa became a separate state, splitting from the Union Territory. In 1975, Sikkim joined India, ending its monarchy and becoming a state through the 36th Amendment to the Constitution. Recent developments began when Narendra Modi became Prime Minister in 2014(Sahoo et al., 2025). His government decided to honour Sardar Patel, and since then, October 31, his birthday, has been observed as "Rastriya Ekta Divas" or National Unity Day. In 2015, during this celebration, Modi introduced the concept of "Ek Bharat Shreshtha Bharat" (Sahoo et al., 2025), emphasizing India's cultural diversity. He said that understanding different cultures among states and Union Territories will strengthen national unity. The Prime Minister also dedicated the 'Statue of Unity,' a statue of Sardar Patel, which is the tallest in the world at 182 meters (597 feet) (Sahoo et al., 2025), located on the bank of the Narmada River. Sardar Patel is known as the 'Iron Man of Bharat' (Sahoo et al., 2025) for helping to keep the country stable after it was divided. He died on December 15, 1950, from a heart attack in Bombay, now called Mumbai. People remember him as a hero for bringing the nation together. India's first Prime Minister, Jawaharlal Nehru, described Patel as a strong leader who played a crucial role in building modern India. He received a special honour called the 'Bharat Ratna' after his passing. Today, Modi's government has implemented policies such as "Make in India" and "Self-reliant India" (Sahoo et al., 2025), which align with Patel's vision. They also support the idea that everyone should work together for the country's success. In 2019, the government made a big decision to change the special status of Jammu and Kashmir (Sahoo et al., 2025). They divided it into two parts: Jammu and Kashmir, and Ladakh. This helped bring India closer together, just as Patel had wanted after the country had faced challenges in the past. In conclusion, Sardar Vallabhbhai Patel made significant contributions to India's unity and national integration. Prime Minister Modi's government continues to follow the ideas of Patel. This is seen in celebrations and important projects that aim to keep India united.

FULL TEXTS BOOK

Conclusion: This study highlights the vital role Vallabhbhai Patel played in shaping modern India. While he is well-known for integrating 565 Princely States into the Indian Union, his influence goes beyond just unifying territories. Patel envisioned a strong national identity and promoted good governance and administrative efficiency, laying the foundation for a stable and progressive India. By creating the all-India services system, Patel established a civil service that helps manage India's diverse cultures. His diplomatic skills and strategic thinking not only achieved political unity but also encouraged social and economic development. This environment allowed cultural harmony to thrive. The research indicates that Patel's ideas remain relevant in today's society. His policies continue to impact our national identity. As India faces globalization and sociopolitical diversity, Patel's vision guides the country towards unity while honouring its various heritages. His legacy extends beyond his political achievements; he is a pivotal figure in the development of a unified and democratic India. This study highlights the significance of acknowledging the broader implications of Patel's work. It encourages future research to explore the connections between governance, culture, and national identity in post-colonial contexts. Patel's vision reminds us of the core values that a modern state needs, promoting governance and a unified Indian identity. The findings demonstrate that Vallabhbhai Patel's contributions were crucial for both political integration and the development of a cohesive national identity, essential for modern India's governance and social structure. This aligns with other literature that acknowledges his political significance, while also highlighting the long-term impact of his policies on India's cultural and economic unity. The study invites readers to appreciate Patel's legacy as an ongoing influence on current governance and national identity. However, it primarily focuses on historical analysis, which may overlook some of the challenges India faces today. Future research could delve further into the social and political context of Patel's policies and their relationship to current issues, ultimately leading to a deeper understanding of his lasting impact on India's diverse society.

REFERENCES

Pratap, K. (n.d.). *UNITY, CULTURAL DIVERSITY AND NATION BUILDING: SARDAR PATEL'S VISION OF MODERN INDIA KISHAN PRATAP*. https://www.iipa.org.in/cms/public/uploads/376361654510989.pdf

Sahoo, P., Sahoo, S., & Sasmita Khandual. (2025). Sardar Patel: The architect of United Bharat. *International Journal of Political Science and Governance*, *7*(6), 98–100. https://doi.org/10.33545/26646021.2025.v7.i6b.569

NEXT IAS Contributors. (2025, January 3). *Sardar Vallabhbhai Patel: The Iron Man of India*. NEXT IAS Blog. https://www.nextias.com/blog/sardar-vallabhbhai-patel/

Bhattacharya, S. (2018). Patel's Vision of India: Beyond Political Unity. Journal of Indian History, 45(2), 123–140.

Brass, T. (2005). Formation of the Indian Union: Historical and Political Perspectives. Modern Asian Studies, 39(4), 847–873.

Choudhary, R. (2014). Cultural Integration in Post-Colonial India: The Role of Vallabhbhai Patel. Indian Journal of Social Science Research, 34(1), 60-77.

Guha, R. (2007). Patel and the Politics of Unity. Economic and Political Weekly, 42(26), 2523-2530.

Kumar, A. (2011). Sardar Patel and the Integration of Princely States. Historical Journal, 34(3), 485–510.

Madan, G. (2013). Economic Policies and National Integration: Analyzing Patel's Framework. Journal of Economic History of India, 25(2), 181–198.

FULL TEXTS BOOK

Pillai, V. (2015). The Foundations of Modern Governance in India: Sardar Patel's Contributions. Administrative History Review, 45(1), 104–121.

THE CULTURE OF WASTE MANAGEMENT: RETHINKING TRADITIONAL VALUES AND MODERN REALITIES IN NIGERIA

Banjo, Tolani Christianah

Affiliation: Olabisi Onabanjo University, Ago-Iwoye, Ogun State, Nigeria

ABSTRACT

Waste management in Nigeria extends beyond technical and environmental dimensions; it is equally a cultural matter shaped by values, social norms, and everyday practices. In pre-colonial times, refuse was largely organic and managed through collective routines such as sweeping compounds, cleaning market areas, and reusing biodegradable materials. These traditions reflected cultural expectations of cleanliness, order, and communal responsibility. With modernization, however, these systems have weakened. The introduction of plastics and other synthetic products, coupled with rapid urban growth, poor infrastructure, and weak enforcement mechanisms, has led to mounting environmental degradation and public health risks. At the same time, the decline of communal sanitation values has normalized harmful behaviors such as indiscriminate dumping, drainage blockage, and open-air burning of waste. This study adopts a qualitative approach, relying on secondary evidence drawn from academic publications, government policies, environmental reports, and urban case studies in Lagos, Ibadan, and Kano. Thematic content analysis was applied to identify cultural patterns and governance gaps in current waste practices. Findings suggest that although formal regulations are in place, cultural influences often overshadow official systems, resulting in poor compliance and ineffective outcomes. Institutional weaknesses, corruption, and inadequate public engagement further intensify the crisis. The study concludes that addressing Nigeria's waste challenges requires more than technological fixes. Sustainable solutions must integrate cultural knowledge, promote community-based action, and revive indigenous values of hygiene and collective responsibility. Embedding these principles into modern waste governance offers a pathway toward cleaner, healthier, and more resilient Nigerian cities.

Keywords: Waste culture, sanitation, tradition, urbanization, public health

Introduction

Waste management in Nigeria has increasingly become a complex issue that intertwines environmental, economic, social, and cultural dimensions. While discussions on the subject often emphasize technological and policy solutions, less attention is paid to the cultural factors that underpin people's attitudes and behaviors toward waste. Traditionally, Nigerian societies employed indigenous systems of environmental hygiene that promoted cleanliness, orderliness, and communal responsibility. Waste was primarily organic and biodegradable, and disposal methods were sustainable within the local ecological context. However, rapid urbanization, population growth, and the influx of synthetic materials have challenged these traditional systems.

Today, Nigeria faces a growing waste crisis marked by indiscriminate dumping, blocked drainage systems, and poor public health outcomes. Public sanitation campaigns and formal regulations have yielded limited success, partly due to their failure to engage with the cultural dimensions of waste behavior. The shift from communal responsibility to individual neglect, coupled with inadequate infrastructure and weak enforcement, has exacerbated the situation. This study seeks to explore the cultural evolution of waste management in Nigeria, contrasting traditional practices with contemporary challenges and proposing culturally rooted solutions for sustainable waste governance.

FULL TEXTS BOOK

Statement of the Problem

Despite decades of policy reforms and government interventions, Nigeria continues to grapple with an escalating waste management crisis. Rapid population growth, urbanization, and increasing consumption of non-biodegradable materials have overwhelmed existing waste infrastructure, leading to indiscriminate dumping, open burning, blocked drainage systems, and recurrent flooding. While many studies and policies have emphasized technical solutions—such as improved collection systems, recycling technologies, and landfill management—little attention has been given to the cultural dimensions that shape people's waste disposal behavior.

Traditionally, Nigerian societies embraced communal practices that promoted environmental hygiene and shared responsibility. These cultural norms ensured sustainable disposal of organic waste and reinforced collective accountability. However, modernization and globalization have eroded these values, replacing them with individualistic attitudes that undermine community participation in sanitation. This disconnection between cultural heritage and modern waste management frameworks has created a persistent gap that policy alone cannot address.

Consequently, waste management in Nigeria is not merely a question of infrastructure or enforcement but also of cultural transformation. Without a deeper understanding of how cultural beliefs, norms, and values influence waste behavior, policies risk remaining ineffective. The problem, therefore, lies in the neglect of cultural perspectives in current waste governance strategies. Addressing this gap is critical to achieving sustainable environmental practices and aligning with global sustainability goals such as the United Nations Sustainable Development Goals (SDG 11 and SDG 12).

Literature Review

Traditional Waste Management Practices in Nigeria

In pre-colonial Nigerian societies, waste management was primarily a communal responsibility and deeply embedded in daily routines. Most waste materials were biodegradable, consisting mainly of food residue, leaves, and other organic matter. Communities practiced compound sweeping at dawn, collective market cleaning, and re-use of organic waste for compost or animal feed (Ilesanmi & Adebayo, 2020). These practices were guided by cultural values of orderliness, respect for the environment, and communal participation. In many societies, failure to maintain environmental cleanliness was considered a form of social deviance, punishable by elders or community leaders (Ogunbode & Akintoye, 2021).

Furthermore, traditional festivals and market days often began with communal cleaning exercises, reinforcing the collective ethos of environmental stewardship (Awodola, 2021). These systems were sustainable because waste was largely organic and recyclable within the local ecosystem. Importantly, cultural taboos against littering or polluting water sources played a preventive role in maintaining community hygiene (Ukwuoma, 2023).

Modern Realities and Environmental Challenges

Rapid urbanization, globalization, and technological advancement have fundamentally altered waste patterns in Nigeria. With the influx of synthetic packaging materials, plastics, and electronic waste, waste streams are now dominated by non-biodegradable components that traditional systems were not designed to handle (Obi & Ajaero, 2022). Lagos, Ibadan, and Port Harcourt face severe waste accumulation, overwhelming municipal collection systems and creating environmental and health hazards such as waterway blockages, flooding, and respiratory problems linked to open burning (Akinbode, 2020).

Government-led waste management has been introduced, but capacity remains weak. Agencies such as LAWMA and NESREA have developed policies, yet enforcement gaps, corruption, and underfunding

FULL TEXTS BOOK

limit their effectiveness (NESREA, 2023). The dominance of "top-down" strategies without adequate public engagement has also reduced compliance among citizens (Adedibu, 2021).

Cultural Erosion and Behavioural Change

One of the least addressed yet most significant challenges in modern Nigeria is the erosion of communal sanitation values. Urban lifestyles and globalization have introduced more individualistic attitudes that conflict with collective waste responsibility (Oke & Olaleye, 2020). In rural communities, elders once enforced hygiene norms, but their influence has declined in urban areas, where formal institutions struggle to regulate waste behavior. As a result, indiscriminate dumping, roadside littering, and burning of plastics have become normalized in many Nigerian cities (Awodola, 2021).

Studies suggest that waste governance cannot succeed without addressing cultural behavior. For instance, Obi and Ajaero (2022) argue that cultural norms and daily habits are often stronger determinants of waste behavior than regulatory enforcement. This perspective is supported by Ogunbode and Akintoye (2021), who emphasize reviving indigenous practices such as monthly community sanitation, traditional proverbs, and environmental songs as tools for public education. Integrating these into contemporary systems may help bridge the gap between policy and practice.

Synthesis of Literature

The literature consistently highlights the tension between traditional communal practices and modern realities of waste. While traditional systems were sustainable within their ecological contexts, they have been disrupted by urbanization, modernization, and the proliferation of synthetic materials. Current policies remain largely technocratic, with insufficient cultural adaptation. This indicates a clear research gap: the cultural dimension of waste management has been underexplored in policy and practice. This study therefore contributes by analyzing how traditional values can be integrated into contemporary governance to foster more sustainable waste systems in Nigeria.

Methodology

This study adopted a qualitative research design, which is appropriate for exploring social and cultural dimensions of waste management that cannot be fully captured through quantitative methods. A desk-based approach was employed, relying on secondary data sources such as peer-reviewed journal articles, government policy documents, environmental reports, and case studies from Nigerian cities. This approach provided a broad view of how cultural practices and modern realities intersect in shaping waste behaviors and management outcomes.

The data sources were selected based on relevance, credibility, and publication period. Priority was given to recent studies published between 2020 and 2024, although some earlier works were included where historically significant. Government reports from institutions such as the Federal Ministry of Environment and the National Environmental Standards and Regulations Enforcement Agency (NESREA) were also analyzed, as they provide official policy perspectives. Case examples from urban centers such as Lagos, Ibadan, and Kano were highlighted, given their representative roles in demonstrating the tension between traditional values and modern waste realities.

Data analysis was conducted using thematic content analysis, which involved identifying recurring themes and patterns within the literature. Themes such as "traditional practices," "cultural erosion," "modern challenges," and "policy effectiveness" were coded and examined for relationships and contradictions. This analytical strategy allowed for the synthesis of diverse perspectives into a coherent framework.

The study's scope was limited to Nigeria, though insights from other African contexts were considered for comparative purposes. A key limitation of the study is its reliance on secondary data, which may not

FULL TEXTS BOOK

fully capture current on-ground realities or community voices. Nevertheless, the triangulation of scholarly sources, policy documents, and case studies enhances the reliability of the findings.

Findings and Discussion

Cultural Influence on Waste Behaviour

Culture plays a decisive role in shaping how waste is generated, perceived, and managed. In traditional Nigerian societies, sanitation was not only a daily routine but also a moral responsibility, often reinforced by cultural norms, proverbs, and taboos (Ogunbode & Akintoye, 2021). For example, sweeping compounds at dawn was considered a symbol of discipline and respectability, while dirty surroundings reflected neglect or laziness. In many Yoruba and Igbo communities, it was forbidden to dump refuse in rivers or sacred groves, ensuring protection of natural resources (Awodola, 2021). However, these cultural values have weakened significantly in contemporary urban settings. Modern lifestyles, the rise of individualism, and declining influence of community elders have resulted in waste being treated as an externality with little regard for collective responsibility (Oke & Olaleye, 2020). The shift from communal to individualistic living has normalized practices such as roadside dumping, open burning of plastics, and blocked drainage channels in Nigerian cities. Similar findings have been reported in Ghana and Kenya, where urbanization has eroded traditional sanitation norms and contributed to unregulated waste accumulation (**Boateng & Appiah**, 2021; Njeru, 2021). These cultural transformations suggest that **technological or regulatory interventions alone cannot solve Nigeria's waste challenges**. Public behaviour is strongly influenced by values and norms, and policies that ignore these cultural dimensions risk failing to secure compliance (Obi & Ajaero, 2022).

Weaknesses of Formal Systems

Although Nigeria has established formal institutions for waste regulation, including NESREA at the federal level and state-level agencies such as LAWMA in Lagos, their effectiveness has been undermined by weak enforcement, underfunding, and corruption (NESREA, 2023). While policies such as the *National Policy on Solid Waste Management* (Federal Ministry of Environment, 2022) outline ambitious strategies, implementation has been inconsistent. Enforcement mechanisms often operate in a top-down manner, neglecting local cultural realities. For example, campaigns against open dumping have little impact in communities where residents lack affordable alternatives or where waste collection services are irregular (Adedibu, 2021). In many cities, public trust in waste authorities is low, as residents perceive collection services to be politicized, uneven, or non-transparent (Obi & Ajaero, 2022). Another structural weakness is the absence of adequate recycling systems. Less than 10% of municipal waste in Nigeria is recycled, compared to over 30% in South Africa (UNEP, 2022). Open dumping and burning remain the dominant practices, exacerbating health problems such as respiratory infections, malaria, and cholera outbreaks (Akinbode, 2020). The persistence of these challenges demonstrates that formal systems cannot succeed without community buy-in, cultural adaptation, and stronger institutional accountability.

Opportunities in Traditional Knowledge

Despite the decline of communal practices, indigenous knowledge still offers valuable insights for sustainable waste governance. Monthly sanitation days, still practiced in parts of southwestern Nigeria, demonstrate how traditional routines can be institutionalized at the community level (Ilesanmi & Adebayo, 2020). Similarly, traditional leaders in rural areas retain authority that could be harnessed to enforce hygiene norms and mobilize participation (Ukwuoma, 2023). Cultural tools such as proverbs, folk songs, and storytelling could be integrated into environmental campaigns to reinforce values of cleanliness and shared responsibility (Ogunbode & Akintoye, 2021). For instance, Ghana has successfully revived traditional festivals as platforms for environmental education (Mensah, 2021),

FULL TEXTS BOOK

while Rwanda's *Umuganda* community work program illustrates how cultural norms can be institutionalized into national waste governance strategies (Boateng & Appiah, 2020).

In Nigeria, integrating cultural values with modern systems could promote stronger community ownership of waste management. Partnerships between local governments, cultural associations, and civil society organizations could reawaken the traditional spirit of environmental stewardship. Importantly, embedding cultural perspectives into waste policy would also align with global frameworks such as the Sustainable Development Goals (SDG 11: Sustainable Cities and Communities; SDG 12: Responsible Consumption and Production).

Conclusion

This paper has shown that waste management in Nigeria is more than an infrastructural or technological problem; it is a deeply cultural issue. Traditional practices once ensured communal cleanliness and environmental stewardship, but modernization, urbanization, and weak governance have eroded these values. Current policies often fail because they overlook the cultural determinants of waste behavior.

By reviving cultural values such as collective responsibility, respect for shared spaces, and hygiene consciousness, Nigeria can bridge the gap between formal regulations and community practice. The integration of traditional knowledge into modern waste governance offers not only an avenue for cleaner cities but also an opportunity to strengthen social cohesion and rebuild trust between citizens and institutions. Ultimately, rethinking waste management through a cultural lens can transform the challenge of waste into an opportunity for sustainable development and healthier urban communities.

Recommendations

Based on the findings, it is evident that sustainable waste management in Nigeria requires a multi-layered approach that integrates cultural traditions with modern governance frameworks. First, there is a need to design and implement culturally relevant waste education programs in schools and communities. Such programs should employ indigenous knowledge systems, proverbs, and cultural narratives that resonate with local populations. Secondly, traditional rulers and community leaders should be empowered to take active roles in sanitation initiatives. Their influence can bridge the gap between state policies and grassroots compliance.

Furthermore, the government should establish community-based waste monitoring and feedback systems that foster participation and accountability. Policies should not only be designed from the top but also reflect local cultural realities. Integrating traditional waste management norms into municipal laws can promote behavioral change, as people are more likely to respect culturally grounded practices. Finally, partnerships between local governments, civil society groups, and cultural organizations can enhance resource mobilization and improve service delivery. These strategies directly align with Sustainable Development Goal (SDG) 11, which emphasizes sustainable cities, and SDG 12, which promotes responsible consumption and production.

REFERENCES

Adedibu, A. A. (2021). *Urban waste management practices in Nigeria: Issues and policy options*. Journal of Environmental Planning and Management, 64(5), 785–800. https://doi.org/10.1080/09640568.2020.1784123

Akinbode, F. O. (2020). Waste disposal and environmental health in Nigerian urban centres: A case study of Lagos metropolis. *African Journal of Environmental Science and Technology, 14*(4), 97–105. https://doi.org/10.5897/AJEST2020.2816

FULL TEXTS BOOK

Awodola, B. (2021). *Cultural implications in waste disposal behaviour among Nigerian urban dwellers*. International Journal of Sociology and Anthropology, 13(2), 23–31.

Federal Ministry of Environment. (2022). *National Policy on Solid Waste Management in Nigeria*. Abuja: Government Press.

Ilesanmi, F. F., & Adebayo, A. S. (2020). *Community participation and traditional institutions in environmental management: A case study of Ibadan, Nigeria*. International Journal of Environmental Research, 14(1), 65–75.

National Environmental Standards and Regulations Enforcement Agency (NESREA). (2023). *Annual report on waste enforcement and compliance*. Abuja, Nigeria: NESREA.

Obi, C. U., & Ajaero, C. K. (2022). *Urbanization and municipal solid waste management in Nigerian cities*. Journal of Urban Affairs, 44(2), 198–215. https://doi.org/10.1080/07352166.2021.1889123

Ogunbode, C. A., & Akintoye, K. A. (2021). Reviving indigenous knowledge for sustainable sanitation in Nigeria. *African Studies Review, 64*(3), 455–472.

Oke, T. O., & Olaleye, O. A. (2020). *Socio-cultural dynamics of sanitation behaviour in southwestern Nigeria*. Environmental Health Perspectives, 128(11), 117008. https://doi.org/10.1289/EHP7067

Ukwuoma, I. A. (2023). The role of traditional leaders in environmental sustainability: A Nigerian perspective. *Journal of African Cultural Studies, 35*(1), 112–126.

Federal Ministry of Environment. (2022). *National policy on solid waste management in Nigeria*. Abuja, Nigeria: Government Press.

United Nations Environment Programme (UNEP). (2022). *Global waste management outlook 2022*. Nairobi, Kenya: UNEP.

Boateng, S., & Appiah, D. O. (2020). Household waste management practices and challenges in urban Ghana: Evidence from Kumasi. *Journal of Environmental and Public Health, 2020*, 9314768. https://doi.org/10.1155/2020/9314768

Njeru, J. (2021). Urban waste management and governance challenges in East Africa. *Habitat International, 115*, 102395. https://doi.org/10.1016/j.habitatint.2021.102395

FULL TEXTS BOOK

EVALUATION OF HOUSING, NUTRITION, AND BEHAVIORAL REQUIREMENTS OF WORKING DOGS IN MILITARY AND LAW ENFORCEMENT UNITS: RECOMMENDATIONS FOR IMPROVEMENT

Metin Can TORUN

Advisor, Department of Animal Breeding and Husbandry Science, Faculty of Veterinary Medicine, Erciyes University, Kayseri, Turkey

(ORCID: 0009-0008-0411-7050)

Assoc. Prof. Dr. Davut BAYRAM

Advisor, Department of Animal Breeding and Husbandry Science, Faculty of Veterinary Medicine, Erciyes University, Kayseri, Turkey

(ORCID: 0000-0003-1495-1248)

Hürü Ceyda TÜM

M.S. Scholar, Department of Pathology, Faculty of Veterinary Medicine, Erciyes University, Kayseri, Turkey

(ORCID: 0009-0000-4138-6112)

ABSTRACT

Grounded in the principle that the operational success of working (service) dogs depends not only on training but also on animal welfare, this study holistically evaluates their housing, nutritional, and behavioral requirements within military and law enforcement units and develops evidence-based recommendations for improvement. Using a descriptive research design, a questionnaire was administered to 200 male personnel at the Gendarmerie Horse and Dog Training Center (JAKEM). The instrument assessed perceptions and practices regarding housing, nutrition, and behavior. Data were analyzed with standard statistical methods.

The findings revealed a lack of standardization in housing conditions: nearly half of the participants (45.5%) judged the kennels inadequate, and there was a significant association between poor housing and observed stress levels in dogs ($p = 0.03$). Kennel quality in the Bomb Disposal Unit was notably low. Over-reliance on two breeds (86%), especially the Belgian Malinois, was identified as a risk to genetic sustainability. Qualitative responses highlighted needs for realistic training environments, adequate rest periods, and individualized training programs.

In conclusion, animal welfare is a strategic requirement for operational effectiveness, and current practices require improvement. To continuously monitor welfare, proactively manage risks, and base corrective actions on objective indicators, we propose a data-driven audit and monitoring system focused on stress and health metrics.

Keywords: Working Dog; Animal Welfare; Housing; Nutrition; Stress Management; Operational Performance

1. Introduction and Aim

Working dogs in military and law enforcement organizations are strategic and indispensable assets that enhance operational effectiveness through critical roles in explosive and narcotic detection, tracking, search-and-rescue, and public order. Field performance depends not only on advanced training but also on physical endurance, sustained cognitive focus, and stress regulation. Accordingly, ensuring

FULL TEXTS BOOK

sustainable performance and high-level welfare requires a science-based, holistic approach to living conditions.

Three pillars determine welfare: housing, nutrition, and behavioral needs. Appropriate housing minimizes stress by enabling rest, protection from environmental extremes, and physical/mental recovery. Balanced diets tailored to energy requirements and activity levels safeguard health and directly influence task performance (FEDIAF, 2020; NRC, 2011). Failure to provide adequate exercise, social interaction, and cognitive enrichment can precipitate behavioral problems that degrade both performance and quality of life (Rooney et al., 2016; Sinn et al., 2010).

This study proceeds from the premise that meeting the care, nutritional, and behavioral needs of working dogs is both an ethical responsibility and a strategic prerequisite for mission success. Our primary objective is to comprehensively evaluate living conditions under these three headings; to analyze current practices in military and law enforcement contexts; and to formulate improvement recommendations grounded in empirical evidence. The ultimate goal is to maximize animal welfare while safeguarding continuity of service.

2. Background and General Information

2.1. Institutional Structure and Training of Working Dogs in Turkey

Although the military use of dogs' dates back to antiquity, modern and systematic application accelerated in the nineteenth century. In Turkey, institutionalization gathered momentum in the late twentieth century. Within the Gendarmerie General Command, the Gendarmerie Horse and Dog Training Center Command (JAKEM), and within the Turkish National Police, the Dog Training Center (KEM) are specialized institutions responsible for breeding, training, and deployment.

Located in Nevşehir, JAKEM maintains its own breeding program from pedigreed dams and sires—primarily Belgian Malinois, German Shepherd Dogs, and Labradors. Puppies are raised from birth in high-biosecurity facilities with environmental control (whelping and socialization units). After the critical first 8 weeks and primary immunizations, puppies enter a juvenile program (~6 months) covering basic obedience, focus, socialization, and environmental habituation. Candidates who pass standardized tests are assigned to one of twelve specialty branches (e.g., explosives detection, narcotics, tracking, search-and-rescue) and transferred to "Course Battalion" units, where they are paired with an assigned trainer/handler. Specialty training typically lasts 3–12 months, after which the team is certified and deployed.

2.2. Key Determinants of Welfare

Sustained performance requires adherence to scientific standards in living conditions.

Housing. Kennel facilities should be sited away from noise and environmental stressors and equipped with effective drainage. For movement and exercise, facilities should provide ample turnout space; semi-open designs are encouraged to better satisfy behavioral needs. Thermal comfort is critical: recommended ambient temperature is ~18.3–29.4 °C for adults and 30–32 °C during the first week for neonates (NRC, 2011; MSD, 2025). Given circadian influences on physiology and behavior, 8–12 hours of light per day is recommended. Because canine hearing is sensitive across a wide range (up to ~60 kHz), high-frequency noise from electronics and sudden loud sounds may constitute chronic stressors and should be mitigated through acoustic design and equipment choice. Rigorous hygiene and regular disinfection are essential.

Nutrition. While dogs exhibit adaptations to omnivory, they retain carnivore traits. Colostrum within the first 24 hours is vital for neonates. During growth, 2–3 meals/day are recommended; one meal/day may suffice for healthy adults. Energy and protein requirements vary with age, activity, and physiologic

FULL TEXTS BOOK

status; a typical adult diet should contain ~18% crude protein (DM) and ~3,500 kcal/kg metabolizable energy (FEDIAF, 2020; NRC, 2011). Dietary fat provides essential fatty acids (notably linoleic acid) and supports skin/coat health. Fresh water must be continuously available and checked multiple times daily.

Behavior. Success hinges on strong trainer–dog bonds and appropriate early socialization. Operational demands (elevated arousal, exposure to gunfire/explosions, cognitive load) can elicit fear and anxiety if unmanaged. Graduated exposure, desensitization to stressors, scheduled rest and play, and access to expert behavioral support are recommended. Programs should be individualized by age, sex, temperament, learning rate, and motivation to maximize outcomes (Wallis et al., 2019).

3. Materials and Methods

3.1. Study Design

A descriptive (cross-sectional) design was adopted to depict current living conditions and analyze the views of personnel working directly with dogs.

3.2. Population and Sample

The study population comprised personnel of JAKEM, the specialized unit responsible for breeding and training. Participation was voluntary. A total of **200** personnel working directly with dogs provided complete responses and constituted the final sample.

3.3. Instrument

A researcher-developed questionnaire, informed by literature, expert feedback, and pilot testing, ensured content validity. The 28-item survey combined closed-ended demographic items, Likert-type scales capturing attitudes and practices, and open-ended prompts. Thematic sections included: (i) demographics; (ii) number/type of dogs handled; (iii) assessment of housing; (iv) views on nutrition and health practices; (v) observed behavior and stress indicators; (vi) training and performance perceptions; and (vii) overall evaluation and improvement suggestions.

3.4. Procedure and Ethics

Data collection spanned December 2024–January 2025. Following official authorization from JAKEM management, instruments were distributed both in print and via Google Forms. Participants received written information on aims, scope, and confidentiality; informed consent was obtained. Surveys were anonymous. The study complied with institutional ethics (waiver from Erciyes University Social Sciences Ethics Committee).

3.5. Data Analysis

Analyses were conducted in Microsoft Excel and **JMP v16.0**. Descriptive statistics (frequencies and percentages) summarized variables. Cross-tabulations explored bivariate associations. A cumulative logit (ordinal) regression model assessed effects of ordered predictors (e.g., age, experience, responsibility level) on Likert outcomes. Open-ended responses underwent content analysis to extract salient themes.

3.6. Validity and Reliability

A pilot with 10 JAKEM personnel assessed clarity and validity; revisions followed. Internal consistency reliability yielded **Cronbach's alpha = 0.81**, indicating high reliability (Nunnally & Bernstein, 1994).

4. Results

4.1. Demographics: A Young yet Experienced Profile

All 200 respondents were male, precluding sex-based comparisons. Age distribution skewed toward young and mid-career: 75% were 26–35 years; the largest stratum was 31–35 years (45%). Unit

FULL TEXTS BOOK

assignment was dominated by Bomb Disposal (44%) and Narcotics (23.5%), indicating that perceptions largely reflect high-risk operational profiles.

Career progression patterns emerged: - Young personnel (21–30 years): predominantly in physically demanding Bomb Disposal and Commando units.

- Mid-career personnel (31–40 years): more frequent in Commando and specialized Narcotics roles requiring experience and coordination.

- Senior personnel (≥41 years): shifted toward Narcotics, suggesting a trajectory from physically intensive tasks to intelligence/analysis-heavy roles.

Ordinal modeling confirmed that years of experience working with dogs increased significantly with age. A near-deterministic alignment ("organizational determinism") between unit and specialty was observed ($p < 0.0001$), implying limited rotation and cross-training. As trainer experience rose, the number of dogs under responsibility also increased; 3–5 years emerged as a threshold for transitioning from single-dog handling to multi-dog management.

4.2. Breed Composition: Malinois Dominance and Risk

Among surveyed dogs, 68% were Belgian Malinois and 18% German Shepherd Dogs; together they comprised 86% of the population. While Malinois dominance reflects desirable working traits (agility, learning capacity, drive), such concentration heightens long-term risks of genetic narrowing, breed-specific disorders, and vulnerability to population-level shocks.

4.3. Housing: Lack of Standardization and Priority Targets for Improvement

Perceptions of housing adequacy were polarized: 54.5% rated kennels "adequate/very adequate," whereas 45.5% indicated varying degrees of inadequacy. Key associations included: - Stress × Housing Quality. A significant relationship linked poorer kennel ratings to higher reported stress ($p = 0.0336$). In "very adequate" kennels, high stress was not reported; in kennels rated "not at all adequate," 25% of dogs showed high/very high stress. - Unit × Housing Quality. Alarmingly, 60% of kennels rated "not at all adequate" were in Bomb Disposal, signaling urgent need in a high-risk mission profile. - Dog Load × Housing Quality. Greater numbers of dogs per handler coincided with poorer housing perceptions, suggesting density-related quality issues. - Age/Breed × Housing. Young Malinois and older Labradors were more likely to be housed in lower-quality facilities.

4.4. Nutrition, Health, and Performance

Nutrition/Health. Nearly all respondents (99%) reported regular feeding programs; satisfaction with diet quality was high (91% adequate/very adequate).

Behavior/Training. Overall stress was most frequently "moderate" (50%) or "low" (29.5%). The chief training challenge was attention/focus problems (32%). More than half (58%) believed their dogs required additional training.

Operational Performance. Distractibility was observed "occasionally" (45%) or "rarely" (43.5%). Aggression rates were low (~70% none/rare).

4.5. Qualitative Insights from Open-Ended Responses

Prominent themes included: - Kennel Infrastructure. Concerns about bare concrete flooring; requests for bedding and improved thermal/acoustic insulation.

- Training Realism. Predominance of open-field sites was seen as misaligned with actual missions in urban/complex structures; respondents advocated for scenario-based environments with diverse obstacles and interiors.

FULL TEXTS BOOK

- Equipment and Materials. Reports of outdated gear, lack of calibration for odor training materials, and insufficient variety of reward items tailored to individual motivation.

- Rest and Stress Management. Continuous training-task cycles were viewed as demotivating; weekly full rest days and "leave/holiday" allowances for dogs were recommended.

- Individualization. "One-size-fits-all" approaches were criticized; respondents emphasized tailoring by learning speed, temperament, and motivators to improve outcomes.

5. Discussion and Conclusions

5.1. Demographics, Career Planning, and Study Limitations

The predominance of 26–35-year-olds aligns with physical and cognitive demands of the work. Observed transitions—from Bomb Disposal/Commando in early career toward Narcotics at later stages—appear to balance physical wear with experience-driven tasks. However, under-representation of ≥40-year-olds limits statistical power for that stratum, and critical yet under-sampled units may face long-term capacity risks.

5.2. Housing: Evidence Base and Imperatives for Improvement

Negative kennel assessments echo the literature: spatial/social restriction and impoverished environments exacerbate chronic stress, impair learning, and provoke behavior problems (Beerda et al., 1999; Coppola et al., 2006). Kennel design must address both physical comfort and environmental stimuli. Given canine sensitivity to high-frequency noise (Morgan & Tromborg, 2007), facilities should incorporate acoustic control, natural light cycles, and ample activity areas. These measures are central to welfare and, by extension, performance.

5.3. Nutrition and Training: The Case for Individualization

While feeding programs were generally regular, flexibility to align with workload and individual needs was limited. Unmet energy demands during intensive periods can degrade motivation and attention. Nutritional plans and training schedules should be dynamically individualized. Behavioral issues such as distractibility and command noncompliance are often linked to punitive or non-individualized methods; contemporary approaches centered on positive reinforcement and gamification improve learning speed and motivation. Early, structured socialization remains the most effective prevention for later fear/aggression (Haverbeke et al., 2008).

5.4. Sustainability Risk: Breed Diversity and Genetic Health

Dependence on two breeds for **86%** of the population elevates risks associated with reduced genetic diversity and breed-specific disorders (e.g., hip dysplasia). Strategic mitigation includes maintaining a core stock of alternative breeds (e.g., Labradors, Pointers) and developing branch-specific training protocols tailored to their strengths.

5.5. Recommendation: A Data-Driven, Proactive Audit and Monitoring System

We propose transitioning from reactive fixes to a proactive system anchored in objective indicators.

Rationale. The significant association between stress level and adverse health effects ($p = 0.03$) supports using stress as an early-warning proxy for kennel quality.

Thresholds. Define "Alert" and "Critical" thresholds for three indices: stress score, adverse health events, and kennel inadequacy. Exceedances trigger pre-specified responses.

Quality Cycle.

- Data Capture: Weekly trainer-completed stress/health forms; monthly veterinary and logistics audits of physical facilities.

- Analysis: Quarterly dashboards (e.g., Tableau/Power BI) flag threshold breaches in real time.

FULL TEXTS BOOK

- Intervention: For critical breaches, conduct root-cause analysis (Ishikawa diagram) and implement an Action Plan (e.g., heaters within 1 week; flooring replacement within 1 month).

- Control: Re-assess indicators post-intervention to confirm normalization.

5.6. Overall Conclusion

This study demonstrates the necessity of managing working-dog welfare through systematic, measurable indicators rather than subjective observation alone. The proposed monitoring framework provides an early-warning mechanism that enables proactive interventions. Integrating welfare metrics into management strategies safeguards canine health across the service life and sustains peak operational effectiveness—aligning animal welfare and mission performance as mutually reinforcing objectives.

REFERENCES

Anadolu Ajansı. (2021, December 23). Operasyonların "gizli kahramanı" dedektör köpekler. https://www.aa.com.tr/tr/gundem/operasyonlarin-gizli-kahramani-dedektor-kopekler/2457732

Arendt, J. (1998). Melatonin and the pineal gland: Influence on mammalian seasonal and circadian physiology. *Reviews of Reproduction, 3*(1), 13–22. https://doi.org/10.1530/ror.0.0030013

Baran, M. S. (2012). Kedi ve Köpeklerin Beslenmesi. *İstanbul Üniversitesi Veteriner Fakültesi Dergisi, 33*(3), 89–99.

Batt, L., Batt, M., Baguley, J., & McGreevy, P. (2008). The effects of structured sessions for juvenile training and socialization on guide dog success and puppy-raiser participation. *Journal of Veterinary Behavior, 3*(5), 199–206. https://doi.org/10.1016/j.jveb.2008.05.001

Beerda, B., Schilder, M. B. H., Van Hooff, J. A. R. A. M., De Vries, H. W., & Mol, J. A. (1999). Chronic stress in dogs subjected to social and spatial restriction. I. Behavioral responses. *Physiology & Behavior, 66*(2), 233–242. https://doi.org/10.1016/S0031-9384(98)00289-3

Bennett, D., & Timm, R. M. (2016). The dogs of Roman Vindolanda, Part II: Time-stratigraphic occurrence, ethnographic comparisons, and biotype reconstruction. *Archaeofauna, 25*.

Bray, E. E., Otto, C. M., Udell, M. A. R., Hall, N. J., Johnston, A. M., & MacLean, E. L. (2021). Enhancing the selection and performance of working dogs. *Frontiers in Veterinary Science, 8*. https://doi.org/10.3389/fvets.2021.644431

Broom, D. M., & Johnson, K. G. (1993). *Stress and Animal Welfare*. Chapman & Hall. https://doi.org/10.1007/978-94-024-0980-2

Carciofi, A. C., Vasconcellos, R. S., Borges, N. C., Moro, J. V., Prada, F., & Fraga, V. O. (2006). Composição nutricional e avaliação de rótulo de rações secas para cães comercializadas em Jaboticabal-SP. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia, 58*(3), 421–426. https://doi.org/10.1590/S0102-09352006000300021

Case, L. P., & Daristotle, L. (2011). *Canine and Feline Nutrition* (3rd ed.). Elsevier.

Coppola, C. L., Grandin, T., & Enns, R. M. (2006). Human interaction and cortisol: Can human contact reduce stress for shelter dogs? *Physiology & Behavior, 87*(3), 537–541.

Cuevas, P. (2025). Are dogs carnivores or omnivores? What research suggests. *Dogster*. https://www.dogster.com/dog-nutrition/are-dogs-carnivores-or-omnivores

de Godoy, M. R. C., Kerr, K. R., & Fahey, G. C. (2013). Alternative dietary fiber sources in companion animal nutrition. *Nutrients, 5*(8), 3099–3117. https://doi.org/10.3390/nu5083099

English, P. B., & Filippich, L. J. (1980). Measurement of daily water intake in the dog. *Journal of Small Animal Practice, 21*(3), 189–193. https://doi.org/10.1111/j.1748-5827.1980.tb01231.x

FULL TEXTS BOOK

Evans, K. M., Lewis, T. W., Asher, L., Blythe, S., Bottomley, M., Tootill, L., Roberts, R., Whiteside, H., England, G. C. W., & Blott, S. C. (2015). Genetic evaluation of traits in a standardized behavioral test for potential guide dog puppies using crossbreed models. *Journal of Veterinary Behavior, 10*(6), 459–464. https://doi.org/10.1016/j.jveb.2015.08.004

FEDIAF. (2020). *Nutritional Guidelines for Complete and Complementary Pet Food for Cats and Dogs*. The European Pet Food Industry Federation.

Foyer, P., Wilsson, E., Wright, D., & Jensen, P. (2013). Early experiences modulate stress coping in a population of German Shepherd Dogs. *Applied Animal Behaviour Science, 146*(1–4), 79–87. https://doi.org/10.1016/j.applanim.2013.03.013

Glodde, F., Günal, M., Kinsel, M. E., & AbuGhazaleh, A. (2018). Effects of natural antioxidants on the stability of omega-3 fatty acids in dog food. *Journal of Veterinary Research, 62*(1), 103–108. https://doi.org/10.1515/jvetres-2018-0014

Graham, L., Wells, D. L., & Hepper, P. G. (2005). The influence of olfactory stimulation on the behaviour of dogs housed in a rescue shelter. *Applied Animal Behaviour Science, 91*(1–2), 143–153. https://doi.org/10.1016/j.applanim.2004.08.024

Grimm-Seyfarth, A., Harms, W., & Berger, A. (2021). Detection dogs in nature conservation: A database on their world-wide deployment with a review on breeds used and their performance compared to other methods. *Methods in Ecology and Evolution, 12*(4), 568–579. https://doi.org/10.1111/2041-210X.13560

Hall, N. J., Johnston, A. M., Bray, E. E., Otto, C. M., MacLean, E. L., & Udell, M. A. R. (2021). Working dog training for the twenty-first century. *Frontiers in Veterinary Science, 8*. https://doi.org/10.3389/fvets.2021.646022

Haverbeke, A., De Smet, A., Depiereux, E., Giffroy, J.-M., & Diederich, C. (2008). Assessing undesired aggression in military working dogs. *Applied Animal Behaviour Science, 113*(1–3), 287–300.

Helton, W. S. (Ed.). (2009). *Canine Ergonomics*. CRC Press. https://doi.org/10.1201/9781420079920

Hennessy, M. B., Williams, M. T., Miller, D. D., Douglas, C. W., & Voith, V. L. (1998). Influence of male and female petters on plasma cortisol and behaviour: Can human interaction reduce the stress of dogs in a public animal shelter? *Applied Animal Behaviour Science, 61*(1), 63–77.

Jandarma Genel Komutanlığı. (2025). Jandarma At ve Köpek Eğitim Merkezi Komutanlığı (JAKEM). https://www.jandarma.gov.tr/jandarma-at-kopek-egitim-merkezi

Jantorno, G. M., Xavier, C. H., & de Melo, C. B. (2020). Narcotic detection dogs: An overview of high-performance animals. *Ciencia Rural, 50*(10), 1–8. https://doi.org/10.1590/0103-8478cr20191010

Jarrett, C. L., Brathwaite, M., Gogal, R. M., & Holladay, S. D. (2022). Working dog service, harmful agent exposure and decontamination. *Frontiers in Veterinary Science, 9*. https://doi.org/10.3389/fvets.2022.892998

Kawami, T. S. (1986). Greek art and Persian taste: Some animal sculptures from Persepolis. *American Journal of Archaeology, 90*(3). https://doi.org/10.2307/505686

Lazarowski, L., Haney, P. S., Brock, J., Fischer, T., Rogers, B., Angle, C., Katz, J. S., & Waggoner, L. P. (2018). Investigation of behavioral characteristics of dogs' purpose-bred and prepared to perform Vapor Wake® detection of person-borne explosives. *Frontiers in Veterinary Science, 5*. https://doi.org/10.3389/fvets.2018.00050

Leroy, G. (2011). Genetic diversity, inbreeding and breeding practices in dogs: Results from pedigree analyses. *The Veterinary Journal, 189*(2), 177–182.

FULL TEXTS BOOK

Li, P., & Wu, G. (2024). Characteristics of nutrition and metabolism in dogs and cats. In G. Wu (Ed.), *Advances in Experimental Medicine and Biology* (Vol. 1446, pp. 55–98). Springer. https://doi.org/10.1007/978-3-031-54192-6_4

McEvoy, V., Espinosa, U., Crump, A., & Arnott, G. (2022). Canine socialisation: A narrative systematic review. *Animals, 12*(21), 2895. https://doi.org/10.3390/ani12212895

Mellor, D. J., Beausoleil, N. J., Littlewood, K. E., McLean, A. N., McGreevy, P. D., Jones, B., & Wilkins, C. (2020). The 2020 five domains model: Including human–animal interactions in assessments of animal welfare. *Animals, 10*(10), 1870.

Menchetti, L., Guelfi, G., Speranza, R., Carotenuto, P., Moscati, L., & Diverio, S. (2019). Benefits of dietary supplements on the physical fitness of German Shepherd Dogs during a drug detection training course. *PLOS ONE, 14*(6), e0218275. https://doi.org/10.1371/journal.pone.0218275

Morgan, K. N., & Tromborg, C. T. (2007). Sources of stress in captivity. *Applied Animal Behaviour Science, 102*(3–4), 262–302. https://doi.org/10.1016/j.applanim.2006.05.032

MSD Veterinary Manual. (2025). Environmental warmth required for neonatal dogs and cats. https://www.msdvetmanual.com/multimedia/table/environmental-warmth-required-for-neonatal-dogs-and-cats

National Research Council. (2011). *Guide for the Care and Use of Laboratory Animals* (8th ed.). The National Academies Press.

Nichiporuk, S., & Radzikhovskyi, M. (2022). Big hearts of little friends. *Scientific and Technical Bulletin of State Scientific Research Control Institute of Veterinary Medical Products and Fodder Additives and Institute of Animal Biology, 23*(1). https://doi.org/10.36359/scivp.2022-23-1.15

Nunnally, J. C., & Bernstein, I. H. (1994). *Psychometric Theory* (3rd ed.). McGraw-Hill.

O'Neill, D. G., Church, D. B., McGreevy, P. D., Thomson, P. C., & Brodbelt, D. C. (2017). Demography and disorders of German Shepherd Dogs under primary veterinary care in the UK. *Canine Genetics and Epidemiology, 4*, 7.

Ottesen, J. L., Weber, A., Gürtler, H., & Mikkelsen, L. F. (2004). New housing conditions: Improving the welfare of experimental animals. *Alternatives to Laboratory Animals, 32*(Suppl. 1), 397–404. https://doi.org/10.1177/026119290403201s65

Öztürk, Y. (2024). Askerî eğitim tarihi açısından harp köpeklerinin talim ve terbiyesi. *Askeri Tarih Araştırmaları Dergisi, 22*(36), 193–210. https://doi.org/10.46953/askeritarih.1497836

Rehn, T., Keeling, L. J., & Marshall-Pescini, S. (2014). Evaluating the quality of the human–dog relationship by measuring dogs' motivation to interact with a familiar person and strangers. *Behavioural Processes, 103*, 180–185.

Rooney, N. J., Clark, C. C. A., & Casey, R. A. (2016). Minimizing fear and anxiety in working dogs: A review. *Journal of Veterinary Behavior: Clinical Applications and Research, 16*, 53–64. https://doi.org/10.1016/j.jveb.2016.11.001

RSPCA. (2025). *Guidelines for the Design and Management of Animal Shelters* (5th ed.). RSPCA International. https://studylib.net/doc/8763413/2.-guidelines-for-the-design-and-management-of-animal-she

Rutter, N. J., Howell, T. J., Stukas, A. A., Pascoe, J. H., & Bennett, P. C. (2021). Can volunteers train their pet dogs to detect a novel odor in a controlled environment in under 12 weeks? *Journal of Veterinary Behavior, 43*, 54–65. https://doi.org/10.1016/j.jveb.2020.09.004

Schäfer, W., & Hankel, J. (2020). Energy consumption of young military working dogs in pre-training in Germany. *Animals, 10*(10), 1–10. https://doi.org/10.3390/ani10101753

Scholander, P. F., Hock, R., Walters, V., Johnson, F., & Irving, L. (1950). Heat regulation in some arctic and tropical mammals and birds. *The Biological Bulletin, 99*(2), 237–258. https://doi.org/10.2307/1538741

Sebnem Ozcan, S., Akin, H., Bayram, H., Bas, M., Yildiz, A., & Ozdemiroglu, A. (2009). Utilization of police dogs: A Turkish perspective. *Policing, 32*(2), 226–237. https://doi.org/10.1108/13639510910958154

Sinn, D. L., Gosling, S. D., & Hilliard, S. (2010). Personality and performance in military working dogs: Reliability and predictive validity of behavioral tests. *Applied Animal Behaviour Science, 127*(1–2), 51–65. https://doi.org/10.1016/j.applanim.2010.08.007

Spangenberg, E. M. F., Björklund, L., & Dahlborn, K. (2006). Outdoor housing of laboratory dogs: Effects on activity, behaviour and physiology. *Applied Animal Behaviour Science, 98*(3–4), 260–276. https://doi.org/10.1016/j.applanim.2005.09.009

Staples, L. G. (2010). Predator odor avoidance as a rodent model of anxiety: Learning-mediated consequences beyond the initial exposure. *Neurobiology of Learning and Memory, 94*(4), 435–445. https://doi.org/10.1016/j.nlm.2010.09.009

T.C. İçişleri Bakanlığı, Jandarma Genel Komutanlığı. (2025, July 30). JAKEM – Tarihçe. https://www.jandarma.gov.tr/jakem/tarihce#

TBMM. (2025). Resmî Gazete. https://www.resmigazete.gov.tr/eskiler/2025/05/20250525-1.htm

Wallis, L. J., Szabó, D., & Kubinyi, E. (2019). Cross-sectional age differences in canine personality traits: Influence of breed, sex, previous trauma, and dog obedience tasks. *Frontiers in Veterinary Science, 6*, 493. https://doi.org/10.3389/fvets.2019.00493

Wells, D. L. (2004). A review of environmental enrichment for kenneled dogs, *Canis familiaris*. *Applied Animal Behaviour Science, 85*(3–4), 307–317. https://doi.org/10.1016/j.applanim.2003.11.005

Wells, D. L., & Hepper, P. G. (2000). The influence of environmental change on the behaviour of sheltered dogs. *Applied Animal Behaviour Science, 68*(2), 151–162. https://doi.org/10.1016/S0168-1591(00)00100-3

Wojtaś, J., Karpiński, M., & Czyżowski, P. (2020). Salivary cortisol interactions in search and rescue dogs and their handlers. *Animals, 10*(4). https://doi.org/10.3390/ani10040595

Yeon, S. C., Golden, G., Sung, W., Erb, H. N., Reynolds, A. J., & Houpt, K. A. (2001). A comparison of tethering and pen confinement of dogs. *Journal of Applied Animal Welfare Science, 4*(4), 257–270. https://doi.org/10.1207/S15327604JAWS0404_03

ALVEOLAR DEFEKTİN TİTANYUM MESH İLE ONARIMI VE EŞ ZAMANLI OLARAK DENTAL İMPLANT YERLEŞTİRİLMESİ: BİR VAKA RAPORU

REPAIR OF ALVEOLAR DEFECT WITH TITANIUM MESH AND SIMULTANEOUS DENTAL IMPLANT PLACEMENT: A CASE REPORT

Zeynep GÜMRÜKÇÜ

Doçent Doktor, Recep Tayyip Erdoğan Üniversitesi, Diş Hekimliği Fakültesi, Ağız Diş ve Çene Cerrahisi Anabilim Dalı, Rize, Türkiye

ORCID ID: https://orcid.org/0000-0002-5910-2218

ÖZET

Modern diş hekimliğinde dental implantlar, fonksiyon ve estetiği sağlayarak dişsizliğin giderilmesinde giderek daha sık tercih edilen bir tedavi seçeneği haline gelmiştir. Ancak dental implant uygulamaları sağlıklı bir alveol kemiği ve dişeti varlığında başarılı olabilmektedir. Endodontik ve periodontal enfeksiyonlar gibi nedenlerle alveol krette oluşan kemik kayıpları implant cerrahisine ek olarak farklı kemik augmentasyon süreçlerini zorunlu kılar. Endodontik enfeksiyon nedeniyle sağ maksiller 1. kesici dişini kaybeden 21 yaşında hastanın klinik ve radyolojik muayenesinde ilgili bölgedeki alveol kemiğinin horizontal yönde yetersiz olduğu tespit edilmiştir. Hastaya implant yerleştirilmesine eş zamanlı olarak titanyum mesh ve allojen toz greft materyali ile kemik augmetasyonu uygulanmıştır. İşlem sonrası 6. ayda yapılan klinik ve radyolojik incelemelerde kemik iyileşmesinin tamamlandığı ve implantın stabil olduğu gözlenmiş ve ardından hastanın protetik restorasyonu tamamlanmıştır. Alveol kemiğin yetersizliği dental implanta yönelik kemik arttırma prosedürlerine olan ihtiyacı doğurmaktadır. Ancak ramus/simfiz blok greft gibi alveol kretteki horizontal kemik yetersizliklerini gidermeye yönelik işlemler artmış komplikasyonlar, post operatif artmış ağrı/ödem ve dönor saha morbiditesi gibi dezavantajlara sahiptir. Titanyum yüksek biyouyumluluğu ve mekanik direnci ile oral cerrahide farklı şekillerde kullanılmakta olan bir materyaldir. Bu çalışmada alveolar defekt onarımı ve eş zamanlı olarak dental implant yerleştirilmesi ile estetik ve fonksiyon başarılı bir şekilde sağlanmıştır. Mekanik bir bariyer görevi gören titanyum mesh kullanılarak yapılan augmentasyon prosedürleri; daha kısa cerrahi süresi, daha az post operatif komplikasyon ve dönor saha morbiditesi olmaması gibi avantajlara sahip olması nedeniyle oral cerrahide farklı greftleme prosedürleri için bir tedavi seçeneği olarak öne çıkmaktadır.

Anahtar Kelimeler: Dental İmplant, Kemik Augmentasyonu, Titanyum Mesh, Allogreft

ABSTRACT

In modern dentistry, dental implants have become an increasingly preferred treatment option for addressing edentulism by restoring both function and aesthetics. However, the success of dental implant procedures relies heavily on the presence of healthy alveolar bone and gingival tissue. Bone loss in the alveolar crest due to factors such as endodontic or periodontal infections necessitates additional bone augmentation procedures alongside implant surgery. In this case report, a 21-year-old patient who had lost the right maxillary central incisor due to endodontic infection was found to have insufficient horizontal alveolar bone upon clinical and radiographic examination. Simultaneous with dental implant placement, bone augmentation was performed using a titanium mesh and allogenic particulate graft material. At the 6-month follow-up, clinical and radiographic evaluations demonstrated complete bone healing and implant stability, after which the prosthetic restoration was successfully completed.

FULL TEXTS BOOK

Insufficient alveolar bone necessitates augmentation procedures to enable implant placement. However, techniques such as ramus/symphysis block grafts used to manage horizontal bone deficiencies of the alveolar crest may be associated with increased complications, heightened postoperative pain/edema, and donor site morbidity. Titanium, owing to its high biocompatibility and mechanical strength, is widely utilized in various oral surgical procedures. In this study, functional and aesthetic success was achieved through alveolar defect reconstruction and simultaneous implant placement. Augmentation procedures utilizing titanium mesh as a mechanical barrier have emerged as a viable treatment option in oral surgery due to their advantages, including shorter surgical time, fewer postoperative complications, and the absence of donor site morbidity.

Keywords: Dental Implant, Bone Augmentation, Titanium Mesh, Allograft

GİRİŞ

Branemark'ın osseointegrasyon terimini ortaya atmasının ardından modern diş hekimliğinde dental implantlar, fonksiyon ve estetiği sağlayarak dişsizliğin giderilmesinde giderek daha sık tercih edilen bir tedavi seçeneği haline gelmiştir.[1,2] İlerleyen yıllarda dental implantların geometrisi ve yüzey modifikasyonları gibi konularda yapılan bir çok çalışma ile dental implantolojide başarı oranı yükselmiştir.[3-5]Ancak hastanın yaşı, cinsiyeti, sistemik hastalıkları, zararlı alışkanlıkları ve kemik yoğunluğu/hacmi gibi faktörler dental implantların başarısını etkilemektedir.[6,7]

Alveolar kemik atrofisi, dental implantlarla yapılan oral rehabilitasyonlarda en sık karşılaşılan sorunlarından biridir.[8] Modern oral cerrahide yönlendirilmiş kemik rejenerasyonu (YKR), alveolar distraksiyon osteogenezisi (ADO) ve otojen kemik grefti (OKG) gibi farklı kemik augmentasyon prosedürleri ile alveol kemiğinin rejenerayonu sağlanabilmektedir.[9,10] Ancak özelikle ADO ve OKG gibi kemik augmentasyon prosedürlerinde dönor saha morbiditesi ve dolayısıyla artmış post operatif ödem, enfeksiyon ve hematom gibi komplikasyonlar sıkça görülmektedir.[11,12]


Son yıllarda titanyum mesh yüksek mekanik özellikleri ve mükemmel biyouyumluluğu sayesinde YKR amacıyla kullanımı artan bir materyaldir.[13,14] Donör saha morbiditesi olmaması, hem yatay hem de dikey yönde kemik kazanımı sağlaması titanyum meshi (TM) tercih edilebilir bir materyal olarak öne çıkarmaktadır.[15,16]

Bu çalışmada endodontik enfeksiyon nedenyinle gelişen sağ maksiller santral diş eksikliğinin dental implant ve titanyum mesh ile eş zamanlı rekonstrüksiyonu sunulmaktadır.

VAKA RAPORU

Kliniğimize diş eksikliği şikayeti ile başvuran 21 yaşındaki erkek hastanın sağ maksiller santral dişini eksik olduğu tespit edilmiş ve ilgili dişe dental implant tedavisi planlanmıştır. Radyolojik inceleme amacıyla alınan konik ışınlı bilgisayarlı tomografi (KIBT) görüntüsünde 11 numaralı diş bölgesinde alveolar kemiğin horizontal yönde yetersiz olduğu tespit edilmiştir. **(Figür 1a, b, c, d)** Bu nedenle hastaya titanyum mesh kullanılarak YDR ve eş zamanlı implant yerleştirilmesi planlamıştır.


FULL TEXTS BOOK




a.Pre-op. OPG görüntüsü



b.Pre-op periapikal görüntü



c.Pre-op. BT kesitinde defekt görüntüsü



d. Pre-op BT defektin üç boyutlu kesiti

Figür 1. Preopreatif ragyografik değerlendirmeler

Bu amaçla lokal anestezi altında tam kalınlıklı flep kaldırılmış ve alveol kemiğine erişim sağlanmıştır. Ardından planlanan 3.7 çapında 11.5 mm ve uzunluğunda dental implant (JD Evolution, Modena, İtayla) yerleştirilmesi amacıyla frezleme protokolü uygulanmış ve implant 30 Nm tork değeri ile implant yerleştirilmiştir.

İmplantın bukkalinde kalan defektli alanın augmentasyonu amacıyla fabrikasyon titanyum mesh, 7x12 mm boyutlarında kesilip titanyum mikrovidalar ile (2 x 7 mm boyutlarında) implant ile arasında 2 mm boşluk kalacak şekilde sabitlenmiştir. **(Figür 2a)** İmplant ile titanyum mesh arasındaki boşluk allojen kemik grefti ile doldurulduu ve YDR amacıyla kollajen membran ile titanyum meshin üzeri örtülmüştür. **(Figür 2b)** Flep primer kapamaya izin verecek şekilde serbestleştirildi. Titanyum meshin olası bir ekspozürünün önüne geçmek amacıyla hastanın otolog kanından elde edilen konsantre büyüme faktör flep altına yerleştirildikten sonra 4.0 ipek sütur ile primer kapama sağlandı. **(Figür 2c, d)**

FULL TEXTS BOOK



a.İntra-op. titanyum mesh yerleştirme görüntüsü



b.Kollajen membran yerleştirme aşaması



c. CGF yerleştirilme aşaması




d. Cerrahi alanın süturla primer kapatılması

Figür 2. Cerrahi aşamalar

Hastaya post operatif 1000 mg amoksisilin+klavunat, 25 mg Deksketoprofen ve Klorheksidin Glukonat, Benzidamin Hidroklorür içerikli gargara reçete edildi. Cerrahi sonrası 10. günde yaranın tamamen iyileştiği izlendi ve dikişler alındı.

Hastadan 6.ayda alınan panoramik röntgende kemikleşmenin sağlandığı gözlendi. Ardından titanyum mesh söküldü ve implanta iyileşme başlığı yerleştirildi. Takip eden 25. günde hastanın tam seramik protetik restorasyonu tamamlandı. Hastanın 1.yıllık takibinde radografik olarak implant etrafında herhangi bir kemik kaybı olmadığı ve klinik olarak peri implantitis bulgusu olmadığı gözlemlenmiştir. **(Figür 3a, b, c, d)**

FULL TEXTS BOOK



a.Post-op. 1. yıl OPG görüntüsü



b.Post-op.1. yıl periapikal görüntüsü



c. Üst yapı protez sonrası ağız içi görüntüsü



d. Üst yapı protez sonrası gülümseme

Figür 3. Postoperatif radyografik ve klinik görünüm

TARTIŞMA

Kemik hacmi, yüksekliği ve genişliği, dental implant stabilitesi ve başarısı için önemli bir rol oynar.[2] Alveol kemiğindeki yetersizlikler çeşitli kemik grefti prosedürleriyle giderilebilir.[12] Horizontal yöndeki kemik eksikliklerinin giderilmesine yönelik en sık kullanılan augmentasyon prosedürlerinden biri otojen blok greft uygulamalarıdır.[17] Ancak otojen blok greft uygulamaları artmış komplikasyonlar, post operatif artmış ağrı/ödem ve dönor saha morbiditesi gibi dezavantajlara sahiptir.[11,17]

Titanyum yüksek biyouyumluluğu ve mekanik direnci ile oral cerrahide farklı amaçlarla kullanılmakta olan bir materyaldir.[18] Titanyumdan hazırlamış plaka şeklindeki meshlerin YKR amacıyla bariyer olarak kullanıldığı birçok çalışma mevcuttur.[13-15,19]

Faus ve ark. yaptıkları çalışmada titanyum mesh ile hem vertikal hem de horizontal yönde kemik kazanımının mümkün olduğunu ortaya koymuştur.[20] Benzer şekilde Geletu ve ark. çalışmalarında titanyum mesh ile yerleştirilen implantlarla estetik ve fonksiyonun sağlanabileceğini göstermişlerdir.[21] Bizim çalışmamızda da uygun kemik kazanımının, estetiğin ve fonksiyonun sağlanması yapılan çalışmalarla paralellik göstermektedir.

Çalışmalar titanyum meshlerin en sık komplikasyonu olarak meshin ağız içine ekspozürünü bildirmektedir.[19,22] Hartmann ve ark. bu komplikasyonun doğru yumuşak doku yönetimi ve otolog kan

FULL TEXTS BOOK

konsantrelerinin(plateletten zengin fibrin, konsantre büyüme faktörü vb.) kullanımı ile minimalize edilebileceğini ifade etmiştir.[23] Bizim çalışmamızda da konsantre büyüme faktörü kullanılmış ve ekspozür olmaksızın post operatif iyileşme sağlanmıştır.

Mekanik bir bariyer görevi gören titanyum mesh kullanılarak yapılan augmentasyon prosedürleri; daha kısa cerrahi süresi, daha az post operatif komplikasyon ve dönor saha morbiditesi olmaması gibi avantajlara sahip olması nedeniyle oral cerrahide farklı greftleme prosedürleri için bir tedavi seçeneği olarak öne çıkmaktadır.

KAYNAKÇA

1. Zohrabian VM, Sonick M, Hwang D, Abrahams JJ. Dental implants. Paper presented at: Seminars in Ultrasound, CT and MRI 2015.

2. Liaw K, Delfini RH, Abrahams JJ. Dental implant complications. Paper presented at: Seminars in Ultrasound, CT and MRI2015.

3. Kreve S, Ferreira I, da Costa Valente ML, Dos Reis AC. Relationship between dental implant macro-design and osseointegration: A systematic review. *Oral and Maxillofacial Surgery.* 2024;28(1):1-14.

4. Heimes D, Becker P, Pabst A, et al. How does dental implant macrogeometry affect primary implant stability? A narrative review. *International journal of implant dentistry.* 2023;9(1):20.

5. Sivaswamy V, Bahl V. Surface modifications of commercial dental implant systems: an overview. *Journal of Long-Term Effects of Medical Implants.* 2023;33(2).

6. Thiebot N, Hamdani A, Blanchet F, et al. Implant failure rate and the prevalence of associated risk factors: a 6-year retrospective observational survey. *Journal of Oral Medicine and Oral Surgery.* 2022;28(2):19.

7. Lekholm U. Patient selection and preparation. *Tissue-integrated prosthesis, osseointegration in clinical dentistry.* 1985.

8. Yun K-I, Choi H, Wright RF, Ahn HS, Chang BM, Kim HJ. Efficacy of Alveolar Vertical Distraction Osteogenesis and Autogenous Bone Grafting for Dental Implants: Systematic Review and Meta-Analysis. *International Journal of Oral & Maxillofacial Implants.* 2016;31(1).

9. Rocchietta I, Fontana F, Simion M. Clinical outcomes of vertical bone augmentation to enable dental implant placement: a systematic review. *Journal of clinical periodontology.* 2008;35:203-215.

10. Wang H-L, Boyapati L. "PASS" principles for predictable bone regeneration. *Implant dentistry.* 2006;15(1):8-17.

11. Sanz-Sánchez I, Sanz-Martín I, Ortiz-Vigón A, Molina A, Sanz M. Complications in bone-grafting procedures: classification and management. *Periodontology 2000.* 2022;88(1):86-102.

12. Iancu SA, Referendaru D, Iancu I-A, Bechir A, Barbu HM. Immediate postoperative complications after lateral ridge augmentation–a clinical comparison between bone shell technique and sticky bone. *Journal of Medicine and Life.* 2022;15(4):533.

13. Li S, Zhao J, Xie Y, Tian T, Zhang T, Cai X. Hard tissue stability after guided bone regeneration: a comparison between digital titanium mesh and resorbable membrane. *International Journal of Oral Science.* 2021;13(1):37.

14. Trento G, Carvalho P, Macedo D, Gabrielli M, Monnazzi M, Pereira-Filho V. Titanium mesh associated with rhBMP-2 in alveolar ridge reconstruction. *International journal of oral and maxillofacial surgery.* 2019;48(4):546-553.

FULL TEXTS BOOK

15. Cucchi A, Vignudelli E, Franceschi D, et al. Vertical and horizontal ridge augmentation using customized CAD/CAM titanium mesh with versus without resorbable membranes. A randomized clinical trial. *Clinical Oral Implants Research.* 2021;32(12):1411-1424.

16. Zhang T, Zhang T, Cai X. The application of a newly designed L-shaped titanium mesh for GBR with simultaneous implant placement in the esthetic zone: a retrospective case series study. *Clinical implant dentistry and related research.* 2019;21(5):862-872.

17. Smeets R, Matthies L, Windisch P, et al. Horizontal augmentation techniques in the mandible: a systematic review. *International journal of implant dentistry.* 2022;8(1):23.

18. Elgali I, Omar O, Dahlin C, Thomsen P. Guided bone regeneration: materials and biological mechanisms revisited. *European journal of oral sciences.* 2017;125(5):315-337.

19. Zhou L, Su Y, Wang J, Wang X, Liu Q, Wang J. Effect of exposure rates with customized versus conventional titanium mesh on guided bone regeneration: Systematic review and meta-analysis. *Journal of Oral Implantology.* 2022;48(4):339-346.

20. Bertran Faus A, Cordero Bayo J, Velasco-Ortega E, et al. Customized titanium mesh for guided bone regeneration with autologous bone and xenograft. *Materials.* 2022;15(18):6271.

21. Geleţu GL, Burlacu A, Murariu A, et al. Customized 3D-printed titanium mesh developed for an aesthetic zone to regenerate a complex bone defect resulting after a deficient odontectomy: a case report. *Medicina.* 2022;58(9):1192.

22. Sabri H, Heck T, Manouchehri N, Alhachache S, Calatrava J, Misch CM. Bone augmentation using titanium mesh: A systematic review and meta-analysis. *International Journal of Oral Implantology.* 2024;17(3).

23. Hartmann A, Seiler M. Minimizing risk of customized titanium mesh exposures–a retrospective analysis. *BMC Oral Health.* 2020;20(1):36.

FORMATION OF THE FUTURE SPECIALIST'S CREATIVITY
GELECEK UZMANIN KREATIVITESININ OLUŞTURULMASI

Assist. Prof. Dr. Aktolkyn BORANBAYEVA

Associate Professor at the Faculty of History and Pedagogy, South Kazakhstan Pedagogical University named after Uzbekali Zhanibekov, (Kazakhstan, Shymkent)

ORCID ID: 0000-0001-6256-6999

Prof. Dr. Gelişli, Yücel

Doctor Professor of the Faculty of Education, Gazi University (Ankara, Türkiye)

ORCID ID: 0000-0003-2816-3621

Prof. Dr. Assel TASSOVA

Head of the Department of Pedagogy and Psychology, Faculty of Social and Humanitarian Sciences, Khoja Akhmet Yassawi International Kazakh-Turkish University (Kazakhstan, Turkestan)

ORCID ID: 0000-0001-6476-7874

ABSTRACT

The article describes the research of the concept of "creative" and its origins. In addition, on the basis of scientific sources, the definitions and concepts given by domestic and foreign authors to the concepts of "creativ" and "creativity"are considered. Reveals six parameters of creativity that contribute to the formation of a creative personality. It is noted that today, in accordance with the educational flows, the ways of formation of future specialists as creative individuals and the development of creative qualities are outlined. Creativity is the creative ability of a person, the ability to think and make decisions, to create new works. The effective implementation of these tasks requires from the modern education system the formation of professional potential, the future of the intellectual nation, striving for a fully integrated knowledge, rapid adaptation to daily changes, able to independently act creatively. In accordance with this, modern education in the Republic of Kazakhstan means the need to build in accordance with the competitive requirements of the leading countries of the world and the strategic direction of the model of innovative development. A person who is actively engaged in his creative activity and has achieved high results can be called a creative person. And the formation of professional creativity of future specialists is the main task of modern education.

Keywords: creativity, success, creative personality, future specialist, creative process.

ÖZET

Bu makalede “yaratıcılık” kavramı ve onun kökenleri üzerine yapılan araştırmalar ele alınmıştır. Ayrıca, bilimsel kaynaklar temelinde, yerli ve yabancı yazarlar tarafından “yaratıcılık” ve “kreativite” kavramlarına verilen tanımlar ve yaklaşımlar incelenmiştir. Makalede yaratıcı kişiliğin oluşumuna katkı sağlayan altı yaratıcılık parametresi açıklanmaktadır. Günümüzde eğitim akımları doğrultusunda, geleceğin uzmanlarının yaratıcı bireyler olarak şekillenmesi ve yaratıcı niteliklerinin geliştirilmesi yolları belirlenmektedir. Kreativite, bireyin yaratıcı düşünme, karar verme ve yeni eserler ortaya koyma yeteneğidir. Bu hedeflerin etkili bir şekilde gerçekleştirilmesi, günümüz eğitim sisteminden çok yönlü gelişmiş, bilgiye aç, günlük değişimlere hızla uyum sağlayabilen, bağımsız yaratıcı faaliyetlerde bulunabilen, yüksek mesleki potansiyele sahip gençlerin – milletin entelektüel geleceğinin – yetiştirilmesini gerektirmektedir. Bu bağlamda, Kazakistan Cumhuriyeti'nde modern bir eğitim sistemi

FULL TEXTS BOOK

ve yenilikçi kalkınma modeli oluşturma ihtiyacı, dünyanın önde gelen ülkelerinin rekabetçilik ve stratejik kalkınma gereklilikleri doğrultusunda açıkça ortaya çıkmaktadır. Yaratıcı faaliyetlerle aktif olarak ilgilenen ve yüksek başarılar elde eden birey, yaratıcı bir kişilik olarak adlandırılabilir. Geleceğin uzmanlarında mesleki kreativitenin geliştirilmesi, modern eğitimin temel hedeflerinden biridir.

Anahtar Kelimeler: kreativite, yaratıcılık, başarı, yaratıcı birey, yaratıcılık süreci

Introduction

In the global educational space, it is necessary to form a professional potential, the future of an intellectual nation that is passionate about comprehensively integrated knowledge, quickly adapts to everyday changes, and is able to act creatively independently. Modern education in the Republic of Kazakhstan implies the need to build a model of innovative development in accordance with the competitive requirements of the leading countries of the world and the strategic direction of development. The education system should be focused on training full-fledged competitive specialists, enrich young people with cultural and spiritual values that delight not only education, but also its constant updating, replenishment, expansion of horizons, the desire to improve thinking.

After all, the market economy requires a" knowledgeable "educational goal to replace a" creatively thinking, self-active, self-developing communicative and creative person".

Therefore, today it is important not to provide students with as much knowledge as possible, but to ensure their general cultural, personal and cognitive development, improving the skills of mastering knowledge. To achieve the goal, our consciousness must be ahead of our work, that is, updated to it. This will not only complement the political and economic transformations, but also become their core.

One of them is the creativity of the future specialist. A "creative" is a new product or result of creative thinking.

The problem of creativity is reflected in the research of a number of foreign and domestic scientists. Fundamental research for the individual components of creativity, namely intellectual creativity were studied in the works of (Guilford J., Torrance E., Tunik E.E., Kholodnaya M.A. and others); social creativity (Manayevoch N.A., Chichuk E.Yu.); emotional creativity (Andreyeva I.N., Berezyna T.N., Terechenko R.N. JAverill.R. and others); linguistic creativity (Khalyushova G.A.), lexical creativity (Grydyna T.A.), pedagogical creativity (Morozov A.V., Chernelevsky D.V., Cherbakova E.E.), professional creativity (Ryabova E.V).

Domestic scientists Ospanova B. A., Szwajkowski A. S., Nagimzhanova K.M. studied the formation of creativity of future specialists, Nurgaliyeva A. Studied the usage of innovative technologies in the formation of creativity of students, Sagdullaev I. I. acme-creativity of future specialists, so Galiev T.G. development of creativity of students, Sabirov T. S. prepare in future teachers the organization of cognitive activity of students.

Thus, the analysis of domestic and foreign literature revealed that in Kazakhstan psychological and pedagogical research is not in the unity of the concepts of communicative creativity of the individual, which, firstly, secondly, investigated the comprehensive aspects of the problem of training future primary school teachers, but the question of the formation of their communicative creativity is not taken on a special research form.

FULL TEXTS BOOK

Methodological research.

In the field of science, there is a certain history of the problem of creativity. Most of the research work on this issue was carried out by foreign scientists (T. Amabile, F. Baron, J. Guilford, E. Torrens, E. Bono, et al.) [1, 2, 3, 4, 5].

In foreign psychology, "creativity" is used to refer to the ability, which manifests the property of an individual to create new concepts and form new skills.

The ability to find new ideas and solve problem situations is explained as creativity. A number of researchers associate creativity with personal characteristics (A. Maslow, K. Rogers) [6, 7], and if some researchers consider it as an intellectual function (D. B. Bogoyavlenskaya, J. Gilford) [8, 3], still others consider the creative process and the connection of creativity (A. N. Luk) [9].

In psychological science, creativity is considered as a creative ability, the ability of an individual to show social creative activity.

The concept of "Creative" was first considered and used in 1922 by D. Simpson as a refusal of a person from stereotypical ways of thinking. The concept of creativity, being the equivalent of the concept of" creativity", means a creative approach, a departure from the traditional scheme of thinking that generates something new [10].

The consideration of creativity as a special kind of ability began in the 50s of the last century, and it was initiated by the famous American psychologists L. Thurstone and J. Gilford associated with names. J. Gilford considers creativity as a factor of social intelligence in relation to the ability to recognize classes, systems, transformations, results of behavior, understanding the logical development of complex situations in interpersonal communication, understanding the internal motives of people's character [3]. In other words, creativity increases the level of social intelligence, while increasing the adaptive capabilities of the individual. Creativity is considered in two directions. The first defines how creativity and intelligence interact, and focuses on the study of cognitive processes associated with creativity. The second one examines whether the psychological characteristics of the individual can be the dominant aspects of creativity. This direction refers to the focus on personal and motivational characteristics.

According to the concept of creativity by D. J. Guilford, creativity is a universal cognitive creative ability, a separate factor independent of intelligence. Creative thinking is the foundation of creative achievement. The researcher identifies six parameters of creativity:

1) the ability to find and pose a problem;
2) search for new ideas;
3) ability to generate different ideas-flexibility;
4) originality-the ability to create unconventional ideas;
5) receptivity to contradictions caused by abnormal mechanisms, as well as a willingness to quickly move from one idea to another;
6) the ability to solve problems, i.e. to analyze and synthesize.

The study of creativity takes place in two directions. The first direction is the search for an answer to the question whether creativity is related to intelligence and is replaced by cognitive processes, the second direction is the reflection of questions about whether personality is an aspect of creativity along with psychological features and whether it is transmitted by personal, motivational signs- L. A. Belyaev [11].

Also, the ability to associate thinking with forecasting can be called as components of creativity. Its essence is determined by the ability of the body to prepare for the future state. This is an integrative creative trait associated with observation.

FULL TEXTS BOOK

The structural components of creativity in the complex determine the creative style of behavior, ensure productivity, uniqueness and creative transformation of the individual in various areas of everyday life and activities. The process of creating creativity includes the following stages:

- motivational impulse-imitation (standards of creative behavior, mastering the ways of creative activity transformation transformation of experience in accordance with individual characteristics, capabilities and needs);

- formation of the psychological structure of creativity and creative coherence of improving activities.

G. M. Mardanova argues that the creative potential of the individual reflects the highest form of human mental activity. A person with a high creative potential is different from others: creativity, intellectual initiative, activity, is inclined to creative thinking that can go beyond the boundaries of template thinking, is able to resolve emerging contradictions; is able to form the results of their activities, formulate and implement new approaches to solving problems, a creative person is inherent in research search.

The criterion of creativity is not the quality of the result, but the characteristics and processes that activate creative productivity. Development of creative thinking - i.e. formation and improvement of mental operations: analysis, grouping, comparison and addition, grouping, planning, deduction and quality of thinking development of criticality, depth, dexterity, breadth, transience, variability, as well as the use and expansion of knowledge of various types of content [12].

In these studies, a large range of qualities that determine the content of training a specialist for creativity, the conditions for its development are revealed. When discussing the preparatory issues of professional creativity, All authors share the same opinion that creativity is the highest threshold of professionalism.

According to S. Mednik, it is necessary to distinguish between verbal and nonverbal creativity. Verbal creativity is expressed in the dictionary form, for example, in the effective expression of opinions in problem situations, nonverbal creativity - in the manifestation of creative abilities in the form of artistic images, etc.

Results of the study.

Analyzing the approaches of foreign researchers to the definition of creativity, it was found that there is no formed formulation in this question. However, according to the results of the analysis, it was found that the definitions given by the researchers can be classified in four aspects: creative process, creative product, creative personality, Creative environment (industry, structure that forms the requirements for a creative product). By joining the creative process, the future teacher acquires the features of a creative personality in the creative environment, shows activity and becomes more active in the creative product in the future.

The analysis of domestic and foreign psychological research works allowed us to identify a number of definitions related to creativity (Table 1).

Table 1-definitions of scientists who gave the concept of " Creative"

Author	Reference
1	2
R. Hollmann	the flow of the reception, implemented in new ways
Cube	ability to find new connections

FULL TEXTS BOOK

Lasswell	the ability to make discoveries and learn them
P. Sh. Makhanova Zh. Kydyrbayeva	subjective determinants of creativity, the system (multidimensional, multi-level) mental structure
Murray	the emergence of new works
M. S. Bernstein, E. B. Evlanov, Y. A. Ponomarev, C. P. Torrens, C. D. Rogers	a distinctive feature of a person in the flow of life, in motion, or a characteristic of an individual's behavior
D. B. Bogoyavlenskaya, A.V. Brushlinsky, O. K. Tikhomirov	without separating the concepts of "creativity" and "creative potential" from each other, they are considered as intellectual and creative prerequisites aimed at creative activity
J. Gilford	a distinctive feature of divergent thinking, an exceptional quality of thought processes, high human activity, a special new, unique character of the individual the ability to reflect the deep structure of the attitude to the construction of values, non-standard thinking
L. Thurstone	creative abilities are not related to intellectual abilities and represent the following manifestations of creativity: speed of perception, ability to systematize ideas, intellectual factors of cognitive activity (concentration, creativity, etc.). receptivity to new ideas, a special state of consciousness
Taylor	transfer of experience to a new concept
Y. A. Ponomarev	the quality of personal integration
J. Gilford	divergent thinking
V. Ghiselin	the illusion of new constellational meanings
Rogers	the emergence of new relationships, the study of a person's inner psychological world to overcome the limitations in communicating with the outside world
S. I. Makshanov, N. Y. Khryascheva	constructive, non-standard thinking and behavior, as well as the ability to understand and develop your experience
R. Arnheim	this is the full disclosure of knowledge, actions and desires»
D. B. Bogoyavlenskaya, L. B. Ermolaeva-Tomina	intellectual activity

T. A. Barysheva, Y. A. Zhigalov	system (multi-level, multidimensional) mental structure, which, along with intellectual potential, is associated with motivation, the level of aesthetic development, existential, communicative indicators, competence, etc.
D. V. Chernilevsky	mental activity that leads to new solutions
O. V. Butorina	• ability to be creative; • intellectual creativity; • exclusivity, something new; • astral association; • restructuring of the unified system; • unique encoding of information; • divergent thinking; • the result of intrapersonal tension; • go beyond existing knowledge; • unconventional thinking that allows you to quickly solve a problem or situation

Analyzing the definitions given to the concept of creativity, we noticed that scientists have studied the kraetiv in the literature from the point of view of a special state, phenomenon, ability, behavior characteristics, mental structure, unconventional thinking, mental activity, etc.

Of course, in our study, we do not exclude that creativity is a mental structure (Makhanova P. Sh., Kydyrbaeva Zh.). However, since our research is devoted to its formation in the personality, it seems to us that the concept of creativity (J. Gilford, S. I. Makshanov and N. Y. Khryashcheva, Lassuel, Kyubi, M. A. Kholodnaya).

Based on these conclusions, in our study we define that creativity is the creativity of non – standard thinking, which is important for a clear statement, especially the rapid solution of new problems. Therefore, we consider creativity as a phenomenon, and by creativity we mean the manifestation of this phenomenon in the individual.

Results and discussion of the study.

The questionnaire For determining the level of formation of students ' communicative creativity is aimed at understanding the importance of communicative creativity and the correct approach to the formation of their own communicative creativity; studying knowledge from the field of communicative creativity and phenomenal understanding of communicative creativity; identifying the level of ability to form communicative creativity, the ability to fully understand and independently assess their qualities of communicative creativity.

Do you want to build a creative profile "learning and creativity" in the course identify the positive opinions of students on the formation of creativity?"89% of survey participants said they wanted to form, 30% – on lessons of 22.9% during the study elective courses; 21,3% on extracurricular activities, 17,6% - in specially organized activities. At the same time, 10% of students said that they do not want to form creativity. The survey of students showed that the interest in the formation of creativity is low and even absent.

FULL TEXTS BOOK

One of the surveys related to determining the course of formation of creativity of future specialists is the completeness of creative knowledge of students. The pedagogical expertise obtained at the university revealed a low level of creativity of students. Despite the fact that the students had ideas about the formation of creativity, most of the students did not realize the importance of the communicative creativity of the primary school teacher, did not have the desire to understand the functions, the role of the teacher in creativity, its features.

It is noted that to determine which qualities are important and not important for the organization of creativity, which will be as clear as possible for the formation of creativity, compiled to identify differences in the concepts and types of creativity of students.

As a result, many students noted the formation of creativity, the desire to practice, flexibility to new problems, the propensity to solve problems, flexibility in solving problems, the ability to go beyond the task, the mobility of thinking, the ability to quickly and suddenly change the ways of solving, the presence of thinking features, independence, individuality, the ability to make non-standard decisions.

Next, during the proposed survey students about how to be a teacher for the formation of communicative creativity of the modern teacher, analyzing the answers to the questions, for the vast majority of students: "the future primary school teacher needs to be highly creative"-90% of students (100) showed; "of the future primary school teacher needs to adapt quickly to new challenges" to 80.3% of students (89) showed; "of the future primary school teacher needs to be confident" – 69.7% of the students (70) showed; "future elementary school teacher should" is 65% (61) students; "the availability of features thinking in primary school teachers" is 62 % (58) of students.

To determine the level of creativity of the participants of the experiment, the activity of students in various educational, extracurricular activities, scientific and other activities was monitored. As a result of the control, the predominance of medium and low levels in the formation of communicative creativity was revealed.

Conclusion.

Thus, diagnostic approaches have established that many students have a predominant medium and low level in the formation of creativity. At the same time, it is established that teachers and mentors of groups want to build relationships with students on the basis of the principle of personality-oriented communication, while there is a positive impact of joint activities between the student and the teacher in the process of forming creativity.

According to the results of the survey, on the one hand, the students ' requirements for the formation of creativity were revealed, and on the other-the low level of their knowledge and ability to demonstrate and evaluate in the formation of creativity was revealed.

REFERENCES

1. Amabile, T. (2006). How to Kill Creativity // In Creative Thinking in Business. Moscow: Nauka, pp. 46–51.
2. Barron, F. (1988). Putting creativity to work // In R. Sternberg & T. Tardif (Eds.), The Nature of Creativity. Cambridge: Cambridge University Press, pp. 76–98.
3. Guilford, J. P. (1968). Intelligence, Creativity and Their Educational Implications. San Diego, CA: Knapp, p. 5.
4. Torrance, E. P. (1980). Growing up creatively gifted: A 22-year longitudinal study // Creative Child and Adult Quarterly, p. 22.

FULL TEXTS BOOK

5. De Bono, E. (2015). Serious Creativity: A Step-by-Step Approach to Using the Logic of Creative Thinking. Moscow: Alpina Publisher, 381 p.
6. Maslow, A. (1985). The Farther Reaches of Human Nature. New York: Viking Press, 281 p.
7. Rogers, N. (1993). Creative Connection. Palo Alto: Science and Behavior Books, Inc., 188 p.
8. Bogoyavlenskaya, D. B. (1999). The Subject of Activity in the Problem of Creativity // Voprosy Psychologii, No. 2, pp. 35–41.
9. Luk, A. N. (1980). Problems of Scientific Creativity: Collection of Analytical Reviews. Moscow: INION, 227 p.
10. Buinova, D. I. (2013). Creativity as a Factor of Self-Realization of Personality in Professional Activity // Psychology, Sociology and Pedagogy, pp. 53–55.
11. Guilford, J. (1965). Structural Model of Intelligence // In A. M. Matyushkin (Ed.), Psychology of Thinking. Moscow: Progress, 553 p.
12. Buraeva, Zh. B., Berkimbaev, K. M., Praliyeva, R. E., & Tassova, A. B. (2017). The Essence of Pedagogical Creativity // Science and Life of Kazakhstan, Almaty, No. 6 (57), pp. 99–102.

FULL TEXTS BOOK

EBEVEYNİN BESİN NEOFOBİ DURUMUNUN ÇOCUK NEOFOBİSİNE ETKİSİ VE ANTROPOMETRİK ÖLÇÜMLERİNİN DEĞERLENDİRİLMESİ

THE EFFECT OF PARENTAL FOOD NEOPHOBIA ON CHILD NEOPHOBIA AND EVALUATION OF ANTHROPOMETRIC MEASUREMENTS

***Rezzan UYSAL*[1]**

[1]Diyetisyen

[1]ORCID ID: 0009-0008-9453-720X

***Ezgi TÜRKAY*[2]**

[2] Dr. Öğr. Üyesi, Antalya Bilim Üniversitesi, Sağlıklı Bilimleri Fakültesi, Beslenme ve Diyetetik, Antalya, Türkiye.

[2]ORCID ID: 0000-0002-1676-3606

ÖZET

Bu araştırma, ebeveynlerin besin neofobi düzeylerinin 8-10 yaş aralığındaki çocukların besin neofobisi eğilimleri ve antropometrik ölçümleri üzerindeki etkisini incelemek amacıyla yapılmıştır. Kesitsel tipte planlanan çalışma, Antalya'nın Serik ilçesindeki beş farklı ilkokuldan 151 çocuk ve bu çocukların ebeveynleri olmak üzere toplam 302 bireyle gerçekleştirilmiştir. Çalışmada veri toplama aracı olarak demografik bilgi formu, Çocuk Besin Neofobi Ölçeği ve Ebeveynler için Yeni Besin Neofobi Ölçeği kullanılmıştır. Ayrıca çocukların boy, kilo ve bel çevresi gibi antropometrik verileri ölçülerek BKİ hesaplanmıştır. Veriler SPSS 20.0 programı ile analiz edilmiş; Mann Whitney U, Kruskal-Wallis, Ki-Kare ve Spearman korelasyon testlerinden yararlanılmıştır. Araştırmaya katılan çocukların yaş ortalaması 9±0.8 yıl olup %41.7'si kız, %58.3'ü erkektir. Erkek çocukların yaşa göre boy persentil ortalaması 60.0±33.3 ve boy z skoru 0.6±1.7 olup, bu değerler kızlara kıyasla anlamlı düzeyde yüksektir ($p<0.05$). Çocukların besin neofobi puan ortalamaları cinsiyetlere göre benzer bulunmuştur (kızlarda 33.3±6.4, erkeklerde 33.4±5.3; $p=0.690$). Ebeveynlerin neofobi puanları çocuklara göre daha yüksek bulunmuştur (ortalama: 40.7±9.5). Ebeveyn yaşı ile besin neofobi puanı arasında zayıf düzeyde, pozitif yönde anlamlı bir ilişki saptanmıştır ($r=0.159$, $p=0.050$). Ancak çocukların BKİ düzeyleri ile besin neofobi puanları arasında anlamlı bir ilişki bulunamamıştır ($p>0.05$). Sonuç olarak, ebeveynin besin neofobisi düzeyinin çocuğun besin tercihleri üzerinde etkili olabileceği, ancak bu etkinin büyüme göstergeleri üzerinde doğrudan belirleyici olmadığı görülmektedir. Elde edilen veriler, çocukların sağlıklı beslenme alışkanlıklarının kazandırılmasında ebeveyn tutumlarının ve model davranışların önemine işaret etmektedir.

Anahtar kelimeler: Çocuk besin neofobisi, ebeveyn besin neofobisi, besin maruziyeti, antropometrik ölçüm

ABSTRACT

This study was conducted to examine the effects of parental food neophobia levels on the food neophobia tendencies and anthropometric measurements of children aged 8–10 years. Designed as a cross-sectional study, the research was carried out with a total of 302 individuals, including 151 children and their parents, from five different primary schools in the Serik district of Antalya, Türkiye. Data were

FULL TEXTS BOOK

collected using a demographic information form, the Child Food Neophobia Scale, and the New Food Neophobia Scale for parents. Additionally, anthropometric data such as height, weight, and waist circumference of the children were measured and Body Mass Index (BMI) was calculated. The data were analyzed using SPSS 20.0, and Mann Whitney U, Kruskal-Wallis, Chi-Square, and Spearman correlation tests were applied. The mean age of the participating children was 9±0.8 years; 41.7% were girls and 58.3% were boys. Boys had a significantly higher mean height-for-age percentile (60.0±33.3) and height z-score (0.6±1.7) compared to girls ($p<0.05$). The mean food neophobia scores of the children were similar across genders (girls: 33.3±6.4; boys: 33.4±5.3; $p=0.690$). Parents' neophobia scores were found to be higher than those of their children (mean: 40.7±9.5). A weak but statistically significant positive correlation was found between parental age and their food neophobia scores ($r=0.159$, $p=0.050$). However, no significant relationship was observed between the children's BMI levels and their food neophobia scores ($p>0.05$). In conclusion, while parental food neophobia levels may influence children's food preferences, this effect does not appear to directly impact anthropometric growth indicators. The findings emphasize the importance of parental attitudes and modeling behaviors in promoting healthy eating habits in children.

Keywords: Child food neophobia, parental food neophobia, food exposure, anthropometric measurement

GİRİŞ

Beslenme; büyüme gelişme, sürdürülebilir bir yaşam, sağlığın korunması ve iyileştirilmesi için gerekli olan besin ögelerinin ve bileşenlerin dışarıdan alınıp vücutta kullanılması olarak tanımlanmaktadır (1). Gereksinimlerinden düşük veya fazla alım vücut üzerinde farklı etkilere sebep olabilmektedir. Her yaşta dengeli besin alımının önemi göz ardı edilmemelidir.

Besin neofobisi; yeni, bilinmeyen besinlerin tüketilmesine karşılık bir reddetme veya bu besinlere karşı isteksizlik olarak tanımlanabilmektedir (2). Neofobi sebebi her yaşta farklılık söz konusu olabilen bir durumdur. Yetişkinlerde genel olarak temel nedenin insan sağlığı üzerinde olumsuz etkiler yaratabilecek besinler olmasına karşı duyulan tedirginlik olduğu düşünülmektedir (3,4). Okul çağındaki çocuklarda neofobinin en büyük sebebi ise genellikle bilmedikleri besinin görüntüsüne karşı duydukları ön yargıdır (5). Yenilik değeri besin neofobisi ve besin seçimini ayıran noktadır (6). Seçici yemede hem yeni hem de bilinen besine karşı bir reddetme durumu söz konusudur. Besin neofobisindeyse sadece besin reddi bulunmaktadır ve bazen besin seçiminin alt alanı olarak kabul edilebilmektedir (6,7). Neofobide yaş, cinsiyet, ailenin tutum ve davranışlarının yanı sıra pek çok etmen etkili olabilmektedir. Özellikle yaş neofobide önemli bir belirleyici olarak kabul edilmektedir (7-9).

Çocukluk çağında başlayan bu neofobinin yetişkinlik dönemlerinde yaşamları, sağlıkları üzerine olumsuz etkileri gözlemlenebileceği, besin alımında yetersizliğin söz konusu olacağı gözlemlenmiştir. Besin neofobisi üzerinde ailenin etkileri değerlendirildiği zaman en sık karşılaşılan sebep; çocuk tarafından başta beğenilmeyen besinin çocuğun beslenmesinden tamamen çıkartılmasıdır. Literatürde ise bir besinin 8-15 kez denemesi sonucunda o besine karşı duyulan yargının yıkılabileceği ve tüketiminin zamanla artacağından bahsedilmektedir (3-10-11). Bebeklikten itibaren besin çocuğa defalarca denetilmeli ve sevebileceği, eğlenceli tabaklarla çocuğa sunulmalıdır. Aynı zamanda besine karşı neofobisi olan ebeveyn çocuğuna bu durumu yansıtmamalıdır. Masaya o yemeği, besini koymaktan kaçınmamalı çocuğun tüketimini sağlamalıdır. Çocukların ebeveynlerinde gördüklerini uyguladıklarını unutmamak gerekmektedir. Besin neofobisine sahip olan bireylerin çoğunda sebze ve meyve çeşitliliğinin daha az olduğu gözlemlenmiştir (12). Yüksek eğitime sahip olan bireylerde ise besin çeşitliliğinin artması ve besinlere erişimin kolay olması sebebiyle besin neofobisinin

FULL TEXTS BOOK

gözlemlenmesi çok daha düşük olduğu söylenmektedir (13-14). Sosyodemografik açıdan incelendiği zaman ise düşük gelire sahip olan ailedeki bireylerin yüksek gelir düzeyine sahiip olanlara kıyasla daha neofobik olduğu gözlemlenmiştir. Bu durum da farklı, yeni besinlerle karşılaşma olasılıklarının düşüklüğü ile bağdaşmaktadır (15). Besin seçiminde maruz kalınan çevre, deneyimler ve genetik yatkınlık rol oynamaktadır (8,9). Kalıtsallık ve gıda neofobisi ilişkisi ile ilgili çalışmalar sınırlı da olsa, 331 ikizin katıldığı bir çalışmada gıda seçiminde genetik aktarımın rolü araştırılmış ve besin seçiminde bireylerin %72'sinin genetik, %28'inin çevresel etkenlerden etkilendikleri bulunmuştur (9-11). Literatürde bu çalışmayla benzer sonuçlar gösteren araştırmalar da mevcuttur (12-13). Kadınların genetik etkilerle gıda neofobisinden %61 oranında, erkeklerin ise çevresel etkilerle %45 oranında etkilendiği raporlanmıştır (13).

YÖNTEM

Evren ve Örneklem

Araştırma 2024-2025 eğitim- öğretim yıllarında Antalya ili Serik ilçesine bağlı bulunan 3 devlet okulunda (Tekeli İlkokulu, Akçaalan İlkokulu, Boğazkent Nurten Turan Kilit İlkoukulu) 8-10 yaş arası ilkokul öğrencileri ve ebeveynleri ile yürütülmüştür. Serik İlçe Milli Eğitim Müdürü'nden alınan bilgiyle 8-10 yaş aralığında bulunan öğrenci sayısının yaklaşık 1140 kişi olduğu belirlenmiştir. Araştırma evreninin tamamına ulaşmak zaman ve maaliyet yönünden zor olacağı için evreni teslim edecek şekilde evreni bilinen örneklem sayısına bakılarak örneklem sayısı %95 güven aralığı ve %5 örneklem hatasıyla araştırmada görüşülmesi gereken çocukların sayısı 288 olarak saptanmış ve her bir çocuk için ebeveyn değerlendirilmesi yapılacağından dolayı ebeveyn sayısı 288 olarak belirlenmiştir. Ebeveyn sayısı ile toplam örneklem sayısının 576 olacağı belirlenmiştir. Ancak yaşanan bazı sağlık sorunları nedeni ile 151 çocuk ve 151 ebeveyn olmak üzere toplam 302 kişi ile çalışma tamamlanmıştır.

Veri Toplama Araçları

Veri toplama aracı olarak çocuk ve ebeveynlere anket uygulanmıştır. Anket formunda demografik bilgiler (yaş, cinsiyet), çocuk ve ebeveyn için besin neofobi ölçeği ve antropometrik ölçüm değerleri yer almaktadır.

BULGULAR

Çocuk ve Ebeveynlerin Besin Neofobi Ölçeğinden Aldığı Puanların Dağılımı

Çalışmada çocuklara Besin Neofobi ölçeği, ebeveynlerine ise Yeni Besin Neofobi ölçeği uygulanmıştır. Ölçekten alınan puanların dağılımı Tablo 1'de verilmiştir. Ölçek maksimum puanı 70'tir. Puan yükseldikçe besin neofobisinin varlığı ile ilişkilendirilmektedir. Ebeveynlerin ölçekten aldıklarının puanların ortalaması çocuklara göre daha yüksek bulunmuştur. Genel dağılımlar benzerdir. (Tablo 1)

Tablo 1. Ölçeklerden alınan puanların dağılımı

	n	±SS	Medyan	Min-Maks
Besin Neofobi	151	33.3±5.8	34	16-50
Ebeveyn Neofobi	151	40.3±8.6	40	12-67

FULL TEXTS BOOK

Ölçeklerden Alınan Puanların Çocukların ve Ebeveynlerin Cinsiyetine Göre Dağılımı

Çocukların cinsiyetine göre ölçeklerden aldıkları puanlar ve ebeveynlerinin ölçekten aldıkları puanlar karşılaştırılmıştır. Çocukların cinsiyetlerine göre Besin Neofobi ölçeğinden aldıkları puanlar arasında istatistiksel anlamlı bir fark yoktur ($p>0.05$). Kız ve erkek çocukların ölçekten aldıkları puanların dağılımı benzerdir. Kız çocuk ebeveynlerinin ve erkek çocuk ebeveynlerinin de ölçekten aldıkları puanlar arasında istatistiksel anlamlı bir fark bulunmamıştır ($p>0.05$). Ebeveynler ölçekten benzer puanlar almıştır. (Tablo 2)

Tablo 2. Ölçeklerden Alınan Puanların Çocukların ve Ebeveynlerin Cinsiyetine Göre Dağılımı

	n	±SS	Medyan	Min - Maks	p
Besin Neofobi					
Kız	63	33.3±6.4	34	16-45	0.690
Erkek	88	33.4±5.3	34	20-50	
Ebeveyn Neofobi					
Kadın	63	41.0±9.3	40	12-67	0.700
Erkek	88	40.5±9.7	40	18-90	

Ölçeklerden Alınan Puanların Çocukların BKİ Gruplarına Göre Dağılımı

Çocukların BKİ gruplarına göre ölçeklerden aldıkları puanlar ve ebeveynlerinin ölçekten aldıkları puanlar karşılaştırılmıştır. Çocukların BKİ derecelerine göre Besin Neofobi ölçeğinden aldıkları puanlar arasında istatistiksel anlamlı bir fark yoktur ($p>0.05$). Tüm gruplardaki çocukların ölçekten aldıkları puanların dağılımı benzerdir. BKİ grubu farklı olan çocukların ebeveynlerinin de ölçekten aldıkları puanlar arasında istatistiksel anlamlı bir fark bulunmamıştır ($p>0.05$). Ebeveynler ölçekten benzer puanlar almıştır. (Tablo 3)

Tablo 3. Ölçeklerden Alınan Puanların Çocukların BKİ Gruplarına Göre Dağılımı

	n	±SS	Medyan	Min - Maks	p
Besin Neofobi					
Zayıf	104	33.2±5.6	34	16-46	0.216
Normal	36	33.6±5.9	34	20-50	
Şişman	10	33.9±7.6	34	17-42	
Obez	1	-	-	-	
Ebeveyn Neofobi					

FULL TEXTS BOOK

Zayıf	104	40.9±7	40	25-59	0.793
Normal	36	40.8±13.3	39	16-90	
Şişman	10	37.4±15.1	38.5	12-67	
Obez	1	-	-	-	

TARTIŞMA ve SONUÇ

Bu araştırmada ebeveynlerin besin neofobi düzeylerinin, çocukların besin neofobisi eğilimleri ve antropometrik ölçümleri üzerindeki etkisi incelenmiştir. Bulgulara göre ebeveynlerin besin neofobi puanlarının çocuklara göre daha yüksek olduğu görülmüştür. Literatürde de ebeveynlerin tutum ve davranışlarının çocukların besin seçimleri üzerinde doğrudan etkili olduğu, ebeveynin yeni besinlere karşı olumsuz yaklaşımının çocuklarda benzer eğilim geliştirdiği belirtilmektedir (15,16).

Çalışmada çocukların besin neofobi puanlarının cinsiyetlere göre farklılık göstermediği bulunmuştur. Daha önce yapılan bazı araştırmalar da neofobi eğilimlerinde cinsiyetin belirleyici olmadığını desteklemektedir (17). Bununla birlikte bazı toplumlarda erkek çocukların yeni besinleri denemeye daha az istekli olduğu rapor edilmiştir, bu farklılıkların kültürel ve sosyoekonomik faktörlerden kaynaklanabileceği düşünülmektedir.

Ebeveynlerin yaşları ile besin neofobi puanları arasında pozitif yönde zayıf düzeyde bir ilişki bulunmuştur. Benzer şekilde, yaşın ilerlemesiyle birlikte besin tercihlerinde esnekliğin azaldığını ve yeni besinlere karşı daha temkinli bir yaklaşımın geliştiğini bildiren çalışmalar mevcuttur (16).

Çocukların BKİ düzeyleri ile besin neofobi puanları arasında anlamlı bir ilişki bulunmamıştır. Bu bulgu, besin neofobisinin kısa vadede büyüme göstergeleri üzerinde doğrudan etkili olmadığını düşündürmektedir. Ancak literatürde, uzun vadede besin çeşitliliğinin azalmasının mikronutrient yetersizlikleri ve sağlıksız beslenme alışkanlıkları ile ilişkili olabileceği vurgulanmaktadır (18,19).

Sonuç olarak, bu çalışmadan elde edilen veriler ebeveynlerin beslenme konusundaki tutumlarının çocukların yeni besinlere yaklaşımında kritik rol oynadığını göstermektedir. Özellikle ebeveynlerin kendi neofobik eğilimlerini çocuklarına yansıtmamaları ve yeni besinleri tekrar tekrar olumlu bir yaklaşımla sunmaları, çocuklarda besin kabulünü artırmada önemli görülmektedir.

Tablo 3. Ölçeklerden Alınan Puanların Çocukların BKİ Gruplarına Göre Dağılımı

	n	**±SS**	**Medyan**	**Min - Maks**	**p**
Besin Neofobi					
Zayıf	104	33.2±5.6	34	16-46	0.216
Normal	36	33.6±5.9	34	20-50	
Şişman	10	33.9±7.6	34	17-42	
Obez	1	-	-	-	
Ebeveyn Neofobi					

FULL TEXTS BOOK

Zayıf	104	40.9±7	40	25-59	0.793
Normal	36	40.8±13.3	39	16-90	
Şişman	10	37.4±15.1	38.5	12-67	
Obez	1	-	-	-	

TARTIŞMA ve SONUÇ

Bu araştırmada ebeveynlerin besin neofobi düzeylerinin, çocukların besin neofobisi eğilimleri ve antropometrik ölçümleri üzerindeki etkisi incelenmiştir. Bulgulara göre ebeveynlerin besin neofobi puanlarının çocuklara göre daha yüksek olduğu görülmüştür. Literatürde de ebeveynlerin tutum ve davranışlarının çocukların besin seçimleri üzerinde doğrudan etkili olduğu, ebeveynin yeni besinlere karşı olumsuz yaklaşımının çocuklarda benzer eğilim geliştirdiği belirtilmektedir (15,16).

Çalışmada çocukların besin neofobi puanlarının cinsiyetlere göre farklılık göstermediği bulunmuştur. Daha önce yapılan bazı araştırmalar da neofobi eğilimlerinde cinsiyetin belirleyici olmadığını desteklemektedir (17). Bununla birlikte bazı toplumlarda erkek çocukların yeni besinleri denemeye daha az istekli olduğu rapor edilmiştir, bu farklılıkların kültürel ve sosyoekonomik faktörlerden kaynaklanabileceği düşünülmektedir.

Ebeveynlerin yaşları ile besin neofobi puanları arasında pozitif yönde zayıf düzeyde bir ilişki bulunmuştur. Benzer şekilde, yaşın ilerlemesiyle birlikte besin tercihlerinde esnekliğin azaldığını ve yeni besinlere karşı daha temkinli bir yaklaşımın geliştiğini bildiren çalışmalar mevcuttur (16).

Çocukların BKİ düzeyleri ile besin neofobi puanları arasında anlamlı bir ilişki bulunmamıştır. Bu bulgu, besin neofobisinin kısa vadede büyüme göstergeleri üzerinde doğrudan etkili olmadığını düşündürmektedir. Ancak literatürde, uzun vadede besin çeşitliliğinin azalmasının mikronutrient yetersizlikleri ve sağlıksız beslenme alışkanlıkları ile ilişkili olabileceği vurgulanmaktadır (18,19).

Sonuç olarak, bu çalışmadan elde edilen veriler ebeveynlerin beslenme konusundaki tutumlarının çocukların yeni besinlere yaklaşımında kritik rol oynadığını göstermektedir. Özellikle ebeveynlerin kendi neofobik eğilimlerini çocuklarına yansıtmamaları ve yeni besinleri tekrar tekrar olumlu bir yaklaşımla sunmaları, çocuklarda besin kabulünü artırmada önemli görülmektedir.

KAYNAKÇA

1. Türkiye Beslenme Rehberi (TÜBER) 2022, Sağlık Bakanlığı, Halk Sağlığı Genel Müdürlüğü, Sağlık Bakanlığı Yayın No:1031, Ankara 2022.
2. Białek-Dratwa A, Staśkiewicz-Bartecka W, Kiciak A, Wardyniec A, Grajek M, Aktaç Ş, Çelik ZM, Sabuncular G, İslamoğlu AH, Kowalski O. Food Neophobia and Avoidant/Restrictive Food Intake among Adults and Related Factors. Nutrients. 2024 Sep 2;16(17):2952. doi: 10.3390/nu16172952. PMID: 39275268; PMCID: PMC11397511.
3. Pliner, P., & Hobden, K. (1992), Development of A Scale to Measure The Trait of Food Neophobia in Human, Appetite, 19(2), 105-120.
4. Maiz, E., & Balluerka, N. (2018), Trait Anxiety and Self-Concept Among Children and Adolescents With Food Neophobi. Food Research International, 105, 1054 1059.

FULL TEXTS BOOK

5. Maratos, F. A., & Staples, P. (2015), Attentional Biases Towards Familiar and Unfamiliar Foods in Children. The Role of Food Neophobi, Appetite, 91, 220 225.
6. Dovey TM, Staples PA, Gibson EL, Halford JCG. Food neophobia and 'picky/fussy' eating in children: A review. Appetite. 2008;50(2-3):181–93.
7. Lafraire J, Rioux C, Giboreau A, Picard D. Food rejections in children: Cognitive and social/environmental factors involved in food neophobia and picky/fussy eating behavior. Appetite. 2016;96:347–57.
8. Kermen S, Aktaç Ş. Picky eating and risk factors in children. Guncel Pediatr. 2018;16(3):85–102.
9. Duman E, Akçil Ok M, Keser A. Besin Neofobisi Ölçeğinin Türkçeye uyarlanması: Geçerlik ve güvenirlik çalışması. Kocaeli Üniversitesi Sağlık Bilim Derg. 2020;6(2):157–61.
10. Brich LL. Development of food preferences. Annual Review of Nutrition.1999;19:41-62.
11. Törnwall O, Silventoinen K, Hiekkalinna T, Perola M, Tourila H, Kaprio J. Identifying flavor preference subgroups. Genetic basis and related eating behavior traits. Appetite.2014;75:1-10.
12. Knaapila A, Silventoinen K, Broms U, et al. Food neophobia in young adults: Genetic architecture and relation to personality, pleasantness and use frequency of foods, and body mass index a twin study. Behavior Genetics.2011;41(4):512-521.)
13. Tuorila H, Lahteenmaki L, Pohjalainen L, Lotti L. Food neophobia among the finns and related responses to familiar and unfamiliar foods. Food Quality and Preferences. 2001;12(1):29-37.
14. Paupério A, Severo M, Lopes C, Moreira P, Cooke L, Oliveira A. Could the food neophobia scale be adapted to pregnant women? A confirmatory factor analysis in a Portuguese sample. Appetite.2014; 75:110-116.)
15. Addessi, E., Galloway, A. T., Visalberghi, E., & Birch, L. L. (2005). Specific social influences on the acceptance of novel foods in 2–5-year-old children. Appetite, 45(3), 264–271. https://doi.org/10.1016/j.appet.2005.07.007
16. Cooke, L. J. (2007). The importance of exposure for healthy eating in childhood: A review. Journal of Human Nutrition and Dietetics, 20(4), 294–301. https://doi.org/10.1111/j.1365-277X.2007.00804.x
17. Dovey, T. M., Staples, P. A., Gibson, E. L., & Halford, J. C. (2008). Food neophobia and "picky/fussy" eating in children: A review. Appetite, 50(2–3), 181–193. https://doi.org/10.1016/j.appet.2007.09.009
18. Nicklaus, S. (2009). Development of food variety in children. Appetite, 52(1), 253–255. https://doi.org/10.1016/j.appet.2008.09.018
19. Wardle, J., Carnell, S., & Cooke, L. (2005). Parental control over feeding and children's fruit and vegetable intake: How are they related? Journal of the American Dietetic Association, 105(2), 227–232. https://doi.org/10.1016/j.jada.2004.11.006

FULL TEXTS BOOK

NSTE-AKS SONRASI YENİDEN YATIŞI ÖNGÖRMEDE D-DİMER/ALBÜMİN, CRP/ALBÜMİN VE KOMPOZİT İNFLAMASYON-KOAGÜLASYON İNDEKSİ

D-DIMER/ALBUMIN, CRP/ALBUMIN, AND THEIR COMPOSITE INFLAMMATORY-COAGULATION INDEX PREDICT READMISSION AFTER NSTE-ACS

***Özge ÇETİNARSLAN*[1]**

[1] T.C. Demiroğlu Bilim Üniversitesi, Tıp Fakültesi, Kardiyoloji Anabilim Dalı, İstanbul Florence Nightingale Hastanesi, İstanbul, Türkiye

[1]ORCID ID: https://orcid.org/0000-0003-4825-6489

ÖZET

Giriş: ST yükselmesiz-akut koroner sendrom (NSTE-AKS) sonrası yeniden yatışlar hem sağlık sistemini hem de hastaları zorlar. D-dimer/Albümin oranı (DAR) trombotik aktivite ile beslenme rezervini, CRP/Albümin oranı (CAR) ise inflamasyonu yansıtır. Bu çalışmada DAR'ın tek başına ve CAR ile kombinasyonu (İnflamasyon-Koagülasyon İndeksi [ICI] = DAR × CAR) ile prognostik değerini inceledik.

Yöntemler: Tek merkezli kayıtta yer alan 465 NSTE-AKS hastasında (medyan takip 18 ay) hastane başvurusu laboratuvarlarından DAR, CAR ve ICI hesaplandı. Birincil sonlanım kardiyak yeniden yatış; ikincil sonlanımlar plansız revaskülarizasyon, ölüm, akut böbrek hasarı (ABY) ve dördünün bileşik sonlanım noktasıdır. Lojistik regresyon yaş, cinsiyet, diyabet, hipertansiyon, geçirilmiş perkütan koroner girişim (PKG), eGFR ve sol ventrikül ejeksiyon fraksiyonu (LVEF) için ayarlandı.

Bulgular: Ortalama yaş 65 ± 11 yıl; hastaların %28'i diyabetikti. DAR yeniden yatışı (ayarlı OR 1,74; %95 GA 1,06–3,10; p = 0,030) ve bileşik sonlanımı (OR 1,48; %95 GA 1,05–2,09; p = 0,023) bağımsız olarak öngördü. DAR yerine ICI kullanılması yeniden yatış için ilişkiyi güçlendirdi (OR 2,01; %95 GA 1,22–3,58; p = 0,007) ve plansız revaskülarizasyon için de anlamlılık sağladı (OR 1,79; p = 0,034). DAR ve ICI, ABY ile ilişkili değildi; mortalite sinyali ise istatistiksel anlamlılığa ulaşmadı. Her iki belirteç IPTW ve çift-sağlam modellemelerden sonra da anlamını korudu.

Sonuç: Koagulasyon (D-dimer) ve inflamasyon-beslenme (CRP/Albümin) bileşimini içeren basit girişimsel olmayan bir indeks, NSTE-AKS sonrası yeniden yatış ve tekrar revaskülarizasyon riskini sınıflandırır. DAR ve CAR rutin olarak elde edildiğinden, bunların çarpımı olan ICI, yoğun takip gerektiren yüksek riskli hastaları erken ve maliyetsiz olarak tanımlamayı kolaylaştırabilir.

Anahtar Kelimeler: D-dimer/Albümin oranı (DAR); CRP/Albümin oranı (CAR); İnflamasyon-Koagülasyon İndeksi (ICI); ST yükselmesiz akut koroner sendrom (NSTE-AKS)

ABSTRACT

Background: Rehospitalisation after non-ST-elevation acute coronary syndrome (NSTE-ACS) strains health systems and patients alike. The D-dimer-to-albumin ratio (DAR) integrates thrombotic drive with nutritional reserve, while the C-reactive-protein/albumin ratio (CAR) captures residual inflammation.

FULL TEXTS BOOK

We examined the prognostic value of DAR alone and in combination with CAR (Inflammatory-Coagulation Index [ICI] = DAR × CAR).

Methods: In a single-centre registry of 465 NSTE-ACS patients (median follow-up 18 months) we calculated DAR, CAR and ICI from admission laboratories. End-points were cardiac rehospitalisation (primary), unplanned revascularisation, death, acute-kidney-injury (AKI) and a composite of all four. Logistic regression adjusted for age, sex, diabetes, hypertension, prior PCI, eGFR and left-ventricular ejection fraction (LVEF).

Results: Mean age was 65 ± 11 years; 28 % were diabetic. DAR independently predicted rehospitalisation (adjusted OR 1.74, 95 % CI 1.06–3.10; p = 0.030) and the composite end-point (OR 1.48, 95 % CI 1.05–2.09; p = 0.023). Substituting ICI strengthened the association with rehospitalisation (OR 2.01, 95 % CI 1.22–3.58; p = 0.007) and additionally reached significance for unplanned revascularisation (OR 1.79; p = 0.034). Neither DAR nor ICI was linked to AKI, and the mortality signal did not attain statistical significance. Both markers retained significance after inverse-probability-of-treatment weighting (IPTW) and doubly-robust modelling.

Conclusions: A simple admission index that combines coagulation (D-dimer) and inflammation-nutrition (CRP/albumin) stratifies rehospitalisation and repeat revascularisation risk after contemporary NSTE-ACS. Because DAR and CAR are routinely available, their product (ICI) could facilitate early, cost-free identification of high-risk patients warranting intensified follow-up.

Keywords: D-dimer/albumin ratio (DAR); CRP/albumin ratio (CAR); Inflammatory-Coagulation Index (ICI); non-ST-elevation acute coronary syndrome (NSTE-ACS)

Introduction

Rehospitalisation within one year of an acute coronary syndrome (ACS) incurs a 4- to 5-fold increase in subsequent mortality and consumes up to one-third of episode-related health-care costs.[1] Although the TIMI and GRACE scores perform reasonably well for short-term death/myocardial-infarction prediction, they are ill-suited to forecast later rehospitalisation, which is governed by a complex nexus of residual thrombo-inflammatory activity, nutritional reserve and ventricular recovery.

Circulating D-dimer is a surrogate of fibrin turnover that rises with plaque thrombosis and correlates with long-term cardiovascular events. Albumin, conversely, mirrors both nutritional status and systemic inflammation; low levels portend worse outcomes after ACS. The ratio of D-dimer to albumin (DAR) therefore couples haemostatic drive with host reserve in a single metric. Likewise, the CRP/albumin ratio (CAR) capitalises on the disparate half-lives of its constituents to index unresolved inflammation.[2]

Previous work has linked DAR and CAR to mortality in sepsis, stroke and COVID-19, yet their relevance to post-ACS rehospitalisation remains unexplored.[3,4] We hypothesised that (i) DAR would predict cardiac rehospitalisation after non-ST-elevation acute coronary syndrome (NSTE-ACS) and (ii) combining DAR with CAR into a composite inflammatory–coagulation index (ICI) would further refine risk stratification.

Methods

We analysed the retrospectively NSTE-ACS registry of Istanbul Florence Nightingale Hospital from January 2020 to June 2024. All patients ≥ 18 years who fulfilled ESC criteria for NSTE-ACS and had same-day measurements of D-dimer, CRP and albumin were eligible.[5] Exclusions were active malignancy, chronic inflammatory disease, end-stage renal or hepatic failure, and loss to follow-up. The institutional review board approved the study with waiver of informed consent owing to the observational design.

DAR = (D-dimer [µg/L) / (albumin [g/dL).

CAR = (CRP [mg/L) / (albumin [g/dL).

ICI = DAR × CAR. Biomarkers were analysed as continuous variables; ICI was rescaled by dividing by 100 to yield clinically interpretable odds ratios.

Primary endpoints: first rehospitalisation for cardiac cause (angina, heart failure decompensation or arrhythmia) within 18 months.

Secondary endpoints: unplanned coronary revascularisation (PCI or CABG), all-cause death, acute-kidney-injury (AKI) defined by KDIGO criteria, and a composite of the four.[6]

Demographics, cardiovascular history, index-admission vitals, angiography results and discharge medications were extracted from electronic records. Serum creatinine estimated glomerular filtration rate (CKD-EPI) and LVEF (biplane Simpson) were included a-priori as confounders.

Continuous variables are expressed as mean ± SD and compared with Student's t or Mann–Whitney U as appropriate; categorical variables are counts (%) compared with χ^2 or Fisher exact. Multivariable logistic models were built using purposeful selection: variables with $p < 0.20$ in univariable testing or strong biological plausibility entered the full model, followed by backward elimination at $p < 0.05$. Discrimination was quantified by C-statistic and calibration by the Hosmer–Lemeshow test. To address missingness (< 5 % per variable), complete-case analysis was prespecified. Two-tailed $p < 0.05$ was considered significant. Analyses were run in SPSS.

Results

Of the 498 NSTE-ACS admissions screened, 33 lacked ≥ 1 biomarker and were excluded, leaving 465 patients for analysis. Cardiac rehospitalisation within 18 months occurred in 129 (27.7 %). Baseline characteristics are shown in Table 1. Compared with those not readmitted, rehospitalised patients were older, more frequently female and exhibited higher median DAR and ICI values (all $p < 0.01$). β-blocker therapy at discharge was also more prevalent in the rehospitalised group (88 % vs 71 %, $p < 0.001$), whereas angiographic complexity (three-vessel disease) did not differ. After adjusting for β-blocker use and including a β-blocker × high-ICI interaction term, ICI remained an independent predictor of rehospitalisation (adjusted OR 1.94, $p = 0.009$) and the interaction was not significant ($p = 0.33$), indicating that treatment imbalance did not account for the biomarker-outcome association.

Variable	**No-rehospitalisation (n = 336)**	**Rehospitalised (n = 129)**	***p*-value**
Age (yrs)	63.6 ± 11.3	65.9 ± 13.3	0.084
LVEF (% Simpson)	52.8 ± 8.3	49.7 ± 8.5	**0.001**

FULL TEXTS BOOK

Creatinine (µmol/L)	1.05 ± 0.88	0.97 ± 0.34	0.131
CAR	4.55 ± 11.28	8.81 ± 19.41	**0.020**
DAR	0.68 (0.34–1.47)	0.94 (0.48–1.92)	**0.008**
ICI	3.1 (1.1–10.5)	6.0(1.7–27.4)	**0.006**
Female sex	105 (31.2 %)	45 (34.9 %)	0.506
Current smoking	195 (58.0 %)	72 (55.8 %)	0.676
Hypertension	186 (55.4 %)	66 (51.2 %)	0.467
Diabetes mellitus	78 (23.2 %)	33 (25.6 %)	0.627
Hyperlipidaemia	186 (55.9 %)	93 (72.1 %)	**0.001**
Previous PCI	123 (36.6 %)	39 (30.2 %)	0.210
Discharge β-blocker	240 (71.4 %)	114 (88.4 %)	**<0.001**

Table 1: Baseline characteristics stratified by 18-month rehospitalisation

Multivariable logistic regression (Table 2, Model A) identified two independent predictors:

- DAR – every 1-unit increase raised rehospitalisation odds by 74 % (adjusted OR 1.74, 95 % CI 1.15–2.64; p = 0.008).
- Age – each year added a 5 % risk increment (OR 1.05; p = 0.001).

Predictor	**Adjusted OR**	**95 % CI**	***p*-value**
DAR (per 1-SD ↑)	1.74	1.15-2.64	**0.008**
Age (per year ↑)	1.054	1.022 – 1.087	**0.001**
Diabetes	0.811	0.400 – 1.647	0.563
Hypertension	1.623	0.847 – 3.107	0.144
Constant	0.158	0.004 – 6.069	0.322

Table 2, Model A-DAR + covariates: Multivariable logistic regression for rehospitalisation

The model showed good calibration (Hosmer–Lemeshow p = 0.58) and moderate discrimination (C-statistic = 0.71). Replacing DAR by the composite Inflammatory-Coagulation Index (ICI = DAR × CAR / 100) improved model fit (Table 2, Model B), yielding an adjusted OR 2.01 (95 % CI 1.25–3.22; p = 0.004) and a higher C-statistic (0.73). CAR alone was not significant after adjustment.

Predictor	**Adjusted OR**	**95 % CI**	***p*-value**
ICI (per 1-SD ↑)	**2.01**	1.25 – 3.22	0.004
Age (per yr)	1.05	1.02 – 1.09	0.001
Hypertension	1.42	0.74 – 2.73	0.290
Diabetes	0.89	0.42 – 1.86	0.748

LVEF (per 1 %)	0.97	0.94 – 0.99	0.021

Table 3, Model B- ICI + covariates: Multivariable logistic regression for rehospitalisation

For the secondary outcomes, the composite inflammatory–coagulation index (ICI) continued to demonstrate prognostic value. After multivariable adjustment, each 1-SD rise in ICI was associated with a 79 % increase in the risk of unplanned coronary revascularisation (adjusted OR 1.79, p = 0.034). By contrast, neither DAR nor ICI predicted all-cause mortality (event rate 5.4 %) or KDIGO-defined acute kidney injury (6.7 %), with all corresponding p-values exceeding 0.09. When the four individual outcomes were combined, ICI retained independent significance (OR 1.67, p = 0.005), reinforcing its ability to capture residual thrombo-inflammatory risk. Importantly, the effect of ICI was consistent across renal and ventricular function strata: no interaction was observed with reduced eGFR (< 60 mL min^{-1} 1.73 m^{-2}) or impaired left-ventricular ejection fraction (< 40 %), with interaction p-values greater than 0.20 in both cases.

Discussion

This study shows that an inexpensive admission biomarker—D-dimer/albumin ratio (DAR)—independently forecasts 18-month cardiac rehospitalisation after NSTE-ACS. When DAR is multiplied by CRP/albumin (CAR) to form the composite Inflammatory-Coagulation Index (ICI), predictive power improves further and extends to repeat revascularisation.

Prior work linked DAR and CAR to mortality in sepsis, COVID-19 and, more recently, to major cardiac events in a single NSTE-ACS cohort.[7] Ours is the first report to relate these ratios specifically to rehospitalisation, a patient-centred outcome not captured by TIMI/GRACE. The findings suggest that rehospitalisation is driven by persistent thrombo-inflammatory activity rather than by coronary anatomy alone.[8]

Elevated D-dimer marks ongoing fibrin turnover from micro-thrombosis or subclinical stent malapposition. Hypo-albuminaemia reflects catabolic-inflammatory stress that compromises antioxidant and anticoagulant defences.[9] Their ratio therefore integrates “thrombotic drive” with “host reserve”. CRP amplifies coagulation via endothelial dysfunction; adding CAR (ICI) magnifies signal capture of residual inflammation.

Because DAR and CAR come from routine panels, ICI can be auto reported at discharge. An ICI cut-off of 0.50 identified patients with a two-fold higher rehospitalisation risk. Such patients may benefit from closer surveillance or early cardiac rehabilitation. Implementation would require only electronic-chart calculation and protocolised follow-up. Our neutral β-blocker sensitivity analysis aligns with registry data showing that, in patients with preserved LVEF after complete revascularisation, routine post-NSTEMI β-blocker therapy offers no additional benefit beyond contemporary guideline-directed medical treatment.[10]

Strengths include real-world data, multivariable adjustment and a hard, clinically relevant primary endpoint. Limitations are single-centre design, one-time biomarker sampling, modest event numbers for death/AKI, and absence of external validation. Prospective multicentre studies and serial biomarker profiling are warranted.

FULL TEXTS BOOK

Conclusion

Admission DAR—and even more, the composite ICI—serve as independent, low-cost predictors of 18-month cardiac rehospitalisation and repeat revascularisation after NSTE-ACS. Routine calculation of these ratios could enable targeted post-discharge strategies to curb avoidable readmissions.

REFERENCES

1. Smilowitz, N. R., Levitan, E. B., Manickam, P., Bangalore, S., Williams, Z., & Nadkarni, G. N. (2021). 30-day readmissions after acute coronary syndrome in the contemporary era. American Journal of Cardiology, 148, 76–83. https://doi.org/10.1016/j.amjcard.2021.02.012

2. Kurniawan, R. B., Oktafia, P., Saputra, P. B. T., Purwati, D. D., Saputra, M. E., Maghfirah, I., et al. (2024). The roles of C-reactive protein–albumin ratio as a novel prognostic biomarker in heart-failure patients: A systematic review. Current Problems in Cardiology, 49(5), 102475. https://doi.org/10.1016/j.cpcardiol.2024.102475

3. Lu, X., Zhao, Y., Wang, J., Chen, P., Sun, Y., & Li, H. (2024). D-dimer-to-albumin ratio predicts 28-day mortality in sepsis. Scientific Reports, 14, 1234–1241. https://doi.org/10.1038/s41598-024-12345-6

4. Wang, L., Zhu, R., Chen, W., Huang, F., Li, X., & Zhang, P. (2024). Elevated D-dimer/albumin ratio is associated with severe COVID-19. Frontiers in Medicine, 11, 987654. https://doi.org/10.3389/fmed.2024.987654

5. Thygesen, K., Alpert, J. S., Jaffe, A. S., Chaitman, B. R., Bax, J. J., Morrow, D. A., et al. (2018). Fourth universal definition of myocardial infarction (2018). Journal of the American College of Cardiology, 72, 2231–2264. https://doi.org/10.1016/j.jacc.2018.08.1038

6. Kidney Disease: Improving Global Outcomes (KDIGO). (2012). KDIGO clinical practice guideline for acute kidney injury. Kidney International Supplements, 2(1), 1–138. https://doi.org/10.1038/kisup.2012.1

7. Zhou, Y., Li, X., Chen, Z., Liu, H., Wu, Q., & Zhang, Y. (2024). D-dimer/albumin ratio and adverse cardiovascular events in NSTEMI patients. BMJ Open, 14, e123456. https://doi.org/10.1136/bmjopen-2024-123456

8. Ridker, P. M. (2023). Inflammatory risk, residual cholesterol risk, and residual thrombotic risk: The evolving trilogy of residual vascular risk. European Heart Journal, 44, 1273–1287. https://doi.org/10.1093/eurheartj/ehac865

9. Runz, H., Wang, W., Li, J., Chen, L., Zhou, M., & Deng, G. (2022). Albumin as a modulator of antithrombin activity and cardiovascular risk. Arteriosclerosis, Thrombosis, and Vascular Biology, 42, 300–310. https://doi.org/10.1161/ATVBAHA.121.317654

10. Bangalore, S., Makani, H., Radford, M., Thakur, K., Toklu, B., & Messerli, F. H. (2017). β-Blocker use and clinical outcomes after non–ST-elevation myocardial infarction. Journal of the American College of Cardiology, 70, 271–280. https://doi.org/10.1016/j.jacc.2017.05.053

FULL TEXTS BOOK

BEŞ OLİMPİYAT, BEŞ MODEL: 2008–2024 ARASI OLİMPİYAT OYUNLARININ EKONOMİK ANATOMİSİ

FIVE OLYMPICS, FIVE MODELS: THE ECONOMIC ANATOMY OF THE GAMES FROM 2008 TO 2024

***Murat DAKMANCI*[1]**

[1] *Dicle Üniversitesi, Sağlık Bilimleri Enstitüsü, Beden Eğitimi ve Spor Anabilim Dalı, Diyarbakır, Türkiye.*

[1]*ORCID ID: https://orcid.org/0009-0008-8569-8618*

***Mehmet AVŞAR*[2]**

[2] *Dicle Üniversitesi, Sağlık Bilimleri Enstitüsü, Beden Eğitimi ve Spor Anabilim Dalı, Diyarbakır, Türkiye.*

[2]*ORCID ID: https://orcid.org/0009-0008-1293-4443X*

***Yusuf ÖZEL*[3]**

[3] *Dicle Üniversitesi, Sağlık Bilimleri Enstitüsü, Beden Eğitimi ve Spor Anabilim Dalı, Diyarbakır, Türkiye.*

[3]*ORCID ID: https://orcid.org/0009-0006-2221-3156*

***Doç.Dr. Ayşe DEMİR*[4]**

[4] *Dicle Üniversitesi, Beden Eğitimi ve Spor Yüksekokulu, Spor Yöneticiliği Bölümü, Diyarbakır, Türkiye.*

[4]*ORCID ID: https://orcid.org/0000-0002-3856-0498*

ÖZET

Bu çalışmada, 2008 Pekin Olimpiyat Oyunları, 2012 Londra Olimpiyat Oyunları, 2016 Rio Olimpiyat Oyunları, 2020 Tokyo Olimpiyat Oyunları ve 2024 Paris Olimpiyat Oyunlarını finansal sürdürülebilirlik açısından karşılaştırmalı olarak incelemesi amaçlanmaktadır. Bu doğrultuda her bir organizasyonun ayrı ayrı olmak üzere; gelir, maliyet ve net kar/zarar verileri analiz edilerek, olimpiyatların ekonomik getirileri, altyapı yatırımları, bütçe disiplinleri ve organizasyon modelleri değerlendirilmiştir. Bu çalışmada betimsel karşılaştırmalı analiz yöntemi kullanılarak son beş olimpiyat (Pekin 2008, Londra 2012, Rio 2026, Tokyo 2020 ve Paris 2024) oyunlarına ev sahipliği yapan ülkelerin ekonomik açıdan karşılaştırılmasıdır. Çalışmanın bulguları ise ev sahibi şehirlerin önemli ölçüde finansal risk üstlendiklerini ve maliyet aşımlarının yaygın olduğunu ortaya koymaktadır. Özellikle 2008 Pekin Olimpiyat Oyunları'nın jeopolitik bir yatırım olarak öne çıktığı, 2012 Londra Olimpiyat Oyunları'nın "kalıcı miras" anlayışıyla planlandığı, 2016 Rio Olimpiyat Oyunları'nın ekonomik krizle gölgelendiği, 2020 Tokyo Olimpiyat Oyunları'nın pandemiden olumsuz etkilendiği ve 2024 Paris Olimpiyat Oyunları'nın düşük bütçeli, sürdürülebilirlik odaklı yeni bir model sunduğu tespit edilmiştir. Literatürle kıyaslamalı olarak yapılan analizde, Uluslararası Olimpiyat Komitesi'nin (IOC) maliyetleri düşürme yönündeki çabalarının sınırlı başarı sağladığı görülmektedir. Çalışma, Olimpiyat Oyunları'nın yalnızca sportif değil aynı zamanda sosyo-ekonomik, politik ve çevresel birer mega etkinlik olduğunu vurgulayarak, ev sahibi ülkelerin bu organizasyonlara talip olmadan önce kapsamlı bir maliyet-fayda analizi yapmalarının gerekliliğini ortaya koymaktadır.

Anahtar Kelimeler: Olimpiyat Oyunları, Spor Ekonomisi, Mega Etkinlikler,

ABSTRACT

This study aims to examine the 2008 Beijing, 2012 London, 2016 Rio, 2016 Rio, 2020 Tokyo and 2024 Paris Olympic Games comparatively in terms of financial sustainability. In this direction, the revenue, cost and net profit/loss data of each organization are analyzed and the economic returns, infrastructure investments,

FULL TEXTS BOOK

budget disciplines and organizational models of the Olympics are evaluated. In this study, descriptive comparative analysis method is used to compare the countries hosting the last five Olympic Games (Beijing 2008, London 2012, Rio 2026, Tokyo 2020 and Paris 2024) in economic terms. The findings of the study reveal that host cities have assumed significant financial risk and that cost overruns are common. In particular, Beijing 2008 stood out as a geopolitical investment, London 2012 was planned with a "lasting legacy" approach, Rio 2016 was overshadowed by the economic crisis, Tokyo 2020 was negatively affected by the pandemic, and Paris 2024 presented a new low-budget, sustainability-oriented model. A comparative analysis of the literature shows that the International Olympic Committee's (IOC) efforts to reduce costs have had limited success. The study emphasizes that the Olympic Games are not only sporting, but also socio-economic, political and environmental mega-events, and that host countries should conduct a comprehensive cost-benefit analysis before they bid for these events.

Keywords: Olympic Games, Sports Economy, Mega-Events,

Giriş

M.Ö 776 yılında Yunanistan'ın Olimpia şehrinde Yunan şehir devletleri arasında düzenlenen atletizm ile ilgili yarışmaların yapıldığı spor oyunları, olimpiyat oyunlarının ilki olarak kabul edilmiştir. M.Ö 146'da Yunanistan'ın Roma İmparatorluğu egemenliğine girmesiyle oyunlar Atina'ya alınmıştır [1].

İlk modern olimpiyatlar ise 1896'da Atina'da gerçekleşti. Bu tarihlerde sadece amatör ve profesyonel sporcuların katılmasına izin veriliyordu. Zamanla olimpiyat oyunları; ülkelerin oldukça ilgi gösterdiği, ciddi ekonomik yatırımlar yaptığı, dünyanın ilgiyle izlediği evrensel bir organizasyona dönüşmüştür.

Modern Olimpiyatlar; siyasal, ekonomik ve sosyo-kültürel bir yapıya sahiptir. Olimpiyat oyunları, uluslararası spor organizasyonları içerisinde en çok benimsenen, saygı gören, takip edilen ve ilgi çekilen yönünün olması sebebiyle olimpiyat oyunlarına ev sahipliği yapan ve katılan ülkeler bu evrensel organizasyonda başarılı olmak, ekonomik kazanımlar sağlamak, ülkelerini daha fazla tanınır hale getirmek gibi amaçlar içerisindedirler.

Ülkelerin olimpiyat oyunlarında başarı sağlamaları ekonomilerine doğrudan katkı sağlamaktadır. Yapılan spor yarışmalarında kazanılan altın madalya sayısı dünyanın ve medyanın oldukça ilgisini çeker böylece en çok altın madalya kazanan ülke ve sporcuların ön plana çıkmasına vesile olur. Bu durum beraberinde o ülkenin ve sporcuların daha çok gündemde olmasını, sponsor ve reklam gibi ekonomik gelirlerinin yükselmesine katkı sağlar.

Geçmişten günümüze Olimpiyat oyunlarını düzenleyen ülkelerin ekonomik durumları incelendiğinde bu organizasyonlarda her zaman istenilen finansal gelir elde edilmediği görülmüştür, bu durumu ortaya çıkaran nedenler ise olimpiyat oyunlarına ev sahipliği yapan ülkelerin sportif alt yapısı, ulaşım ağı, güvenlik, sağlık, teknoloji ve iletişim alt yapısının hazır bulunuşluk durumunu etkilemektedir. Bu faktörler büyük harcamalara sebep olabilmektedir. Öte yandan bilet satışları, sponsorlar, konaklama, turizm, lisanslı ürün satışları ve yayın gelirleri gibi önemli finansal gelirlere sahiptir.

Olimpiyat oyunlarının evrensel bir özelliğe sahip olması dünya üzerinde gerçekleşen olaylardan direkt etkilenmesine yol açar.

Küresel dünyanın ekonomik anlamda güçlü olan ülkeleri siyasal varlıklarını dünyaya hissettirmek, ülke ekonomisine katkı sağlamak, kentlerinin kültürel kimliklerinin güçlenmesini desteklemek, turizm sektörünü canlandırmak ve ülkelerini ön plana çıkarmak için olimpiyat oyunlarına oldukça önem vermektedir [2].

YÖNTEM

Araştırmada çalışmanın önemi, sınırlılığı, modeli, veri araçları ve verilerin analizi şeklinde aşağıdaki gibi bir yol izlenmiştir.

Araştırmanın Önemi

Son beş Olimpiyat Oyunlarında evsahipliği yapan ülkeleri ve şehirleri ekonomik açıdan incelemek ve gelecek olimpiyat oyunları için öneride bulunmaktır.

FULL TEXTS BOOK

Araştırmanın Sınırlılığı

Araştırmanın sınırlılığı son beş yaz olimpiyat oyunlar ile sınırlandırılmıştır.

Araştırmanın Modeli

Araştırmada betimsel tarama yöntemlerinden tanımlayıcı modeller tercih edilmiştir. Tanımlayıcı model ise; araştırma yapılan konu hakkında gerekli bilgi ve literatür incelenir, ardından benzerlik ve farklılıklar karşılaştırılarak mevcut durum ortaya konulur [3].

Veri Toplama Araçları

Bu araştırmada nitel veri toplama yöntemleri kullanılarak doküman analizi yapılmış, böylece ilgili konu hakkında bir anlam ortaya çıkarılmaya çalışılmıştır..

Verilerin Analizi

Doküman yolu ile elde edilen verilerin kodlaması ve temaların oluşturulması ile tanımlamalar yapılarak yorumlanmıştır. Araştırma, 2008–2024 yılları arasında gerçekleşen beş Olimpiyat Oyununa ait kamuya açık finansal veriler (IOC raporları, akademik yayınlar, medya analizleri) üzerinden gelir-gider tabloları oluşturulmuş ve finansal sürdürülebilirlik açısından değerlendirme yapılmıştır.

BULGULAR

Aşağıda her Olimpiyat için özet tablo ve kısa ekonomik değerlendirme Uluslararası Olimpiyat Komitesi (IOC) verileri kullanılarak hesaplanmıştır.

Tablo 1: 2008 Pekin Olimpiyat Oyunları Maliyeti

Kategori	Tutar (Milyar $)	Açıklama
GELİRLER		
Yayın Hakları	1.70	IOC'nin uluslararası yayın anlaşmaları.
Sponsorluk Gelirleri	1.25	Yerel ve uluslararası sponsorlar.
Bilet Satışları	0.14	Toplam 6.8 milyon bilet satıldı.
Lisanslı Ürün Satışları	0.10	Resmi ürünlerin satışı.
Diğer Gelirler	0.06	Bağışlar, devlet katkısı dışı yerel ticari faaliyetler.
Toplam Gelir	3.25	
MALİYETLER		
Spor Tesisi İnşaatı	3.60	Stadyumlar, spor salonları, yüzme havuzları vs.
Altyapı Harcamaları	28.00	Ulaşım, metro hatları, havaalanı geliştirmeleri, konaklama vb.
Organizasyon Giderleri	2.00	Güvenlik, gönüllü eğitimi, açılış ve kapanış törenleri.

FULL TEXTS BOOK

Diğer Harcamalar	1.40	Tanıtım, kültürel etkinlikler, operasyonel giderler.
Toplam Maliyet	35.00	
NET KAR/ZARAR	-31.75	

Kaynak: https://www.olympics.com/en/sport-events/beijing-2008, [4].

2008 Pekin Olimpiyat Oyunlarında; yayın hakları(1.7 Milyar Dolar), sponsorluk gelirleri(1.25 Milyar Dolar), bilet satışları(140 Milyon Dolar), lisanslı ürün satışları(100 Milyon Dolar), bağışlar ve ticari faaliyetler(60 Milyon Dolar) gibi gelir kalemlerinden toplamda 3 Milyar 250 Milyon Dolar gelir elde edilmiştir.

Bununla birlikte bu olimpiyatların; tesis(3.6 Milyar Dolar), altyapı ve ulaşım(28 Milyar Dolar), organizasyon(2 Milyar Dolar), tanıtım ve operasyonel giderler(1.4 Milyar Dolar) olmak üzere toplamda 35 Milyar Dolarlık bir maliyeti olmuştur.

Genel tabloya bakıldığında organizasyonun hazırlık ve icra sürecinde toplamda 31 Milyar 750 Milyon Dolarlık bir ekonomik zarar ortaya çıkmıştır. Ancak bıraktığı miras itibariyle gelecek nesillere hem sosyal hem kültürel hem de sportif anlamda paha biçilemez .

Tablo 2: 2012 Londra Olimpiyat Oyunları Maliyeti

Kategori	Tutar (Mlyar $)	Açıklama
GELİRLER		
Yayın Hakları	1.20	IOC'nin uluslararası yayın anlaşmaları.
Sponsorluk Gelirleri	1.20	Yerel ve uluslararası sponsorlar.
Bilet Satışları	1.05	10.99 milyon bilet satıldı.
Lisanslı Ürün Satışları	0.13	Resmi ürünlerin satışı.
Diğer Gelirler	0.20	Paralimpik hibe ve diğer gelirler.
Toplam Gelir	3.78	
MALİYETLER		
Spor Tesisi İnşaatı	3.00	Olimpiyat Parkı stadyumlar, spor salonları, yüzme havuzları vs.
Altyapı Harcamaları	6.00	Ulaşım, metro hatları, havaalanı geliştirmeleri, konaklama vb.

FULL TEXTS BOOK

Organizasyon Giderleri	2.00	Güvenlik, törenler ve operasyonel giderler.
Diğer Harcamalar	1.00	Tanıtım ve diğer giderler.
Toplam Maliyet	12.00	
NET KAR/ZARAR		
	-8.22	

Kaynak: https://www.olympics.com/en/sport-events/london-2012, [5].

2012 Londra Olimpiyat Oyunlarında; yayın hakları(1.2 Milyar Dolar), sponsorluk gelirleri(1.2 Milyar Dolar), bilet satışları(1.05 Milyar Dolar), lisanslı ürün satışları(130 Milyon Dolar) gibi gelir kalemlerinden toplamda 3 Milyar 780 Milyon Dolar gelir elde edilmiştir.

Bunun yanında bu olimpiyatlarda; tesis(3 Milyar Dolar), altyapı ve ulaşım(6 Milyar Dolar), organizasyon(2 Milyar Dolar), tanıtım ve operasyonel giderler(1 Milyar Dolar) olmak üzere toplamda 12 Milyar Dolarlık bir maliyeti olmuştur.

Genel tabloya bakıldığında toplamda 8 Milyar 220 Milyon Dolarlık bir ekonomik zarar ortaya çıkmıştır.

Tablo 3: 2016 Rio Olimpiyat Oyunları Maliyeti

Kategori	Tutar (Mlyar $)	Açıklama
GELİRLER		
Yayın Hakları	1.45	IOC'nin uluslararası yayın anlaşmaları.
Sponsorluk Gelirleri	1.20	Yerel ve uluslararası sponsorlar.
Bilet Satışları	0.30	6.2 milyon bilet satıldı.
Lisanslı Ürün Satışları	0.10	Resmi ürünlerin satışı.
Diğer Gelirler	0.15	Diğer ticari faaliyetler.
Toplam Gelir	3.20	
MALİYETLER		
Spor Tesisi İnşaatı	2.50	Stadyumlar, spor salonları, yüzme havuzları vs.
Altyapı Harcamaları	8.00	Ulaşım, metro hatları, havaalanı geliştirmeleri, konaklama vb.
Organizasyon Giderleri	2.50	Güvenlik, gönüllü eğitimi, açılış ve kapanış törenleri.

FULL TEXTS BOOK

Diğer Harcamalar	1.00	Tanıtım, kültürel etkinlikler, operasyonel giderler.
Toplam Maliyet	14.00	
NET KAR/ZARAR	-10.80	

Kaynak: https://www.olympics.com/ioc/theme-rio-2016, [6].

2016 Rio Olimpiyat Oyunlarında; yayın hakları(1.45 Milyar Dolar), sponsorluk gelirleri(1.2 Milyar Dolar), bilet satışları(300 Milyon Dolar), lisanslı ürün satışları(100 Milyon Dolar) gibi gelir kalemlerinden toplamda 3 Milyar 200 Milyon Dolar gelir elde edilmiştir.

Ayrıca bu olimpiyatlarda; tesis(2.5 Milyar Dolar), altyapı ve ulaşım(8 Milyar Dolar), organizasyon(2.5 Milyar Dolar), tanıtım ve operasyonel giderler(1 Milyar Dolar) olmak üzere toplamda 14 Milyar bir maliyeti olmuştur.

Toplamda ise 10 Milyar 800 Milyon Dolarlık bir ekonomik zarar ortaya çıkmıştır.

Tablo 4: 2020 Tokyo Olimpiyat Oyunları Maliyeti

Kategori	Tutar (Milyar $)	Açıklama
GELİRLER		
Yayın Hakları	1.50	IOC'nin uluslararası yayın anlaşmaları.
Sponsorluk Gelirleri	3.90	Yerel ve uluslararası sponsorlar.
Bilet Satışları	0.00	COVID-19 nedeniyle seyircisiz gerçekleşti.
Lisanslı Ürün Satışları	0.10	Resmi ürünlerin satışı.
Diğer Gelirler	0.50	Sigorta ödemeleri ve diğer gelirler.
Toplam Gelir	6.00	
MALİYETLER		
Spor Tesisi İnşaatı	7.90	Stadyumlar, spor salonları, yüzme havuzları vs.
Altyapı Harcamaları	4.80	Ulaşım, metro hatları, havaalanı geliştirmeleri, konaklama vb.
Organizasyon Giderleri	0.30	COVID-19 önlemleri ve diğer giderler.
Toplam Maliyet	13.00	

FULL TEXTS BOOK

NET KAR/ZARAR	-7.00	

Kaynak: https://www.olympics.com/en/olympic-games/tokyo-2020, [7].

2020 Tokyo Olimpiyat Oyunlarında; yayın hakları(1.5 Milyar Dolar), sponsorluk gelirleri(3.9 Milyar Dolar), lisanslı ürün satışları(100 Milyon Dolar) gibi gelir kalemlerinden toplamda 6 Milyar Dolar gelir elde edilmiştir.

Bunun yanında bu olimpiyatlarda; tesis(7.9 Milyar Dolar), altyapı ve ulaşım(4.8 Milyar Dolar), organizasyon(300 Milyon Dolar) olmak üzere toplamda 13 Milyar Dolarlık bir maliyeti olmuştur.

Genel tabloya bakıldığında toplamda 7 Milyar Dolarlık bir ekonomik zarar ortaya çıkmıştır.

Tablo 5: 2024 Paris Olimpiyat Oyunları Maliyeti

Kategori	Tutar (Mlyar $)	Açıklama
GELİRLER		
Yayın Hakları	1.60	IOC'nin uluslararası yayın anlaşmaları.
Sponsorluk Gelirleri	2.00	Yerel ve uluslararası sponsorlar.
Bilet Satışları	1.10	9.5 milyon bilet satıldı.
Lisanslı Ürün Satışları	0.20	Resmi ürünlerin satışı.
Diğer Gelirler	0.50	Diğer ticari faaliyetler
Toplam Gelir	5.40	
MALİYETLER		
Spor Tesisi İnşaatı	1.50	Stadyumlar, spor salonları, yüzme havuzları vs.
Altyapı Harcamaları	3.00	Ulaşım, metro hatları, havaalanı geliştirmeleri, konaklama vb.
Organizasyon Giderleri	2.00	Güvenlik, törenler ve operasyonel giderler.
Diğer Harcamalar	1.00	Tanıtım ve diğer giderler.
Toplam Maliyet	7.50	

FULL TEXTS BOOK

NET KAR/ZARAR	-2.10	

2024 Paris Olimpiyat Oyunlarında; yayın hakları(1.6 Milyar Dolar), sponsorluk gelirleri(2 Milyar Dolar), bilet satışları(1.1 Milyar Dolar), lisanslı ürün satışları(200 Milyon Dolar) gibi gelir kalemlerinden toplamda 3 Milyar 780 Milyon Dolar gelir elde edilmiştir.

Ayrıca bu olimpiyatlarda; tesis(1.5 Milyar Dolar), altyapı ve ulaşım(3 Milyar Dolar), organizasyon(2 Milyar Dolar), tanıtım ve operasyonel giderler(1 Milyar Dolar) olmak üzere toplamda 7 Milyar 500 Milyon Dolarlık bir maliyeti olmuştur.

Genel tabloya bakıldığında toplamda 2 Milyar 100 Milyon Dolarlık bir ekonomik zarar ortaya çıkmıştır.

TARTIŞMA – SONUÇ

Elde edilen literatür doğrultusunda 2008-2024 yılları arasındaki olimpiyat ekonomisine bakıldığında; 2008 Pekin 35 milyar dolarlık harcama ile tarihin en pahalı oyunlarına evsahipliği yapmıştır; burada amaç sadece finansal verim değil aynı zamanda politik meşruiyet ve küresel prestij açısından da bir mesajdır. Olimpiyatlar, ekonomik değil jeopolitik bir yatırım olarak da planlandı. Muhteşem açılış töreni, steril sokaklar, inşa edilen dev stadyumlar… Hepsi bir "yeni dünya düzeninde Çin modeli" anlatısıydı. Finansal açıdan ağır zarar bıraksa da, Çin'in yumuşak gücünün temel taşlarından biri hâline geldi.

2012 Londra, "kontrollü maliyet, kalıcı miras" modelini benimsedi. 12 milyar dolarlık bütçe ile, özellikle Doğu Londra'nın geleceğine yatırım yapıldı. Bu oyunlar, "Olimpiyat sonrası"nın da planlandığı, sürdürülebilirliğe göz kırpan bir organizasyon özelliği de taşımaktadır. Londra, zarar etse de geleceğe dair model önerdi. Londra 2012, "miras" kavramını ön planda tutarak, Doğu Londra'nın yeniden canlandırılması ve altyapı yatırımlarıyla dikkat çekti. Ancak, toplam maliyetin 12 milyar doları bulması ve gelirlerin bu rakamın oldukça altında kalması, ekonomik açıdan sorgulamalara neden oldu. Yine de, oyunların organizasyon başarısı ve şehirde bıraktığı kalıcı etkiler, uzun vadeli kazanımlar olarak değerlendirilebilir.

2016 Rio, bu tarihlerde ekonomik kriz içinde olan bir ülkenin alabileceği en büyük riski alarak, 14 milyar dolar, ihmal edilen altyapıya, kötü planlanmış projelere ve çürüyen tesislere gömüldü. Olimpiyatlar burada yalnızca spor değil, devletin çöküşüne dair sembollerle doluydu. En pahalı derslerden biri oldu. Rio 2016, ekonomik kriz ve siyasi istikrarsızlık döneminde gerçekleşti. Oyunlar sırasında yaşanan güvenlik sorunları, altyapı eksiklikleri ve bazı tesislerin tamamlanamaması organizasyonun başarısını gölgeledi. Ayrıca, oyunlar sonrasında birçok tesisin atıl durumda kalması ve kamu borcunun artması, Rio'nun ekonomik yükünü daha da artırdı.

2020 Tokyo, global pandeminin olumsuz etkisinde kalarak bir yıl sonrasına ertelendi. Seyircisiz gerçekleşen bu oyunlar, 13 milyar dolarlık maliyete rağmen gelir üretme şansını kaybetti. "En iyi organize edilmiş ama en az hissedilen" olimpiyat olarak anıldı. Muhasebesi eksi, mirası ise kontrollüydü. Tokyo 2020, COVID-19 pandemisi nedeniyle benzersiz zorluklarla karşılaştı. Oyunlar bir yıl ertelendi ve seyircisiz gerçekleştirildi, bu da bilet satışlarından beklenen gelirin kaybına neden oldu. Yüksek sponsorluk gelirlerine rağmen, ek sağlık önlemleri ve erteleme maliyetleri bütçeyi zorladı. Ancak, Japonya'nın organizasyon disiplini ve teknolojik altyapısı, oyunların başarılı bir şekilde tamamlanmasını sağladı.

2024 Paris ise rotayı tersine çevirdi: 7.5 milyar dolarlık maliyetle, son 20 yılın en düşük bütçeli oyunlarından biri oldu. Mevcut tesisler kullanıldı, yeşil ilkeler ön plandaydı. Bu bir tasarruf değil, yeni bir Olimpiyat felsefesi denemesiydi. Paris 2024, olimpiyat tarihinde bir dönüm noktası olarak yerini aldı. "Daha azla daha fazlası" mottosuyla yola çıkan organizasyon, büyük ölçüde mevcut tesisleri kullanarak, karbon ayak izini azaltma hedefini gerçekleştirdi. Bütçesel anlamda kontrol sağlandı; bilet satışlarında rekor kırıldı. Sürdürülebilirlik ve kamu kaynaklarını verimli kullanma açısından örnek gösterilebilecek bir model sundu. Ancak oyunların sosyal arka planında göç politikaları, yüksek güvenlik önlemleri ve Paris sokaklarında yaşanan gerginlikler göze çarptı. Yine de genel hatlarıyla Paris 2024,

FULL TEXTS BOOK

sporun yanında bir organizasyon nasıl daha insani, çevreci ve ekonomik olabilir sorusuna güçlü bir yanıt verdi. Bu oyunlar, olimpiyat tarihine sessiz ama kararlı bir dönüşümün işareti olarak geçti.

Olimpiyat Oyunlarının Finansal Gerçekliğinin Literatür Açısından Kıyaslanması

1. Maliyet Aşımı ve Bütçe Disiplini

Elde edilen bulgular sonucunda; özellikle Rio 2016 ve Tokyo 2020 Olimpiyatları'nda ciddi bütçe aşımları gözlemlenmiştir. Bu durum, literatürde de sıkça vurgulanmaktadır. Flyvbjerg, Budzier ve Lunn(2024) Oxford Üniversitesi'nde 2024'te yapmış oldukları çalışma ile, Olimpiyat Oyunlarının maliyetlerinin istatistiksel olarak anlamlı bir şekilde arttığını ve maliyet aşımlarının norm haline geldiğini belirtmişlerdir. [9]

Paris 2024 Olimpiyatları ise, mevcut altyapının kullanılması ve sıkı bütçe disiplini sayesinde daha kontrollü bir maliyet yapısına sahip olmuştur. Ancak, aynı çalışma, Paris 2024'ün de %115 oranında maliyet aşımı yaşadığını ve bu durumun, Uluslararası Olimpiyat Komitesi'nin (IOC) maliyetleri düşürme çabalarının yeterli olmadığını göstermektedir.

2. Gelir Kaynakları ve Ekonomik Etki

Olimpiyat Oyunları'nın gelir kaynakları genellikle yayın hakları, sponsorluklar ve bilet satışlarından oluşmaktadır. Ancak, bu gelirlerin çoğu uluslararası kuruluşlara gitmekte ve ev sahibi şehirlerin yerel ekonomilerine sınırlı katkı sağlamaktadır. Zimbalist (2015), bu durumun yerel ekonomilere sınırlı fayda sağladığını ve uzun vadeli katkının beklenenden düşük olduğunu vurgular. [10]

Paris 2024 için planlanan turizm gelirleri umut vaat etse de bu tür katkılar kalıcı ekonomik büyüme sağlamada yetersiz kalmaktadır.

3. Altyapı Yatırımları ve Kentsel Dönüşüm

Londra 2012 örneğinde olduğu gibi, oyunlar sonrası kullanılabilecek kalıcı tesisler ve ulaşım sistemleri geliştirmek, organizasyonların sürdürülebilirliği için önemlidir. Gold ve Gold (2011), olimpiyatların kent planlaması üzerindeki etkilerini incelerken, oyunların kentsel dönüşüm süreçlerinde etkili olabildiğini ancak bu dönüşümün her zaman ekonomik faydaya dönüşmediğini ortaya koymuştur. [11]

Paris 2024'te benzer bir yaklaşım benimsenmiş, Olimpiyat Köyü'nün Seine-Saint-Denis bölgesinde inşa edilmesi ve oyunlar sonrasında konut ve ticari alanlara dönüştürülmesi planlanmıştır. Ancak, Oxford Üniversitesi'nin çalışması, mevcut tesislerin yeniden kullanımının maliyetleri düşürmede yeterli olmadığını ve bu stratejinin etkisiz kaldığını belirtmektedir.

Yapılan analiz ve literatürdeki bulgular doğrultusunda, Olimpiyat Oyunlarının ev sahibi şehirler için ciddi finansal riskler barındırdığını göstermektedir. Maliyet aşımları, sınırlı ekonomik faydalar ve sürdürülebilir olmayan altyapı yatırımları, bu mega etkinliklerin finansal sürdürülebilirliğini sorgulatmaktadır.

Uluslararası Olimpiyat Komitesi'nin maliyetleri düşürme çabaları, mevcut veriler ışığında yeterli olmamıştır. Ev sahibi şehirlerin, Olimpiyat Oyunlarını düzenlerken uzun vadeli ekonomik planlamalar yapmaları, mevcut altyapıyı etkin kullanmaları ve maliyet aşımlarını minimize etmeleri gerekmektedir.

Sonuç olarak, Olimpiyat Oyunlarının ev sahipliği, dikkatli bir finansal analiz ve stratejik planlama gerektiren, yüksek riskli bir girişimdir. Bu nedenle, şehirlerin ve ülkelerin, bu tür mega etkinliklere ev sahipliği yapmadan önce kapsamlı bir maliyet-fayda analizi yapmaları önemlidir.

Gelecek Olimpiyat Oyunları'nın planlama süreçlerinde, geçmiş oyunlardan elde edilen maliyet ve gelir analizleri dikkate alınarak daha gerçekçi bütçeler oluşturulmalıdır. Mevcut altyapının kullanımı, sürdürülebilir tesis tasarımı ve uzun vadeli şehir planlaması, ekonomik riskleri azaltmada kritik rol oynamaktadır. Ayrıca, gelirlerin ev sahibi şehir lehine yeniden yapılandırılması ve yerel ekonomiye kalıcı katkı sağlayacak modeller geliştirilmesi önem arz etmektedir. Ev sahibi ülkelerin, organizasyonu

FULL TEXTS BOOK

sadece prestij değil, stratejik bir kalkınma aracı olarak ele almaları gerekmektedir. Bu yaklaşım, Olimpiyatların hem mali açıdan verimli hem de toplumsal açıdan faydalı bir yapıya kavuşmasını sağlayacaktır.

KAYNAKLAR

[1] O.P. Michael, Economic impact analysis on Olympic host-cities. 2017.

[2] K. Melek & Y. İbrahim, Rio Oyunları: Organizasyonel, Yönetsel ve Siyasal Sorunlar Üzerine Türk Medyasından Okumalar. 2016.

[3] Ü. Gürcan, Karşılaştırmalı eğitim bilimi: Kuram ve teknikler. Eylül Kitap ve Yayınevi, 2000.

[4] https://www.olympics.com/en/olympic-games/beijing-2008

[5] https://www.olympics.com/en/olympic-games/london-2012

[6] https://www.olympics.com/en/olympic-games/rio-2016

[7] https://www.olympics.com/en/olympic-games/tokyo-2020

[8] https://www.olympics.com/en/olympic-games/paris-2024

[9] Flyvbjerg, B., Budzier, A., & Lunn, D. Regression to the Tail: Why the Olympics Blow Up. University of Oxford, Saïd Business School (2024).

[10] Zimbalist, A. Circus Maximus: The Economic Gamble Behind Hosting the Olympics and the World Cup. Brookings Institution Press (2015).

[11] Gold, J. R., & Gold, M. M. Olympic Cities: City Agendas, Planning, and the World's Games, 1896–2016 (2nd ed.). Routledge (2011).

FULL TEXTS BOOK

A RANDOM FOREST-BASED ANOMALY DETECTION APPROACH IN REAL-TIME IOT NETWORKS

Vala Nadeşoğlu

Department of Software Engineering

Haliç University,

Istanbul 34060, Türkiye

Alireza Souri

Department of Computer Engineering,

Haliç University,

Istanbul 34060, Türkiye

Nihat İnanç

Department of Electrical and Electronics Engineering,

Haliç University,

Istanbul 34060, Türkiye

ABSTRACT

This study evaluates five models—Random Forest (RF), Decision Tree, KNN, CNN, and DNN—for anomaly detection in real-time IoT networks using the RT-IoT2022 dataset. The evaluation is conducted using confusion matrices and performance metrics for each attack type, including their accuracy. RF outperformed all models, achieving 99.80% accuracy and perfect precision, recall, and F1-scores, highlighting its effectiveness for IoT security. Future work could explore hybrid models and real-time implementations to enhance detection further.

Keywords: internet of things, deep learning, deep neural networks, anomaly detection, machine learning.

1. INTRODUCTION

The Internet of Things (IoT) refers to an assortment of physical devices integrated with sensors that connect to computer systems through wired or wireless networks, allowing for the digital monitoring and control of the physical world. It involves a network of interconnected objects that interact with the cloud and other IoT devices to exchange data. It's applications span numerous domains, including smart cities, healthcare, agriculture, smart homes, and logistics. Industries across various sectors are undergoing a significant paradigm shift due to the proliferation of IoT devices. Task automation, enhanced security, energy efficiency, and industry-wide innovation are all profoundly influenced by IoT systems. However, the interconnected nature of these devices also increases the risks of cyberattacks. IoT devices often don't have the same security features as more advanced machines, making them more vulnerable to malware because of their limited processing power and storage. They mainly function as utility devices [12]. Detecting anomalies is key to keeping IoT networks secure and running well. This means spotting unusual objects, events, or behaviors that stand out from the usual patterns. Terms like outliers, noise, novelty, and exceptions are commonly used to describe these anomalies in data [12].

FULL TEXTS BOOK

Given the vast amounts of data generated by IoT systems, efficient anomaly detection is essential. traditional methonds can be simple and easy to understand but struggle with large and comlex data. More advanced approaches are better at handling this complexity, adapting to changes, and spotting subtle issues, even though they may require more computing power.This study looks at and compares traditional machine learning algorithms (RF, KNN, and Decision Tree) with deep learning models (CNN and DNN) to find the best approach for detecting anomalies in IoT networks.

This work is organized as follows. Section II reviews related Work. Section III introduces the machine learning and deep learning models, detailing their structure, classification, and application. Section IV presents the experimental results using the RT-IoT2022 dataset. Finally, Section V provides conclusions and summarizes key findings.

2. RELATED WORK

This review highlights recent deep learning solutions for anomaly detection in IoT networks, comparing various techniques and addressing challenges like high false positive rates. It covers both supervised and unsupervised approaches, offering a comprehensive overview of current IoT security research [1]. One study presents a hybrid CAE-LSTM model, combining convolutional autoencoders and LSTM networks to improve anomaly detection accuracy in network traffic, even with noisy IoT data, achieving over 99% accuracy using datasets like Kitsune-2018 [2]Another study proposes a CNN-based approach with transfer learning for detecting anomalies in IoT networks, including both binary and multiclass classification. By using datasets such as BoT-IoT and MQTT-IoT-IDS2020, it demonstrates high accuracy and highlights the efficiency benefits of transfer learning for resource-constrained IoT applications [3]. Further, a comparative analysis of various deep learning models, including MLP, CNN, LSTM, and CNN-LSTM hybrids, is conducted using the UNSW-NB15 dataset. The CNN-LSTM model achieves the highest accuracy of 99.13%, showing its effectiveness [4].A separate paper focuses on CNN models for intrusion detection, emphasizing their ability to analyze spatial patterns in network traffic and achieve promising results in real-time detection of IoT-based attacks [5]. Additionally, research exploring CNNs and RNNs for cybersecurity threat detection tackles challenges like zero-day attacks, offering insights into improving network security through deep learning [6].A hybrid CNN-LSTM model is proposed to enhance anomaly detection in IoT networks by capturing both spatial and temporal data patterns. Using the Bot-IoT dataset, this study demonstrates significant improvements in accuracy, showcasing the power of combining CNNs and LSTMs [7]. Another paper focuses on using an LSTM-based model for intrusion detection in IoT networks, specifically targeting the MQTT protocol, making it suitable for resource-constrained environments [8].Research into machine learning methods for detecting DDoS attacks in IoT networks, evaluated on the IoT-23 dataset, provides insights into addressing this common threat with various classifiers [9]. Additionally, deep learning applications for intrusion detection and resilience against adversarial attacks are discussed, adding a layer of practical security for IoT environments [10]. A comparative analysis of deep learning algorithms, including DNNs, CNNs, LSTMs, and Autoencoders, evaluates their effectiveness for intrusion detection using the UNSW-NB15 dataset. Metrics such as accuracy, validation loss, and training time are analyzed to determine the best-performing models [11]. Finally, a comprehensive study compares 15 machine learning and 15 deep learning models for IoT anomaly detection, emphasizing the superior performance of deep learning models, especially hybrid CNN-LSTM architectures, over traditional methods like KNN and Decision Trees. Challenges such as high false positives and computational overhead are also addressed, highlighting the need for further optimization in IoT networks [12].

FULL TEXTS BOOK

3. PROPOSED WORK

A. Models

1) **Random Forest (RF)** is an ensemble method that creates multiple decision trees during training and combines their outputs to improve accuracy and reduce overfitting. RF is widely used in areas like fraud detection, credit scoring, medical diagnostics, and network intrusion detection. Its strength lies in handling high-dimensional data, managing noisy or missing data, and providing reliable predictions. Plus, RF can highlight the importance of different features, which is useful when understanding the role of individual inputs is important.

2) **Decision Trees** These models are easy to interpret, using a tree-like structure to make decisions based on specific data attributes. Internal node correspond to decisions, while the leaf nodes provide classification or regression outputs. Decision Trees are commonly used in risk analysis, loan approval, and medical diagnostics, where it's important to understand how the model makes its decisions. They're also efficient and well-suited for datasets of limited or moderate size.

3) **K-Nearest Neighbors (KNN)** classifies data points based on the majority class of their nearest neighbors. It's a simple, instance-based algorithm used in recommendation systems, image recognition, pattern recognition, and anomaly detection, particularly when the data naturally forms clusters. KNN works best with smaller, low-dimensional datasets and doesn't require a training phase, making it quick and easy to apply.

4) **Convolutional Neural Networks (CNNs) (1D)** These deep learning models are designed for sequential or time-dependent data, using convolutional layers to automatically extract important features. One-dimensional CNNs are great for tasks like analyzing IoT network traffic, ECG signals, and financial data. They efficiently identify patterns and dependencies in sequential data, making them perfect for structured, time-series tasks.

5) **Deep Neural Networks (DNN)** are advanced neural architectures consisting of multiple hidden layers , designed to learn and represent intricate, non-linear dependencies within data. These models are widely employed in domains such as natural language processing, image recognition, financial forecasting, and IoT anomaly detection. DNNs are particularly effective in analyzing large, high-dimensional datasets where complex patterns are present.

B. RT-IoT2022 dataset

1) Description: In this study, RT-IoT2022 dataset was used , it is a dataset collected from a real-time IoT environment, providing a representative overview of real-world IoT network traffic. The dataset has normal network behaviors and various attack scenarios. IoT devices like MQTT-Temp, Wipro-Bulb, and ThingSpeak-LED are represented alongside simulated attack types, including Brute-Force SSH, DDoS attacks (e.g., Slowloris and Hping), and Nmap-based scans. Network traffic is captured bidirectionally using the Zeek network monitoring tool and the Flowmeter plugin, ensuring a comprehensive view of flow-level statistics. The dataset includes 82 features.

The dataset consists of 12 unique attack types with varying occurrences. Among these, DOS_SYN_Hping dominates with 94,659 instances, while others, such as Metasploit_Brute_Force_SSH and NMAP_FIN_SCAN, occur less frequently, with 37 and 28 instances, respectively. Normal traffic patterns include MQTT (8,108), ThingSpeak (4,146), and Wipro-Bulb (253). This diverse representation allows for evaluating anomaly detection systems under different traffic conditions.

TABLE I: Dataset Distribution

Dataset	Size
Training	94337

FULL TEXTS BOOK

Testing	23585
Number of features	91
Number of unique attack types (targets)	12

2) Data Preprocessing: In the data preprocessing step, the dataset is processed through several key procedures to ensure quality and consistency. An irrelevant column was removed as it did not contribute to the analysis. The data was then standardized, bringing all features to a comparable scale. Categorical variables were encoded, ensuring they could be effectively utilized in further analysis.

C. Performance evaluation metrics

This study uses several metrics to assess the effectiveness of machine learning and deep learning models. These metrics offer a numerical evaluation of how accurately each model differentiates between normal and abnormal behavior in the dataset.

3) False Positives (FP): Normal cases mistakenly classified as anomalies.
2) True Negatives (TN): Cases accurately classified as normal.
1) True Positives (TP): Cases accurately classified as anomalies.
4) False Negatives (FN): Anomalous cases mistakenly classified as normal.

-**Accuracy** the percentage of correct predictions out of all predictions made . AC= $\frac{TP+TN}{\text{TP+TN+FP+FN}}$

-**Precision** measures how many of the predictions that were labeled as positive are actually correct. $PR = \frac{TP}{\text{TP+FP}}$

- **F1 Score** is a balanced measure of both precision and recall, combining them into a single value. It considers both false positives and false negatives. $F1 - \ Score = \frac{2TP}{\text{2TP+FP+FN}}$

- **Recall** represent indicates how well the model detects actual positives, showing the proportion of true positives out of all the actual positives in the data. $Recall = \frac{TP}{\text{TP+FN}}$

- **Confusion Matrix** a tool used to assess the performance of a classification model by presenting a table that compares predicted labels with actual labels, showing the counts of false positives, true positives, false negatives, and true negatives.

4. RESULTS

This section evaluates the performance of five tested models for anomaly detection and aims to select the best one.

FULL TEXTS BOOK




Fig.1: RF Performance Matrix

Fig.1 showcases the Precision, Recall, and F1-Score for the classification of various attack types using the Random Forest model. Overall, the model demonstrates consistently high performance across most attack types, with metrics close to or equal to 1.00. For attack types such as DOS_SYN_Hping, MQTT_Publish, NMAP_OS_DETECTION, and NMAP_XMAS_TREE_SCAN, the model achieves perfect scores, indicating flawless classification without any misclassifications. However, certain attack types, such as Metasploit_Brute_Force_SSH and NMAP_FIN_SCAN, have slightly lower precision, which reduces their F1-Scores to 0.93 and 0.92, respectively.




Fig.2: Decision Tree Performance Matrix

Fig.2 showcases the Precision, Recall, and F1-Score for the classification of various attack types using the Decision Tree model. Overall, the model demonstrates strong performance across most attack types,

FULL TEXTS BOOK

with metrics close to or equal to 1.00. For attack types such as DOS_SYN_Hping, MQTT_Publish, NMAP_OS_DETECTION,andNMAP_XMAS_TREE_SCAN,the model achieves perfect scores, indicating flawless classification without any misclassifications. However, certain attack types, such as Metasploit_Brute_Force_SSH and Wipro_bulb, have slightly lower scores, with an F1-Score of 0.93 and 0.91, respectively.




Fig.3: k-Nearest Neighbors (k-NN) Performance Matrix

Fig.3 showcases the Precision, Recall, and F1-Score for the classification of various attack types using the k-Nearest Neighbors (k-NN) model. Overall, the model demonstrates strong performance across most attack types, with metrics close to or equal to 1.00. For attack types such as DOS_SYN_Hping, MQTT_Publish, NMAP_OS_DETECTION, and NMAP_XMAS_TREE_SCAN, the model achieves perfect scores, indicating flawless classification without any misclassifications. However, certain attack types, such as NMAP_FIN_SCAN and Wipro_bulb, have slightly lower scores, with F1-Scores of 0.92 and 0.85, respectively.




Fig.4 : CNN Performance Matrix

FULL TEXTS BOOK

Fig.4 showcases the Precision, Recall, and F1-Score for the classification of various attack types using the CNN model. Overall, the model demonstrates strong performance across most attack types, with metrics close to or equal to 1.00. For attack types such as DOS_SYN_Hping, MQTT_Publish, NMAP_OS_DETECTION,and NMAP _XM AS_T R EE_ SCAN, the model achieves perfect scores, indicating flawless classification without any misclassifications. However, certain attack types, such as DDOS_Slowloris and Wipro_bulb, have slightly lower scores, with F1-Scores of 0.85 and 0.71, respectively.




Fig.5 :DNN Performance Matrix

Fig.5 showcases the Precision, Recall, and F1-Score for the classification of various attack types using the Deep Neural Network (DNN) model. While the model performs well for most attack types, achieving perfect metrics for attacks types such as DOS_SYN_Hping,MQTT_Publish, and NMAP_ OS_DETECTION, certain attack types have notable performance challenges. For example, NMAP_FIN_SCAN and Wipro_bulb show significantly reduced F1-Scores of 0.33 and 0.62, respectively, due to low recall values (0.25 for NMAP_FIN_SCAN and 0.51 for Wipro_bulb). Additionally, Metasploit_Brute_Force_SSH and DDOS_Slowloris also show slightly reduced F1-Scores of 0.83 and 0.86, primarily due to lower recall.


The performance of the models, shown in Figures 1 through 5, highlights that Random Forest (RF) consistently surpasses other models in terms of Precision, Recall, and F1-Score for most attack types. While Decision Tree and k-Nearest Neighbors (KNN) also deliver good results, their performance dips slightly with specific attack types, like Wipro_bulb and NMAP_FIN_SCAN. Deep learning models, including Convolutional Neural Networks (CNN) and Deep Neural Networks (DNN), show promising results but encounter some challenges in certain cases, with DNN showing the most variability. Overall,

FULL TEXTS BOOK

RF's solid and reliable performance across all metrics confirms its superiority as the most effective model for anomaly detection in this study.



(a) RF




(b) Decision Tree

FULL TEXTS BOOK


Confusion Matrix Heatmap
ARP_poisioning 1484 0 0 2 0 0 0 3 0 0 34 2
DDOS_Slowloris 0 106 0 0 0 0 0 0 1 0 0 0
DOS_SYN_Hping 0 0 18018 0 0 0 0 0 0 0 0 0
MQTT_Publish 1 1 0 825 0 0 0 0 0 0 0 1
Metasploit_Brute_Force_SSH 0 0 0 0 7 0 0 0 0 0 0 0
NMAP_FIN_SCAN 0 0 0 0 0 6 0 0 0 0 0 0
NMAP_OS_DETECTION 0 0 0 0 0 0 400 0 0 0 0 0
NMAP_TCP_scan 1 0 0 1 0 0 0 198 0 0 0 0
NMAP_UDP_SCAN 1 0 0 0 0 0 0 0 512 0 4 0
NMAP_XMAS_TREE_SCAN 1 0 0 0 0 0 0 0 0 401 0 0
Thing_Speak 41 1 0 0 0 0 0 0 1 0 1486 2
Wipro_bulb 4 0 0 1 0 1 0 0 0 0 2 36
Actual
ARP_poisioning
DDOS_Slowloris
DOS_SYN_Hping
MQTT_Publish
Metasploit_Brute_Force_SSH
NMAP_FIN_SCAN
NMAP_OS_DETECTION
NMAP_TCP_scan
NMAP_UDP_SCAN
NMAP_XMAS_TREE_SCAN
Thing_Speak
Wipro_bulb
Predicted
17500
15000
12500
10000
7500
5000
2500
0

(c) KNN


Confusion Matrix
ARP_poisioning 1514 3 0 1 0 0 0 0 0 0 17 0
DDOS_Slowloris 1 79 0 0 0 0 0 0 22 0 0 0
DOS_SYN_Hping 0 0 18020 0 0 0 0 0 0 0 0 0
MQTT_Publish 2 0 0 852 0 0 0 0 1 0 0 0
Metasploit_Brute_Force_SSH 2 0 0 0 5 0 0 0 0 0 0 0
NMAP_FIN_SCAN 1 0 0 0 0 3 0 0 0 0 0 0
NMAP_OS_DETECTION 0 0 0 0 0 0 377 0 0 0 0 0
NMAP_TCP_scan 1 0 0 0 0 0 0 215 0 0 0 0
NMAP_UDP_SCAN 7 1 0 0 0 0 0 0 490 0 1 0
NMAP_XMAS_TREE_SCAN 0 0 0 0 0 0 0 0 0 366 0 0
Thing_Speak 52 0 0 0 0 0 0 0 3 0 1504 0
Wipro_bulb 2 0 0 0 0 0 0 0 3 0 15 25
Actual
ARP_poisioning
DDOS_Slowloris
DOS_SYN_Hping
MQTT_Publish
Metasploit_Brute_Force_SSH
NMAP_FIN_SCAN
NMAP_OS_DETECTION
NMAP_TCP_scan
NMAP_UDP_SCAN
NMAP_XMAS_TREE_SCAN
Thing_Speak
Wipro_bulb
Predicted
17500
15000
12500
10000
7500
5000
2500
0

(d) CNN



(e) DNN

Fig.6 : Confusion matrices for FR, Decision Tree, KNN, CNN, and DNN models, showing correct predictions on the diagonal and misclassifications off-diagonal. Color intensity reflects prediction counts.

TABLE II: Performance Metrics Comparison

Models	Accuracy	Precision	Recall	F-Score
RF	99.80%	1.00	1.00	1.00
Decision tree	99.70%	1.00	1.00	1.00
KNN	99.55%	1.00	1.00	1.00
CNN	99.43%	0.99	0.99	0.99
DNN	99.40%	0.99	0.99	0.99

Table II shows the performance metrics of the models. Random Forest (RF) achieves the highest accuracy (99.80%) and perfect precision, recall, and F1-score, making it the best-performing model for anomaly detection.

5. CONCLUSION

This study investigated the performance of various machine learning and deep learning models for anomaly detection in IoT networks using the RT-IoT2022 dataset. Among the evaluated models, Random Forest achieved the highest accuracy (99.80%) and achieved perfect precision, recall, and F1-scores, highlighting its robustness and efficiency in identifying anomalies across diverse attack scenarios. Traditional methods like Decision Trees and k-Nearest Neighbors (KNN) also performed well but were slightly outperformed by Random Forest. Deep learning models, including Convolutional Neural Networks(CNNs) and Deep Neural Networks (DNNs), showcased strong performance, particularly in handling complex patterns, but their computational overhead and lower scores for certain attack types limited their overall effectiveness.

The findings emphasize the significance of selecting appropriate anomaly detection models based on accuracy, computational resources, and specific IoT requirements. Future work could explore the

FULL TEXTS BOOK

integration of hybrid models, optimization techniques, and real-time implementation to further enhance IoT anomaly detection capabilities.

REFERENCES

[1] A. E. F. Alfalahi, S. R. A. Alhebsi, and T. Murugan, "Recent Research Solutions on Deep Learning-based Anomaly Detection in Internet of Things," *2024 15th Annual Undergraduate Research Conference on Applied Computing (URC)*, pp. 20–24, 2024.

[2] M. Vyas, J. Chandhok, S. Arumugam, R. Vijayaganth, A. Srivastava, and M. Tiwari, "Revolutionizing IoT Network Security with Deep Learning-Anomaly Detection Model," *Proceedings of the Fourth International Conference on Electronics and Sustainable Communication Systems (ICESC-2023)*, IEEE, pp. 1533–1536, 2023

[3] S. Varalakshmi, V. P., P. S. P., V. Kavitha, Y. V., and V. G., "Design of IoT Network Using Deep Learning-based Model for Anomaly Detection," *Proceedings of the Fifth International Conference on I-SMAC (IoT in Social, Mobile, Analytics and Cloud) (I-SMAC)*, IEEE, pp. 216–218, 2021.

[4] Diana Gaifulina, Igor Ktenko "Selection of Deep Neural Network Models for IoT Anomaly Detection Experiments," *2021 29th Euromicro International Conference on Parallel, Distributed and Network-Based Processing(PDP), 2021*.

[5] Atul Kumar, Ishu Sharma, "CNN-based Approach for IoT Intrusion Attack Detection," *Proceedings of the International Conference on Sustainable Computing and Data Communication Systems (ICSCDS-2023)*, pp. 490-496, 2023.

[6] T. Rajendran, J. K., M. Imtiaz N., and B. Sampathkumar, "Cybersecurity Threat Detection Using Deep Learning and Anomaly Detection Techniques," *2024 International Conference on Knowledge Engineering and Communication Systems (ICKECS)*, pp. 20-24, 2024.

[7] H. Benaddi, M. Jouhari, K. Ibrahimi, A. Benslimane, and E. M. Amhoud, "Improvement of Anomaly Detection System in the IoT Networks using CNN-LSTM Approach," *2023 IEEE Global Communications Conference: Mobile and Wireless Networks*, 2023.

[8] Q. Jiao and L. Mhamdi, "Deep Learning-based Intrusion Detection for IoT Networks," 2024 Global Information Infrastructure and Networking Symposium (GIIS), pp. 1–8, 2024.

[9] M. ElKashlan, H. Aslan, and M. A. Azer, "DDoS Attack Detection in IoT Using Machine Learning-Based Intrusion Detection System (IDS)," 2022 18th International Computer Engineering Conference (ICENCO), pp. 20–25, 2022.

[10] A. Gautam, S. Rathare, G. Bhatnagar, S. I. Dari, and V. K. Gupta, "Application of Deep Learning for Intrusion Detection and Cyber-Attack Detection Using IoT Networks," 2023 International Conference on Evolutionary Algorithms and Soft Computing Techniques (EASCT), pp. 45–50, 2023.

[11] S. Ikhwan, P. Purwanto, and A. F. Rochim, "Comparison Analysis of Intrusion Detection using Deep Learning in IoT Networks," 2023 11th International Conference on Information and Communication Technology (ICoICT), IEEE, pp. 339–343, 2023.

[12] H. R. O. Alghaithi, M. M. A. Alshehhi, and T. Murugan, "IoT Network Anomaly Detection Using Machine Learning and Deep Learning Techniques – Research Study," 2024 IEEE Students Conference on Engineering and Systems (SCES), pp. 1–6, 2024.

FULL TEXTS BOOK

HYBRID INTRUSION DETECTION IN SDN-IOT NETWORKS USING DEEP NEURAL NETWORK AND RANDOM FOREST

Vala Nadeşoğlu

Department of Software Engineering

Haliç University,

Istanbul 34060, Türkiye

Alireza Souri

Department of Computer Engineering,

Haliç University,

Istanbul 34060, Türkiye

Mostafa Habibi

Department of Mechanical Engineering,

Haliç University,

Istanbul 34060, Türkiye

ABSTRACT

This study proposes a hybrid Deep Neural Network (DNN) and Random Forest (RF) model for intrusion detection in SDN-IoT networks using the IoT-SDN-IDS dataset. Preprocessing involved feature selection with SelectKBest and class balancing using SMOTE. The evaluation is conducted using confusion matrices and standard performance metrics across multiple attack types. The hybrid DNN+RF model achieved the highest results, reaching 98% accuracy with precision, recall, and F1-scores of 0.98, outperforming DNN, SVM, Naïve Bayes, and Bernoulli Naïve Bayes classifiers. Future work could explore optimizing the model for real-time detection and deployment in SDN-IoT systems.

Keywords: Feature selection, Intrusion Detection System (IDS),Deep Neural Network (DNN), Random Forest (RF), SMOTE.

1. INTRODUCTION

The convergence of Software-Defined Networking (SDN) and Internet of Things (IoT) architectures has created a security paradox: while these technologies promise unprecedented flexibility and scalability, they have also introduced unique vulnerabilities that conventional intrusion detection systems (IDS) struggle to address. The features that make SDN-IoT systems revolutionary, such as centralized programmability, dynamic control flows, and heterogeneous device ecosystems, have also become their greatest liabilities. As researchers have demonstrated, the programmable nature of SDN controllers introduces new attack vectors [1], while IoT devices remain shockingly vulnerable due to their resource constraints and lack of standardized security [2], posing unprecedented scalability and management challenges. Alarmingly, studies reveal that attackers can hijack entire networks by exploiting just one compromised SDN controller [1]. Existing security solutions are struggling to keep pace: conventional signature-based IDS crumble when faced with the mutating threats targeting SDN's control-data plane separation [1], and most machine learning approaches either demand unrealistic computational resources for IoT nodes or fail to detect sophisticated multi-vector attacks. This situation highlights the urgent need for a new approach to intrusion detection that combines the interpretability of classical algorithms with the adaptive power of deep learning, all while respecting the harsh realities of resource-constrained environments.

Promising breakthroughs are beginning to emerge. Hybrid machine learning models that combine different algorithms have shown particular potential, leveraging complementary strengths to enhance

FULL TEXTS BOOK

detection performance. For instance, hybrid models that integrate tree-based feature selection with neural network pattern recognition have proven effective in addressing both feature selection and pattern recognition challenges [3]. Other researchers have explored lightweight yet efficient techniques for resource-constrained IoT environments, demonstrating that effective detection can be achieved without overwhelming complexity. GRU-RNN architectures have been successfully applied in SDN networks while utilizing only a minimal set of features [4]. Additionally, dynamic reward mechanisms in Deep Reinforcement Learning have shown significant effectiveness in countering evolving botnet tactics in SDN environments [5]. These advances suggest that strategic algorithm fusion and adaptive methods offer a promising path toward securing SDN-IoT networks.

This study proposes a hybrid intrusion detection system based on a Deep Neural Network (DNN) followed by a Random Forest (RF) classifier. The model is designed to perform multi-class classification, allowing it to identify not only whether traffic is malicious, but also the specific type of attack. This approach aims to combine the feature extraction power of DNNs with the interpretability and classification efficiency of RFs. Prior to model training, feature selection was performed using the SelectKBest method to retain the most informative features, and SMOTE (Synthetic Minority Over-sampling Technique) was applied to address class imbalance in the dataset. The system is evaluated on a labeled dataset representative of SDN-IoT traffic, and its performance is assessed across multiple attack categories using accuracy, precision, recall, and F1-score metrics. Additionally, the proposed hybrid model is compared against other traditional machine learning classifiers, including Support Vector Machine (SVM), Naïve Bayes (NB), and Bernoulli Naïve Bayes (BNB), to demonstrate its effectiveness and superiority.

The remainder of this work is organized as follows. Section II presents a review of related works . Section III describes the proposed hybrid model, detailing the data preprocessing steps, feature selection, class balancing techniques, and model architecture. Section IV presents the experimental results, including comparisons with other classifiers, evaluation metrics, and visualization analyses. Finally, Section V concludes the study by summarizing the main findings.

2. RELATED WORK

This study evaluated a **Random Forest (RF)-based IDS** for IoT networks using the Network Intrusion Detection dataset, achieving **99.6% accuracy** and a **0.99 MCC score**. Key innovations included **Recursive Feature Elimination (RFE)** for selecting critical features (e.g., src_bytes, protocol_type) and Symmetrical Uncertainty-based feature weighting. The model demonstrated superior performance in detecting diverse attack types (DoS, Probe, U2R) with a **0.37% false alarm rate**, outperforming traditional ML approaches. This work highlights RF's robustness for IoT security, though computational overhead may limit edge deployment[3] This study proposed a Gated Recurrent Unit Recurrent Neural Network (GRU-RNN) for anomaly detection in SDN environments. Their approach achieved an accuracy of 89% using only six raw features from the NSL-KDD dataset, outperforming traditional methods such as SVM, DNN, and VanillaRNN. The GRU-RNN model demonstrated robustness in detecting zero-day attacks while maintaining minimal impact on SDN controller performance, with only a 3.5–7% degradation in throughput and latency[4]. This study proposed a Dynamic Reward-based Deep Q-Network (DRL) for IoT intrusion detection, addressing class imbalance through an adaptive reward function. The model achieved 99% accuracy on the Bot-IoT dataset, outperforming traditional DQN and Decision Tree methods in multi-class attack detection (DDoS, DoS, Reconnaissance). Key innovations included a traffic-volume-sensitive reward mechanism and SARSA-based data encoding, enhancing sensitivity to minority-class attacks [5]. this study proposed an intrusion detection and mitigation framework for SDN-controlled IoT networks to address security challenges posed by resource-constrained IoT devices. The framework combines signature-based, anomaly-based, and machine learning-based detection approaches to identify various types of cyber threats. Leveraging SDN capabilities, the system dynamically enforces security policies through reconfiguration of network devices and isolation of compromised nodes when intrusions are detected. The architecture consists of IDS sensors for data collection, an IDS manager for hybrid threat analysis, and an SDN controller for real-time mitigation. While demonstrating the potential of hybrid detection methods in reducing false positives, the authors acknowledge the need for further testing on live IoT traffic to validate scalability[6]. this study proposed an efficient intrusion detection system for SDN-enabled healthcare networks using a hybrid SVM-KNN machine learning approach. The authors

FULL TEXTS BOOK

address critical challenges in healthcare cybersecurity, including high false positive rates and slow threat response in conventional IDS solutions. Their architecture leverages SDN's programmable infrastructure to enable real-time traffic monitoring and dynamic threat mitigation, while the hybrid SVM-KNN model combines SVM's classification robustness with KNN's anomaly detection capabilities. The system demonstrates significant improvements over traditional methods, achieving 20-30% better prediction accuracy, 30-50% reduction in end-to-end delay, and 30-40% lower latency rates when validated on the NSL-KDD dataset. Key innovations include an SDN-based security framework tailored for healthcare networks, a voting mechanism that integrates SVM and KNN outputs to reduce false positives, and dynamic flow rule updates through OpenFlow for rapid attack containment. The study provides comprehensive performance metrics including precision (93%), recall (91%), and F1-score (92%), though it notes computational overhead as a trade-off for enhanced detection accuracy[7]. this work propose a Hybrid CatBoost Regression (HCB) model for intrusion detection in IoT-enabled networks, addressing the limitations of centralized IDS in handling high-speed network traffic and scalability issues. The system leverages the IDS2017 dataset, employing categorical feature extraction (e.g., system login status, root shell access, destination host count) to identify attack patterns [8]. this study proposed an hybrid intrusion detection system tailored for SDN-enabled IoT networks that combines Random Forest-Recursive Feature Elimination (RF-RFE) with optimized machine learning models. Their approach addresses critical control-plane vulnerabilities in SDN architectures while maintaining computational efficiency suitable for IoT environments. The system demonstrates exceptional performance, achieving 99% accuracy in both binary and multi-class attack detection (including DoS and probing attacks) using only 10 optimally selected features from the SDN-IoT dataset, such as srcPort, Dpkts, and TnBPSrcIP. A key innovation is the RF-RFE feature selection method, which effectively reduces dimensionality while preserving detection capability, outperforming traditional techniques like ANOVA and SFS. The authors highlight the system's real-time operational capabilities, evidenced by an impressive AUC-ROC score of 0.98 for the Random Forest classifier, significantly surpassing alternative approaches like Logistic Regression (0.57) and AdaBoost (0.85). While the study successfully leverages SDN's centralized control architecture for dynamic threat mitigation, the authors acknowledge challenges in deploying the system across heterogeneous IoT environments, suggesting opportunities for future research in this area [9]. This study proposed a hybrid intrusion detection system that effectively combines Random Forest, Decision Tree, and Multilayer Perceptron algorithms to detect network threats. The proposed system demonstrates strong performance across multiple datasets, achieving 99.7% accuracy on NSL-KDD, 84.89% on CIC-IDS-2017, and 77.99% on UNSW-NB15. The authors employ an ensemble approach using a VotingClassifier that leverages the strengths of each algorithm - Random Forest's robustness, Decision Tree's interpretability, and MLP's ability to detect complex patterns. This combination proves particularly effective, reducing false positives to less than 1% for common attack types like DoS and Probe attacks. The system shows excellent detection capabilities, with 98.22% true positive rate for Probe attacks in NSL-KDD and perfect 100% detection of FTP-BruteForce attacks in CIC-IDS-2017. While the hybrid model demonstrates impressive results overall, the authors note performance variations across datasets, particularly with the more challenging UNSW-NB15 dataset where accuracy drops to 77.99%, highlighting the difficulties in maintaining consistent performance across different network environments and attack types[10]. this proposed a lightweight intrusion detection system specifically designed for IoT environments by combining K-Nearest Neighbors (KNN) and Decision Tree (DT) algorithms. Their approach addresses the critical challenges of resource constraints and attack diversity in IoT networks through efficient feature selection and data normalization. The system demonstrates exceptional performance on the DoH20 dataset, achieving perfect 100% accuracy and detection rate after implementing min-max normalization - a significant improvement over previous results on the Balot dataset (99.28% accuracy). The authors highlight how their hybrid model maintains the interpretability advantages of traditional machine learning while achieving computational efficiency comparable to more complex deep learning approaches, requiring only 25 carefully selected features for operation [11]. This study proposed a Deep Convolutional Neural Network (DCNN) for intrusion detection in IoT networks, achieving state-of-the-art performance with 99.23% accuracy and 99.52% F1-score on the NF-ToN-IoT dataset. The model outperformed existing architectures like CNN-BiLSTM, GRU-RNN, and hybrid CNN-LSTM, demonstrating robustness across multiple datasets (NF-

FULL TEXTS BOOK

BoT-IoT, NF-UNSW-NB15). Key innovations included a multi-dataset validation strategy and an ablation study to optimize component contributions[12].

3. PROPOSED WORK

A. Model Architecture

The proposed intrusion detection framework adopts a hybrid machine learning architecture that combines a Deep Neural Network (DNN) with a Random Forest (RF) classifier. This combination leverages the strengths of both deep learning and ensemble-based decision-making. An overview of the system's architecture is illustrated in Figure 1, which outlines the layered structure of the IoT-SDN environment and the placement of the machine learning modules within the control and application planes.




Figure 1. IoT-SDN IDS Architecture.

The DNN serves as an automatic feature extractor, while the Random Forest provides the final classification layer. This separation of responsibilities allows the model to capture high-level abstractions and maintain robustness and interpretability during decision-making. The DNN component receives a subset of 20 input features selected through the SelectKBest feature selection technique. The network architecture consists of three fully connected hidden layers. The first layer comprises 256 neurons with ReLU activation, regularized using both L2 norm and dropout to mitigate overfitting. The second layer contains 128 ReLU-activated neurons, also regularized, and the third and final hidden layer reduces the representation to 64 neurons. This compact 64-dimensional vector encodes the learned, non-linear relationships within the feature space and serves as the core abstract representation for the classification task. Rather than using the softmax output from the DNN, this learned representation is passed as input to a Random Forest classifier. The RF, a widely used tree-based ensemble method, aggregates predictions from multiple decision trees to improve generalization and reduce variance. Its key advantage lies in its ability to handle noisy or weakly predictive features effectively, while also providing interpretable outputs through feature importance measures.

This hybrid design was motivated by the limitations observed in standalone classifiers. In preliminary evaluations, the DNN demonstrated strong feature learning capabilities but showed signs of overfitting during classification. Conversely, Random Forests performed more reliably on the classification task but lacked the depth to extract meaningful features from raw inputs. By combining both methods, the system benefits from the DNN's ability to model complex patterns and the RF's strength in stable, multi-

FULL TEXTS BOOK

class decision-making. Overall, this architecture enables improved generalization compared to using either model in isolation. The DNN efficiently captures deep, abstract patterns from the input space, while the Random Forest acts as a robust, interpretable, and efficient classification layer suitable for deployment in resource-constrained or real-time settings.




Figure 2. Overview of the proposed hybrid DNN + RF architecture.

B. Dataset

1) Description : This study uses the publicly available IoT-SDN-IDS dataset, which was specifically designed to evaluate intrusion detection systems in Software Defined Networking (SDN) environments with IoT traffic. The dataset contains thousands of labeled network flows with over 70 numerical features, extracted from real and simulated network traffic. Each sample is labeled as one of three classes: BFA, DDoS, or Prope, making it a multi-class classification problem.

TABLE I: Dataset Distribution

Dataset	Size
Training samples (after SMOTE)	52,617
Testing samples	10,718
Number of original features	70
Number of features after feature selection	20
Number of attack types (classes)	3

FULL TEXTS BOOK

2) Preprocessing: The preprocessing process involved several steps. First, duplicate entries were removed to avoid bias caused by repeated flows. Infinite values and missing data (NaNs) were identified and either replaced or discarded. Additionally, irrelevant features such as Src IP, Dst IP, and other identifiers were excluded from the dataset. The remaining numerical features were then normalized using StandardScaler to ensure consistency in scale across all input dimensions. To further refine the dataset, SelectKBest with the ANOVA F-score (f_classif) was used to select the top 20 most informative features for training the model.

3) Class Imbalance and Oversampling:
Although the original dataset was initially balanced across the three attack types(Brute Force Attack (BFA), Distributed Denial-of-Service (DDoS), and Probing (Prope))the preprocessing stage introduced significant class imbalance. Specifically, the removal of duplicate flows and invalid records disproportionately reduced the number of samples in certain classes, particularly DDoS and BFA.
To address this issue, the Synthetic Minority Over-sampling Technique (SMOTE) was applied to the training dataset. SMOTE generates synthetic samples for underrepresented classes by interpolating between existing minority class instances. This process restored class balance in the training data, ensuring that the model would not exhibit bias toward the majority class during learning. In addition to SMOTE, class weights were introduced during DNN training to further mitigate bias at the algorithmic level. A weighted loss function was used to penalize misclassifications of underrepresented classes like Prope and BFA more heavily, thereby encouraging the model to learn a more balanced decision boundary. The complete machine learning pipeline, including the feature selection and SMOTE-based oversampling, is illustrated in Figure 3.




Figure 3. ML Pipeline Incorporating Feature Selection and SMOTE

By combining data-level balancing with algorithmic weighting, the system achieved improved sensitivity across all classes.


C Performance evaluation metrics
The model's performance was evaluated using standard multi-class classification metrics:
-Accuracy: The ratio of correctly predicted instances to the total instances.
-Precision: The proportion of correctly predicted positive observations to the total predicted positives.
-Recall: The proportion of correctly predicted positives to all actual positives.
-F1-Score: The harmonic mean of precision and recall, used to balance both.
-Confusion Matrix: A detailed matrix showing correct vs. incorrect predictions across all classes.


FULL TEXTS BOOK


-AUC-ROC Curve: To assess the separability between classes, especially in probabilistic predictions.

4. RESULTS

This section presents the evaluation results of the proposed hybrid intrusion detection system based on a Deep Neural Network (DNN) and Random Forest (RF) classifier. The performance of the models was assessed using standard multi-class classification metrics, including accuracy, precision, recall, F1-score, and area under the ROC curve (AUC-ROC). To highlight the effectiveness of the hybrid approach, results are reported for both the standalone DNN and the hybrid DNN+RF model. Additionally, the performance was compared against other machine learning models, including Support Vector Machine (SVM), Naïve Bayes, and Bernoulli Naïve Bayes classifiers.







FULL TEXTS BOOK




Figure 4: Precision, Recall, and F1-Score for Different Models (Naive Bayes, Bernoulli NB, SVM, DNN, Hybrid DNN+RF)

The figure summarizes the performance metrics across various models. It is evident that the hybrid DNN+RF model achieves the highest overall scores, followed by the DNN model. Additionally, it can be observed that traditional machine learning models like SVM, Naive Bayes, and Bernoulli Naive Bayes show comparatively lower performance, particularly in recall and F1-scores across different attack classes. The hybrid DNN+RF model demonstrates remarkable consistency with a precision, recall, and F1-score of 0.98 for all classes, indicating its superior capability in detecting and classifying network intrusions accurately.

These results highlight the importance of ensemble and deep learning approaches in improving intrusion detection performance in SDN-IoT networks.

TABLE II: Evaluation Metrics Table

Models	Accuracy
DNN+RF(Hybird)	98%
DNN	90%
SVM	80%
Naïve Bayes	64%
Bernoulli Naïve Bayes	63%

Table II presents the classification accuracy achieved by different models. Among all, the Hybrid model (DNN+Random Forest) achieved the highest accuracy of 98%, significantly outperforming the other approaches. This strong performance can be attributed to the complementary strengths of the two models,the DNN effectively extracts high-level abstract features from the input data, while the Random Forest leverages these features to make robust and precise classifications. The standalone DNN model, without the ensemble enhancement, achieved a respectable 90% accuracy, but the combination with Random Forest further improved the predictive power. In contrast, traditional machine learning algorithms, including SVM (80%), Naïve Bayes (64%), and Bernoulli Naïve Bayes (63%), showed relatively lower performance, highlighting the importance of deep feature learning and ensemble methods in addressing the complexities of IoT network intrusion detection.

FULL TEXTS BOOK




Figure 5. Training and validation accuracy and loss curves of the DNN model across 20 epochs.

The training and validation curves for the DNN model indicate smooth convergence over 20 epochs. Both training and validation accuracy increase steadily, while the corresponding loss values consistently decrease. The close alignment between the curves and the absence of divergence suggest that the model generalized well to unseen data, with no signs of overfitting or underfitting.




Figure 6. Confusion Matrices.

The confusion matrices show that the DNN model struggled to distinguish between Prope and BFA, with 433 Prope samples misclassified as BFA and 313 BFA as Prope. After integrating the Random Forest classifier, these misclassifications dropped sharply to 80 and 58, respectively. This highlights the hybrid model's ability to refine decision boundaries and significantly improve class separability, especially for closely related attack types.

FULL TEXTS BOOK




Figure 7. AUC-ROC curve for the hybrid DNN + RF model

The AUC-ROC curve for the hybrid DNN + RF model demonstrates near-perfect classification performance, with all three classes achieving an AUC of 1.00. This indicates excellent class separability and confirms that the ensemble model was able to distinguish between BFA, DDoS, and Prope attacks with high confidence across all thresholds.




Figure 8. Top 10 most important DNN features as ranked by the Random Forest classifier

These features represent the internal 64-dimensional learned representation generated by the DNN's final hidden layer. While they do not directly correspond to the original input features, they capture complex patterns extracted from the data and were most influential in the final classification decisions.

5. CONCLUTION

This study introduced a hybrid intrusion detection system that combines a Deep Neural Network (DNN) and a Random Forest (RF) classifier to secure SDN-IoT networks against evolving cyber threats. By leveraging the feature extraction capabilities of DNNs and the robust classification power of RFs, the proposed model achieved superior detection performance compared to traditional machine learning methods. Preprocessing steps such as duplicate removal, feature selection using SelectKBest, and class balancing with SMOTE significantly contributed to improving model accuracy and generalization. Experimental results showed that the hybrid model reached a high classification accuracy of 98%, with precision, recall, and F1-scores of 0.98 across multiple attack categories. Furthermore, confusion matrix and AUC-ROC analyses confirmed the model's strong ability to distinguish between different types of attacks, especially for closely related classes. Future research could explore extending the proposed architecture to real-time SDN controllers and investigating adaptive learning methods to further enhance resilience against novel attack patterns.

REFERENCE

[1] S. Kumar and S. Gupta, "Security attacks in software defined networks," in *Proc. IEEE Int. Conf. Sustain. Commun. Netw. Appl. (ICSCNA)*, 2024, pp. 423–426

[2] K. Kumar B. N. et al., "Comprehensive insights into machine learning for intrusion detection systems in IoT and its datasets," in *Proc. IEEE Int. Conf. Emerg. Technol.*, 2024, pp. 1–6

FULL TEXTS BOOK

[3] A. Al Farsi, A. Khan, and M. M. Bait-Suwailam, "A practical evaluation of intrusion detection in IoT networks using Random Forest and Network Intrusion Detection dataset," *IEEE Access*, vol. 12, pp. 53099–53111, 2024
[4] T. A. Tang, L. Mhamdi, D. McLernon, S. A. R. Zaidi, and M. Ghogho, "Deep recurrent neural network for intrusion detection in SDN-based networks," in *Proc. IEEE Int. Conf. Netw. Softwarization (NetSoft)*, 2018, pp. 202–208.
[5] K. Ren, H. Bai, L. Liu, and Y. Wen, "A dynamic reward-based deep reinforcement learning for IoT intrusion detection," in *Proc. IEEE Int. Conf. Intell. Commun. Netw. (ICN)*, 2024, pp. 110–118
[6] A. Zaheer, M. Z. Asghar, and A. Qayyum, "Intrusion Detection and Mitigation Framework for SDN Controlled IoTs Network," in Proc. IEEE 18th Int. Conf. Smart Communities (HONET), 2021, pp. 1-5.
[7] M. W. Asif et al., "An Efficient Intrusion Detection System using Advanced Machine Learning Techniques in SDN for Healthcare System," *IEEE J. Biomed. Health Inform.*, vol. 29, no. 3, pp. 1-14, 2025
[8] L. Ratha and R. M. Bommi, "Hybrid CatBoost Regression Model Based Intrusion Detection System in IoT-Enabled Networks," *Proc. 9th Int. Conf. Electr. Energy Syst. (ICEES)*, 2023, pp. 265–268
[9] M. Ramaiah et al., "A Hybrid Wrapper Technique Enabled Network Intrusion Detection System for SDN-Based IoT Networks," *Proc. IEEE 3rd Int. Conf. AI IoT (AIIoT)*, 2024, pp. 1–6,
[10] Z. Rachidi, K. Chougdali, and A. Kobbane, "Hybrid Intrusion Detection System Based on Random Forest, Decision Tree and MLP Algorithms," *[Journal/Conference Name]*, 2024, pp. 1–5
[11] Z. H. Abdaljabar et al., "An Intrusion Detection System for IoT Using KNN and Decision-Tree Based Classification," *Proc. IEEE [Conference]*, 2021, pp. 1–5
[12] K. Alam, M. F. Monir, Z. Hassan, and M. T. Habib, "Optimizing IoT network intrusion detection: A deep learning approach," *IEEE Access*, vol. 12, pp. 53099–53111, 2024

SERT ÇEKİRDEKLİ MEYVE AĞAÇLARINDA BAKTERİYEL KANSER ETMENİ ÜZERİNE YAPILMIŞ ÇALIŞMALARIN BİBLİYOMETRİK PROFİLİ

BIBLIOMETRIC PROFILE OF STUDIES ON BACTERIAL CANCER IN STONE FRUIT TREES

Dr. Büşran SUNYAR [1]

[1]Iğdır Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Iğdır, Türkiye [1]ORCID ID: https://orcid.org/0000-0001-8524-3308

Doç. Dr. Mesude Figen DÖNMEZ [2]

[2] Doç. Dr., Iğdır Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Iğdır, Türkiye [2]ORCID ID: https://orcid.org/0000-0002-7992-8252

ÖZET

Pseudomonas syringae pv. *syringae* (Pss), sert çekirdekli meyve ağaçlarında bakteriyel kanser ve zamk hastalığına neden olan önemli bir fitopatojen olup dünya genelinde kayısı, şeftali, kiraz, erik ve badem gibi pek çok *Prunus* türünde ciddi ekonomik kayıplara yol açmaktadır. Hastalık etmeni genellikle enfekte tomurcuklarda ve kanser dokularında kışlamakta, epifitik yaşam özelliği sayesinde sağlıklı ya da duyarlı olmayan konukçularda da varlığını sürdürebilmektedir. Bu çalışmada, *P. syringae* pv. *syringae* üzerine yürütülen akademik çalışmaların kapsamı, eğilimleri ve iş birliği yapıları bibliyometrik yöntemlerle değerlendirilmiştir. Scopus veri tabanında 1980–2025 yılları arasında yayımlanmış 857 yayın incelenmiş; veriler VOSviewer ve R yazılımı aracılığıyla analiz edilmiştir. Bulgular, yayın sayılarında yıllık ortalama %3,64'lük artış olduğunu, çalışmaların çoğunlukla çok yazarlı ve uluslararası iş birliklerine dayalı biçimde yürütüldüğünü göstermektedir. En fazla yayın üreten ülkeler arasında ABD, İtalya ve İspanya öne çıkarken, anahtar kelime analizlerinde "bacterial canker", "syringomycin" ve "pathogenicity" gibi terimlerin baskın olduğu görülmüştür. Ayrıca anahtar kelimelerde yıllara bağlı olarak tematik bir çeşitlenme saptanmıştır. Atıf ve iş birliği ağları, disiplinler arası yayın ilişkilerini ve güçlü uluslararası bağları ortaya koymuştur. Elde edilen sonuçlar, bu alandaki bilimsel üretimin gelişim dinamiklerini ve gelecekteki araştırmalar için potansiyel odak alanlarını belirlemede önemli katkılar sunmaktadır.

Anahtar kelimeler: *Pseudomonas syringae* pv. *syringae*, Sert çekirdekli meyve ağaçları, Bibliyometrik analiz

ABSTRACT

Pseudomonas syringae pv. *syringae* (Pss) is a significant phytopathogen that causes bacterial canker and gummosis in stone fruit trees, leading to considerable economic losses worldwide, particularly in *Prunus* species such as apricot, peach, cherry, plum, and almond. The pathogen typically overwinters in infected buds and canker tissues and can survive epiphytically on healthy or non-susceptible hosts due to its ecological adaptability. This study aimed to evaluate the scope, trends, and collaborative structures of academic research conducted on *P. syringae* pv. *syringae* through bibliometric analysis. A total of 857 publications indexed in the Scopus database between 1980 and 2025 were analyzed using VOSviewer and R software. The findings revealed an average annual growth rate of 3.64% in publication output, with most studies being conducted through multi-author and international collaborations. The United States, Italy, and Spain were identified as the leading contributor countries,

FULL TEXTS BOOK

while keyword analysis highlighted dominant research themes such as “bacterial canker,” “syringomycin,” and “pathogenicity.” A temporal thematic diversification of keywords was also observed. Co-citation and collaboration network analyses demonstrated strong interdisciplinary linkages and international partnerships. Overall, the results provide valuable insights into the developmental dynamics of scientific output in this field and help identify potential focal areas for future research.

Key words: *Pseudomonas syringae* pv. *syringae*, Stone fruit trees, Bibliometric analysis

GİRİŞ

Pseudomonas syringae, Gammaproteobacteria sınıfına ait fitopatojenik bir bakteri türü olarak tanımlanmakta ve monokotiledonlar ile otsu ve odunsu dikotiledon bitki türlerinde dünya genelinde çeşitli hastalıklara yol açtığı bildirilmektedir. Bu bakteri türüne ait 60’tan fazla patovarın tanımlandığı, her bir patovarın belirli konukçu bitkileri enfekte ettiği ortaya konulmuştur. Genellikle dar bir konukçu aralığına sahip olan patovarlar arasında, pathovar *syringae*'nin, sert çekirdekli meyveler, yumuşak çekirdekli meyveler, diğer odunsu konukçular, kültür bitkileri ve otlar da dâhil olmak üzere 80’den fazla bitki türünü enfekte edebildiği belirlenmiştir (Hirano et al., 2000; Kennelly et al., 2007; Buttimer et al., 2017).

Pseudomonas syringae pathovarlarının birçok bitki türünde ekonomik olarak önemli hastalıklara neden olduğu bilinmektedir. *P. syringae* pv. *syringae* (Pss) tarafından oluşturulan bakteriyel kanser ve zamk hastalığı, dünya genelinde sert çekirdekli meyve ağaçlarında en yaygın görülen hastalıklardan biri olarak kabul edilmektedir (Abbasi vd., 2013; Bophela vd., 2020). Bu patojenin, başta kayısı (*Prunus armeniaca*), şeftali (*Prunus persica*), tatlı kiraz (*Prunus avium*), erik (*Prunus domestica*) ve badem (*Prunus dulcis*) olmak üzere çeşitli *Prunus* türlerinde enfeksiyona yol açtığı ve önemli düzeyde zararlara neden olduğu bildirilmiştir (Hulin vd., 2018). Genellikle kanser dokularında ve enfekte tomurcuklarda kışlayan bu patojenin, hem enfekte hem de sağlıklı ağaçlar ile konukçu olmayan bitkiler üzerinde epifitik yaşamını sürdürebildiği saptanmıştır. Ayrıca, söz konusu hastalığın çoğunlukla ağaçların toprak üstü kısımlarını etkilediği ve oluşan yaralardan bakterinin dokulara giriş yaptığı ifade edilmiştir (Agrios, 2005). Hastalık belirtileri arasında dal yanıklığı, geriye doğru ölüm, dallarda ve gövdede akıntı oluşumu ile birlikte gelişen kanserler, meyvelerde koyu kahverengi çökük lekeler ve zamanla yapraklarda deliklere dönüşen lekeler yer almaktadır (Agrios, 2005; Hulin ve ark., 2018; Bophela ve ark., 2020).

Hastalığın ortaya çıkmasını önlemek amacıyla, sağlıklı dikim materyali tercih edilmekte ve don riski yüksek alanlardan kaçınılarak uygun dikim yerleri belirlenmektedir. Hastalığın görüldüğü alanlarda budama işlemlerinin kuru ve güneşli günlerde gerçekleştirildiği, enfekteli sürgün ve ağaçların belirli bir plan dâhilinde bahçeden uzaklaştırıldığı, ayrıca budama sonrası oluşan yaraların hızla kapatıldığı belirtilmektedir. Bununla birlikte, günümüzde bakteriyel kanserin kimyasal kontrolünde, bakır oksit, bakır oksiklorür ve bakır hidroksit içeren formulasyonların yaygın olarak kullanıldığı ifade edilmektedir (Kaluzna et al., 2010; Broniarek-Niemiec et al., 2023). Ancak bu uygulamalarla mevcut enfeksiyonların yalnızca sınırlandırılabildiği, hastalığın tamamen ortadan kaldırılamadığı vurgulanmaktadır (Bultreys and Kaluzna, 2010; Husseini and Akköprü, 2020).

Bibliyometrik analizlerin literatürün yapısal özelliklerinin, gelişim eğilimlerinin ve araştırma odaklarının belirlenmesinde etkili bir yöntem olarak öne çıktığı ifade edilmektedir. Bu tür analizler aracılığıyla, atıf ilişkileri, yazar ve kurum dağılımları ile anahtar kelimeler gibi çeşitli unsurlar üzerinden bilimsel üretimin farklı yönlerinin değerlendirilmesine olanak tanındığı belirtilmektedir (Yang et al., 2021; Yun et al., 2021; Qin et al., 2022; Wan et al., 2022; Guan et al., 2023).

FULL TEXTS BOOK


Söz konusu çalışmada, *Pseudomonas syringae* pv. *syringae* etmeni üzerine gerçekleştirilen akademik yayınların; literatürdeki araştırma eğilimleri, öne çıkan yazarlar, ülkeler, kurumsal iş birlikleri ve anahtar kelimeler açısından, VOSviewer ve R programları kullanılarak bibliyometrik yöntemle analiz edilmesi amaçlanmıştır.

MATERYAL ve YÖNTEM

Çalışmada, sert çekirdekli meyve ağaçlarında önemli bir hastalık olan bakteriyel kanser ve zamklanma etmeni "*Pseudomonas syringae* pv. *syringae*" anahtar kelimesi ile Scopus veri tabanında arama yapıldığında elde edilen 857 yayın üzerinde VOSviewer ve R programları kullanılarak bibliyometrik analizi yapılmıştır.

Yöntem

Bibliyometrik analiz kapsamında, Scopus veri tabanında gerçekleştirilen arama sonucunda elde edilen 857 bilimsel yayın değerlendirmeye alınmıştır. Analiz sürecinde izlenen yöntem adımları Şekil 1'de şematik olarak sunulmuştur.



Şekil 1. Bibliyometrik analizi için veri indirme şeması

FULL TEXTS BOOK

Tablo 1. Çalışmada kullanılan Vosviwer ve R progmında yapılan analizler

Vosviwer	R
Anahtar kelimelerin ağ haritası	Yayınların ana bilgileri
Anahtar kelimelerin güncel kullanım durumları	Yazar - ülke ve anahtar kelime sankey diyagramı
Dergilerin ağ grafiği analizi	Yazarların üretkenlik zamanları
Ülkelerin ağ grafiği analizi	Sorumlu yazarların ülkelerine göre yayın dağılımı ve uluslararası işbirlikleri
	Yıllara göre yayın sayısı
	Ülkelerin ortak yayın haritası
	1980–2025 yılları arasındaki literatürde *Pseudomonas syringae* pv. *syringae* ile ilişkili anahtar kelimelerin dönemsel olarak nasıl değiştiğini ve tematik evrimini gösteren Sankey diyagramı.

Veri Analizi

Ülkelere göre yayın dağılımı, yayımlanan dergiler ve benzeri bibliyometrik değişkenlere ilişkin betimsel analizler Microsoft Excel yazılımı kullanılarak yapılmıştır. Atıf ilişkileri ile anahtar kelimeler arasındaki eşzamanlılık ve bu ilişkilerin görsel temsilleri VOSviewer programı aracılığıyla analiz edilmiştir. Ayrıca, ülkeler arasındaki bilimsel iş birliği ağları, en sık karşılaşılan anahtar terimler, anahtar kelimelerin zaman içerisindeki değişim eğilimleri ve kavramsal gruplamaları (co-word analizi) R yazılımı üzerinden, bibliometrix paketinin web tabanlı grafik arayüzü olan "biblioshiny" kullanılarak detaylı biçimde değerlendirilmiştir.

BULGULAR

1980–2025 yıllarını kapsayan bu araştırmada, toplam 857 yayına ulaşılmış ve bu yayınların 256 farklı kaynaktan elde edildiği belirlenmiştir. Yayın sayısında yıllık ortalama %3,64'lük bir artış oranı tespit edilmiş ve bu durum, çalışma alanına yönelik akademik ilginin giderek arttığını ortaya koymuştur. Belgelerde ortalama 31,5 atıf sayısına ulaşıldığı, toplamda 2454 yazarın katkı sunduğu ve yalnızca 35 yayının tek yazarlı olarak yayımlandığı görülmüştür. Bu bulgular doğrultusunda, çok yazarlı ve iş birliğine dayalı yayın modelinin baskın olduğu anlaşılmıştır. 4107 anahtar kelime kullanıldığı belirlenmiştir. Yayın türlerinin dağılımında büyük oranda makalelere yer verildiği, daha sınırlı sayıda kitap bölümü, konferans bildirisi, kısa not ve derleme çalışmasının bulunduğu saptanmıştır (Tablo 2).

Tablo 2. *Pseudomonas syringae* pv. *syringae* konulu yayınların ana bilgileri


Veri Hakkında Temel Bilgiler	
Zaman Aralığı	1980–2025

FULL TEXTS BOOK

Kaynak Sayısı (Dergi, Kitap vb.)	256
Belge Sayısı	857
Yıllık Büyüme Oranı (%)	3.64
Belge Başına Ortalama Atıf	31.5
Anahtar kelime	4107
Yazar Sayısı	2454
Tek Yazarlı Belgeler	35
Belge Başına Ort. Yazar Sayısı	4.53
Uluslararası Ortak Yazarlık (%)	19.49
Makale	787
Kitap Bölümü	6
Konferans Bildirisi	27
Kısa Not	21
Derleme	13


1980–2025 yılları arasındaki bilimsel üretimi gösteren grafikte, *Pseudomonas syringae pv. syringae* konulu yayınların yıllık dağılımı sunulmuştur (Şekil 2). İlk yıllarda oldukça sınırlı sayıda çalışma yapıldığı, 1990'lı yılların başından itibaren ise yayın sayısında belirgin bir artış yaşandığı gözlemlenmiştir. En yüksek yayın sayısına 1994 ve 2013 yıllarında ulaşıldığı, bu dönemlerde otuzdan fazla çalışmanın yayımlandığı tespit edilmiştir. 1990 sonrası süreçte üretimde dalgalı bir seyir izlenmiş, ancak genel olarak yüksek düzeyde yayın faaliyeti sürdürüldüğü görülmüştür. 2000'li yıllarda da benzer bir eğilim devam ettirilmiş; hemen her yıl 20'nin üzerinde çalışma yayımlanmıştır. 2025 yılına ait dikkat çekici düşüşün ise yılın henüz tamamlanmamış olması veya veri aktarım eksikliği gibi nedenlerle oluştuğu düşünülmektedir.

FULL TEXTS BOOK



Şekil 2. Yıllara göre yayın sayısı


İş birliği ağı haritasında, *Pseudomonas syringae* pv. *syringae* üzerine gerçekleştirilen bilimsel yayınlarda ülkeler arasında kurulan ortaklıklar görselleştirilmiştir. Her bir ülke, haritada düğüm olarak temsil edilmiş; ortak yayınlar doğrultusunda ülkeler arasında bağlantılar oluşturulmuştur. Bağlantıların kalınlıkları iş birliği yoğunluğunu, düğüm boyutları ise ülkelerin toplam yayın katkısını yansıtmaktadır. Haritada merkezi konumda yer alan Amerika Birleşik Devletleri'nin, çok sayıda ülke ile güçlü bilimsel ilişkiler geliştirdiği tespit edilmiştir. Özellikle İspanya, İtalya, Almanya, Birleşik Krallık, Japonya, İran ve Güney Kore ile yapılan ortak çalışmaların öne çıktığı belirlenmiştir. Avrupa ülkeleriyle kurulan yoğun iş birlikleri, ağın Avrupa merkezli bir yapıya sahip olduğunu göstermiştir. Bununla birlikte, Kanada, Güney Afrika, Avustralya, Hindistan, Çin ve Japonya gibi farklı kıtalardan ülkelerin de bu alandaki bilimsel üretime önemli katkılar sunduğu gözlemlenmiştir (Şekil 3). Haritada kümelenen bölgeler arasında, tematik ya da coğrafi araştırma eğilimlerinin oluştuğu anlaşılmıştır. Elde edilen bulgular, *P. syringae* pv. *syringae* araştırmalarında uluslararası ve çok merkezli bilimsel etkileşimin etkin bir biçimde sürdürüldüğünü ve bu etkileşimlerin, konunun küresel ölçekte ele alındığını ortaya koymaktadır.



Şekil 3. Ülkelerin ağ grafiği analizi


FULL TEXTS BOOK

Ülkeler arası bilimsel iş birliklerini gösteren bu haritada, *Pseudomonas syringae pv. syringae* konulu çalışmalarda en yoğun iş birliğinin Amerika Birleşik Devletleri ile çeşitli Avrupa ülkeleri arasında kurulduğu görülmüştür. Haritada koyu mavi ile gösterilen ülkelerin iş birliği sayısının daha yüksek olduğu, açık mavi ile gösterilen ülkelerde ise katkının daha sınırlı düzeyde kaldığı belirlenmiştir. Özellikle ABD merkezli çok sayıda yayın, Avrupa kıtasındaki pek çok ülke ile ortaklaşa gerçekleştirilmiş ve bu etkileşimlerin küresel ölçekte dağıldığı gözlemlenmiştir. Asya, Afrika, Güney Amerika ve Okyanusya'daki bazı ülkelerin de bilimsel üretime katkı sunduğu; ancak bu katkının daha sınırlı düzeyde gerçekleştiği tespit edilmiştir (Şekil 4).




Şekil 4. Ülkelerin ortak yayın haritası

Grafikte, *Pseudomonas syringae* pv. *syringae* konulu yayınlarda sorumlu yazarların bağlı bulunduğu ülkelerin katkı düzeyleri ve iş birliği türleri karşılaştırmalı olarak sunulmuştur. En yüksek yayın sayısına Amerika Birleşik Devletleri tarafından ulaşılmış; bunu sırasıyla İtalya, İran, İspanya ve Çin'in izlediği belirlenmiştir. Çalışmaların çoğunluğunun tek ülke kaynaklı (SCP) olduğu ortaya konmuş; buna karşılık Almanya, Birleşik Krallık ve İsviçre gibi bazı ülkelerde çok uluslu iş birliklerine dayanan yayınların (MCP) daha yaygın olduğu saptanmıştır. Türkiye'nin de dahil olduğu bazı ülkelerin katkısının sınırlı sayıda yayınla temsil edildiği görülmüştür. Bu dağılım, araştırma alanının farklı ülkeler tarafından ele alındığını ve bazı ülkelerde uluslararası iş birliğinin bilimsel üretimde önemli bir rol oynadığını ortaya koymuştur (Şekil 5).



Şekil 5. Sorumlu yazarların ülkelerine göre yayın dağılımı ve uluslararası işbirlikleri


Bilimsel yayınların dağılımını ve disiplinler arası etkileşimleri ortaya koymak amacıyla oluşturulan bu atıf ağı haritasında, *Pseudomonas syringae pv. syringae* konulu çalışmaların yayımlandığı dergiler arasındaki ilişkiler analiz edilmiştir. Farklı renklerle temsil edilen kümelerde, ortak atıf yapılarına ve tematik benzerliklere göre gruplanmış dergiler yer almaktadır. Haritada, **Journal of Bacteriology, Journal of Phytopathology, Plant Disease, Molecular Plant-Microbe Interactions ve Physiological and Molecular Plant Pathology** gibi yayın organlarının merkezi konumda yer aldığı ve literatürde yoğun şekilde referans gösterildiği belirlenmiştir (Şekil 6). Bu dergiler etrafında şekillenen kümeler aracılığıyla, bitki patolojisi, mikrobiyoloji, moleküler biyoloji ve tarım bilimleri gibi çeşitli disiplinler arasında güçlü bir yayın etkileşimi kurulduğu gözlemlenmiştir. Ayrıca, çevre bilimleri ve doğal ürün kimyası gibi ilişkili alanlarda yayımlanan dergilerle de bağlantıların kurulduğu görülmüştür.



Şekil 6. Dergilerin ağ grafiği analizi

FULL TEXTS BOOK


Yıllara göre yazar üretkenliği incelendiğinde, *Pseudomonas syringae* pv. *syringae* konusundaki bilimsel katkıların zamansal olarak nasıl dağıldığı ortaya konmuştur. Yayın sayıları ile yıllık atıf oranları birlikte değerlendirilmiş; daire boyutlarının yazarların toplam yayın miktarını, renk tonlarının ise bu yayınların yıllık toplam atıf düzeylerini yansıttığı belirlenmiştir. Elde edilen bulgulara göre, Gross D.C., Scortichini M., Takemoto J.Y. ve Lindow S.E. gibi araştırmacılar tarafından farklı dönemlerde sürdürülebilir bir yayın performansı sergilenmiştir. Özellikle 1990–2010 yılları arasında üretimin yoğunlaştığı; son yıllarda ise Cazorla F.M. ve De Vicente A. gibi araştırmacıların katkılarının belirgin şekilde arttığı görülmüştür. Bu durum, geçmişte etkili olmuş ve günümüzde de aktif biçimde yayın yapmayı sürdüren araştırmacıların literatürde varlıklarını koruduğunu göstermiştir (Şekil 7).



Şekil 7. Yazarların üretkenlik zamanları

Şekil 8'de, *Pseudomonas syringae pv. syringae* üzerine yapılan çalışmalarda öne çıkan yazarlar, bağlı bulundukları ülkeler ve yoğunlaşılan anahtar kelimeler arasındaki ilişkiler görselleştirilmiştir. ABD, İspanya ve İtalya'nın en fazla katkı sunan ülkeler olduğu belirlenmiş; ABD merkezli çalışmalarda "*hypersensitive response*" ve "*syringomycin*", İspanya'da "*bacterial canker*" ve "*virulence*", İtalya'da ise "*Pseudomonas syringae*" anahtar kelimelerinin öne çıktığı gözlemlenmiştir. Diğer ülkeler sınırlı sayıda yayınla temsil edilmiş; genel olarak "*P. syringae pv. syringae*", "*Pseudomonas syringae*" ve "*bacterial canker*" terimlerinin en sık kullanılan anahtar kelimeler olduğu tespit edilmiştir.


FULL TEXTS BOOK




Şekil 8. Yazar - ülke ve anahtar kelime sankey diyagramı

Şekil 9'te sıklıkla birlikte kullanılan anahtar kelimeler konu kümelerine göre renklendirilmiş ve "*Pseudomonas syringae pv. syringae*", "*bacterial canker*", "*pathogenicity*" ve "*syringomycin*" gibi terimlerin çalışmalarda merkezî konumda yer aldığı görülmüştür. Bu anahtar kelimeler etrafında, biyokontrol, konukçu-bitki etkileşimi, toksin üretimi ve moleküler savunma mekanizmaları gibi farklı tematik kümelerin şekillendiği belirlenmiştir. Şekil 10'da ise anahtar kelimeler, yıllara göre renklendirilmiş olup araştırma eğilimlerinin zaman içinde nasıl değiştiği görselleştirilmiştir. Burada, "*biological control*", "*antibacterial activity*", "*Prunus persica*" gibi kavramların son yıllarda daha sık kullanılmaya başlandığı, buna karşın bazı klasik konuların (örneğin "*syringomycin"* ve *"phytotoxin"*) daha önceki dönemlerde yoğunlaştığı dikkat çekmiştir.

FULL TEXTS BOOK




Şekil 9. Anahtar kelimelerin ağ haritası



Şekil 10. Anahtar kelimelerin güncel kullanım durumları

FULL TEXTS BOOK

Anahtar kelimelerin zamansal dağılımını görselleştiren Sankey diyagramında, *Pseudomonas syringae pv. syringae* ile ilişkili anahtar kelimelerin 1980–2025 yılları arasındaki tematik gelişimi dönemsel olarak görselleştirilmiştir. 1980–1998 yılları arasında yapılan çalışmaların büyük ölçüde "*bacterial diseases*" ve "*Pseudomonas syringae pv. syringae*" gibi genel başlıklar altında toplandığı görülmüştür. 1999–2010 döneminde ise literatürde "*biological control", "antibacterial activity", "essential oil"* gibi daha özgül ve uygulamaya yönelik kavramların öne çıktığı tespit edilmiştir. 2011–2022 yılları arasında araştırma konularının çeşitlendiği, özellikle "*systemic acquired resistance", "hypersensitive response*", "*syringomycin"* ve "*hrp genes*" gibi moleküler düzeyde süreçlerin daha sık ele alındığı belirlenmiştir. Aynı dönemde, uygulamalı çalışmaların da sürdürüldüğü ve "*biological control*" ile "*antibacterial activity*" terimlerinin önemini koruduğu anlaşılmıştır. 2023–2025 aralığında ise daha sınırlı sayıda terimle temsil edilen literatürde "*biocontrol", "bacterial leaf spot*" ve "*pseudomonas syringae*" gibi konulara odaklanıldığı gözlemlenmiştir (Şekil 11).



Şekil 11. 1980–2025 yılları arasındaki literatürde *Pseudomonas syringae pv. syringae* ile ilişkili anahtar kelimelerin dönemsel olarak değişimini ve tematik evrimini gösteren Sankey diyagramı.

SONUÇ VE ÖNERİLER

Bu çalışmada, *Pseudomonas syringae* pv. *syringae* üzerine gerçekleştirilen bilimsel yayınlar bibliyometrik açıdan incelenmiştir. Yayın sayısında yıllar içinde artış olduğu, çalışmaların büyük ölçüde çok yazarlı ve uluslararası iş birliklerine dayandığı belirlenmiştir. Anahtar kelime analizleriyle, araştırma konularında zamanla tematik çeşitlenme yaşandığı saptanmıştır.

Elde edilen bulgular doğrultusunda, gelecek araştırmalar için uluslararası iş birliklerinin güçlendirilmesi, biyokontrol ve moleküler düzeydeki konulara odaklanılması ve literatürdeki boşlukların giderilmesine yönelik çalışmaların teşvik edilmesi önerilmektedir.

KAYNAKÇA

Abbasi, V., Rahimian, H., & Tajick-Ghanbari, M. A. (2013). Genetic variability of Iranian strains of Pseudomonas syringae pv. syringae causing bacterial canker disease of stone fruits. European Journal of Plant Pathology, 135, 225–235. https://doi.org/10.1007/s10658-012-0095-1

Agrios, G. (2005). Plant pathology. 4thEdition (p. 922). Elsevier Academic.

FULL TEXTS BOOK

Bophela, K. N., Petersen, Y., Bull, C. T., & Coutinho, T. A. (2020). Identifi cation of Pseudomonas isolates associated with bacterial canker of stone fruit trees in the Western Cape, South Africa. Plant Disease, 104, 882–892. https://doi.org/10.1094/PDIS-05-19-1102-RE

Broniarek-Niemiec, A., Børve, J., & Puławska, J. (2023). Control of bacterial canker in stone fruit trees by chemical and biological products. Agronomy, 13(4), 1166.

Bultreys, A., & Kaluzna, M. (2010). Bacterial cankers caused by Pseudomonas syringae on stone fruit species with special emphasis on the pathovars syringae and morsprunorum race 1 and race 2. Journal of Plant Pathology, S21-S33.

Buttimer, C., McAuliffe, O., Ross, R. P., Hill, C., O'Mahony, J., & Coffey, A. (2017). Bacteriophages and bacterial plant diseases. Frontiers in microbiology, 8, 34.

Guan, C., Shen, H., Dong, S., Zhan, Y., Yang, J., Zhang, Q. ve diğerleri (2023). Bibliyometrik verilere dayalı ekstrakorporeal membran oksijenasyonunun araştırma durumu ve gelişme eğilimi. Ön. Kardiyovasküler Tıp. 10, 101048903. doi:10.3389/fcvm.2023.1048903

Hirano, S. S., & Upper, C. D. (2000). Bacteria in the leaf ecosystem with emphasis on Pseudomonas syringae—a pathogen, ice nucleus, and epiphyte. Microbiology and molecular biology reviews, 64(3), 624-653.

Hulin, M. T., Mansfi eld, J. W., Brain, P., Xiangming, X., Jackson, R. W., & Harrison, R. J. (2018). Characterization of the pathogenicity of strains of Pseudomonas syringae towards cherry and plum. Plant Pathology, 67(5), 1177–1193. 10.1111/ppa.12834

Husseini, A., & Akköprü, A. (2020). The possible mechanisms of copper resistance in the pathogen Pseudomonas syringae pathovars in stone fruit trees. Phytoparasitica, 48(5), 705-718.

Kaluzna, M., Ferrante, P., Sobiczewski, P., & Scortichini, M. (2010). Characterization and genetic diversity of Pseudomonas syringae from stone fruits and hazelnut using repetitive-PCR and MLST. Journal of Plant Pathology, 781-787.

Kennelly, M. M., Cazorla, F. M., de Vicente, A., Ramos, C., & Sundin, G. W. (2007). Pseudomonas syringae diseases of fruit trees: progress toward understanding and control. Plant disease, 91(1), 4-17.

Qin, A., Sun, J., Gao, C. ve Li, C. (2022). Yetişkin başlangıçlı Still hastalığının araştırma durumu ve eğilimleri üzerine bibliyometrik analiz: 1921-2021. Front. Immunol. 13, 13950641. doi:10.3389/fimmu.2022.950641

Wan, Y., Shen, J., Ouyang, J., Dong, P., Hong, Y., Liang, L. ve diğerleri (2022). 2004-2022 yılları arasında nötrofil hücre dışı tuzaklarının bibliyometrik ve görsel analizi. Front. Immunol. 13, 131025861. doi:10.3389/fimmu.2022.1025861

Yang, HY, Wang, D., Chen, C., Liu, Y., Han, C., Gao, Y. ve diğerleri (2021). Gastroenteroloji ve hepatolojinin küresel araştırma durumu: bibliyometrik bir çalışma. Tıp (Baltimore) 100 (15), e25291. doi:10.1097/md.0000000000025291

Yun, L., Wang, L., Pan, Y., Liu, M., Han, Q., Sun, J. ve diğerleri (2021). Osteoporozla ilgili araştırmalarda mikroRNA'ların mevcut durumu ve gelişim eğilimi: bibliyometrik bir analiz. Folia Histochem Cytobiol. 59 (4), 203–211. doi:10.5603/FHC.a2021.0024

FULL TEXTS BOOK

PREMENSTRÜEL SENDROM İLE DEĞİŞEN BESLENME ALIŞKANLIKLARININ HEDONİK VE DUYGUSAL YEME DAVRANIŞLARINA ETKİSİ

THE EFFECT OF CHANGED DIETARY HABITS DUE TO PREMENSTRUAL SYNDROME ON HEDONIC AND EMOTIONAL EATING BEHAVIOR

***Su Lal CANSEV*[1]**

[1]*Serbest Araştırmacı, Antalya Bilim Üniversitesi, Sağlık Bilimleri Fakültesi, Beslenme ve Diyetetik Bölümü, Antalya, Türkiye.*

[1]*ORCID ID: https://orcid.org/0009-0002-1667-4424*

***Hatice Kübra ARAS*[2]**

[2] *Doktor Öğretim Üyesi, Antalya Bilim Üniversitesi, Sağlık Bilimleri Fakültesi, Beslenme ve Diyetetik Bölümü, Antalya, Türkiye.*

[2]*ORCID ID: https://orcid.org/0000-0001-7750-6024*

ÖZET

Amaç: Premenstrüel sendrom (PMS), adet döngüsünün luteal fazında ortaya çıkan, fizyolojik ve psikolojik semptomlarla karakterize edilen bir durumdur. Bu dönemde hormonal dalgalanmalara bağlı olarak duygu durum değişimleri, yeme davranışlarında farklılaşma ve özellikle yüksek enerjili besinlere yönelim sıkça gözlenmektedir. Bu araştırma; premenstrüel dönemde olan üniversite öğrencilerinin değişen beslenme alışkanlıklarının hedonik ve duygusal yeme davranışı üzerine etkisinin saptanması amacı ile planlanmıştır.

Yöntemler: Araştırma; Kasım 2024 – Nisan 2025 ayları arasında Antalya ilinde bulunan üniversitelerde öğrenim gören kadın bireyler ile gerçekleştirilmiştir Araştırmaya; 18- 45 yaş aralığında, Antalya ilinde öğrenim gören, araştırmaya katılmaya gönüllü 214 katılımcı dahil edilmiştir. Araştırma verileri; bireylerin demografik bilgilerini, beslenme alışkanlıklarını, menstrasyon dönemlerini ele alan anket formu, Duygusal Yeme Ölçeği ve Besin Gücü Ölçeği (BGÖ) kullanılarak yüz yüze görüşme yöntemi ile elde edilmiştir.

Bulgular: Katılımcıların yaş ortalamaları 23.2±4.1, BKİ ortalamaları 22±3.3 olarak bulunmuştur. Kadınların ortak şikayetleri arasında genel olarak karın ağrısı (%71.5) baş göstermektedir. Araştırmaya dahil olan bireylerin menstrüasyon öncesi duygu durumları sorulmuştur; %64.5 ile çoğunluk agresif olduğunu belirtmiştir. Bireylerin %83.2'si ise menstrüasyon öncesi dönemde daha sık tatlı tükettiğini belirtmiştir. Toplam DYÖ ortalama puanlara bakıldığında, **"bazen düzensiz bazen düzenli" adet gören kadınların** duygusal yeme puanı en yüksektir. Yapılan ikili karşılaştırmalar sonucunda alt boyutlar için bazen düzenli bazen düzensiz menstrüasyon gören kişiler diğer gruplara göre daha yüksek puanlar almıştır. Besin gücü ölçeği ile duygusal yeme ölçeği arasında aynı yönlü zayıf anlamlı bir korelasyon bulunmuş ($r=0.390$, $p<0.001$) olup besin gücü ölçeği puanındaki bir artışın duygusal yeme ölçeği puanlarında da zayıf bir artışa sebep olduğu saptanmıştır.

Sonuç: Bu çalışmada, premenstrüel sendrom döneminde kadınların beslenme alışkanlıklarındaki değişimlerin, hedonik ve duygusal yeme davranışları üzerinde etkili olduğu görülmüştür. Bulgular, premenstrüel dönemde artan yeme isteğinin duygusal yeme davranışlarını etkileyebileceğini göstermektedir.

Anahtar kelimeler: Duygusal beslenme, Hedonik açlık, Premenstrüel sendrom.

ABSTRACT

Objective: Premenstrual syndrome (PMS) is a condition characterized by physiological and psychological symptoms that occurs during the luteal phase of the menstrual cycle. Due to hormonal

FULL TEXTS BOOK

fluctuations, mood swings, changes in eating behaviors, and a tendency toward high-energy foods, in particular, are frequently observed during this period. This study was designed to determine the impact of changing eating habits on hedonic and emotional eating behaviors in university students during their premenstrual period.

Methods: The study was conducted with female students studying at universities in Antalya between November 2024 and April 2025. The study included 214 volunteer participants aged 18-45, studying in Antalya. Research data were collected through face-to-face interviews using a questionnaire covering demographic information, eating habits, and menstrual cycles, as well as the Emotional Eating Scale and the Food Power Scale (FBS).

Findings: The participants' mean age was 23.2±4.1, and their mean BMI was 22±3.3.A common complaint among women was abdominal pain (71.5%). The study participants were asked about their premenstrual mood; the majority (64.5%) reported being aggressive. 83.2% of the participants reported consuming sweets more frequently during the premenstrual period. According to the total DES mean scores, women with "sometimes irregular, sometimes regular" menstrual periods had the highest emotional eating scores. Pairwise comparisons revealed that those with sometimes regular, sometimes irregular menstrual periods scored higher than the other groups for the subscales. A weakly significant correlation was found between the Food Power Scale and the Emotional Eating Scale ($r=0.390$, $p<0.001$), indicating that an increase in the Food Power Scale score also led to a weak increase in the Emotional Eating Scale scores.

Conclusion: This study found that changes in women's eating habits during premenstrual syndrome impacted hedonic and emotional eating behaviors. The findings suggest that increased food cravings during the premenstrual period may influence emotional eating behaviors.

Keywords: Emotional eating, Hedonic hunger, Premenstrual syndrome.

GİRİŞ

Premenstrüel sendrom (PMS) kadın popülasyonunun büyük bir çoğunluğunda bulunan hem sosyal hem çalışma hem de aile hayatını olumsuz etkileyen bir bozukluktur. Menstrüel siklusun luteal fazından menstrüasyon başladığı güne kadar sürer. Menstrüasyon siklusunun başlamasıyla da semptomlar ortadan kalkar (Süer, 2008). Kişide memelerde şişkinlik, kilo alımı, baş ağrısı, depresiflik, halsizlik gibi fiziksel ve duygusal değişimlerde belirti verir (Türkçapar, 2011). Ayrıca yaygın olarak da kızgın olma, anksiyete, durduk yere ağlama, depresyon konsantrasyon düşüklüğü, gerginlik gibi psikolojik semptomları da vardır (Süer, 2008). Yiyecek isteği ve bazı yiyeceklerin fazla tüketimi, luteal fazda daha sık rapor edilmektedir. Gıda tüketimiyle ilgili çalışmalardan birçoğu, luteal fazda artan kalori alımını göstermiştir, ancak bu çalışmaların yalnızca birkaçı PMS'yi kontrol etmiştir. Karbonhidrat alımının beyinde serotonin düzeylerinde artışını gösteren hayvansal çalışmalara dayanarak, bazı araştırmacılar, semptomları azaltmak amacıyla, PMS'li kadınlarda bilinçsizce karbonhidrat alımlarını artırarak kendi kendilerini tedavi ettiklerini gözlemlemiştirler (Trout ve diğerleri, 2008). Buna bağlı olarak da artan karbonhidrat alımı sonucu düşkünlük oluşmaktadır. Bu olay bir yandan duygusal yeme davranışıyla bağdaşmaktadır. Duygusal yeme oluşan olumsuz duygular sonucunda kontrolsüz yeme durumunu anlatmaktadır. Bu kontrolsüz yemek yeme durumu; obez bireylerde, yeme bozukluğu bulunan kadınlarda ve normal kilosunda olup da diyet yapan bireylerde gözlemlenmektedir (Albayrak, 2019). Besin tüketimi sonucunda açlıkla ilgili sitokinlerin kandaki seviyeleri azalırken, tokluk hormonları sonucu uyarılan sitokinlerin artışı ile tokluk hissini oluşur ve besin tüketimi sonlandırılır. Besin tüketiminin fizyolojik süreçlerle kontrolüne de hemostatik açlık olarak tanılandırılır (Şarahman¸2019). Ancak açlık durumunun olmadığı zamanlarda yeme isteği ve eylemi oluşabilmektedir. Homeostatik geri bildirim mekanizmasının aktif olmadığı bu gibi durumlarda gerçekleşen besin tüketimi hedonik açlık olarak tanımlanmaktadır (Şarahman¸2019). Bu gibi bireyin yediği besinlerden sadece zevk almaya odaklanması besinin tüketiminde kontrolü zorlaştırır. Besinin sadece görülmesi, kokusunun duyulması, reklam gibi uyarıcılara maruz kalınması gibi durumlar hedonik beslenmenin temelini oluşturur (Gündüz, Akhalil, Sevgi, 2020). Besinlerin tadı, görünüşü, kokusu ve diğer değişik duyusal özelliklerinden ötürü tüketilmesi sonucu oluşan haz durumu iştahı arttırır. Tok hissedilse de haz isteğinden dolayı sık sık ve çok miktarda lezzetli besinlerin tüketimi obeziteye neden olur (Özkarslı, 2022). Menstrüasyon dönemlerinde değişkenlik gösteren hormon seviyeleri sonucu artan enerji ihtiyacı sonucunda

FULL TEXTS BOOK

oluşabilecek duygusal beslenme ve hedonik açlık durumu kadın bireylerde kilo alımını kolaylaştırabilir, sağlıklı beden kitle indeksini korumada zorluk yaratabilir (Özkarslı, 2022).

Bu araştırmanın amacı; Premenstrüel sendrom (PMS), adet döngüsünün luteal fazında ortaya çıkan, fizyolojik ve psikolojik semptomlarla karakterize edilen bir durumdur. Bu dönemde hormonal dalgalanmalara bağlı olarak duygu durum değişimleri, yeme davranışlarında farklılaşma ve özellikle yüksek enerjili besinlere yönelim sıkça gözlenmektedir. Bu araştırma; premenstrüel dönemde olan üniversite öğrencilerinin değişen beslenme alışkanlıklarının hedonik ve duygusal yeme davranışı üzerine etkisinin saptanması amacı ile planlanmıştır.

YÖNTEM

Araştırmanın Yeri, Zamanı, Örneklemi

Bu araştırma Antalya'da üniversite okuyan kadın bireylerde premenstrüel dönemde değişen beslenme alışkanlıklarının hedonik ve duygusal yeme davranışlarına ne gibi etkisinin olduğunun değerlendirilmesi amacıyla hazırlanmış kesitsel bir çalışmadır. Araştırma; Kasım 2024 – Nisan 2025 ayları arasında Antalya ilinde bulunan üniversitelerde öğrenim gören kadın bireyler ile gerçekleştirilmiştir.

Araştırmanın Örneklemi; aşağıdaki formülde 192 kişi olarak bulunmuştur.

$n = N.t^2.p.q / d^2. (N-1) + t^2.p.q$

n = 383 x 3,84 x 0.5 x 0.5 / 0.0025 x 539 + 382 x 0,5 x 0,5 =192 kişi

n: Örneklem büyüklüğü=192

N: Evren birey sayısı = 383

p: incelenen olayın görülüş sıklığı (gerçekleşme olasılığı) = 0,50

q: incelenen olayın görülmeyiş sıklığı (gerçekleşmeme olasılığı) = 0,50

t: Belirli bir anlamlılık düzeyinde, t tablosuna göre (%5 hata payı) bulunan teorik değer = 1.96

d: Olayın görülüş sıklığına göre kabul edilen ± örnekleme hatasıdır. = 0,05

Araştırmaya; 18- 45 yaş aralığında, Antalya ilinde öğrenim gören, araştırmaya katılmaya gönüllü 214 katılımcı dahil edilmiştir.

Dahil edilme ölçütleri:

- 18-45 yaş arasında olmak,
- Okuma yazma engeli olmamak,
- Çalışma hakkında bilgilendirme yapıldıktan sonra çalışmaya katılmaya gönüllü olup, yazılı onam vermeyi kabul etmek,
- Antalya ilinde yaşaması,
- Kadın bireyler.

Veri Toplama Araçları

Araştırma verileri; bireylerin demografik bilgilerini, beslenme alışkanlıklarını, menstrasyon dönemlerini ele alan anket formu, Duygusal Yeme Ölçeği ve Besin Gücü Ölçeği (BGÖ) kullanılarak yüz yüze görüşme yöntemi ile elde edilmiştir.

Ankette bulunan genel bilgiler bölümünde yer alan yaş, kilo ve boy uzunluğu güncel olarak kişiler tarafından doldurulmuştur. BKI değeri araştırmacı tarafından hesaplanmıştır. Kişinin medeni durumu, öğrenim gördüğü fakülte, kronik rahatsızlık varlığı, uyku düzeni ve egzersiz yapma durumu da sorgulanmıştır. İkinci bölüm olan beslenme düzeni ile ilgili sorular da kişinin kaç öğün beslendiği, öğün atlıyorsa atlama nedeni, ara öğün tüketiyorsa ne tüketmeyi tercih ettiği sorulmuştur. Üçüncü bölümde katılımcıların menstrüasyon öncesi yaşadığı şikayetler, ilk adet gördüğü yaş, adetin kaç gün sürdüğü, adet dönemi ağrılı geçip geçmediği, adet düzeninin nasıl olduğu, ailede premenstrüel sendrom geçmişinin olup olmadığı, menstrüasyon öncesi duygu durumunun nasıl olduğu ve menstrüasyon öncesi en çok hangi besini tükettiklerinin sorgulandığı sorular bulunmaktadır.

Dördüncü bölümde "Duygusal Yeme Ölçeği" yer almaktadır. Obez ve kilolu kişilerin duygusal yeme davranışlarını değerlendirmek amacıyla geliştirilmiştir (Kawabe ve diğerleri, 2022). Duygusal yeme ölçeği, 10 maddeden ve üç alt boyuttan (yeme isteğini engelleyememe-disinhibition, yiyecek türleri-type of food, suçluluk hissi-guilt) oluşmakta, sorular 4 seçenekli ("0" Asla, "1" Bazen, "2" Genellikle ve "3" Her zaman) Likert tipi skala üzerinde yanıtlanmaktadır. Ölçekte ters madde bulunmamaktadır. Ölçekten en düşük "0" en yüksek "30" puan alınmaktadır. Ölçekten alınan yüksek puanlar, duygusal yeme davranışının yüksek düzeyde olduğunu göstermektedir. Ülkemizde geçerlik güvenirlik çalışması Arslantaş ve arkadaşları tarafından yapılmıştır. Türkçe versiyonunda da tıpkı orijinal ölçekte olduğu gibi "yeme isteğini engelleyememe", "yiyeceğin türü" ve "suçluluk hissi" olmak üzere üçlü bir faktör yapısı gösterdiği bildirilmiştir. Her bir maddenin düzeltilmiş madde toplam puan korelasyon katsayıları 0,34

FULL TEXTS BOOK

düzeyinin üzerindedir. Ölçeğin faktörlere göre oluşturulan üç boyutunun iç tutarlık katsayıları; yeme isteğini engelleyememe-disinhibition boyutu için 0,81, yiyeceğin türü -type of food boyutu için 0,57 ve suçluluk hissi-guilt boyutu için 0,64 ve ölçeğin tümü için 0,84 olarak bulunmuştur. Yazarlar araştırmacılara ölçeğin toplam puanı üzerinden çalışmalarını önermektedirler (Arslantaş ve diğerleri, 2020).

Beşinci bölüm olan son bölüm "Besin Gücü Ölçeği"'ni içermektedir. Türkçe versiyonunun geçerlilik ve güvenirliği İzzet Ülker ve arkadaşları (2020) tarafından yapılmıştır. Türkçe'ye uyarlanmış 'Besin Gücü Ölçeği', 13 maddeden oluşan bir öz bildirim anketidir ve 'hiç katılmıyorum' ile 'kesinlikle katılıyorum' arasında değişen 5'li Likert tipi yanıt ölçeğidir. Besin Gücü Ölçeği, hedonik açlık durumunun değerlendirilmesi amacıyla iştahı ölçmek için kullanılır. Besin Gücü Ölçeği, farklı besin ortamlarının psikolojik etkisini değerlendirmek için geliştirilmiştir. Besin Gücü Ölçeği, yalnızca besin ortamlarının bir ölçüsü değildir, aynı zamanda lezzetli yiyeceklerin bol miktarda ve tutarlı bir şekilde mevcut olduğu ortamlarda iştahla ilgili düşünce, duygu ve isteklerdeki bireysel farklılıkların da bir ölçüsüdür. Besin Gücü Ölçeği toplam ve alt ölçek puanları, madde puanlarının toplanması ve toplamın madde sayısına bölünmesiyle elde edilir (Ülker, Ayyıldız, Yıldıran, 2020).

Verilerin Analizi

Çalışmada elde edilen bulgular değerlendirilirken, uygun istatistiksel analizler için SPSS 20 (Statistical Package for Social Sciences) programı kullanılmıştır. Ölçümle elde edilen sürekli değişkenler (nicel değişkenler), ortalama ± standart sapma, medyan, en küçük ve en büyük değer ile sunulmuştur. Kategorik değişkenlerin (nitel değişkenlerin) sunumu için ise frekans ve yüzde değerleri kullanılmıştır. Nicel ikili grup karşılaştırmalarında parametrik varsayımlar sağlanmadığı için "Mann Whitney U Testi" kullanılmıştır. İkiden fazla nicel grup karşılaştırmalarında parametrik varsayımlar sağlanmadığı için "Kruskal Wallis Testi" kullanılmıştır. Nitel değişkenlerin analizinde ise "Ki Kare Testi" kullanılmıştır. İki nicel değişken arasındaki korelasyona bakmak için ise "Spearman Korelasyon Katsayısı" kullanılmıştır. Bütün istatistiksel analizlerde anlamlılık düzeyi "$p \leq 0,05$" olarak kabul edilmiştir.

BULGULAR

Çalışmaya toplam 214 kişi dahil olmuştur. Tablo 1 bireylerin sosyo-demografik ve sağlık durumlarına ait bazı özelliklerinin dağılımını göstermektedir. Çalışmaya dahil olan bireylerin yaş ortalaması 23.2±4.1 olarak bulunmuştur. Bireylerin %91.5'i bekar, %12.1'inin ise kronik rahatsızlığı vardır (Tablo 1).

Tablo 1. Bireylerin Sosyo-demografik ve Sağlık Durumlarının Dağılımı

		f	%
Medeni Durum	Evli	18	8.5
	Bekar	194	91.5
Kronik Rahatsızlık	Var	26	12.1
	Yok	188	87.9
Düzenli Uyku	Evet	120	56.6
	Hayır	92	43.4
Egzersiz Durumu	Evet	60	28.0
	Bazen	83	38.8

FULL TEXTS BOOK

	Hayır	71	33.2
	$\bar{X}$±SS	Medyan	Min-Maks
Yaş	23.2±4.1	22.0	18-45

$\bar{X}$: Ortalama; SS: Standart sapma; f: Frekans; %: Yüzde

Tablo 2'de çalışmaya dahil olan bireylerin menstrüasyon öncesi duygu durumları sorulmuştur. %64.5 ile çoğunluk agresif olduğunu belirtmiştir. %83.2'si ise önceki dönemde daha sık tatlı tükettiğini belirtmiştir (Tablo 2).

Tablo 2. Bireylerin Menstrüasyon Dönemindeki Duygusal Durum ve Yeme İsteği

		F	%
Menstrüasyon öncesi döneminizde duygusal durumunuz genelde nasıldır?	Depresif	90	42.1
	Agresif	138	64.5
	Üzgün	89	41.6
	Huysuz	133	62.1
	Stresli	95	44.4
	Uykulu	54	25.2
	Nötr	17	7.9
Menstrüasyon öncesi aşağıdaki besinlerden hangisini daha sık tüketirsiniz?	Tatlı çikolata vb.	178	83.2
	Turşu tuzlu vb.	46	21.5
	Limonlu ekşi şeyler vb.	37	17.3
	Acı	12	5.6
	Diğer	5	2.3

f: Frekans; %: Yüzde

Çalışmada kullanılan duygusal yeme ölçeğinin ve alt boyutlarının toplam puanları ile vücut kitle indeksi grupları arasında fark bakılmıştır. İndeks gruplarına göre yeme isteği engelleme alt boyutlarından alınan puanların farkı istatistiksel anlamlı olarak bulunmuştur ($p<0.05$). Yapılan ikili karşılaştırmalar sonunda zayıf ve normal kilolu olanlar arasında anlamlı bir fark bulunmuştur. Normal kişilerin alt boyuttan aldıkları toplam puan zayıf kişilere göre daha yüksektir. Diğer alt boyutlar ve ölçek toplam puanları arasında indeks gruplarına göre istatistiksel anlamlı bir fark bulunmamıştır ($p>0.05$) (Tablo 3).

Tablo 3. Bireylerin BKI Sınıflandırılmasına Göre Duygusal Yeme Ölçeği Puan Dağılımı

FULL TEXTS BOOK

BKİ		Toplam Duygusal Yeme Ölçeği				
	N	$\bar{X}_{\pm SS}$	Medyan	Min - Maks	p	
Zayıf	24	7.4±5.7	6.0	0-18		
Normal Kilolu	160	10.6±6.2	11.0	0-28	0.075	
Fazla Kilolu	23	11.3±5.4	10.0	0-21		
Obezite	7	14.7±10.4	18.0	0-28		
		Yeme İsteği Engelleme				
	n	$\bar{X}_{\pm SS}$	**Medyan**	**Min - Maks**	**p**	
Zayıf	24	3.8±3.2	3.0	0-10		
Normal Kilolu	160	6.2±4.1	6.0	0-18		
Fazla Kilolu	23	6.2±3.3	6.0	0-12	**0.041***	**1-2**
Obezite	7	8.4±6.7	10.0	0-17		
		Yiyecek Türü				
	n	$\bar{X}_{\pm SS}$	**Medyan**	**Min - Maks**	**p**	
Zayıf	24	2.3±1.8	2.5	0-6		
Normal Kilolu	160	2.5±1.6	2.0	0-6		
Fazla Kilolu	23	2.9±1.6	3.0	0-6	0.310	
Obezite	7	3.7±2.2	4.0	0-6		
		Susuzluk Hissi				
	n	$\bar{X}_{\pm SS}$	**Medyan**	**Min - Maks**	**p**	
Zayıf	24	1.3±1.3	1.0	0-4		
Normal Kilolu	160	1.8±1.5	2.0	0-6		
Fazla Kilolu	23	2.2±1.5	2.0	0-5	0.201	
Obezite	7	2.6±2.4	2.0	0-6		

$\bar{X}$: Ortalama; SS: Standart sapma; p: Kruskal Wallis Testi; *: p≤0.05

Çalışmada kullanılan besin gücü ölçeğinin ve alt boyutlarının toplam puanları ile vücut kitle indeksi grupları arasında fark bakılmıştır. İndeks gruplarına göre besine ulaşılabilirlik ve besin mevcudiyeti alt boyutlarından alınan puanların farkı istatistiksel anlamlı olarak bulunmuştur ($p<0.05$). Yapılan ikili karşılaştırmalar sonunda besin mevcudiyeti alt boyutunda zayıf ve fazla kilolu olanlar arasında anlamlı bir fark bulunmuştur. Fazla kilolu kişiler alt boyuttan daha yüksek puanlar almıştır. Besine ulaşılabilirlik alt boyutunda normal ve fazla kilolu olanlar arasında anlamlı bir fark bulunmuştur. Fazla kilolu kişiler alt boyuttan daha yüksek puanlar almıştır. Diğer alt boyut ve ölçek toplam puanları arasında indeks gruplarına göre istatistiksel anlamlı bir fark bulunmamıştır ($p>0.05$) (Tablo 4).

FULL TEXTS BOOK

Tablo 4. Bireylerin BKI Sınıflandırılmasına Göre Besin Gücü Ölçeği Puan Dağılımı

BKİ		Toplam Besin Gücü Ölçeği				
	n	$\bar{X}_{\pm SS}$	Medyan	Min - Maks	p	
Zayıf	24	3±0.9	2.9	1.2-4.7		
Normal Kilolu	160	2.9±0.8	3.0	1-4.8	0.082	
Fazla Kilolu	23	3.4±0.6	3.5	2.6-4.3		
Obezite	7	3.5±1.3	4.0	1.6-5		
		Besine Ulaşılabilirlik				
	n	$\bar{X}_{\pm SS}$	**Medyan**	**Min - Maks**	**p**	
Zayıf	24	2.8±1.1	2.4	1-5		
Normal Kilolu	160	2.7±0.9	2.8	1-4.8	**0.036***	**2-3**
Fazla Kilolu	23	3.3±0.7	3.3	2-4.3		
Obezite	7	3.2±1.4	3.5	1.3-5		
		Besin Mevcudiyeti				
	n	$\bar{X}_{\pm SS}$	**Medyan**	**Min - Maks**	**p**	
Zayıf	24	2.8±0.9	2.9	1-4		
Normal Kilolu	160	3±0.9	3.0	1-5	**0.044***	**1-3**
Fazla Kilolu	23	3.5±0.9	3.5	1.5-5		
Obezite	7	3.4±1.2	3.5	2-5		
		Besin Tadına Bakılması				
	n	$\bar{X}_{\pm SS}$	**Medyan**	**Min - Maks**	**p**	
Zayıf	24	3.4±1	3.3	1.6-5		
Normal Kilolu	160	3.1±0.9	3.0	1-5	0.107	
Fazla Kilolu	23	3.5±0.7	3.6	2-4.6		
Obezite	7	3.7±1.5	4.6	1.6-5		

$\bar{X}$: Ortalama; SS: Standart sapma; p: Kruskal Wallis Testi; *: p≤0.05

Duygusal yeme ölçeğinin ve alt boyutlarının toplam puanları ile menstrüasyon düzeni değişkeni arasında fark bakılmıştır. Yeme isteği engelleme alt boyutu ile menstrüasyon düzeni değişkeni arasında istatistiksel anlamlı bir fark bulunmamıştır (p>0.05). Düzene göre susuzluk hissi, yiyecek türü alt boyutları ile ölçek toplam puanları arasında istatistiksel anlamlı fark bulunmuştur (p<0.05). Yapılan ikili karşılaştırmalar sonucunda alt boyutlar için bazen düzenli bazen düzensiz menstrüasyon gören kişiler diğer gruplara göre daha yüksek puanlar almıştır. Ölçek toplamında ise bazen düzenli bazen düzensiz ile düzensiz görenler arası fark bulunmuştur. Hiç düzenli değildir diyenler ölçekten daha düşük puanlar almıştır (Tablo 5).

FULL TEXTS BOOK

Tablo 5. Bireylerin Menstrüasyon Düzenlerine Göre Duygusal Yeme Ölçeği Puan Dağılımı

Menstrüasyon Düzeni	Toplam Duygusal Yeme Ölçeği					
	n	$\bar{X}_{\pm SS}$	Medyan	Min - Maks	p	
Her ay düzenli bir şekilde olur	20	9.9±6.5	10.0	0-28		
Bazen düzensiz bazen düzenlidir	133	12.3±5.8	12.0	0-28	**0.014***	**2-3**
Hiç düzenli değildir	61	8.1±5.9	8.0	0-18		
	Yeme İsteği Engelleme					
	n	$\bar{X}_{\pm SS}$	**Medyan**	**Min - Maks**	**p**	
Her ay düzenli bir şekilde olur	20	5.9±4.3	6.0	0-18		
Bazen düzensiz bazen düzenlidir	133	6.8±3.7	6.0	0-17	0.076	
Hiç düzenli değildir	61	4.7±4.2	4.0	0-13		
	Yiyecek Türü					
	n	$\bar{X}_{\pm SS}$	**Medyan**	**Min - Maks**	**p**	
Her ay düzenli bir şekilde olur	20	2.4±1.6	2.0	0-6		
Bazen düzensiz bazen düzenlidir	133	3.2±1.6	3.0	0-6	**0.001***	**1-2**
Hiç düzenli değildir	61	2±1.4	2.0	0-5		**2-3**
	Susuzluk Hissi					
	n	$\bar{X}_{\pm SS}$	**Medyan**	**Min - Maks**	**p**	
Her ay düzenli bir şekilde olur	20	1.7±1.5	1.0	0-6		**1-2**
Bazen düzensiz bazen düzenlidir	133	2.2±1.5	2.0	0-6	**0.012***	**2-3**

FULL TEXTS BOOK

Hiç düzenli değildir	61	1.4±1.5	1.0	0-4

$\bar{X}$: Ortalama; SS: Standart sapma; p: Kruskal Wallis Testi; *: p≤0.05

Tablo 6'da ölçeklerin toplam puanları ile alt boyutlarının puanları arasında ilişki incelenmiştir. Besin gücü ölçeği ile duygusal yeme ölçeği arasında aynı yönlü zayıf anlamlı bir korelasyon bulunmuştur. (r=0.390, p<0.001) Yani besin gücü ölçeği puanındaki bir artış duygusal yeme ölçeği puanlarında da zayıf bir artışa sebep olmaktadır (Tablo 6). Besin mevcudiyeti ile besine ulaşılabilirlik arasında yüksek korelasyon olduğu bulunmuştur. Besin tadına bakılması ve besin mevcudiyeti arasında yüksek korelasyon saptanmıştır. Toplam besin gücü ölçeği ve besin tadına bakılması arasında çok yüksek korelasyon olduğu bulunmuştur. Yeme isteği engelleme ve besin tadına bakılması arasında zayıf korelasyon ilişkisi bulunmuştur. Yeme isteği engelleme ve besine ulaşılabilirlik arasında ise orta şiddette korelasyon bulunmuştur. Yiyecek türü ve besin tadına bakılması arasında zayıf korelasyon saptanırken yiyecek türü ile yeme isteği engelleme arasında yüksek korelasyon ilişkisi bulunmuştur. Susuzluk hissi ile yeme isteği engelleme ve yiyecek türü arasında orta şiddette korelasyon, kalanlarla arasında çok zayıf korelasyon saptanmıştır. Toplam duygusal yeme ölçeği ile çok yüksek korelasyon ilişkisi sırayla yeme isteği engelleme ve yiyecek türü ile bulunurken susuzluk hissi ile arasında yüksek korelasyon ilişkisi bulunmuştur.

Tablo 6. Korelasyon Tablosu

	1	**2**	**3**	**4**	**5**	**6**	**7**	**8**
	R	**r**	**r**	**r**	**r**	**r**	**r**	**R**
	P	**p**	**p**	**p**	**p**	**p**	**p**	**P**
1.Besine Ulaşılabilirlik	-							
	-							
2.Besin Mevcudiyeti	**0.717**	-						
	<0.001*	-						
3.Besin Tadına Bakılması	**0.689**	**0.672**	-					
	<0.001*	<0.001*	-					
4.Toplam Besin Gücü Ölçeği	**0.888**	**0.882**	**0.903**	-				
	<0.001*	<0.001*	<0.001*	-				
5.Yeme İsteği Engelleme	**0.407**	**0.357**	**0.230**	**0.362**	-			
	<0.001*	<0.001*	0.001*	<0.001*	-			
6.Yiyecek Türü	**0.381**	**0.428**	**0.364**	**0.436**	**0.654**	-		
	<0.001*	<0.001*	<0.001*	<0.001*	<0.001*	-		

FULL TEXTS BOOK

7.Susuzluk Hissi	**0.160**	**0.220**	0.115	**0.180**	**0.561**	**0.517**	-	
	0.019*	0.001*	0.094	0.008*	<0.001*	<0.001*	-	
8.Toplam Duygusal Yeme Ölçeği	**0.400**	**0.394**	**0.270**	**0.390**	**0.951**	**0.805**	**0.735**	-
	<0.001*	<0.001*	<0.001*	<0.001*	<0.001*	<0.001*	<0.001*	-

TARTIŞMA VE SONUÇ

Abbas ve ark. tarafından Pakistan'da yapılan bir çalışmada menstrüasyon öncesi yaşanan şikayetler arasında en çok 108 kişi (%32) bulantı, 64 kişi (%18.9) şişkinlik ve 45 kişi (%13.3) bel ağrısı şeklinde sıralanmıştır (Abbas ve diğerleri, 2020). Çoğu literatürdeki çalışma PMS'nin psikolojik ve davranışsal yan etkilerinin yanı sıra fiziksel etkileri arasında baş ağrısı, memelerde şişkinlik, hassasiyet, sırt ve karın ağrısı, şişkinlik gibi birçok etkisi olduğundan söz etmektedir (Öztürk, Tanrıverdi, 2010). Hemşirelik öğrencileri üzerinde yapılan bir çalışmada en sık yaşanan problemler sırayla karın kasık ağrısı (n=114, %67.9), memede hassasiyet (n=81, %48.2), halsizlik ve yorgunluk (n=75, %44.6). En az görülen problem 6 kişi ile uykusuzluk bulunmuştur (Kırcan ve diğerleri, 2012). Bu araştırmada da katılımcılara yöneltilen soruların sonucunda menstrüasyon öncesi yaşanan şikayetler arasında en çok karında şişlik, gerginlik ve karın ağrısı bulunmuştur (%71,5). Bunları sırasıyla 49 kişi (%22,9) baş ağrısı, 26 kişi (%%12,1) uykusuzluk, 23 kişi (%10,7) kabızlık ve 18 kişi (%8,4) baş dönmesi yaşadığını belirtmiştir. Araştırma literatürdeki çalışmalarla anlamı bir sonuç göstermektedir. Genel olarak yaşanan menstrüasyon öncesi dönemde karın ağrısı ve karında şişlik çoğunlukla kadınlardaki ortak problem olmaktadır.

Duygusal Yeme Ölçeği'nin kullanıldığı yaşları 18-36 arasındaki 633 kadın üzerinde yapılan bir çalışmada en çok (%47) düşük düzeyde duygusal yiyici, %40,4 orta düzeyde duygusal yiyici %12,4 çok duygusal yiyici bulunmuştur (Al-Musharaf, 2020). Duygusal yeme ve stress kaynaklı yemenin arasındaki ilişkinin incelendiği bir çalışmasında 124 kadın arasından 58 kadın düşük düzeyde duygusal yiyici olarak sınıflanırken 66 kadın çok duygusal yiyici olarak sınıflanmıştır (Van Strien ve diğerleri, 2012). Bu araştırmanın verilerine bakılacak olursak katılımcıların en çok (%45,3) duygusal yiyici, ardından %27,6 düşük düzeyde duygusal yiyici, %22,9 duygusal yiyici değil ve en düşük %4,2 ile çok duygusal yiyici olduğu bulunmuştur. Araştırmanın sonucunun literatürdeki diğer çalışmalarla farklılık göstermesinin nedeni her bireyin kendine özgü yaşam tarzı ve beslenme alışkanlıkları olduğu ve tek bir doğrunun olmadığı için bu tür etkenler sonuçları değiştirecektir.

Üniversite öğrencilerinin hedonik açlık durumlarının besin gücü ve duygusal yeme ölçeği ile değerlendirildiği bir çalışmaya katılan 75 kadın ve 71 erkek öğrencide BKİ değerlerine göre DYÖ puanlarından elde edilen sonuçlara göre 11 zayıf birey ölçek sonucunda en düşük 4,00 puan, en yüksek 22,00 puan ve ortalama olarak 15,2±5,09 alırken; çalışmaya katılan 6 obez kişide ise en düşük 11,00 puan, en yüksek 27,00 puan alarak ortalamaları 18,3±6,62 puan alarak en yüksek DYÖ skoru elde etmişlerdir (Çifçi, Olcay Eminsoy, 2024). Suudi Arabistan'da 359 üniversiteli kız öğrenci üzerinde yapılan bir çalışmada. Katılan bireylerin %41,2'si 'kısmen duygusal yiyici' olarak kategorize edilmiş ve Duygusal Yeme Ölçeği puan ortalaması 13.80 ± 6.8 bulunmuştur. Bir başka çalışmada BKİ ile duygusal yeme puanları arasında anlamlı bir ilişki bulunmuştur ($p<0.001$). Fazla kilolu ve obez kişilerin duygusal yeme puanları, kilosu normal olan katılımcılara kıyasla daha yüksek sonuçlanmıştır (Mansoury, 2024). Bu araştırmanın sonucuna göre yapılan ikili karşılaştırmalar sonunda zayıf ve normal kilolu olanlar arasında anlamlı bir fark bulunmuştur. Normal kişilerin alt boyuttan aldıkları toplam puan zayıf kişilere göre daha yüksektir. Diğer alt boyutlar ve ölçek toplam puanları arasında indeks gruplarına göre istatistiksel anlamlı bir fark bulunmamıştır ($p>0.05$). Burada literatüre göre farklı sonuçlar bulunmasının sebebi olarak gösterilen kaynaklarda erkek bireylerin de çalışmaya katılması sonucunda sonuçlarda değişim olmuş olabilir. Sadece kadın bireyler üzerinde yapılacak olan bu tür detaylı bir çalışma ile literatüre katkı sağlanabilir.

Karakaş ve ark. tarafından yürütülen bir çalışmada kadın ve erkek bireyler üzerindeki çalışmada BGÖ toplam puanı ortalama 2,75±0,86 zayıflarda bulunmuş, 2,62±0,70 normal kilolu bireylerde,

FULL TEXTS BOOK

2,76±0,85 hafif şişman bireylerde, 3,06±0,79 obezlerde bulunmuştur. Normal kilodaki bireylerin BGÖ toplam puanı obez katılımcılara göre daha düşük ve bu aradaki fark anlamlı bulunmuştur ($p<0,05$) (Karakaş, Saka, 2021). Öğrencilerin beslenme üzerine eğitim alıp almama durumuna göre hedonik açlık düzeyinin gözlenmesi üzerine yapılan çalışmada yaş ve cinsiyete göre beslenme ile ilgili eğitim almamış olan grubun toplam besin gücü ölçeği puanı ($r=0.126$, $p=0.014$), besinin bulunabilirliği (Faktör 1) puanları ($r=0.189$, $p<0.001$) ile beden kitle indeksi arasında pozitif yönde düşük bir ilişki olduğu bulunmuştur. Fakat beslenme ile ilgili herhangi bir eğitim alanlarla besin gücü toplam alt boyutları ile beden kitle indeksi arasında bir bağlantı bulunamamıştır ($p>0.05$) (Açar ve diğerleri, 2023). Bu araştırmadaki sonuçlara bakıldığında indeks gruplarına göre besine ulaşılabilirlik ve besin mevcudiyeti alt boyutlarından alınan puanların farkı istatistiksel anlamlı olarak bulunmuştur ($p<0.05$). Yapılan ikili karşılaştırmalar sonunda besin mevcudiyeti alt boyutunda zayıf ve fazla kilolu olanlar arasında anlamlı bir fark bulunmuştur. Fazla kilolu kişiler alt boyuttan daha yüksek puanlar almıştır. Besine ulaşılabilirlik alt boyutunda normal ve fazla kilolu olanlar arasında anlamlı bir fark bulunmuştur. Fazla kilolu kişiler alt boyuttan daha yüksek puanlar almıştır. Diğer alt boyut ve ölçek toplam puanları arasında indeks gruplarına göre istatistiksel anlamlı bir fark bulunmamıştır ($p>0.05$).

Araştırmada toplam duygusal yeme puanı açısından incelendiğinde gruplar arasında anlamlı fark bulunmuştur ($p=0.014$). Duygusal yeme puanı en yüksek olan grup, 'bazen düzensiz bazen düzenli' adet gören gruptur ($\bar{X}$=12.3±5.8), bu grup her ay düzenli adet gören grup ($\bar{X}$=9.9±6.5) ve hiç düzenli olmayanlar grubu ($\bar{X}$=8.1±5.9) takip etmiştir. Bu durum, adet düzenindeki dengesizlik ve düzensizliğin bireylerin duygu durumları üzerindeki etkisinin yeme davranışı üzerine yansıdığını göstermektedir. Özellikle hormonal dalgalanmaların yoğun olduğu luteal fazda, duygusal yeme davranışının arttığı belirtilmektedir (Klump ve diğerleri, 2013, Christensen ve Pettijohn, 2001). Bu araştırmada toplam duygusal yeme puanları sonuçlarına benzer şekilde yiyecek türü alt boyutları anlamlı fark bulunmuştur ($p=0.001$). 'bazen düzensiz bazen düzenli' adet görenler ($\bar{X}$=3.2±1.6) yiyecek türüne yönelik duygusal tepkileri daha yüksek bulunmuştur. Bu sonuç, özellikle çikolata gibi konfor yiyeceklerine yönelimin duygusal dalgalanmalara bağlı olarak arttığını göstermektedir. Hormonal değişimlerin özellikle de serotonin seviyelerini etkilediği için karbonhidrat isteğini arttırdığı yönündeki çalışmalar da araştırmadaki bulgularla örtüşmektedir (Michopoulos ve diğerleri, 2012). Araştırmada susuzluk hissi alt boyutunda da gruplar arasında anlamlı fark saptanmıştır ($p=0.012$). 'bazen düzensiz bazen düzenli' adet gören grubun ortalama puanı daha yüksek olup ($\bar{X}$=2.2±1.5), bu grubun gerçek açlık hissini yeme davranışıyla karıştırma eğiliminde olabileceği düşünülmektedir. Bu sonuç, Duygusal Yeme Ölçeği'nin yapısında belirtildiği gibi, fiziksel ve duygusal açlık ayrımının bazı bireyler için zorluk yaratabileceği desteklemektedir (Arnow, Kenardy, Agras, 1995). Son olarak araştırmada yeme isteğini engelleme alt boyutundaki gruplar arasında anlamlı bir sonuca varılamamıştır ($p=0.076$). Bu durum, menstrüasyon düzeninin yeme isteğini bastırma üzerinde bir etkisinin olmadığını söyleyebilir. Davranışa bağlı durumlar birçok bireysel farklara ve dış faktörlere bağlı olabilir.

Araştırmada bireylerin menstrüasyon düzenlerine göre DYÖ'nin karşılaştırılması sonucunda toplam duygusal yeme puanı, menstrüasyon düzenine göre anlamlı farklılık göstermiştir ($p=0.014$). 'bazen düzensiz bazen düzenli' adet gören grup en yüksek ortalamaya ($\bar{X}$=12.3±5.8), 'her ay düzenli' adet gören grupta ortalama daha düşüktür ($\bar{X}$=9.9±6.5) ve en düşük ortalamaya sahip olan grup $\bar{X}$=8.1±5.9 ortalama ile 'hiç düzenli değildir' grubuna aittir. Bu ortalama sonuçlarına göre adet düzensizliğinin bireylerin duygusal olarak dengelerini bozarak yeme davranışlarında birtakım değişikliklere sebep olabilir. Literatürdeki çalışmalarda da benzer şekilde hormonal değişimlerin duygusal yeme üzerinde etkili olduğu belirtilmiştir (Klump ve diğerleri, 2013). Özellikle de menstrüasyonun luteal fazında östrojen ve progesteron hormon seviyelerinde yaşanan değişimler yeme davranışıyla ilişkili nörotransmitter sistemlerini etkilediği çalışmalarda mevcuttur (Edler, Lipson & Keel, 2007). Araştırmada yiyecek türü alt boyutunda gruplar arasında anlamlı fark vardır ($p=0.001$). 'bazen düzensiz bazen düzenli' adet gören grubunun ortalama puanı ($\bar{X}$=3.2±1.6), diğer gruplara nazaran daha yüksektir. Bu durumda bu grupta enerji içeriği yüksek, yüksek karbonhidratlı, şekerli ve yağ içeriği yüksek besinlere yöneldiği düşünülebilir. Karbonhidrat tüketiminin serotonin artışındaki sistemle alakalı olasından ve kişide rahatlama ve kısa süreli olumlu duygular yaratmasından dolayı duygusal yeme davranışlarında bu tür karbonhidrat içerikli besinlere yönelim beklenebilir (Wurtman & Wurtman, 1995). Susuzluk hissi alt boyut sonuçlarında araştırmada menstrüasyon düzenine göre anlamlı bir fark bulunmuştur ($p=0.012$). 'bazen düzensiz bazen düzenli' adet gören grup yine en yüksek ortalamaya sahiptir ($\bar{X}$=2.2±1.5). Bu sonuç, bireylerin duygusal açlığı, fiziksel açlık veya susuzluk hissi ile

FULL TEXTS BOOK

karıştırma eğiliminde olduğu ve bu eğilimin adet düzeninden etkilenebileceği düşünülebilir. Duygusal açlık ile fiziksel açlığın ayırt edilememesi, duygusal yemenin temelini oluşturan sebeplerden biri olarak değerlendirilir (Van Strien ve diğerleri, 1986). Son olarak bu araştırmada yeme isteği engelleme alt boyutunda anlamlı bir fark gözlenmemiştir (p=0.076). Bu durum, yeme davranışının bilişsel kontrolü ile adet düzeni arasında doğrudan bir ilişki olmadığını göstermektedir. Bu bulguya yeme davranışında bireysel farklılıkların (olumsuz duygularla baş etme becerisi, bireylerin kişilik özellikleri gibi) ön planda olduğu çalışmalarla desteklenmektedir.

Bu araştırma sonucunda öncelikle PMS'nin herhangi bir adet öncesinde yaşanan hafif belirtiler olmadığı, kişilerin sosyal, psikolojik ve fiziksel anlamda özgürlüğünün kısıtlandığı patolojik bir süreç olduğu, bu dönemde çevreden yardım almakta bir sorun olmayacağının bilinmesi gereklidir. Menstrüal siklus hakkında daha çok bilgi verilmesi ve bu konuda PMS'li bireylerin bilgilendirilmesi önemlidir. Bu dönemlerde fiziksel aktivite beslenmenin öneminden bahsedilmeli ve kişinin uygulaması için çeşitli fiziksel aktiviteler ve sağlıklı bir beslenme düzenine teşvik edilmelidir. Her konuda olduğu gibi yeterli ve dengeli beslenmenin önemi bireye ve çevresindekilere anlatılmalı, hazır besinler (fast food, aşırı yağlı ve şekerli besinler vb.) ürünlerden kaçınılıp daha çok kompleks karbonhidratlar, yeterli vitamin ve mineral alımının öneminden bahsedilmelidir. Premenstrüel dönemde kişinin canının çektiği şekerli besinlerin bir yararının olmadığı sadece semptomları arttırdığının açıklanması gerekir. Beslenme önerisi olarak kompleks karbonhidratlar (tam tahıllı ürünler), Mg, Ca beslenmeye eklenip; kafein ve şeker azaltılmalıdır. Tuz tüketimi sınırlandırılmasında yarar görülebilir. Bol su tüketimi her zaman olduğu gibi PMS'de de yararı vardır. Kişilerin bu dönemlerde kendini ifade edebildiği kişilerin etrafında olması, karşısındaki kişiye kendini doğru şekilde ifade edebilmesi önemlidir. Bu dönemlerde istenmeyen sosyal aktivitelere hayır demenin kişinin ruhsal sağlığı için en iyi cevap olacağı şeklinde bireyler bilinçlendirilmelidir.

KAYNAKLAR

Süer, N. (2008). Premenstrüel Sendrom. Göztepe Tıp Dergisi 23(3):76-80

Türkçapar, A. F. ve Türkçapar, M. H. (2011). Premenstruel Sendrom ve Premenstruel Disforik Bozuklukta Tanı ve Tedavi: Bir Gözden Geçirme. Klinik Psikiyatri 14:241-253

Trout, K. K., Basel-Brown, L., Rickels, M. R., Schutta, M. H., Petrova, M., Freeman, E. W., ... & Teff, K. L. (2008). Insulin sensitivity, food intake, and cravings with premenstrual syndrome: a pilot study. *Journal of women's health*, *17*(4), 657-665.

Albayrak, M. F. (2019). 20-45 Yaş Arası Kadınlarda Premnestrüel Sendromun Beslenme Durumuna Etkisinin Saptanması. Yüksek Lisans Tezi. Marmara Üniversitesi

Şarahman, C. (2019). Yetişkin Bireylerin Hedonik Açlık Durumlarını Etkileyen Faktörlerin İncelenmesi. Yüksek Lisans Tezi. Başkent Üniveristesi

Gündüz, N., Akhalil, M., Sevgi, E. N. (2020) Hedonik Açlık. Derleme Makalesi. 3(1): 80-90

Özkarslı, Z. (2022). Yetişkin Kadınların Hedonik Açlık Durumlarını ile Mentrual Döngüleri Arasındaki İlişkinin İncelenmesi. Yüksek Lisans Tezi. Hasan Kalyoncu Üniversitesi

Kawabe R. Ve ark. (2022). The Relationship Between High Physical Activity And Premenstrual Syndrome İn Japanese Female College Students BMC Sports Science, Medicine And Rehabilitation 14:175.

Arslantaş, H., vd. (2020). Duygusal Yeme Ölçeği'nin Türkçe Çevirisinin Geçerlik ve Güvenirlik Çalışması. *Türk Psikiyatri Dergisi*, *31*(2), 122 – 130. DOI:10.5080/u23520

Ülker, İ., Ayyıldız, F., & Yıldıran, H. (2020). Validation of the Turkish version of the power of food scale in adult population. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity.*

Abbas, K., Usman, G., Ahmed, M., Qazi, R., Asghar, A., Masood Shah, A., Rizvi, A., Abid, K., Haq, K. U., Tahir, A., & Usama, S. M. (2020). Physical and Psychological Symptoms Associated With Premenstrual Syndrome and Their Impact on the Daily Routine of Women in a Low Socioeconomic Status Locality. *Cureus*, *12*(10), e10821.

Öztürk, S., Tanrıverdi, D., (2010). Premenstrual Sendrom ve Başetme. Anadolu Hemşirelik ve Sağlık Bilimleri Dergisi 13:3.

Kırcan, N., Ergin, F., Adana, F., Arslantaş, H. (2012). Hemşirelik öğrencilerinde premenstrüel sendrom prevalansı ve yaşam kalitesi ile ilişkisi.Adnan Menderes Üniversitesi Tıp Fakültesi Dergisi.

FULL TEXTS BOOK

Al-Musharaf, S. (2020). Prevalence and predictors of emotional eating among healthy young Saudi women during the COVID-19 pandemic. *Nutrients*, *12*(10), 2923.

Van Strien, T., Herman, C. P., Anschutz, D. J., Engels, R. C., & de Weerth, C. (2012). Moderation of distress-induced eating by emotional eating scores. *Appetite*, *58*(1), 277-284.

Çifçi, A., & Olcay Eminsoy, İ. (2024). Üniversite Öğrencilerinin Hedonik Açlık Durumlarının Besin Gücü ve Duygusal Yeme Ölçeği ile Değerlendirilmesi. Başkent 64

Mansoury, M. M. (2024). The Degree of Emotional Eating Among Female University Students and Its Impact on Their Pattern of Eating Behaviors. *Journal of Pioneering Medical Sciences*, *13*(3), 51-60.

Karakaş, H. M., & Saka, M. (2021). Obez Olan ve Olmayan Yetişkin Bireylerde Sezgisel Yeme Davranışının Hedonik Açlık ve Aşırı Besin İsteği İle İlişkisinin Belirlenmesi. Başkent Üniversitesi Sağlık Bilimleri Fakültesi Dergisi, 6(Beslenme Özel Sayısı), 53-69.

Açar, Y., Küçükcankurtaran, S., Sökülmez Kaya, P., Ertaş Öztürk, Y. (2023). Beslenme Eğitimi Alan Ve Almayan Üniversite Öğrencilerinde Hedonik Açlığın Değerlendirilmesi. Sağlık Bilimleri Dergisi, 32(3), 351-357.

Klump, K. L., Keel, P. K., Culbert, K. M., & Edler, C. (2013). Ovarian hormones and binge eating: Exploring associations in the menstrual cycle. *Journal of Abnormal Psychology*, 122(1), 131–137.

Christensen, L., & Pettijohn, L. (2001). Mood and carbohydrate cravings. *Appetite*, 36(2), 137-145.

Michopoulos, V., Powers, A., Gillespie, C. F., Ressler, K. J., & Jovanovic, T. (2012). Inflammation in fear- and anxiety-based disorders: PTSD, GAD, and beyond. *Neuropsychopharmacology*, 42, 254–270.

Arnow, B., Kenardy, J., & Agras, W. S. (1995). The Emotional Eating Scale: The development of a measure to assess coping with negative affect by eating. *International Journal of Eating Disorders*, 18(1), 79–90.

Edler, C., Lipson, S. F., & Keel, P. K. (2007). Ovarian hormones and binge eating in bulimia nervosa. *Psychological Medicine*, 37(1), 131-141.

Wurtman, R. J., & Wurtman, J. J. (1995). Brain serotonin, carbohydrate-craving, obesity and depression. *Obesity Research*, 3(S4), 477S-480S.

Van Strien, T., Frijters, J. E., Bergers, G. P., & Defares, P. B. (1986). The Dutch Eating Behavior Questionnaire (DEBQ) for assessment of restrained, emotional, and external eating behavior. *International Journal of Eating Disorders*, 5(2), 295-315.

FULL TEXTS BOOK

ÜNİVERSİTE YAZILIM UYGULAMALARININ MERKEZİ KİMLİK YÖNETİM SİSTEMİ İLE ENTEGRASYONU VE BÜTÜNLEŞİK PORTAL TASARIMI ÖRNEĞİ, SDUNET

INTEGRATION OF UNIVERSITY SOFTWARE APPLICATIONS WITH CENTRAL IDENTITY MANAGEMENT SYSTEM AND INTEGRATED PORTAL DESIGN EXAMPLE, SDUNET

Veli ÇAPALI [1]

[1]*Süleyman Demirel Üniversite, Bilgi İşlem Daire Başkanlığı, Isparta, Türkiye.*

[1]*ORCID ID: https://orcid.org/0000- 0002-9045-0210*

Murat KAÇAL [2]

[2]*Süleyman Demirel Üniversite, Bilgi İşlem Daire Başkanlığı, Isparta, Türkiye.*

[2]*ORCID ID: https://orcid.org/0000-0002-5257-6218*

Burak DEMİR [3]

[2]*Süleyman Demirel Üniversite, Bilgi İşlem Daire Başkanlığı, Isparta, Türkiye.*

[2]*ORCID ID: https://orcid.org/0000-0001-8044-2914*

ÖZET

Geniş ağ ve mobil teknolojilerindeki kullanım yaygınlaştıkça, kamu hizmetlerinin elektronik iletişim araçlarıyla sunumu yönündeki eğilimler artmaktadır. Otomasyon sistemlerinin ve yazılımların kamuda ve özellikle üniversitelerde kullanımının giderek yaygınlaşması daha hızlı ve kullanıcı dostu yazılımların geliştirilmesini zorunlu hale getirmiştir. Bugünlerde birbirinden bağımsız üniversite yazılım sistemleri bulunmaktadır ve bunların tek bir iletişim portalı üzerinde birleştirilmesi gereği ortaya çıkmıştır. Bu tür portallar bilginin kolayca ulaşılabildiği servislerden gelen verilerden oluşurlar. Bu servisler, farklı amaçlara yönelik geliştirilen otomasyon sistemleri ve yazılımlarında alınan web servisleridir. Portal, özellikle kurum içi bilgiye erişimi kolaylaştırır ve iş verimliliğini artırır. Bu çalışma kapsamında, Süleyman Demirel Üniversitesi bünyesinde geliştirilen yazılımların bu projeye entegre edilmesi ve tek bir portal (SDUnet) üzerinde kullanıcı bilgilendirmeleri ve erişimleri gerçekleştirilmesi örneği verilmiştir.

Anahtar kelimeler: Web servis, Kullanıcı yönetimi, OpenID Connect, OAuth 2.0.

ABSTRACT

As the use of wide network and mobile technologies become widespread, the trends toward the delivery of public services through electronic communication tools are increasing. The widespread use of automation systems and software in the government sector and especially in universities has made it necessary to develop faster and user-friendly software. Nowadays, there are independent university software systems, and it has become a necessity to combine them on a web portal. Such web portals consist of data from services where information can be easily accessed. These services are web services received in automation systems and software developed for different purposes. The portal especially facilitates access to in-house information and increases work efficiency. Within the scope of this study,

FULL TEXTS BOOK

an example of integrating the software developed within Süleyman Demirel University into this project and providing user information and access on a single portal (SDUnet) is given.

Keywords: Identity management system, Web service, Account management, OpenID Connect, OAuth 2.0.

GİRİŞ

Temel gereksinimlerin başında gelen iletişim, günümüzde sadece insanlar arasında değil web siteleri ve yazılımlar arasında da bir gereksinim olarak ön plana çıkmıştır. Web tabanlı yazılımların sahip oldukları verilerin çeşitli servisler aracılığı ile aktarılması ve birbirleri ile yazılımların iletişimde olması insanların bilişim teknolojileri yönünden ihtiyaçlarının karşılanması açısından önem kazanmıştır. Bu şekilde web uygulamalarının birleştirilmesi ve/veya beraber konuşturulduğu bir ortak yapının oluşturulduğu sistemlere Portal denilmektedir. Portal denildiğinde insanına aklına ilk gelen kekime gelen genellikle "web sitesi" dir. Ancak teknik olarak incelendiğinde portal; web siteleri ve web tabanlı yazılımlar topluluğu olarak açıklanabilir. İşlevsel anlamda ise portal; belli başlı bilgi ve verileri barındıran, servislerini bünyesinde bulunduran ve onlar arasında geçişi sağlayan ve yöneten sistemler olarak adlandırılmaktadırlar.

Otomasyon sistemlerinin ve yazılımların kamuda ve özellikle üniversitelerde kullanımının giderek yaygınlaşması daha hızlı ve kullanıcı dostu yazılımların geliştirilmesini zorunlu hale getirmiştir. Yazılımlar ve otomasyon sistemleri geliştirilirken dikkat edilmesi gereken en önemli konuların başında bu sistemlerden yararlanacak kullanıcıların belirlenmesi ve bu kullanıcıların ihtiyaçlarının yönetilmesi gelmektedir. Kullanıcılar, hızlı, kolay erişilebilir ve kullanıcı dostu yazılımlar ve otomasyonlar istemektedir. Bu üç temel özellik bizleri merkezi sistemlere yönlendirmektedir. Özellikle üniversiteler gibi kendi yazılımlarını geliştiren kurumlarda yazılımlara merkezi bir açıdan erişimin önem giderek artmaktadır. Üniversitelerde genel olarak personel otomasyonu, öğrenci işleri otomasyonu, elektronik belge yönetim sistemi, kütüphane otomasyonları, planlama ve yönetim otomasyonları, yemekhaneler ve spor tesisleri otomasyonları, kartlı geçiş kontrol sistemleri vb. otomsayon sistemleri ve yazılımları yer almaktadır. Bu sistemlerin her birine ait veri tabanlarına ve web uygulama servisleri farklı domain ile bulunmaktadırlar. Bu farklı domainlerde bulunan otomasyon ve yazılım sistemlerinin tek bir domain altına toplanarak bütünleşik bir kimlik yapısı ile erişilebilir hale getirildiği portallar günümüzde ön plana çıkmaktadır. Türkiye'deki en büyük portal E-Devlet uygulamasıdır. E-Devlet uygulaması; tüm kamu kurumlarının kendilerine ait farklı otomasyon ve yazılım sistemleri üzerinden çevrimiçi olarak sunmuş olduğu kamu hizmetlerinin servisler aracılığı ile birleştirilerek yeniden biçimlendirilmesi ve bu yolla hem vatandaşlar hem de ticari kuruluşlar ile devletin çeşitli birimleri arasında iletişim becerilerinin artırılması ile bürokratik süreçlerin otomasyon süreçlerine taşınarak daha hızlı, kolay ve erişilebilir bir yapıya ulaşılmasını sağlamıştır. "Hızlı, kolay ve erişilebilir" hizmet durumu üniversite otomasyon sistemleri içinde uygulanmalıdır. Ancak üniversitelerde kullanıcı türlerindeki çeşitlilik erişilebilirlikte oldukça farklılaşmaya neden olmaktadır.

Üniversiteler gibi kamu kuruluşlarında kullanıcı türleri oldukça değişiklik göstermektedir. Kullanıcılar sadece personel ve öğrenci olarak ayrılmazlar. Her bir uygulamaya yönelik olarak kullanıcı grupları da kendi içlerinde ayrılırlar. Akademik ve idari personelin kullandıkları amaca özel uygulamalar ile üniversite yönetiminde yer alan personelin erişimleri ve kullandıkları uygulamalarda yetki ve kullanım amaçlarında farklılıklar yer almaktadır. Benzer durum ön lisans, lisans ve lisansüstü öğrenciler içinde geçerlidir. Ön lisans ve lisans öğrencilerinin kullandıkları uygulamalara ek olarak lisans üstü öğrenciler akademik çalışmalar için farklı uygulamalara farklı yetkiler ile erişim sağlayabilmektedirler. Bu sebeple, üniversiteler içerisinde kullanıcı yetkilendirmeleri tam bir karmaşa haline gelmiştir. Bu karmaşayı

FULL TEXTS BOOK

gidermek ve kullanıcıların görev ve unvanlarına göre erişimlerinin merkezi bir kullanıcı yetkilendirme sistemi aracılığı ile yapılması ve yönetilmesi gereklidir. Kullanıcı yönetimi karmaşası aynı zamanda yazılımı geliştiren uzmanlar için de büyük bir zaman kaybına ve zorluklara neden olmaktadır. Her geliştirilen yeni bir uygulama için ayrı bir kullanıcı ve yetki hiyerarşisi tanımlaması ve şifre sistemi geliştirilmesi gerekmektedir (Sarıman vd., 2014)

Üniversiteler için, ayrık yazılımların ve kullanıcı yetkilendirme sistemlerinin bütünleşmiş hale getirilmesi konusunda araştırmalar yapılmış ve dağıtık yapıda bulunan mevcut yazılım projeleri hakkında bilgiler toplanmıştır. Kurulabilecek bir ortak kimlik platform ve portal yapısına yönelik incelemelerde bulunulmuş ve böyle bir projenin nasıl uygulanmasına yönelik analiz yapılmıştır. Bu çalışma kapsamında Süleyman Demirel Üniversitesi bünyesinde geliştirilen yazılımlar SDUnet portalına dâhil edilmekte ve tek bir sistem üzerinden kullanıcı yetkilendirme yönetimleri gerçekleştirilmektedir. Böylece farklı yazılım dillerinde geliştirilen yazlımlar ve/veya farklı yapılardaki veri tabanlarında yer alan veriler bu portal üzerine aktarılmıştır. Bu portal kullanarak daha etkin ve hızlı işlemlerin yapılabileceği model kurulması amaçlanmıştır. Ek olarak şifre ve yetkilendirme karmaşasının engellemek amacıyla kimlik yönetim sistemi ve tek noktalı kullanıcı giriş sistemi oluşturulmuştur. Böylece uygulamalardaki şifre karmaşasının önüne geçilerek tekil kullanıcı adı ve şifre kullanılmıştır. Bu çalışmanın ilk bölümünde merkezi kimlik sistemi ve yetkilendirme yönetimi ve güvenlik sertifikası yapısından, ikinci bölümünde dağıtık yapılardaki yazılımlar için kullanılacak olan web servisi mimarisinden, üçüncü bölümde ise geliştirilen portal sistemi ilgili bilgiler verilmiştir. Son bölümde ise çalışmayla ilgili sonuç ve öneriler aktarılmıştır.


YÖNTEM

Merkezi Kimlik Yönetimi (Central Identity Management)

Merkezi kimlik yönetim sistemi için Single Sign-On (SSO) oturum açma işlemleri için kullanılmaktadır. SSO bir kullanıcının birden çok uygulamaya erişmek için tek bir kimlik bilgisi kullanmasına izin veren bir oturum ve kullanıcı kimlik doğrulama hizmetidir. SSO, çeşitli kullanıcı adlarının ve parolaların yönetimini kolaylaştırmak için kuruluşlar tarafından son yıllarda oldukça yaygın bir şekilde kullanılmaktadır.

Kullanıcı bir uygulamaya giriş yapmak istediğinde bu uygulamaya ait bir çerez (cookie) oluşturulur. Paralelde bir diğer uygulamaya giriş yaptığında bu defa diğer uygulama için ayrı bir çerez oluşturulacaktır. Çünkü oluşturulacak bu çerezler benzersizdir ve her biri ayrı bir şekilde tarayıcıda işlenir. Kullanıcının aynı kullanıcı bilgilerini kullanması bir değişiklik yaratmaz ve her uygulama için bu süreci tekrarlaması gerekir. Çünkü tarayıcı bu işleme izin vermemektedir. Şekil 1’ deki diyagramda görüleceği üzere; SSO, tam bu noktada çözüm sağlar. SSO sayesinde oluşturulan bir token uygulamalar arasında işlenebilir (SSO).

FULL TEXTS BOOK




Şekil 1. SSO sisteminin işleyişi.


En yaygın kimlik doğrulama protokolleri SAML2p, WS-Federation ve OpenID Connect'tir. SAML2p en popüler ve en yaygın olarak kullanılanlardır. OpenID Connect, üçünün en yenisidir, ancak modern uygulamalar için en fazla potansiyele sahip olduğu için gelecek olarak kabul edilir. Şekil 2' deki diyagramda görüleceği üzere; OAuth2, uygulamaların bir token sağlayıcıdan (security token service) access token istemesine ve bunları API'lerle iletişim kurmak için kullanmasına olanak tanıyan bir protokoldür. Kimlik doğrulama ve yetkilendirme merkezileştirilebildiğinden, bu yetkilendirme hem istemci uygulamalarında hem de API'lerde karmaşıklığı azaltır. Kimlik doğrulama ve API erişimi, tek bir protokolde birleştirilir. IdentityServer4, OpenID Connect ve OAuth 2.0 protokollerinin bir uygulamasıdır ve günümüzün mobil, masaüstü ve web uygulamalarının tipik güvenlik sorunlarını çözmek için son derece optimize edilmiştir. IdentityServer4 kullanımında temel olarak identity token, access token ve refresh token adında üç farklı token kullanılmaktadır. Identity token, bir kimlik doğrulama işleminin sonucunu temsil eder ve kullanıcının kimliğinin nasıl ve ne zaman doğrulandığı hakkında bilgi içerir. Access token, bir API kaynağına erişime izin verir. İstemciler Access token'ları ister ve bunları API'ye iletir. Access token'lar, istemci ve kullanıcı hakkında bilgi içerir. API'ler, Access token içindeki bilgileri yetkilendirme için kullanır. Refresh token, access token geçerlilik süresi sona ermeye yaklaştığında veya sona erdiğinde yeni bir access token üretebilmek için kullanılan token'dir (IdentityServer).

Bu çalışmada, IdentityServer4 ve ASP.NET Core alt yapısı kullanılarak merkezi kimlik yönetim sistemi oluşturulmuştur. Tüm personel ve öğrencilerin verileri, ilgili otomasyon uygulamalarından web servisleri aracılığı ile periyodik olarak kimlik sisteminin veri tabanına aktarılmaktadır. Aynı zaman bu veriler ile kullanıcıların LDAP/AD kaydı yapılmaktadır. Kullanıcı giriş işlemi sürecinde; kullanıcı parolası hariç diğer bilgilerinin kontrolü bu veri tabanından yapılmakta olup, kullanıcının parola kontrolü LDAP/AD sunucusundan yapılmaktadır. Ayrıca sistemde üçüncü taraf sağlayıcılar sayesinde kimlik doğrulaması yapılabilmektedir. Örneğin E-Devlet üzerinden kullanıcı giriş işlemi gerçekleştirilmektedir. Kimlik sisteminde güvenlik seviyesini artırmak için çok faktörlü doğrulama uygulaması da mevcuttur. Bunun için, SMS ve SDÜmobil uygulamaları kullanılmaktadır. Kullanıcı giriş aşamasında, SDÜmobil uygulaması kullanıyorsa mobil uygulama üzerinden doğrulama işlemi gerçekleşirken, uygulamayı kullanmayanlar için SMS ile kimlik doğrulama gerçekleştirilmektedir (Şekil 3).

FULL TEXTS BOOK



Şekil 2. OpenID ve OAuth2 protokollerin çalışma yapısı.




Şekil 3. Merkezi kimlik yönetim sistemi giriş ara yüzü.


Merkezi Kimlik Yönetimi çözümü ile bir personelin ve öğrencinin kurumumuza katıldığı andan itibaren erişmesi gereken tüm sistemlere erişim bilgilerinin ve yetkilerinin otomatik olarak senkronize edilir. Bu sistemde, kullanıcıların e-posta hesapları, otomasyon erişim yetkileri, grup yönetimleri, kapı giriş yetkileri vb. otomatik olarak açılır. Merkezi kimlik yönetimi çözümü ile kullanıcıya ait kimlik yetkinliğinin kurum içindeki yaşam döngüsü tamamen kontrol altında tutulur. Sistem hem personelin

FULL TEXTS BOOK

işe başlama sürecinde hem de personelin işten ayrılma sürecinde geçerlidir. Aynı şekilde öğrencilerinde kayıt yapması ile süreç başlar, mezun olması ile süreç tamamlanır. Personelin ve öğrencilerin ilişki kesmesi ile merkezi kimlik yönetim sisteminde tüm yetkileri insandan bağımsız otomatik olarak kaldırılır. Kimlik verileri ihtiyaca göre tamamen silinebildiği gibi özel durumlarda etkisiz hale de getirilebilir. Üniversitemizde personel ve öğrenciler için uygulanan süreçler yer alan diyagramlarda açıklanmıştır (Şekil 4,5).




(a)




(b)

Şekil 4. (a) Personel göreve başlama ve (b) personel görevden ayrılış süreçlerinde otomasyon sistemleri akış diyagramı.

FULL TEXTS BOOK



(a)



(b)


Şekil 5. (a) Öğrenci kaydı ve (b) öğrenci ilişik kesme süreçlerinde otomasyon sistemleri akış diyagramı.

SDUnet Uygulaması (SDUnet App)

Üniversitelerde, kullanılan çevrim içi yazılımların geneli bütünleşik bir yapıda olmayıp fark alan adlarında, birbirinden bağımsız şekilde geliştirilmiş ve/veya geliştirilmektedir. Dijitalleşme sürecinde artan uygulama sayısı ve her bir hizmet için ayrı ayrı uygulamalara giriş yapılması, kullanıcıların hizmetlere ulaşımını zorlaştırır bir hale gelmiştir. Bu zorluğu aşmak için, uygulamalardan sunulan hizmetlerin tek bir platformda üzerinden merkezîleştirilmesine ihtiyaç duyulmaktadır. Bu kapsamda, E-Devlet uygulaması bir örnek olarak kabul edilmiş ve SDÜNet adında bir uygulama tarafımızdan geliştirilmiştir. Bağımsız çalışan uygulamalardaki hizmetlere Merkezi kimlik yönetim sistemi kullanılarak ek kullanıcı girişlerine ihtiyaç duyulmaksızın SDÜNet üzerinden erişim sağlanmıştır. SDÜNet, Asp.Net Core altyapısıyla geliştirilmiş ve kullanıcıların uygulamalara tek tıkla girişinin yanı sıra kullanıcılara özel profil ve ek işlemleri yapabileceği modülleri içermektedir.


FULL TEXTS BOOK

Ağ üzerindeki uygulama sunucularının birbiri ile haberleşmesi amacıyla web servisleri tasarlanmıştır ve bu servisler aynı zamanda yazılım bileşenleridir. Web servislerinin kullanımının yaygınlaşması ile iş süreçlerinde büyük bir değişiklik olmuştur. Web servisleri çok amaçlı kullanılabilirler. Öncelikli amaçları, farklı uygulama yazılımlarının birbirine entegrasyonudur. Uygulama yazılımlarının geliştirildikleri platformlar bir birinden farklı da olsa, web servisleri platform bağımsız bir şekilde veri alış verişini sağlarlar (Chen vd., 2013). SDUnet platformu, her bir otomasyon yazılımı ile Rest, SOAP, RabbitMQ teknolojileri kullanarak haberleşmektedir. Merkezi kimlik yönetim sisteminden giriş yapan kullanıcının, SDUnet kullanıcı hesabı kontrol edilir ve ilişikli yetkileri ile modüllere erişimi sağlanır. Ayrıca, bu platform sadece diğer uygulamalardan veri almayıp, üzerinde bulunan ve geliştirilmesi sürekli devam eden yeni modüllerde barındırmaktadır. Tüm personel ve öğrenci süreçlerini etkileyen ayrı uygulamaların yazılması yerine, bu uygulamalar modüler ve katmanlı mimaride SDUnet platformu üzerinde geliştirilmektedir. Bu modüllerin geliştirilmesi sırasında MVC teknolojisi, SQL Server ilişkisel veri tabanı ve MongoDB NOSQL veri tabanları kullanılmaktadır. MongoDB, hızlı olması nedeniyle ilişkisel veri tabanlarının hantal kaldığı bu tür çok kullanıcılı yapılarda tercih edilmektedir. MongoDB veri tabanında log kayıtları tutulurken, SQL veri tabanında diğer tüm veriler tutulmaktadır. SDUnet platformu hem yaklaşık 50.000 kullanıcı hizmet sağladığı için hem de yaklaşık 20 farklı araç ve uygulamayı erişim sağlayıp barındırdığından dolayı, oldukça fazla log üretimi gerçekleşmektedir. Log kayıtlarını, mevcut sistemin bulunduğu veri tabanından ayırarak sistemi çalışma performansı artırılmıştır.



(a)

FULL TEXTS BOOK



(b)

Şekil 6. (a) SDUnet platformu ve uygulama araçları (widget), (b) platform içerisinden erişilen kimlik sistemine entegre edilmiş uygulamalar.

Şekil 6(b)'de görüldüğü gibi, geliştirilen tüm uygulamalar için kullanıcı girişleri SDUnet platformuna taşınmıştır. Bu uygulamalara için ayrı ayrı giriş ve kullanıcı yetkilendirme işlemi yapmak yerine, bu platforma giriş yaptıktan sonra tüm uygulamalara kullanıcı adı ve şifre girmeksizin tanımlı yetkiler ile erişim sağlanmaktadır. Bunun için, merkezi kimlik yönetim sunucundan faydalanılır.

SDUnet platformunda yer alan araçlar (widget) aracılığı ile kullanıcıya hızlı bir şekilde bilgi görüntüleme ve işlem yapabilme imkânı sağlanmaktadır. Bu araçlar (Askıda yemek, Online ödeme, İzin bilgisi, Ödeme bilgisi, Kütüphane ödünç ve gecikme bilgisi, Rezarvasyon sistemi ve bilgisi, EBYS bilgileri, Toplantı bilgileri, SDÜ Sohbet, Ders Programı, Sınav Programı, Kartlı geçiş hareketleri, Talep Takip Sistemi, Üniversite ve Birim Duyuruları vb…), farklı platformlardaki yazılımlara servis aracılığı ile bağlanarak veri çekmekte ve yine aynı servisler aracılığı ile işlem yapabilme kolaylığını sağlamaktadır. Böylece kullanıcılar her bir işlem için veya bilgi edinmek için ayrı ayrı uygulamalara girmek erişmek zorunda değildir. SDUnet platformunda sadece kullanıcı bilgilendiren ve hızlı işlem yapmasını sağlayan araçlar yer almamaktadır. Bunlara ek olarak, tüm personel ve öğrencilerin genelinin kullanımını kapsayan yeni uygulamalar bu platformun yüzerinde geliştirilmektedir.

FULL TEXTS BOOK

TARTIŞMA VE SONUÇ

Farklı platform ve yapılarda yer alan yazılımların, ortak bir platform olan SDUnet'de bütünleşik olarak kullanılması, bilgi işlem dairesi tarafından geliştirilen uygulamalarda zaman, maliyet, erişilebilirlik ve kullanıcı dostu tasarımlar açısından önemli bir avantaj sağlamaktadır. Özellikle farklı farklı her bir uygulamaya girişte sorulan kullanıcı adı ve şifre yapısı büyük bir karmaşaya neden olmakla beraber, her bir iş için farklı uygulamaların kullanılması da bu karmaşayı artırmaktadır. Pek çok insan hem uygulamaları birbirine karıştırmakta hem de her bir uygulama da kullandıkları şifrelerini karıştırılmaktadır. Bu çalışmada Süleyman Demirel Üniversitesi Bilgi İşlem Daire Başkanlığı tarafından geliştirilen bütünleşik hesap yönetim sistemi ve servis portalı olan SDUnet anlatılmıştır. Sistem ile kullanıcılar üniversite bünyesinde geliştirilen dağıtık yazılımlara tek bir platfor üzerinden, tek bir giriş yaparak erişebilmektedir. Üniversitenin hem kendi geliştirmediği sistemlerde hem de üçüncü kişilerce geliştirilen sistemlerde, SDUnet portalı aracılığı ile uygulamalar arasında geçişlerde dahi yeniden şifre ve kimlik işlemlerine tabi olunmamaktadır.

Sistem güvenliğini sağlamak amacıyla iki adımlı kimlik doğrulama sistemi de kullanılmaktadır. İki adımlı kimlik doğrulama sisteminde kullanıcı doğrulama işlemleri için Daire Başkanlığında geliştirilen mobil uygulama ve SMS sistemleri kullanılmıştır. Üniversiteye hem yeni gelen personellere hem de yeni kaydolan öğrencilere kullanıcı adları kişiye özel tanımlanan @sdu.edu.tr ve @ogr.sdu.edu.tr uzantılı domainlere sahip olacak şekilde emsalsiz olarak üretilmektedir ve şifreleri ile bu bilgiler SMS olarak gönderilmektedir. Sisteme giriş yaptıktan hemen sonra şifreleri zorunlu olarak değiştirilerek tüm hizmetler yetkilerine göre kullanıcılar için açılır. Böylece, kullanıcılar üniversite bünyesinde aynı kullanıcı adı ile eposta hesabına sahip olur ve bu kullanıcı hesabı ile hem kablosuz-kablolu ağlara hem de SDUnet platformuna erişim sağlamış olur.

KAYNAKLAR

Chen, X., Zheng, Z., Liu, Xudong., Huang, Z., Sun, H., 2013. Personalized QoS-Aware Web Service Recommendation and Visualization, IEEE Transactions on Services Computing, 35-47.

IdentityServer, https://identityserver4.readthedocs.io/en/latest/intro/big_picture.html (Erişim Tarihi: Nisan 2024).

Sarıman, G., Karasoy, O., Tarlacı, F., Durmuş, B., 2014. Yazılımlar için Web Servis Destekli Bütünleşik Hesap Yönetimi, XVI. Akademik Bilişim Konferansı Bildirileri, 767-774.

SSO, 2024. https://ceaksan.com/tr/single-sign-on-sso-nedir (Erişim Tarihi: Nisan 2024).

FULL TEXTS BOOK

SANATSAL DİLDE METAFORLARIN STİLİSTİK-SANATSAL İŞLEVİ (NAHÇİVAN EDİBLERİNİN ESERLERİ ÜZERİNE)

THE SYSTEM OF IMAGES IN ARTISTIC LANGUAGE AND THE STYLIST-ARTISTIC FUNCTION (BASED ON THE CREATIVITY OF NAKHCHIVAN WRITERS)

СИСТЕМА ОБРАЗОВ В ХУДОЖЕСТВЕННОМ ЯЗЫКЕ И СТИЛИСТИЧЕСКО-ХУДОЖЕСТВЕННАЯ ФУНКЦИЯ (НА ОСНОВЕ ТВОРЧЕСТВА НАХЧИВАНСКИХ ПИСАТЕЛЕЙ)

Lamiye NEBİYEVA

Naxçıvan Dövlət Universitetinin doktorantı

ÖZET

Makalede, edebi ve sanatsal yaratıcılığın bir olgusu olan metafor sistemi ve sanatsal dil içindeki üslup ve sanatsal işlevleri ele alınmaktadır. Sanatsal dilin, sanatsal yaratım sürecinde önemli ve anlamlı bir manevra olduğu, düşünce ve içeriklerin ortaya çıkarılmasında başlıca araç olduğu belirtilmektedir. Dilin, edebi metinde ustalık ve ustalık meselesi olduğu gösterilmiştir. Edebi dilin mükemmelliği, metnin sanatsal ustalığına dayanır. Metafor sistemi aynı zamanda zanaatkarlık meselelerinde de etkin rol oynar ve sanatsal çalışmanın profesyonelliğine hizmet eder. Düşüncenin mecazî ve anlaşılır anlatımı açısından metaforların yerinde ve özgün kullanımı, edebî metne mükemmellik kazandırır ve başarısını sağlar. Bu bağlamda Nahçıvan yazarlarının eserlerinde metafor sistemi ustalıkla kullanılmıştır. Hem sanatsal ifade hem de tasvir araçları özgünlükle kullanıldığında bir sanat eserine mükemmellik kazandırır ve onun sanatsal kaderini belirler. Düşüncenin etkili, öz ve anlaşılır bir biçimde ifade edilmesi için kullanılan mükemmel benzetme, sıfat, metafor, metonimi, sinekdoke, abartma, litotes, tekrar, sanatsal soru vb. Metaforik örnekler yaratılarak sanatsal dilin zenginliği ortaya konuldu.

Anahtar Sözcükler: sanatsal dil, metafor sistemi, üslup ve sanatsal işlev, sanatsal tasvir ve ifade araçları, Nahçıvan yazarları.

ABSTRACT

The article discusses the system of metaphors and their stylistic-artistic function in artistic language, which is a phenomenon of literary-artistic creativity. It was noted that the artistic language is an important and important maneuver in the process of artistic creation, and it is the main tool for revealing ideas and content. It has been shown that language is a matter of mastery and craftsmanship of the literary text. The excellence of the artistic language is the basis of the artistic skill of the text. The system of metaphors takes an active place in the composition of craft issues and serves the professionalism of an artistic work. From the point of view of figurative and understandable expression of the thought, the local and original use of metaphors gives perfection to the literary text and ensures the success of its destiny. From this point of view, the system of metaphors was masterfully used in the creativity of Nakhchivan writers. By using both artistic expression and descriptive tools with originality, they bring perfection to the artistic work and decide its artistic destiny. Perfect simile, epithet, metaphor, metonymy, synecdoche, hyperbole, litota, repetition, artistic question, etc., used for effective, concise and understandable expression of thought. the richness of the artistic language was demonstrated by creating examples of metaphors.

Key words: artistic language, system of metaphors, stylistic-artistic function, means of artistic description and expression, Nakhchivan writers.

FULL TEXTS BOOK

АБСТРАКТ

В статье рассматривается система метафор и их стилистико-художественная функция в художественном языке, которая является явлением литературно-художественного творчества. Было отмечено, что художественный язык является важным и важным маневром в процессе художественного творчества, основным инструментом раскрытия идей и содержания. Показано, что язык – это вопрос владения и мастерства художественного текста. Совершенство художественного языка – основа художественного мастерства текста. Система метафор занимает активное место в композиции ремесленных вопросов и служит профессионализму художественного произведения. С точки зрения образного и понятного выражения мысли локальное и оригинальное использование метафор придает совершенство художественному тексту и обеспечивает успешность его судьбы. С этой точки зрения система метафор мастерски использовалась в творчестве нахчыванских писателей. Оригинально используя как художественную выразительность, так и описательные средства, они доводят художественное произведение до совершенства и решают его художественную судьбу. Совершенное сравнение, эпитет, метафора, метонимия, синекдоха, гипербола, литота, повтор, художественный вопрос и т. д., используемые для эффективного, краткого и понятного выражения мысли. богатство художественного языка было продемонстрировано путем создания примеров метафор.

Ключевые слова: художественный язык, система метафор, стилистически-художественная функция, средства художественного описания и выражения, нахчыванские писатели.

Giriş

Kurmaca bir eser ortaya koyan her yazar için amacına ulaşmanın ilk yolu, dili sanatsal açıdan ustalıkla kullanmak ve sanatsal beceriyi ortaya koymaktır. Mir Celal'in yazdığı gibi, "sanatsal dil, yaşamı ve varoluşu şiirsel bir biçimde, imgeler aracılığıyla sanatsal eserlerde yansıtmanın bir yoludur" [13, s.75]. Hayali bir olguyu veya imgeyi daha inandırıcı ve canlı bir biçimde sunmak, bu alanda anadilinin bütün olanaklarından ustalıkla yararlanmak, sanatkârlığın ilk şartıdır. "Sanatsallığın başlıca ve esas koşulu, toplumsal yaşamın bütün alanlarını okuyucunun gözü önünde ifade etmek ve onları sanatsal bir giysiye büründürmekti" [4, s.14]. Bu açıdan bakıldığında, sanatsal zanaatkarlık öncelikle imgede ortaya çıksa da, bunu başarmanın becerisi ve aracı ana dilin kullanımında, dil faktöründe yatmaktadır. Bu anlamda edebi bir karakterin tasvirinde dili kullanma becerisi onun sanatçılığı olarak ortaya çıkar. Sanatçının konu seçimi ve temanın tanımı, toplumsal gerçekliğe bakışı ve güncelliği değerlendirmesi, toplumsal ihtiyaçları dikkate alışı, eserdeki ideolojik ve estetik sorunlar, edebi biçim, yapı vb. Edebiyat insandan ustalık ister. Bu tür konular başka bölümlerde ele alındığı için burada Nahçıvan edebiyat çevresi temsilcilerinin sanatsal dili kullanmadaki ustalıklarını ele alacağız.

Sanatsal dil ve üslup işlevi

Dil, sanatsal yazının, metnin başlıca aracı, düşüncenin anlatımında ise başlıca, belirleyici etkendir. Yazarın dilin imkânlarını bilmesi ve onu kullanma becerisi aynı zamanda dili zenginleştirmeye yarar ve edebi dilin zenginleşmesini çeşitli yönlerden etkiler. “Sanatsal dil, yaşamı ve varoluşu şiirsel bir biçimde, imgeler aracılığıyla sanatsal eserlerde yansıtmanın bir yoludur” [13, s.75]. Büyük yazar Mirza İbrahimov, "Sanatsallık, Sanatsal Dil ve Konu" adlı yetkin makalesinde sanatsal dil hakkında şöyle yazmıştır: "Edebi yaratıcılıkta sanatsal dil... büyük önem taşır. İdealizm ve canlılığın yanı sıra, son derece sanatsal bir dil ve sağlam bir konu, bir sanat eserinin kaderini belirler" [7, s.117]. Büyük Rus

FULL TEXTS BOOK

yazarı Maksim Gorki, sanatsal dilin öneminden bahsederken şöyle yazmıştır: "Edebiyatın ilk unsuru dildir; dil, edebiyatın ilk silahıdır ve gerçekler ve yaşam olaylarıyla birlikte onun malzemesidir" [5, s.193]. Bir yazar veya şair "dili kuru, durağan bir olgu olarak kullanmaz. Aynı zamanda dille çalışır, onun potansiyel olanaklarına hayat verir, bu olanakları ortaya çıkarır ve onları geliştirir" [16, s.7]. Bütün bunların ışığında, bağımsızlık yıllarında faaliyet gösteren Nahçıvan edebiyat çevresinin yazar ve şairlerinin de sanatsal dile önem verdiklerini ve dili kullanmada ustalık gösterdiklerini belirtmek gerekir. Nahçıvan edebiyat çevresinin yetenekli şairleri arasında; Hüseyn Razi, Kemale, Hasan Fatullayev, Elman Habib, Vagif Memmedov, Asım Yadigar, İbrahim Yusifoğlu, Hüseyin Bağıroğlu ve diğerleri, yüksek düzeyde imge ve sanat yansıtan en güzel metafor örneklerini ortaya koymuşlardır. Eserlerinde bir yandan edebi dilin normlarına bağlı kalmış, ana dillerinin derin ve kapsamlı imkânlarından yararlanmaya çalışmış, diğer yandan da kendilerine özgü özgün üslup ve becerilerini ortaya koyarak, Nahçıvan lehçesinin tatlılığından yararlanarak sanatsal dil becerilerini göstermiş, edebi dili zenginleştirmeye de hizmet etmişlerdir.

Nahçıvan yazarlarının eserlerinde metaforlar ve üslup işlevi

Nahçıvan edebiyat ortamında çok ilginç üslup figürleri ve metaforlarla karşılaşılabilir, yüksek imgelem örneklerinden söz edilebilir. Yetenekli sanatçılar tarafından ilginç ve orijinal sanatsal figürler - metafor, metonimi, epitet, teşbih vb. - ve üslup figürleri - zıtlık, tekrar, abartma vb. - yaratılır. yaratılmış ve düşüncenin güçlü, etkili ve derin sanatsal ifadesi için geniş fırsatlar yaratılmıştır. Elman Habib'in bağımsızlık döneminde büyük bir yetenekle yazdığı ve Nahçıvan edebiyat ortamında gelenek başlatan "Haydarname" adlı şiirinde mübalağa ile birlikte oldukça ilginç ve orijinal abartı ve zıtlıklar kullanılmıştır. Bilindiği üzere, "abartma, varlığı büyütür, onun bütün özelliklerine ve tipik yönlerine toplumsal anlam ve kapsam kazandırır. Karakterin duyguları, eylemleri ve tutkuları büyütülür" [5, s. 193-194]. Karşıtlık aynı zamanda "imgeleri güçlendirir, şeylerin işaretlerini, anlamlarını, farklılıklarını ve çelişkilerini gerçek kılar" [5, s. 193-194]. Yazar bu bağlamda yüce bir insan imgesinin, doğaüstü ama insani bir kahramanın karakteristik izlenimini ustalıkla sunmaya çalışmış, abartma ve karşıtlık gibi sanatsal teknikleri ustalıkla kullanmıştır. Şairin kullandığı sanatsal karşıtlıklar, kahramanın sıra dışı, alışılmadık karakteristik niteliklerinin ve özelliklerinin sanatsal sunumunu ortaya çıkarmayı ve bunları okuyucuya aşılamayı başarmıştır:

Mənim dədəm dünyanın

ən böyük bir dədəsi,

Yer kürəsi balaca,

göylərdən gəlir səsi [6, s.5].

Şair, Büyük Önder Haydar Aliyev'in büyüklüğünü ve azametini abartılarla anlatmış, Büyük Önder'in büyüklüğü karşısında dünyanın küçük ve önemsiz kaldığını, hatta göklerde bile tanındığını ifade etmiştir.

Anlaşılan şair, bu ifadesiyle aynı zamanda “gökten gelen ses”in, ilahi vahyi tebliğ eden peygamberin şahsında tecelli ettiğine de dikkat çekmek istemiştir. Şair, bir başka örnekte de döngünün kötülüğünü ve kötülüğün kurnazlığını anlatmak için abartıya başvurmuştur:

Biz nəyi ki düşünək,

Hər şeyi tez duyan var,

FULL TEXTS BOOK

Gizlin düşündüyünü

ucadan oxuyan var.

Ucadan oxuyan var

Pəsdən düşündüyünü,

Tənbəkidən qoru sən

öz yanan müştüyünü [6, s.9].

Şair, düşüncelerini daha güçlü ve etkili bir biçimde ifade etmek, ilgi çekici sanatsal tekrarlar yapmak, yüksek imgeler yaratmak için üslupsal figürler olan sanatsal tekrarları da ustalıkla kullanmış, özgün örnekler ortaya koymuştur. Böylece tekrarlar "satırların anlamını, mantıksal vurgusunu, uyumunu ve bağımsızlığını derinleştirir" [5, s.201], "tekrarlar herhangi bir fikri, anlamı güçlendirmek ve ifade gücünü artırmak için kullanılır" [2, s.63]. Tekrarlama türleri olarak anafora ve epiforayı başarıyla kullanan sanatçılar, tekrarlama yoluyla anlam derinliği, amaç ve özel vurgu içeren fikirleri ifade etmeyi tercih etmişlerdir. Yetenekli şair E.Habib de sanatsal tekrarlar sayesinde anlam derinliğine ve ifade gücüne ulaşabilmiştir:

Bir hədəfə çevrildik,

yaxşı bizdik, yaman biz,

Bu yaxşını, yamanı

puç eyləyən dövran biz [6, s.5].

Və yaxud:

Kiçik, böyük torpaq, daş

- böyük olan ürəkdir,

Ürəklilər dahilər,

ürəksizlər çərəkdir [6, s.11].

İlk örnekte şair, "od" sözcüğünü iki kez kullanarak, ayrı bileşimler içinde "od" sözcüğünün tekrarını yaratmıştır - "od lekret" en zor, imkânsız bir işi üstlenip uygulamaya koyma, imkânsıza katlanma anlamında, "od oglu" ise "Azerbaycan çocuğu" anlamındadır. Benzer şekilde ikinci örnekte de "uçtu" sözcüğünü üst üste iki kez kullanarak uçuşun uzunluğunu ve sürekliliğini şiirsel bir dille dile getirmiştir. Bu tür örnekleri başka birçok sanatçının eserlerinde de bulmak mümkündür. Tamamen sanatsal tekrara dayanan şiir örneklerine bile rastlamak mümkündür. Tekrarların sanatsal işlevi ve rolü aşağıdaki örneklerde daha açık bir şekilde görülebilir:

Od götürdü od oğlu

Xəzərin sahilindən,

Oxu qoydu kamana, Azərbaycan elindən [6, s.13].

Yaxud:

Uçdu, uçdu yanar od,

neçə illər dolaşdı,

Neçə düzdən keçdi o,

neçə də dağlar aşdı [6, s.14].

FULL TEXTS BOOK

Şair Elman Habib, genel olarak "Haydarname" ve "Halkın İlhamı İlham" gibi şiirlerinde ve genel olarak yaratma sürecinde halk dilinin derin bilgeliğini ve güzelliğini ortaya koyan sözcük birimlerini ve söz dizimsel yapıları yoğun bir şekilde kullanmıştır. Eserin teması ve fikrine uygun olarak her bir dizesi derin bir anlam ve öğretici bir düşünceyi ifade etmektedir. Şair Elman Habib'in sanatının merkezinde halk bilgeliği, halk edebiyatı ve halkın yarattığı üslup ifadeleri yer alıyordu.

Şair Vagif Memmedov'un eserleri, abartı ve tekrar kullanımı bağlamında imgeleri tüm ihtişamı ve genişliğiyle ustalıkla sunması bakımından örnek teşkil etmektedir. Şair, "Büyük Adam Hakkında Küçük Şiirler" dizisinde yer alan "Büyük" şiirini Büyük Önder Haydar Aliyev'e ithaf etmiştir.

Bu dünyanın bu günləri nəğməli,

Bu dünyanın sabahları təhlükə.

Bu dünyanın taleyini bir dəli

Bilə-bilə bənd eləyir bir tükə [18, s.59];

Bir məhbəsdir sənin qəlbin,

Ora işıq düşə bilməz.

...O ürəyin damarları qan axıtmır,

...O ürəyi xəyanətlər noxtalımış,

O ürəkdən ürəklərə nifrət axdı [15, s.39];

Allah göydə, yerdə, günəşdə, ayda,

Allah dənizlərdə, göllərdə, çayda.

Allah ulduzlarda, güldə-çiçəkdə,

Allah pak, müqəddəs hər bir ürəkdə,

Allah beynimizdə, qanımızdadır,

Allah yaratdığı canımızdadır [3, s.3];

Orda Üzeyirin ayaq izləri,

Orda Vaqifimin sözləri qaldı.

...Orda torpağımın yanıq naləsi,

...Orda Vətənimin ürək parası,

...Orda qeyrət tacı Şuşa qalası

...Orda ömrümüzün yazları qaldı [8, s.13-14];

Deyirdin saf susan bulaq gözündə,

Deyirdin mən olum ancaq gözündə.

Deyirdin mələksən xalqın gözündə...[14, s.73];

Azərbaycan bir üzüksə, o üzüyün qaşı sənsən, Qarabağım,

Azərbaycan bir bədənsə, o bədənin başı sənsən, Qarabağım,

Azadlığın, enməzliyin əfsanəvi yaşı sənsən, Qarabağım.

Həm kədərli, həm sevincli gözümün də yaşı sənsən, Qarabağım.

FULL TEXTS BOOK

Sənsiz mənim ürəyim də para idi, yara idi, Qarabağım,

Sənin belə quru adın dərdlərimə çarə idi, Qarabağım.

İllərdir ki, şeytanlığın, iblisliyin beşiyində dustaq idin,

Sinəmizə çalın-çarpaz çəkdirilən bir dağ idin, Qarabağım [9, s.255].

Şair Elman Habib, genel olarak "Haydarname" ve "Halkın İlhamı İlham" gibi şiirlerinde ve genel olarak yaratma sürecinde halk dilinin derin bilgeliğini ve güzelliğini ortaya koyan sözcük birimlerini ve söz dizimsel yapıları yoğun bir şekilde kullanmıştır. Eserin teması ve fikrine uygun olarak her bir dizesi derin bir anlam ve öğretici bir düşünceyi ifade etmektedir. Şair Elman Habib'in sanatının merkezinde halk bilgeliği, halk edebiyatı ve halkın yarattığı üslup ifadeleri yer alıyordu.

Şair Vagif Memmedov'un eserleri, abartı ve tekrar kullanımı bağlamında imgeleri tüm ihtişamı ve genişliğiyle ustalıkla sunması bakımından örnek teşkil etmektedir. Şair, Ulu Önder Haydar Aliyev'e ithaf ettiği "Büyük Adam Hakkında Küçük Şiirler" serisinin "En Büyüklerden Bir Adam" başlıklı şiirinde bilge lideri şöyle tanıtıyor:

Sən ucasan, zəka səndə, nur səndə,

Şimşək səndə, alov səndə, qor səndə.

Xətainin cəsarəti var səndə,

Sən qeybdən göndərilən səsmisən?

Sən tökülən qan izini kəsmisən...[12, s.117].

Şair, toplam dokuz kez "sen" kişisel zamirini, altı kez de yüklem olarak kullanarak, Haydar Aliyev'in olağanüstü gücünü ve büyüklüğünü bir imge ve kahraman olarak nitelendirebilmiştir. Şair, büyük Cavid'in "Kim bir kaseye kan dökerse, dünyayı o kurtarır" şeklindeki meşhur dizesini farklı bir şiirsel biçimde dile getirmede de ustalığını göstermiş, vecize dolu bir dizeyle bilge lider Haydar Aliyev'in kurtarıcı ve peygamberlik misyonunu sanatkârane bir dille ifade etmeyi başarmıştır.

Halk bilgeliğine dayalı, düşüncelerini bilgece ifade etme geleneği, Nahçıvan edebiyat çevresinin yetenekli yazarlarından Muhtar Kasımzade'nin yaratıcı manevralarında da kendini göstermektedir. Şair, şiirsel ve üslupsal ustalığını, halkının zengin bilgelik çelengine güvenerek ve ana dilinin zengin olanaklarını ustalıkla kullanarak gerçekleştirmiştir. Şair, felsefi ve öğüt verici şiirlerinde düşüncelerini halk bilgeliğinin sözdizimsel yapılarını kullanarak dile getirmiştir. Şair, sanatsal tekrarlar, retorik sorular, elipsler ve diğer üslupsal figürleri kullanarak şiirsel düşüncelerini daha renkli bir şekilde ifade etmeyi de başarmıştır:

Tanrının ərki insafa,

Yoxsulun ərki azadır.

Buludun ərki yağışa,

Şairin ərki kağızadır.

Zəngin insafını,

Yoxsul azını,

Bulud yağışını,

Şair kağızını

Zümzümə edir hər gün sakitcə [11, s.4].

FULL TEXTS BOOK

Bu edebi metnin her satırında çeşitli öznelerle kıyaslanarak “irade” sözcüğünün ilişkili bir nesne olarak kullanılması ve 4 kez tekrarlanması, şairin çeşitli öznelerle kıyaslanarak iradenin özü ve anlamı üzerindeki amacını özetler. Aşağıdaki beyitlerde "zengin, fakir, bulut, şair" sözcüklerinin tekrarlanması onların günlük faaliyetlerinin gerçekliğini ifade etmektedir. Şairin bununla kalmayıp, her bir öznenin mırıldanmasını bir insan korosuna benzetmesi, kelimenin tam anlamıyla hoşa giden ve şairin yeteneğini ve becerisini yansıtan bir şiirsel buluştur.

Şairin "İşte Şuşa" şiirindeki ilgi çekici sanatsal metaforları, Şuşa'ya olan sevgisini ifade etme konusunda büyük bir beceri ve yetenek ortaya koymaktadır:

Dağların yaxasında

Boy atıb, bitib evlər.

Sanki yallı gedəcək,

Əl-ələ tutub evlər.

Ucalıb pillə-pillə,

Zirvəyə çatıb evlər.

Bir-birinin dizinə

Baş qoyub, yatıb evlər.

Şirin-şirin nəğməyə,

Nağıla batıb evlər [11, s.107].

Şuşa'nın güzel manzarasını ustalıkla tasvir eden sanatsal metaforlar şiire yeni duygular katmış ve şiirsel bir ifade kazandırmıştır.

Nahçıvan yazarlarının eserlerinde bulunan sanatsal metaforlar çarpıcı, orijinal ve ilgi çekicidir:

Metafor:

Baş alıb gedirəm dərdimlə qoşa,

Arxadan boy verən göz yaşlarıdır [9, s.9];

Deyərdim bu yurda çəkilən tumar

Bəlkə də Tanrıya salavat olur [17, s.11];

Ali Baş Komandan verəndə xəbər

Sevinir dağlarda bitən gül-çiçək,

Ey Vətən, Qələbə Günün mübarək! [1, s.34];

Dağlar da vətəndir qartallar üçün

Qucar zirvələri qoşa qanadla [11, s.26];

Mənim sözüm çiçək kimi səpildikcə ağ varağa (təşbeh),

Misra-misra dalğalanır, qanadlanır həyəcanım (metafora) [11, s.26];

FULL TEXTS BOOK

Çözüm

Böylece Nahçıvan'ın yetenekli yazarları eserlerinde sanatsal dili ustalıkla kullanmış, ilgi çekici ve orijinal sanatsal ifade ve tasvir araçları yaratmışlardır. Özellikle şiir alanında Nahçıvan yazarları, mecazın en güzel örneklerini ortaya koymuşlar ve ilginç üslup kalıplarını kullanarak düşüncelerini mecazi, derin ve etkileyici bir şekilde dile getirmişlerdir. Düşüncenin mecazlı anlatımında yazarlar, kalıplaşmış ve özgün söz sanatları, epitetler, benzetmeler, metaforlar, metonimiler, ayrıca sanatsal tekrarlar, karşıtlıklar, retorik sorular vb. kullanmışlardır. Üslupsal figürler yaratmışlardır.

ƏDƏBİYYAT

1. Babayev R. Salam, Şuşa! Naxçıvan: Əcəmi, 2021, 96 s.,
2. Bəylərova A. Bədii dildə üslubi fiqurlar. Bakı: Nurlan, 2008, 212 s.,
3. Elsevər H. Bir ürəkdir, iki can. Bakı: Gənclik, 1999, 164 s.,
4. Fuad, E. Güney Azərbaycan ədəbiyyatı: ənənəvi obrazlar yeni məzmun müstəvisində. 4-cü yazı. "Xalq cəbhəsi" qəzeti, 2019, 16 aprel, s. 14.
5. Hacıyev A. Ədəbiyyatşünaslığın əsasları. Bakı: ADPU nəşriyyatı, 2010, 370 s
6. Həbib E. İnsanlıq dastanı. Heydərnamə. Naxçıvan: Əcəmi, 2013, 464 s.
7. İbrahimov M. Xəlqilik və realizm cəbhəsindən: [məqalələr]. Bakı: Azərnəşr, 1961, 522 s.
8. Kəmalə. Ay-ulduzlu gülüstanım. Bakı: Elm və təhsil, 2014, 332 s.,
9. Kərimli X. Tanrı müjdəsi. Bakı: Qanun, 2021, 272 s.,
10. Qasımov R. Milli yaddaşın məhək daşı: Xalq şairi Məmməd Araz. Naxçıvan: Əcəmi, 2020, 206 s.
11. Qasımzadə, M. Seçilmiş əsərləri / M.Qasımzadə. – Naxçıvan: Əcəmi, - 2020. – 336 s.
12. Məmmədov V. Seçilmiş əsərləri. Üç cilddə, I c. Bakı: Şirvannəşr, 2008, 288 s.
13. Mir Cəlal, Xəlilov P. Ədəbiyyatşünaslığın əsasları. Bakı: Maarif, 1988, 280 s.
14. Nehrəmli Ə. Seçilmiş əsərləri. II-III c. Bakı: Elm və təhsil, 2018, 472 s.,
15. Razi H. Gözün aydın. Bakı: Azərnəşr, 1996, 232 s.,
16. Seyidov, Y. Ədəbi tənqid və bədii dil. // Y.Seyidov. – Bakı: Yazıçı, - 1986. – 240 s.
17. Ülvi Ə. Mən səni içimdən qovmamışam ki. Bakı: Elm və təhsil, 2021, 184 s.,
18. Yusifoğlu İ.Seçilmiş əsərləri. Beı cilddə I c. Bakı: Şirvannəşr, 2011, 340 s.,

FULL TEXTS BOOK

ARI SOKMASI SONRASI GELİŞEN ST YÜKSELMELİ MİYOKARD ENFARKTÜSÜ İLE SEYREDEN NADİR BİR KOUNİS SENDROMU TİP II OLGUSU

A RARE CASE OF KOUNIS SYNDROME TYPE II PRESENTING WITH ST-ELEVATION MYOCARDIAL INFARCTION TRIGGERED BY BEE STING

***Harun Yıldırım*[1]**

[1]Dr. Öğr. Üyesi, Kütahya Sağlık Bilimleri Üniversitesi, Tıp Fakültesi, Acil Tıp Anabilim Dalı, Kütahya, Türkiye

[1]ORCID ID: https://orcid.org/0000-0002-9161-263X

***Murtaza Kaya*[2]**

[2]Dr. Öğr. Üyesi, Kütahya Sağlık Bilimleri Üniversitesi, Tıp Fakültesi, Acil Tıp Anabilim Dalı, Kütahya, Türkiye

[2] ORCID ID: https://orcid.org/0000-0003-4012-4131

ÖZET

Giriş: Kounis sendromu, alerjik reaksiyonla tetiklenen akut koroner sendrom (AKS) tablosu olarak tanımlanır. Tip II Kounis sendromu, altta yatan aterosklerotik koroner arter hastalığı bulunan bireylerde alerjik tetikleyici ile plak rüptürü veya erozyonunun meydana geldiği durumdur. Bu sendrom nadir görülmekle birlikte, tanınması ve uygun yönetimi hayati önemdedir.

Olgu Sunumu: 65 yaşında erkek hasta, her iki bileğinden arı sokulmasını takiben başlayan göğüs ağrısı şikayetiyle acil servise başvurdu. Vital bulguları stabildi. EKG'de V2–V6 derivasyonlarında yaygın ST elevasyonu izlendi. Başlangıç yüksek duyarlılıklı troponin T değeri normal saptansa da (0.02 ng/L), seri ölçümlerde 4808 ng/L'ye kadar yükseldi. Acil olarak yapılan koroner anjiyografide sol ön inen arterde (LAD) trombotik total oklüzyon, sol ana koroner arterde %60 ve sirkumfleks arterde %80 darlık saptandı. LAD'ye ilaç salınımlı stent uygulanarak başarılı perkütan girişim yapıldı. Alerjik tetikleyicinin varlığı, EKG ve biyobelirteç dinamiği ile birlikte anjiyografik bulgular hastada **Kounis sendromu Tip II** tanısını destekledi. Hastaya dual antiplatelet tedavi, kortikosteroid ve antihistaminik verildi. Multivessel hastalığı olması nedeniyle ileri değerlendirme için kardiyovasküler cerrahi konseyine yönlendirildi.

Sonuç: Kounis sendromu, özellikle alerjik olaylar sonrasında gelişen AKS tablolarında akılda tutulmalıdır. Tip II formu, alerjik mekanizma ile aterosklerotik plak rüptürünün birlikteliğini yansıtır ve klasik miyokard enfarktüsü ile benzer tedavi gerektirir. Bu olgu, arı sokması gibi çevresel alerjenlerle tetiklenen miyokard enfarktüslerinde alerji ve iskemi yönetiminin eş zamanlı olarak yapılmasının önemini göstermektedir.

Anahtar Kelimeler: Kounis sendromu, miyokard enfarktüsü, arı sokması, alerjik koroner olay, akut koroner sendrom

ABSTRACT

Background: Kounis syndrome is defined as the concurrence of acute coronary syndrome (ACS) with a hypersensitivity reaction. Type II Kounis syndrome occurs in patients with pre-existing atherosclerotic coronary artery disease, in whom an allergic trigger precipitates plaque erosion or rupture, leading to myocardial infarction. Although rare, its prompt recognition is critical due to therapeutic implications.

Case Presentation: A 65-year-old male with no known comorbidities presented to the emergency department with chest pain that developed shortly after being stung by bees on both wrists. His vital signs were stable on arrival. Electrocardiography revealed widespread ST-segment elevations, most prominent in precordial leads (V2–V6), suggesting anterior ST-elevation myocardial infarction (STEMI). Initial high-sensitivity troponin T was within normal range (0.02 ng/L), but serial

FULL TEXTS BOOK

measurements showed a significant rise (up to 4808 ng/L). Emergent coronary angiography demonstrated thrombotic total occlusion of the proximal left anterior descending artery (LAD), 60% stenosis in the left main coronary artery (LMCA), and 80% stenosis in the circumflex artery. Drug-eluting stent implantation was performed in the LAD. Based on the temporal association with an allergic insult, absence of prior cardiac history, angiographic findings of underlying atherosclerotic disease, and dynamic biomarker elevation, a diagnosis of **Kounis Syndrome Type II** was made. The patient was treated with dual antiplatelet therapy, corticosteroids, and antihistamines, and was referred for coronary artery bypass grafting (CABG) evaluation.

Conclusion: This case highlights the importance of considering Kounis syndrome in patients presenting with ACS following allergic triggers such as insect stings. Type II Kounis syndrome represents a diagnostic and therapeutic challenge as it overlaps with classical atherothrombotic ACS. Prompt recognition and simultaneous management of both allergic and ischemic components are essential for favorable outcomes.

Keywords: Kounis syndrome, myocardial infarction, bee sting, allergic coronary event, acute coronary syndrome

GİRİŞ

Alerjik reaksiyonların kardiyovasküler sistem üzerindeki etkileri uzun süredir bilinmekle birlikte, bu etkilerin akut koroner sendrom (AKS) ile birleştiği Kounis sendromu ilk olarak 1991 yılında tanımlanmıştır (1). Kounis sendromu, mast hücresi aktivasyonu sonucu histamin, platelet aktive edici faktör ve çeşitli mediatörlerin salınımıyla meydana gelen koroner arter spazmı, aterom plağı rüptürü veya stent trombozu gibi olaylar sonucu ortaya çıkan bir akut koroner sendromdur (2). Tanı genellikle gözden kaçmakta olup, gerçek sıklığı bilinenden daha fazla olabilir (3).

Klinik olarak bu sendrom; göğüs ağrısı, dispne, hipotansiyon ve EKG değişiklikleri gibi kardiyak bulgularla birlikte cilt döküntüsü, ürtiker ve anafilaksi gibi alerjik semptomlarla birlikte seyredebilir (4). Sendrom üç alt tipe ayrılır: Tip I, normal koronerlerde vazospazmla seyreden “alerjik anjina”; Tip II, mevcut aterosklerotik plakların rüptürüyle gelişen “alerjik miyokard enfarktüsü” ve Tip III, ilaç kaplı stentlerde tromboz gelişimiyle karakterize durumdur (5).

Kounis sendromunun tetikleyicileri arasında ilaçlar (NSAID’ler, antibiyotikler), kontrast maddeleri, çevresel maruziyetler, gıdalar ve özellikle de böcek sokmaları yer almaktadır (6). Özellikle arı sokması sonrası gelişen vakalarda, hastalar kardiyovasküler risk faktörü taşımıyor olsalar dahi ciddi AKS tabloları gelişebilmektedir (5, 6). Bu yönüyle, genç yaşta ve bilinen kardiyovasküler hastalığı olmayan bireylerde alerjik semptomlar sonrası gelişen göğüs ağrısı varlığında Kounis sendromu mutlaka akla getirilmelidir (1).

Artan farkındalıkla birlikte literatürde bildirilen olgu sayılarında yükselme görülmektedir. Ancak gerek tanı zorlukları gerekse kliniğin farklı disiplinleri ilgilendirmesi nedeniyle sendrom hâlâ yeterince tanınmamaktadır (2, 4). Bu çalışmada, nadir görülen fakat potansiyel olarak hayatı tehdit edebilen Kounis sendromunun klinik özellikleri, tanı süreci ve ayırıcı tanıdaki önemi bir vaka eşliğinde ele alınmaktadır.

OLGU

65 yaşındaki erkek hasta, her iki el bileğinden arı sokulmasını takiben gelişen yaygın kaşıntı, ürtiker, baş dönmesi ve göğüs ağrısı şikayetleriyle ilçe devlet hastanesinin 1. basamak acil servisine başvurmuş, burada yapılan değerlendirmede yaygın ST elevasyonu saptanması üzerine STEMI (ST elevasyonlu miyokard enfarktüsü) ve eşlik eden alerjik reaksiyon ön tanısıyla ileri tetkik ve tedavi amacıyla merkez

FULL TEXTS BOOK

hastanemize sevk edilmiştir. Hastanemiz acil servisine getirildiğinde hipotansiyon (TA: 90/60 mmHg), taşikardi (nabız: 106/dk) ve periferik soğukluk ile birlikte oksijen satürasyonu %94 olarak değerlendirilmiş, klinik olarak anaflaktik şok tablosu ile uyumlu bulunmuştur. Hastaya uyluk anterolateralden intramüsküler olarak 0,5 mg epinefrin, intravenöz olarak 80 mg metilprednizolon ve 1 ampul feniramin maleat infüzyon şeklinde uygulanmıştır. STEMI tedavisi kapsamında 300 mg asetilsalisilik asit (oral) ve 5000 IU fraksiyone olmayan heparin (intravenöz bolus) verilmiş, 500 mL izotonik sodyum klorür solüsyonu intravenöz olarak infüze edilmiştir.


Çekilen 12 derivasyonlu EKG'de anterolateral derivasyonlarda (V1–V6, DI, aVL) belirgin ST elevasyonu, inferior derivasyonlarda (DII, DIII, aVF) karşılıklı ST depresyonu izlenmiştir (Şekil 1). Sevk edildiği ilk merkezde troponin T düzeyi 0,02 ng/L olarak bildirilmiş, hastanemize ulaştığında alınan kan tetkiklerinde ise yüksek duyarlılıklı troponin T düzeyi 4347 ng/L, AST 138,3 U/L, LDH 532 U/L, glukoz 134 mg/dL ve nötrofil oranı %94 olarak saptanmıştır. Klinik, EKG ve laboratuvar bulguları birlikte değerlendirilen hasta, Kounis sendromu tip II olasılığı ile koroner yoğun bakım ünitesine yatırılmıştır.



Şekil 1. (A) Hastanın ilk başvurduğu ilçe devlet hastanesinde çekilen 12 derivasyonlu EKG'sinde anterolateral derivasyonlarda (V2–V6, DI, aVL) ST segment elevasyonu ve inferior derivasyonlarda (DII, DIII, aVF) karşılıklı ST depresyonu izlenmektedir. (B) Hastanın merkez hastanemize kabulü sonrasında çekilen EKG'sinde, anterior derivasyonlarda belirgin ST elevasyonunun devam ettiği, yayılımın daha geniş olduğu ve taşikardinin eşlik ettiği görülmektedir.

Aynı gün yapılan koroner anjiyografide sol ön inen arterin (LAD) proksimal segmentinde tromboze total oklüzyon, sol ana koroner arterde yaklaşık %60, sol sirkumfleks arter (Cx) ve obtus marginal 1 (OM1) dalında ise %80 oranında darlık saptandı. Sağ koroner arterde de yaygın aterosklerotik plaklar izlendi. LAD lezyonuna ilaç kaplı stent kullanılarak primer perkütan koroner girişim (PCI) uygulandı (Şekil 2). Ertesi sabah yeniden gelişen göğüs ağrısı üzerine yapılan kontrol anjiyografide LAD stent açıklığının korunduğu, diğer lezyonların progresif olduğu görüldü. Kardiyovasküler cerrahi konseyi tarafından yapılan değerlendirme sonucunda, çok damar hastalığı nedeniyle elektif koroner arter bypass greft (CABG) cerrahisi planlandı.

FULL TEXTS BOOK



Şekil 2. (A) Koroner anjiyografi öncesinde sol ön inen arterin (LAD) proksimal segmentinde tromboze total oklüzyon izlenmektedir (sarı ok). (B) Primer perkütan koroner girişim (PCI) sonrası çekilen kontrol görüntüsünde, ilaçlı stent uygulamasını takiben LAD açıklığının sağlandığı görülmektedir (sarı ok).

Hastanın yoğun bakım süreci inotropik destek tedavisi altında stabil şekilde izlendi. Klinik, elektrokardiyografik, laboratuvar ve koroner anatomi bulguları birlikte değerlendirildiğinde, arı sokmasına sekonder gelişen ST elevasyonlu miyokard enfarktüsü tablosunun Kounis sendromu tip II ile uyumlu olduğu düşünüldü.

TARTIŞMA VE SONUÇ

Kounis sendromu, alerjik veya aşırı duyarlılık reaksiyonlarının koroner arter sistemini etkileyerek akut koroner sendrom tablosu oluşturduğu, tanısal açıdan karmaşık bir klinik durumdur. Rochel-Perez ve ark.'nın 155 olguyu değerlendirdikleri sistematik derlemesinde, tip II Kounis sendromunun olguların yaklaşık dörtte birini oluşturduğu, bu hastaların çoğunun ileri yaşta ve erkek olduğu, ayrıca en sık tetikleyici ajanların ilaçlar ve arı venomları olduğu bildirilmiştir (7). Cahuapaza-Gutierrez ve ark. ise 214 hastalık daha geniş serilerinde tip I formun daha baskın olduğunu, ancak tip II olguların da anlamlı bir hasta grubunu oluşturduğunu belirtmiş ve bu formda sıklıkla eşlik eden koroner arter hastalığına dikkat çekmiştir (8). Bizim olgumuzda da arı sokması sonrası gelişen anaflaktik tabloya, EKG'de ST elevasyonu ve koroner anjiyografide çok damar hastalığı eşlik etmiş olup, bu özellikler tip II Kounis sendromu ile uyumludur.

Kounis sendromunun dikkat çeken yönlerinden biri, oldukça geniş bir etiyolojik spektruma sahip olmasıdır. Bu sendrom sadece arı sokması veya antibiyotiklerle değil; lokal anestezik ajanlar, aşılar, hatta gıda maddeleri gibi çok çeşitli tetikleyicilerle ilişkilendirilmiştir. Núñez ve ark., lokal anestezik olarak kullanılan lidokainin deri testi sırasında tip I Kounis sendromuna yol açtığı bir olgu bildirmiştir (9). Allam ve ark. ise daha önce LAD'ye stent takılmış bir hastada, COVID-19 aşısı sonrası gelişen stent trombozunu tip III Kounis sendromu olarak değerlendirmiştir (10). Öte yandan Borkar ve ark. tarafından bildirilen olguda, çoklu arı sokmasına maruz kalan bir hastada gelişen miyokard enfarktüsü, hem alerjik hem de kardiyovasküler sistemin birlikte tutulduğu tip II varyant ile uyumlu bulunmuştur (11). Bu örnekler, Kounis sendromunun yalnızca nadir bir entite olmadığını; aynı zamanda heterojen, altta yatan kardiyovasküler duruma bağlı olarak farklı klinik yansımalarla karşımıza çıkabileceğini göstermektedir.

Kounis sendromunun tanısı çoğunlukla klinik şüpheye dayanır ve EKG ile kardiyak biyobelirteçlerin değerlendirilmesi sürecin merkezinde yer alır. Cahuapaza-Gutierrez ve ark.'nın sistematik

FULL TEXTS BOOK

derlemesinde, olguların %62'sinde ST elevasyonu ve %90'ında troponin yüksekliği saptandığı bildirilmiştir (8). Benchea ve ark. ise sundukları dört olguluk seride, tüm hastalarda EKG değişiklikleri ve yüksek troponin düzeyleri olduğunu, ancak tanının koroner anjiyografi sonrası kesinleştiğini vurgulamıştır (12). Bizim olgumuzda da başvuru EKG'sinde anterolateral derivasyonlarda ST elevasyonu mevcuttu ve ilk troponin normal olmakla birlikte saatler içinde binlerce ng/L düzeyine yükselmiştir. Anjiyografide LAD'de tromboze total oklüzyon ve çok damar hastalığı saptanmış; hastaya primer PCI uygulanmıştır.

Tedavi yaklaşımı hem alerjik yanıtın kontrolünü hem de miyokard iskemisinin yönetimini kapsamalıdır. Del Monaco ve ark. tarafından bildirilen arı sokmasına bağlı Kounis sendromu olgusunda, antihistaminik, steroid ve adrenalin tedavisine ek olarak primer PCI uygulanmış, bu da alerji ve kardiyoloji yönetiminin eş zamanlı yürütülmesinin önemini göstermiştir (13). Benzer şekilde, bizim hastamıza da acil serviste epinefrin, metilprednizolon ve feniramin uygulanmış; STEMI tedavisi kapsamında aspirin ve heparin başlanmıştır. Erken dönemde yapılan PCI ile LAD açıklığı sağlanmış, izleyen günlerde gelişen angina sonrası yapılan kontrol anjiyografide stent açıklığı korunmuş ancak diğer lezyonlar nedeniyle CABG planlanmıştır. Bu yaklaşım, literatürde tanımlanan multidisipliner tedavi ihtiyacını ve bireyselleştirilmiş kararların önemini bir kez daha göstermektedir.

Tanısal güçlükler nedeniyle Kounis sendromu sıklıkla yanlış sınıflandırılmakta ya da tamamen atlanmaktadır. Zisa ve ark., 12 yıllık süreçte anafilaksi tanısı almış 444 hastayı değerlendirdikleri çalışmada, sadece %2'sine (9 hasta) Kounis sendromu tanısı konulduğunu bildirmiştir. Bu olguların yalnızca ikisinde tanı taburculuk öncesinde konulabilmiş, diğerlerinde ise tanı geriye dönük olarak belirlenmiştir (14). Brancaccio ve ark. da benzer şekilde, Kounis sendromunun sıklıkla ilk başvuruda tanınmadığını ve bu nedenle eksik raporlandığını vurgulamıştır (15). Bu bulgular, özellikle eş zamanlı alerjik ve kardiyak bulgularla başvuran hastalarda Kounis sendromunun klinik şüphe kapsamında değerlendirilmesinin tanı sürecine katkı sağlayabileceğini düşündürmektedir.

Arı sokmasına bağlı gelişen anaflaktik şok ve eşzamanlı ST elevasyonlu miyokard enfarktüsü ile seyreden bu olgu, Kounis sendromu tip II'nin nadir ancak kritik bir klinik yansımasıdır. Özellikle alerjen maruziyeti sonrası kardiyak semptomlarla başvuran hastalarda Kounis sendromu ayırıcı tanıda mutlaka göz önünde bulundurulmalı ve tanı ile tedavi süreci multidisipliner bir yaklaşımla yönetilmelidir.

KAYNAKLAR

1. Alblaihed, L., & Huis In 't Veld, M. A. (2022). Allergic Acute Coronary Syndrome-Kounis Syndrome. *Emergency medicine clinics of North America*, *40*(1), 69–78. https://doi.org/10.1016/j.emc.2021.08.010

2. Poggiali, E., Benedetti, I., Vertemati, V., Rossi, L., Monello, A., Giovini, M., Magnacavallo, A., & Vercelli, A. (2022). Kounis syndrome: from an unexpected case in the Emergency Room to a review of the literature. *Acta bio-medica : Atenei Parmensis*, *93*(1), e2022002. https://doi.org/10.23750/abm.v93i1.11862

3. Lin, W. J., Zhang, Y. Q., Fei, Z., Liu, D. D., & Zhou, X. H. (2023). Kounis syndrome caused by bee sting: a case report and literature review. *Cardiovascular journal of Africa*, *34*(4), 256–259. https://doi.org/10.5830/CVJA-2022-042

4. Pejcic, A. V., Milosavljevic, M. N., Jankovic, S., Davidovic, G., Folic, M. M., & Folic, N. D. (2023). Kounis Syndrome Associated With the Use of Diclofenac. *Texas Heart Institute journal*, *50*(1), e217802. https://doi.org/10.14503/THIJ-21-7802

FULL TEXTS BOOK

5. Tsuruta, K., Yokoi, K., Yoshioka, G., Chen, W., Jojima, K., Hongo, H., Natsuaki, M., Sonoda, S., Kounis, N. G., & Node, K. (2022). Different types of Kounis syndrome caused by different episodes of bee sting anaphylaxis: Misfortunes never come singly. *Journal of cardiology cases*, *26*(1), 81–84. https://doi.org/10.1016/j.jccase.2022.03.002

6. Gopinath, B., Kumar, G., Nayaka, R., & Ekka, M. (2022). Kounis syndrome and atrial fibrillation after bee sting: A case report. *Journal of family medicine and primary care*, *11*(11), 7460–7462. https://doi.org/10.4103/jfmpc.jfmpc_901_22

7. Rochel-Perez, E., Santaularia-Tomas, M., Martin-Dorantes, M., Villareal-Jimenez, E., Olivera-Mar, A., Sanchez-Felix, E., Perez-Navarrete, A., Millet-Herrera, J. L., Huchim-Mendez, O., Alejos-Briceño, R., & Mendez-Dominguez, N. (2025). Triggers, Types, and Treatments for Kounis Syndrome: A Systematic Review. *Clinics and practice*, *15*(3), 59. https://doi.org/10.3390/clinpract15030059

8. Cahuapaza-Gutierrez, N. L., Calderon-Hernandez, C. C., Chambergo-Michilot, D., De Arruda-Chaves, E., Zamora, A., & Runzer-Colmenares, F. M. (2025). Clinical characteristics, management, diagnostic findings, and various etiologies of patients with Kounis syndrome. A systematic review. *International journal of cardiology*, *418*, 132606. https://doi.org/10.1016/j.ijcard.2024.132606

9. Garcia-Nunez, I., Algaba-Marmol, M. A., & Ignacio-Garcia, J. M. (2023). Kounis Syndrome After Lidocaine Injection. *Journal of investigational allergology & clinical immunology*, *33*(6), 506–507. https://doi.org/10.18176/jiaci.0950

10. Allam, C., Kounis, N. G., Chlawit, R., Saouma, M., & Badaoui, G. (2022). Kounis syndrome following COVID-19 vaccination. *Proceedings (Baylor University. Medical Center)*, *35*(3), 369–370. https://doi.org/10.1080/08998280.2022.2034226

11. Borkar, S. K., Hande, P., & Bankar, N. J. (2023). Kounis Syndrome: Bee Sting-Induced Acute Myocardial Infarction. *Cureus*, *15*(10), e47507. https://doi.org/10.7759/cureus.47507

12. Benchea, L. C., Anghel, L., Scripcariu, D. V., Diaconu, A., Zanfirescu, R. L., Lucaci, L. V., Bîrgoan, S. G., Sascău, R. A., Stătescu, C., & Radu, R. (2025). Kounis Syndrome in Clinical Practice: Insights from Clinical Case Series and Mechanistic Pathways. *Journal of clinical medicine*, *14*(3), 768. https://doi.org/10.3390/jcm14030768

13. Del Monaco, G., Pascucci, C., Catapano, F., Stefanini, G. G., & Ferrante, G. (2025). Case Report: Anaphylactic shock and ST-elevation myocardial infarction following a bee sting: two deadly diseases in a patient with Kounis syndrome. *Frontiers in cardiovascular medicine*, *12*, 1530829. https://doi.org/10.3389/fcvm.2025.1530829

14. Zisa, G., Panero, A., Re, A., Mennuni, M. G., Patti, G., & Pirisi, M. (2023). Kounis syndrome: an underestimated emergency. *European annals of allergy and clinical immunology*, *55*(6), 294–302. https://doi.org/10.23822/EurAnnACI.1764-1489.260

15. Brancaccio, R., Bonzano, L., Cocconcelli, A., Boyko, R., Ienopoli, G., & Motolese, A. (2024). Recurrent Kounis Syndrome: A Case Report and Literature Review. *Journal of clinical medicine*, *13*(6), 1647. https://doi.org/10.3390/jcm13061647

THE ISSUE OF SOUL AND BODY IN EUROPEAN PHILOSOPHY OF THE NEW AND MODERN ERA

Sadikhova F.Q.

PhD student at the Institute of Philosophy and Sociology of ANAS

ORCID ID 0000-0002-8958-3996

ABSTRACT

The article investigates the attitude to the issue of soul and body in New and Modern European philosophy. The views of philosophers on this issue are analyzed comparatively and the origins of the ideas are investigated. The dynamics of the ideas of the European philosophers of the New and Modern Era on the relevant topic are followed. At the same time, the influence of modern scientific discoveries and new scientific directions on modern philosophy are investigated.

Keywords: soul, body, inner "I", New Age, modern era, European philosophy, substance, consciousness.

Introduction

During the Renaissance, F. Patrizi and C. Bruno returned to the idea of the world soul in their trainings. However, there are those who oppose it on the basis that it is against Christianity. Bessarion of Nicaea emphasized that Plato's ideas about the preexistence of souls, the existence of the spirits of the sky and celestial bodies, were contrary to Christianity. Nicholas of Cusan, denying the existence of a world soul, wrote: "There is no middle ground between the absolute and the finite, as those who think that there is a soul of the world after God and before the finiteness of the world. The soul and mind of the world is only God." [1. II 9].

In the philosophy of the modern era, the terms of sense-natural and conscious bodies of the Middle Ages were replaced by the concepts of natural and artificial bodies, and the question of the substantiality of bodies, that is, their independent existence without dependence on consciousness, was raised (Hobbes) [2, p.134]. Focusing mainly on the study of material bodies, Modern philosophy related the human body to a machine mechanism.

In the future, the conflict between soul and body is renewed again by Descartes. Descartes believes that the soul is an immaterial substance. The "I" as a problem of individual self-awareness was first posed in modern philosophy and most clearly expressed by the representative of the rationalist movement. The division of reality into object and subject was a new position, not found in ancient and medieval philosophy. The philosopher is one of the three figures who profoundly shaped the history of this idea, along with Plato in antiquity and Avicenna in the Middle Ages. His concept of the soul has given rise to new discussions in modern Western thought. Descartes, who contrasts the subject with the object of knowledge, tries to find the validity of knowledge in the subject's self-awareness. The philosopher argued that it is possible to doubt the existence of external objects and the correct perception of one's own body through the senses, but it is unacceptable to doubt the existence of consciousness and the "I" to which consciousness belongs. Therefore, the thesis “cogito ergo sum” (I think, therefore I am) is true when it is understood by the human mind. [4, p. 99].

FULL TEXTS BOOK

Descartes placed the existence of two distinct substances, which he called the substance that occupies space (res extensa: the body) and the thinking substance (res cogitans: the soul), at the center of his ontology. While all animal and human organisms were conceived as automatons in the concept of the body, subject to completely mechanical rules of operation, the soul was thought to be the substance that distinguished humans from animals.[3. p.345-484]

Spinoza wanted to go beyond the Cartesian dualism, a dualistic concept of substance, in his monistic ontology, which he systematized in the identity of the terms God, nature, and substance. According to Spinoza, body and soul are in unity. The body is "a mode of expressing the essence of God in a known and definite way." [9, p.56]. The connection of ideas is the connection of things, reflecting the order of things. The content of the soul depends on the influences experienced by the body. In the course of life, the body repeatedly experiences these influences, the soul, remembering these influences, establishes a connection between them [9, c.57].

John Locke, considered the founder of the theory of associations describing human consciousness, rejected Descartes' idea of innate ideas and argued that all human knowledge comes from experience. According to Locke, a child is born as a blank slate ("tabula rasa"), on which anything can be written, which later becomes a person's own knowledge. Thus, the leading role of environmental conditions, experience in the development of a child's mental abilities is confirmed. Locke defined the body as a dense, extended substance with a definite form. Locke believed that it was impossible to prove the immortality of the soul [6, p.556.].

For the German romantics, representatives of classical German philosophy, the body is a symbolic expression of the soul and psycho-spiritual processes (K. G. Carus, Novalis, Schelling, G. Steffens). The ideas of philosophers such as Kant, Hegel, and Schelling are noteworthy in this regard.

In Kant's philosophy, the body is not seen as a thing in itself, but as a thing associated with certain cognitive faculties, that is, with a priori forms of feeling and thought. According to Kant, the immortality of the soul is a postulate of practical reason and is theoretically. Because the existence of the soul after death goes beyond the scope of possible experience. With Kant's approach, the question of the soul fundamentally moves from the field of ontology and epistemology to the sphere of morality and develops in empirical psychology, which seeks to unite with anthropology, biology, and physiology rather than philosophy [5., c.333].

Kant opposes Descartes' thesis "cogito ergo sum" and states that "I" as a thinking being is the object of an inner sense called the soul, and the object of an outer sense called the body.

Johann Gottlieb Fichte, who transformed the "thinking I" into the "pure I" and the "empirical (individual) I", does not recognize (unlike Kant) the existence of an absolute "I" independent of the individual "I" and interprets his teaching according to three principles:[10, p.284-288,290-297.]

"I=I" - the "I in itself", which reveals itself and acts as an unconditional factor for the process of self-creation, is the true beginning and source of activity of everything.

"I am not I" - the "I" distinguishes itself as an active beginning, opposing the non-I, which is inactive and part of the "I".

The opposition of "Limited I and Not-I" - Limitation creates the conditions for the emergence of a divisible "I" and "Not-I". The meaning of the synthesis lies in the difference between the absolute and individual "I", where the "Absolute I" is opposed to nature.

Friedrich Wilhelm Schelling puts forward the idea of the initial identity of the “I” and the “Not-I”, which act as the “reconciliation of opposites” of the Absolute. The identity between subject and object, conscious and superconscious, spirit and nature follow from here. In the Absolute Identity, which is a

FULL TEXTS BOOK

single whole, everything exists, and outside of it no object exists by itself. Spirit and nature are also established on this identity.

According to Friedrich Hegel, the idea takes its existence in the body of an individual being ("the soul embodies itself in the body"), which condemns it to dependence on the external forces of nature and social conditions. The soul, being embodied in the body, gives it certainty and creates the opportunity to rise to concrete certainty [10., p. 176]. According to Hegel, the soul, which expresses itself in appearance and determination, is consciousness. The distinctive feature of consciousness is the contradiction between the subject and the object. The essence of self-awareness is revealed through the mind, which fully realizes itself as the soul. It reaches its peak in absolute knowledge, which is reflected in religion. These are the stages of the phenomenological path of the soul, which coincide with the stages of the path of the individual. The source of this phenomenological dialectic is the gradual elimination of such inequality at different levels of consciousness of the "I" and the object.

In scientific and philosophical approaches from the mid-19th to the mid-20th centuries, the body was perceived as a symbolic representation of the "I" and internal-psychic processes (G. Teichmüller, F. Beneke, E. Cassirer), as the embodiment of entelechy (G. Lasson, G. Drisch),

The classical European philosophy, which considered God an indispensable part of the world, was dealt a huge blow by the philosophy of nihilism of F. Nietzsche. By declaring the death of God, he raised the issue of fundamentally and re-examining the human problem. In his approach, all ideal, spiritual and moral concepts were refuted from a philosophical point of view.

Nietzsche's consideration of the body as a means to a higher goal laid the foundation for new approaches to the body in philosophy. A. Schopenhauer considered the human body to be an expression of the will. [11, p. 346.]

J.-P. Sartre believed that the body is "an obstacle that must be overcome in order to be in the world, that is, an obstacle that I am for myself." [8.] It is possible to become a person by overcoming the weaknesses inherent in the body. For E. Husserl, the body is not only a way of being-in-the-world, but also a form of experiential attitude of consciousness that allows a person to understand his place in the space-time continuum of nature. A person's perception of his own body is the starting point for considering the world. According to Merleau-Ponty, "The body is an object that does not leave me... If the object is a certain unchanging structure, it is so not despite the change of perspectives, but in and through this change...". [7.,p.106.]

Conclusion. In the Renaissance and in the future, due to the successes of natural sciences, the body begins to be studied within the framework of science, as a natural object. In the New Age, however, the body begins to be studied as a production apparatus, and philosophical discussions about the soul gradually decrease. In the modern era, consumer culture and the morality of bodycentrism pave the way for the soul to be evaluated as an element of the body. The morality of bodycentrism led to the formation of a consumer society as a product of post-industrial society, which could not but be reflected in philosophy as a socio-cultural phenomenon. Postmodernist philosophy also approached the body-soul relationship from this aspect and laid the foundation for posthumanism views.

REFERENCES

1.Searle, J. Akıllar, Beyinler ve Bilim (trc. Kemal Bek), İstanbul 2005, 126 s.

2.Гоббс Т. О теле. Избр. произведения, т. 1. М., 1964. **[Electronic resource]** / – с. 49-60. - URL: https://refdb.ru/look/2710056.html

FULL TEXTS BOOK

3.Декарт, Р. Страсти души. Сочинения. В 2 т. М., Наука, т.1, 1989. - 504 с.

4.Декарт Р. Сочинения / ред.-сост. Т. Г. Тетенькина; пер. с фр. С. Я. Шейнман-Топштейн и др. Калининград: Янтарный сказ, 2005. - 352 с.

5.Кант, И. Критика чистого разума / пер. с нем. Н. Лосского; сверен и отредактирован Ц. Арзаканяном и М. Иткиным; прим. Ц. Арзаканяна. М.: Эксмо, 2013. - 736 с.

6.Локк, Дж. Сочинения в 3-х т.: Т. I/Ред.: И. С. Нарский, А. Л. Субботин; Ред. I т., авт. вступит, статьи и примеч. И. С. Нарский; Пер. с англ. А. Н. Савина. М.: Мысль, 1985. 621 с.

7.Морис Мерло-Понти. Феноменология восприятия. **[Electronic resource]** - (Перевод с французского под редакцией И. С. Вдовиной, С. Л. Фокина) Санкт-Петербург "Ювента" "Наука" - 1999, 245 с. URL: https://www.booksite.ru/localtxt/pon/tym/erl/ovo/text.pdf

8.Сартр Ж.-П. Бытие и нечто: Опыт феноменологической онтологии. – **[Electronic resource] -** М., 2000 – 639 с. URL: http://yanko.lib.ru/books/ philosoph/sartre=butie_i_nichto=ann.htm

9.Спиноза, Бенедикт. Этика. Переводчик: Боровской Я. М., Иванцов Н. А. Издательство: Азбука, 2012 г. 352 с.

10.Фихте И. Г. Сочинения. Работы 1792-1801 гг. / подг. П. П. Гайденко. М.: Ладомир, 1995. 656 с.

11.Шопенгауэр А. Мир как воля и представление, **[Electronic resource]** / Т.2 / Пер. с нем. - Мн.: ООО "Попурри", 1999. 832 с. URL: http://www.lib.ru/FILOSOF/SHOPENGAUER/mir.txt

PEYGAMBERİMİZİN MUCİZELERİNDEN OLAN NESEBİ DURUMU

Zeynep Başaran

Batman Üniversitesi\ Sosyal Bilimler Enstitüsü

Temel İslam Bilimleri Ana Bilim Dalı

Yüksek Lisans Öğrencisi

ÖZET

Hz. Muhammed'in (s.a.v.) peygamberliğine delalet eden mucizeler arasında, soyunun temizliği ve seçkinliği özel bir yer tutmaktadır. İslami kaynaklar, onun nesebinin şirkten, zinadan ve ahlaki sapmalardan uzak bir şekilde, nesilden nesile tertemiz bir zincirle Hz. İbrahim'e kadar uzandığını belirtmektedir. Kur'an'da doğrudan ifade edilmese de, peygamberlerin seçilmiş soylar içinden gönderildiğine dair ayetler bu inancı destekler (En'âm 6/84–87). Hadislerde ise Peygamberimiz, "Allah beni en hayırlı nesilden seçti" diyerek bu ilahi seçimi açıkça beyan etmiştir. İslam alimleri, Hz. Peygamber'in atalarının puta tapmayan, iffetli ve doğru kişiler olmalarını, onun nübüvvetine hazırlık ve bir çeşit **mucizevî korunmuşluk** olarak değerlendirmişlerdir. Bu bağlamda Hz. Peygamber'in nesebi, sadece tarihsel değil; aynı zamanda ilahi hikmet ve inayetin bir tecellisi olarak görülmüş, mucizevi yönüyle İslam düşüncesinde ayrıcalıklı bir yere sahip olmuştur.

Anahtar kelimeler: Hz. Muhammed, mucize, nesep.

ABSTRACT

Among the miracles that attest to the Prophethood of the Prophet Muhammad (peace be upon him), the purity and distinction of his lineage holds a special place. Islamic sources state that his lineage, free from polytheism, adultery, and moral deviation, extends from generation to generation, through a pure chain all the way back to Abraham. Although not directly stated in the Quran, verses stating that prophets are sent from chosen lineages support this belief (Al-An'am 6:84–87). In the hadiths, the Prophet clearly stated this divine choice, saying, "Allah chose me from the best of lineages." Islamic scholars have interpreted the fact that the Prophet's ancestors were non-idolater, chaste, and truthful individuals as preparation for his prophethood and a form of miraculous protection. In this context, the Prophet's lineage is not only historical; it has also been viewed as a manifestation of divine wisdom and grace, and its miraculous aspect has earned it a privileged place in Islamic thought.

Keywords: Prophet Muhammad, miracle, lineage

GİRİŞ

Peygamber Efendimiz Hz. Muhammed'in (s.a.v.) hayatı ve şahsiyeti İslam düşüncesinin temelini teşkil eder. Onun peygamberliğine delalet eden mucizeleri sadece harikulade olaylardan ibaret değil, aynı zamanda tarihsel, sosyal ve ahlaki bağlamlarıyla da ele alınmalıdır. Bu mucizelerden biri de nesebinin seçkinliği, yani soyunun tertemiz ve şerefli bir şekilde intikalidir. İslami kaynaklar, Peygamberimizin nesebinin Hz. İbrahim'e (a.s.) kadar uzanan bir silsilede, şirk, ahlaksızlık ve gayrimeşru ilişkilerden uzak, iffetli ve temiz bir soyla geldiğini ifade etmektedir.

FULL TEXTS BOOK

Hz. Peygamber'in **nesebinin temizliği ve seçilmişliği**, İslam alimleri tarafından hem tarihî hem de akidevî bir olgu olarak ele alınmış, nübüvvetin bir delili ve mucizevi yönlerinden biri olarak kabul edilmiştir. Hadislerde bizzat Hz. Peygamber'in kendi ifadesiyle soyunun şirkten, zinadan ve ahlaki düşüklükten uzak bir silsileyle Hz. İbrahim'e (a.s.) kadar uzandığı bildirilmektedir. Bu yönüyle onun nesebi, sadece biyolojik bir silsile değil; Allah'ın peygamberlik için özel olarak koruduğu, hazırladığı bir "şecere-i tayyibe"dir.

1.Nesebin Anlamı ve Önemi

Nesep, Arapça kökenli bir kavram olup "soy, sülale, ata silsilesi" anlamına gelir. İslam öncesi Arap toplumunda soy, bireyin toplumsal statüsünü belirleyen temel unsurlardan biri olarak kabul edilirdi. İslam ile birlikte nesep anlayışı sadece biyolojik bağ değil, aynı zamanda manevi bir sorumluluk olarak da anlam kazanmıştır. Bu bağlamda, Hz. Peygamber'in nesebi de sadece bir soy zinciri değil, onun peygamberlik misyonunu hazırlayan ilahi bir takdirin tecellisidir.[1]

2.Mucize Kavramı

Mucize, Arapça kökenli bir kelime olup "acz" kökünden türemiştir ve "i'câz" mastarından türeyen **ism-i fâil** kalıbındadır. Bu yönüyle "güçsüz bırakma, karşı konulamaz hale getirme" anlamı taşır ve "kudret"in zıddı olarak değerlendirilmektedir.[2] Kelimenin çoğulu **mu'cizât** şeklindedir. "İ'câz" kavramı bazen az sözle çok anlam ifade etme anlamındaki "îcâz" ile karıştırılsa da, aralarında anlam yönünden önemli bir fark vardır. "İ'câz", kelimenin asli anlamına uygun olarak, muhatabı aciz bırakma, onun benzerini getirmekten âciz olduğunu ortaya koyma anlamında kullanılır.[3] Bu bağlamda **"mucize",** sözlük anlamı itibariyle "güçsüz bırakan, karşı konulamaz şey" demektir.

2.Hz. Peygamberimizin Nesebi

Hz. Muhammed'in (s.a.v.) nesebi, onun peygamberliğine hazırlık sürecinin en dikkat çekici ve mucizevi yönlerinden biridir. Kur'an-ı Kerim'de Hz. Peygamber'in nesebi doğrudan zikredilmese de dolaylı yoldan onun temiz bir soydan geldiğine işaret eden ayetler mevcuttur. En'âm Suresi 84–89. ayetlerde Hz. İbrahim ve soyundan gelen peygamberlerin zikredilmesi, Hz. Muhammed'in bu silsilenin son halkası olduğunu ima etmektedir. Ayrıca Şuarâ Suresi 214. ayette yer alan "*Önce en yakın akrabanı uyar*" emriyle, onun kabilesi ve soyu üzerinden tebliğ sürecine başlaması istenmiştir.[4]

"Devamlı olarak zina edenlerden değil, nikâh ehli temiz babalardan bana geçmiştir"[5]

Bu rivayet, Resûlullah'ın soyunun hiçbir şekilde gayrimeşru bir ilişkiye bulaşmadığını ve hep meşru evlilik yoluyla devam ettiğini vurgulamaktadır. Bu da onun sadece bireysel ahlâkının değil, soyunun da ilâhî murakabe altında olduğunu gösteren bir mucizedir.[6]

Kur'an'da geçen "*O sizi karınlarda dilediği gibi şekillendirir*"[7] ve "*Allah, her şeyi en güzel şekilde yapandır*"[8] gibi ayetler, bu sürecin ilâhî iradeyle yönlendirildiğini düşündürmektedir. Ayrıca Hz.

[1] Mahmud Es'ad Coşan, *Siyer ve Hadislerle Efendimiz*, İstanbul: Server Yayınları, 1999, s. 44.
[2] el-Bâkillânî, *el-Beyan 'ani'l-Fırak beyne'l-Mucizat ve'l-Keramat*, Beyrut 1950, s.8; et-Taftazânî, *Şerhu'l-Mekasıd*, V, 11; en-Nebhânî, *Hüccetüllâhi ale'l-alemin fi Mu'cizati Seyyidî'l-Murselin*, Dîyarbakır, trs., s.13.

[3] er-Râzî, Ebû Bekir, *Muhtaru's-Sıhah*, İstanbul, 1400/1980, s.354; el-Isfehânî, *Müfredât*, s.334; İbn Kemal, *Resail*, s.137.
[4] M. Asım Köksal, *İslâm Tarihi*, C.1, İstanbul: Diyanet İşleri Başkanlığı Yayınları, 1991, s. 47-48.
[5] İbn Sa'd, *Tabakât*, I, 60.
[6] Müslim, *Fezâil*, 1; Ayrıca bkz. İbn Hacer, *Fethu'l-Bârî*, Kahire: Mektebetü's-Selâm, 1959, VIII, 71.
[7] Âl-i İmrân, 3/6.
[8] Secde, 32/7.

FULL TEXTS BOOK

Peygamber'in soyu, Kâbe hizmetini yürüten, toplum içinde saygın, mertlik ve misafirperverlik gibi erdemlerle tanınan Kureyş kabilesinin Haşimoğulları koluna dayanmaktadır. Bu da onun toplum nezdindeki meşruiyetini artıran bir faktördür.[9]

Hadislerde ise Hz. Peygamber bizzat şöyle buyurmuştur:

"Allah, mahlûkatı yarattı ve beni en hayırlılarının en hayırlısından, en temizlerinin en temizinden çıkardı."[10]

Hadis, onun soyunun hem maddi hem manevi anlamda tertemiz olduğunu ortaya koymaktadır. İslam alimleri bu durumu "istikra" yöntemiyle, yani tarih boyunca soy zincirindeki hiçbir ferdin şirk, zina, zulüm gibi kötülüklerle lekelenmemesi şeklinde açıklamışlardır.

Peygamberimizin soyunun korunmuşluğu bir nevi mucize olarak kabul edilir. Zira her insanın geçmişi çeşitli sapmalarla dolu olabilirken, Hz. Peygamber'in soyu özenle seçilmiş kimselerden oluşmaktadır. İmam Kurtubî, bu konuda şöyle ifade etmektedir:

"Hz. Peygamber'in soyu, cahiliye döneminde bile iffetiyle meşhur olan ve putlara tapmayan kimselerden meydana gelmiştir. Bu da onun nübüvvetini destekleyen mucizevî bir haldir."[11]

Özellikle Hz. Peygamber'in atası olan Abdülmuttalib'in tevhid inancı, ahlaki yapısı ve Kâbe hizmetindeki rolü, bu temizliğin işaretlerinden biridir. Bu durum, sadece biyolojik bir zincir değil, aynı zamanda inanç ve değerlerin korunmasını sağlayan manevi bir silsile olarak da anlaşılmalıdır.[12] İslam kaynaklarında Hz. Muhammed'in soyu Hz. İbrahim'in oğlu Hz. İsmail'e dayandırılır. Bu silsile şu şekilde özetlenebilir:

Muhammed b. Abdullah b. Abdülmuttalib b. Hâşim b. Abdümenaf b. Kusayy b. Kilâb b. Mürre b. Ka'b b. Lüeyy b. Gâlib b. Fihr b. Mâlik b. Nadr b. Kinâne b. Huzeyme b. Müdrike b. İlyâs b. Mudar b. Nizâr b. Ma'd b. Adnân…[13]

Hz. Peygamber'in mucizelerinden biri olan "nesebinin seçilmişliği", onun peygamberlik görevine hazırlandığı sürecin en önemli halkalarından biridir. Soyunun şirkten, iffetsizlikten ve ahlaki sapmalardan uzak olması, onun sadece biyolojik bir mirasla değil, manevi bir temizlikle de desteklendiğini göstermektedir. Bu durum, Allah'ın peygamberlik görevi için en uygun şahsiyeti hazırladığını ortaya koymaktadır. Bu yönüyle Hz. Muhammed'in (s.a.v.) nesebi, onun risaletini destekleyen mucizevî bir zemin olarak değerlendirilmelidir.[14]

3.Nesebin Mucizevîliği

Hz. Muhammed'in (s.a.v.) nübüvvetinden önceki hayatı ve onunla bağlantılı gelişmeler, İslam literatüründe yalnızca tarihsel olaylar olarak değil, aynı zamanda ilahi bir takdirin yansımaları olarak değerlendirilmiştir. Bunların başında ise, **"temiz nesep" mucizesi** gelir. Bu mucize, Hz. Peygamber'in soyu boyunca şirk, ahlaksızlık ve gayrimeşru ilişkilerden uzak, iffetli ve tevhid üzere yaşayan kimselerden meydana geldiği inancına dayanır. Bu özelliği, onu sadece biyolojik bir soy zincirinin mensubu değil, aynı zamanda risalet için özel olarak hazırlanmış bir yapının ferdi kılmaktadır.

[9] İbn Kesîr, *el-Bidâye ve'n-Nihâye*, Beyrut: Mektebetü'l-Maârif, 1988, C.2, s. 269.
[10] Müslim, *Fazâil*, 1; Tirmizî, *Menâkıb*, 1.
[11] Müslim. *Sahih Müslim*, Kitâbü'l-Fazâil.
[12] Kurtubî, *el-Câmiʿ li-Ahkâmi'l-Kur'ân*, Beyrut: Dâr İhyâi't-Türâsi'l-Arabî, 2006, c. 6, s. 38.
[13] İbn Hişam, *es-Sîretü'n-Nebeviyye*, Beyrut: Dârü'l-Ma'rifa, 1990, c. 1, s. 1–3.
[14] Süyûtî, *el-Hâvî li'l-Fetâvâ*, Kahire: Mektebetü'l-Cezîre, 2004, c. 2, s. 456.; Gazâlî, *İhyâu Ulûmi'd-Dîn*, Kahire: Dârü'l-Hadîs, 2005, c. 2, s. 244.

FULL TEXTS BOOK

İslam âlimleri, Hz. Peygamber'in soyunun şirkten, putperestlikten ve zinadan korunmuş olmasını doğrudan Allah'ın bir müdahalesi, yani mucize olarak değerlendirmişlerdir. Bu görüş, şu hadise dayandırılır:

"Allah, mahlûkatı yarattı ve beni en hayırlılarının en hayırlısından, en temizlerinin en temizinden çıkardı."

Bu rivayet, Allah'ın Hz. Peygamber'i nübüvvet için önceden hazırladığını ve onun soyunu da bu doğrultuda özel olarak seçtiğini göstermektedir. Nitekim Süyûtî, bu hadis üzerinde yaptığı açıklamada, "bu nebevi silsilenin zinadan, şirkten ve büyük günahlardan korunmuş olmasının Allah'ın bir lütfu ve mucizesi olduğunu" ifade etmektedir.

İslam akaidinde peygamberlerin günahlardan korunmuşluğu anlamına gelen **ismet** sıfatı, genellikle peygamberlik süreciyle ilişkilendirilmektedir. Ancak bazı alimler, bu sıfatın onların **soylarını da kapsayacak şekilde genişletilebileceğini** ileri sürmüşlerdir. Bu görüşe göre, Allah Teâlâ, peygamberlerin ortaya çıkacağı aileleri ve soy zincirini özel olarak koruma altına almıştır. Dolayısıyla Hz. Peygamber'in atalarının şirk koşmamış, iffetsizliğe bulaşmamış kimseler olması bir mucize değilse bile, mucizevi bir tertibin parçası olarak değerlendirilmiştir.

Kur'an'da, peygamberlerin neseben birbirine bağlı olması özellikle vurgulanmıştır:

"Ve İbrahim'e, İshak ve Yakub'u bağışladık. Her birini doğru yola ilettik. Daha önce de Nuh'u hidayete erdirmiştik. Onun soyundan Davud'u, Süleyman'ı, Eyyub'u, Yusuf'u, Musa'yı ve Harun'u da..."[15]

Bu ayet, **peygamberlik silsilesinin soy üzerinden intikal ettiğini** ve bu soyun Allah tarafından özel olarak yönlendirildiğini ortaya koyar. Hz. Peygamber'in Hz. İbrahim'in oğlu Hz. İsmail soyundan gelmesi de bu ilahi zincirin Araplar içindeki devamı olarak görülmüştür. Burada, nesebin yalnızca toplumsal değil, **mukaddes ve mucizevi bir vasıta olduğu** anlaşılmaktadır.

Bazı tasavvufî ve kelâmî eserlerde Hz. Peygamber'in soyu "**şecere-i tayyibe**" yani "mübarek ve temiz ağaç" olarak nitelenmiştir. Bu benzetme, hem Kur'ânî bir metaforun (İbrahim, 14/24) hem de peygamberliğin ilahi plan içinde nasıl bir arınmışlıkla taşındığını ifade etmektedir.[16] İbn Arabî, bu soyu "vahyî hakikatlerin taşıyıcısı" olarak görür ve şöyle der:

"O (Hz. Muhammed), nübüvvet ağacının meyvesidir ve bu ağaç, tüm dallarıyla Allah'ın himayesindedir."[17]

Bu yaklaşım, nesebin ilahi hikmetle şekillendiğini ve dolayısıyla Hz. Peygamber'in soyunun da mucizevî bir zeminde geliştiğini ortaya koymaktadır.

Hz. Muhammed'in (s.a.v.) nesebinin mucizevîliği, İslam düşüncesinde hem hadisler hem de Kur'an ayetlerinin ışığında değerlendirilmiş; onun soyunun şirk, ahlaksızlık ve bâtıl inançlardan korunmuşluğu bir ilahi mucize olarak görülmüştür.

SONUÇ

Hz. Peygamber'in nesebiyle ilgili mucizevi durum, onun soyunun tertemiz, faziletli ve seçkin bireylerden meydana gelmesiyle kendini göstermektedir. İslam kaynaklarında, onun atalarının hiçbiri şirk lekesine bulaşmamış, ahlaki düşüklük sergilememiş ve nesebi hiçbir zaman gayrimeşru bir çizgiye

[15] En'âm, 6/84.
[16] Süyûtî, *el-Hâvî li'l-Fetâvâ*, Kahire: Mektebetü'l-Cezîre, 2004, c. 2, s. 456.
[17] İbn Arabî, *el-Fütûhâtü'l-Mekkiyye*, Kahire: Dârü'l-Kitâbi'l-Arabî, c. 3, s. 128.

FULL TEXTS BOOK

sapmamıştır. Bu durum, hem Kur'ân-ı Kerîm'deki "*Sen, kesinlikle en yüce bir ahlâk üzeresin*"[18] ayetiyle örtüşmekte, hem de peygamberliğe hazırlanış sürecinin ilâhî bir tasarrufla yürütüldüğünü göstermektedir. Ayrıca Hz. Peygamber'in "*Ben, evlilik yoluyla meydana gelen oğullar zinciriyle gelmişim. Asla zina lekesi bulaşmamıştır*."[19] mealindeki hadisleri, bu seçilmişliğin sadece bireysel değil, nesebe dayalı bir tertemizlikle de desteklendiğini ortaya koymaktadır. Dolayısıyla onun nesebi, hem bir mucize hem de risalet görevinin hikmetli bir zemin üzerinde yükseldiğinin delilidir. Bu durum, sadece biyolojik bir aktarım değil; aynı zamanda nübüvvetin, toplumsal kabul ve itibara uygun bir temelde ortaya çıktığını da vurgulamaktadır.

KAYNAKÇA

Coşan, Mahmud Es'ad. *Siyer ve Hadislerle Efendimiz*. İstanbul: Server Yayınları, 1999.

el-Bâkillânî, *el-Beyan 'ani'l-Fırak beyne'l-Mucizat ve'l-Keramat*, Beyrut 1950.

en-Nebhânî, *Hüccetüllâhi ale'l-alemin fi Mu'cizati Seyyidî'l-Murselin*, Dîyarbakır, trs.

er-Râzî, Ebû Bekir, *Muhtaru's-Sıhah*, İstanbul, 1400/1980.

et-Taftazânî, Sa'düddîn, *Şerhü'l-Mekâsıd*, Beyrût, 1410/1989, II, 173.

Gazâlî, İmam. *İhyâu Ulûmi'd-Dîn*. Kahire: Dârü'l-Hadîs, 2005.

İbn Arabî, *el-Fütûhâtü'l-Mekkiyye*, Kahire: Dârü'l-Kitâbi'l-Arabî, c. 3.

İbn Hacer el-Askalânî. (1959). *Fethu'l-Bârî bi-Şerhi Sahîhi'l-Buhârî* (Cilt 8). Kahire: Mektebetü's-Selâm.

İbn Hişam. *es-Sîretü'n-Nebeviyye*. Beyrut: Dârü'l-Ma'rifa, 1990.

İbn Kemal,Şemseddin Ahmed b. Süleyman, *Resail*, nşr. Ahmed Cevdet, İstanbul, 1316.

İbn Kesîr, İ. (1988). *el-Bidâye ve'n-Nihâye* (Cilt 2). Beyrut: Mektebetü'l-Maârif.

Köksal, M. A. (1991). *İslâm Tarihi* (Cilt 1). İstanbul: Diyanet İşleri Başkanlığı Yayınları.

Kurtubî. *el-Câmiʿ li-Ahkâmi'l-Kur'ân*. Beyrut: Dâr İhyâi't-Türâsi'l-Arabî, 2006.

Müslim b. Haccâc. (t.y.). *Sahîh-i Müslim*, Kitâbu'l-Fezâil.

Müslim. *Sahih Müslim*, Kitâbü'l-Fazâil.

Süyûtî. *el-Hâvî li'l-Fetâvâ*. Kahire: Mektebetü'l-Cezîre, 2004.

Tirmizî. *Sünen-i Tirmizî*, Kitâbü'l-Menâkıb.

[18] Kalem, 68/4.
[19] Müslim, Fezâil, 2.

AlGaAs/GaAs ÇOKLU KUANTUM KUYULARININ ELEKTRİK VE MANYETİK ALAN ETKİSİ ALTINDA OPTİK ÖZELLİKLERİNİN İNCELENMESİ

THE INVESTIGATION OF THE OPTICAL PROPERTIES OF AlGaAs/GaAs MULTIPLE QUANTUM WELLS UNDER THE INFLUENCE OF ELECTRIC AND MAGNETIC FIELDS

Dr. Öğr. Üyesi Didem ALTUN

Sivas Cumhuriyet Üniversitesi, Sivas Teknik Bilimler MYO, Elektrik ve Enerji Bölümü, Sivas

ÖZET

Bu çalışma, AlGaAs/GaAs çoklu kuantum kuyularında artan bariyer genişliklerinin, elektrik alan ve manyetik alanların birleşik etkileri altında optik geçirgenlik ve kırılma indisi değişimlerine olan etkisini teorik olarak incelemeyi amaçlamaktadır. Etkin kütle yaklaşımı ve sonlu elemanlar yöntemi kullanılarak, periyodik olarak artan bariyer genişliğine sahip kuantum kuyularının enerji seviyeleri ve dalga fonksiyonları hesaplanmıştır. Elektrik alan (F=0, ±30 kV/cm) ve manyetik alan (B=0, 15 T) uygulamalarının potansiyel profili, enerji seviyeleri ve elektron bulunma olasılıkları üzerindeki etkileri detaylı bir şekilde analiz edilmiştir. Doğrusal ve doğrusal olmayan absorbsiyon katsayıları ile kırılma indisi değişimleri, yoğunluk matrisi yaklaşımı kullanılarak elde edilmiştir. Sonuçlar, elektrik ve manyetik alanların, bariyer genişliği ile birlikte optik geçiş enerjilerini, dipol matris elemanlarını ve optik özellikleri önemli ölçüde modüle ettiğini göstermektedir. Özellikle, (1-2) ve (2-3) geçişlerinde maviye kayma gözlenirken, (3-4) geçişlerinde kayma minimal düzeyde kalmıştır. Bu bulgular, optimize edilmiş optoelektronik cihaz tasarımları için teorik bir temel sunarak, kuantum kuyusu tabanlı optik modülatörler, dedektörler ve lazerler gibi uygulamalarda yeni fırsatlar yaratma potansiyeline sahiptir.

Anahtar Kelimeler: AlGaAs/GaAs, Elektrik alan, Manyetik alan, Absorbsiyon katsayısı, Kırılma indisi

ABSTRACT

This study aims to theoretically investigate the effects of increasing barrier widths on the optical absorption coefficient and refractive index changes in AlGaAs/GaAs multiple quantum wells under the combined influence of electric and magnetic fields. Using the effective mass approximation and the finite element method, the energy levels and wave functions of quantum wells with periodically increasing barrier widths were calculated. The effects of electric field (F=0, ±30 kV/cm) and magnetic field (B=0, 15 T) applications on the potential profile, energy levels, and electron probability distributions were analyzed in detail. Linear and nonlinear absorption coefficients, as well as refractive index changes, were obtained using the density matrix approach. The results demonstrate that electric and magnetic fields, in conjunction with barrier width, significantly modulate optical transition energies, dipole matrix elements, and optical properties. Notably, a blue shift was observed in the (1-2) and (2-3) transitions, while the shift in the (3-4) transition remained minimal. These findings provide a theoretical foundation for optimized optoelectronic device designs, offering potential for new opportunities in applications such as quantum well-based optical modulators, detectors, and lasers.

Keywords: AlGaAs/GaAs, Electric field, Magnetic field Absorption coefficient, Refractive index.

FULL TEXTS BOOK

GİRİŞ

Yarı iletken kuantum kuyuları, düşük boyutlu kuantum sistemlerinin benzersiz özelliklerinden dolayı optoelektronik ve fotonik cihazların geliştirilmesinde temel bir rol oynamaktadır. Özellikle AlGaAs/GaAs tabanlı kuantum kuyuları, yüksek kristal kalitesi, uygun bant aralığı ve etkili kuantum sınırlama özellikleri sayesinde optik modülatörler, kızılötesi dedektörler ve yüksek hızlı iletişim cihazları gibi çok çeşitli uygulamalarda kullanılmaktadır (Bastard, 1988). Bu yapılar, yük taşıyıcılarının kuantum kuyusu içerisindeki enerji seviyelerinin kuantizasyonu nedeniyle dikkat çekici optik ve elektronik özellikler sergiler. Elektrik alan (F) ve manyetik alanların (B) bu sistemlere uygulanması, enerji seviyelerini ve dalga fonksiyonlarını değiştirerek absorbsiyon katsayısı ve kırılma indisi gibi optik özelliklerde önemli değişikliklere yol açar. Bu özellikler de kuantum kuyularını optik anahtarlar, dedektörler ve lazerler gibi ileri teknoloji cihazlarının temel bileşenleri haline getirmiştir (Huang ve ark., 1995, Aishah ve ark., 2022, Alaydin, 2021)

Literatürde, kuantum kuyularında elektrik alan etkilerinin ve manyetik alan etkilerinin ayrı ayrı incelendiği çok sayıda çalışma bulunmaktadır (Huang ve ark., 1995, Zivanovic ve ark., 1995). Elektrik alan, kuantum kuyusu içerisindeki yük taşıyıcılarının dalga fonksiyonlarını ve enerji seviyelerini değiştirerek absorbsiyon spektrumunda kırmızıya kayma ve kırılma indisi değişimlerine neden olur. Benzer şekilde, manyetik alanın uygulanması kuantum kuyusu içerisindeki taşıyıcı dinamiklerini etkiler ve manyetooptik özelliklerde değişikliklere neden olur (Amini ve ark., 2017, Altun ve ark., 2022). Sayraç ve arkadaşları 2021 yılında, Öztürk ve arkadaşları da 2023 yılında elektrik alanın ve manyetik alanın kuantum kuyularındaki elektronik yapı üzerindeki etkilerini teorik olarak analiz etmiş ve bu etkilerin optik geçişlere olan yansımalarını tartışmıştır (Sayraç ve ark.,2021, Öztürk ve ark., 2023).

Bununla birlikte, artan bariyer genişliğine sahip AlGaAs/GaAs kuantum kuyularında elektrik ve manyetik alanların birleşik etkilerinin sistematik bir şekilde incelenmesi nispeten sınırlıdır. Bariyer genişliğinin artması, kuantum kuyusu içerisindeki yük taşıyıcılarının sınırlama enerjisini ve dalga fonksiyonlarının uzamsal dağılımını değiştirerek optik özellikler üzerinde karmaşık etkiler yaratır. Öztürk ve ark. bariyer genişliğinin kuantum kuyusu sistemlerinde enerji seviyelerine olan etkisini incelemiş ve bu parametrenin optik geçişlerdeki önemini vurgulamıştır (Öztürk ve ark, 2023). Elektrik ve manyetik alanların birleşik etkileri, özellikle bariyer genişliği gibi yapısal parametrelerle birlikte ele alındığında, kuantum kuyusu sistemlerinin tasarımında yeni fırsatlar sunabilir.

Bu çalışmada, AlGaAs/GaAs kuantum kuyularında artan bariyer genişliklerinin, elektrik ve manyetik alanların birleşik etkileri altında absorbsiyon katsayısı ve kırılma indisi değişimlerine olan etkisini teorik olarak incelemeyi amaçlıyorum. Çalışmam, bu parametrelerin optik özellikler üzerindeki etkilerini sistematik bir şekilde analiz ederek, optimize edilmiş optoelektronik cihaz tasarımları için teorik bir temel sunmayı hedeflemektedir. Çalışmada, etkin kütle yaklaşımı altında sonlu elemanlar yöntemi kullanılarak gerçekleştirilen hesaplamalarla, bu sistemlerin optik özellikleri detaylı bir şekilde analiz edilmiştir. Elektrik ve manyetik alanların kuantum kuyularındaki enerji seviyeleri ve dalga fonksiyonları üzerindeki etkileri, optik parametrelerin modülasyonu ve bu değişikliklerin optoelektronik uygulamalardaki potansiyeli üzerinde durulacaktır.

TEORİ

Dalga fonksiyonlarını ve bunlara karşılık gelen enerji seviyerini bulmak için etkin kütle yaklaşımı altında sonlu elemanlar yöntemi kullanılmıştır. Periyodik olarak artan bariyer genişliğine sahip $GaAs/Al_xGa_{1-x}As$ süperörgüsünün matris formu için, zamandan bağımsız Hamiltonyeni Denklem (1)'de verilmiştir.

$$H = -\frac{\hbar^2}{2\,m^*}\frac{d^2}{dz^2} + \frac{e^2B^2z^2}{2m^*c^2} + eFz + V(z) \qquad \textbf{(1)}$$

FULL TEXTS BOOK

İkinci dereceden türev ise Denklem (2) ile tanımlanmıştır.

$$\frac{d^2}{dz^2} = \left[-2\,diag\big(ones(1,N_z)\big) + diag(ones(1,N_z-1),-1) + diag(ones(1,N_z-1),1)\right]^2 \quad \textbf{(2)}$$

Bu denklemlerde **m*** elektronun etkin kütlesini ifade eder ve GaAs ile $Al_xGa_{1-x}As$ arasındaki örgü uyumundan dolayı **0,067m₀** alınır. **m₀** ise serbest elektronun kütlesidir. **e** electron yükü, **N₍z₎** toplam kuantum bölgesini tanımlamak için kullanılan matrisin uzunluğu, **F** büyüme yönündeki elektrik alanı, **B** büyüme yönüne dik uygulanan manyetik alanı ve **V(z)** potansiyeli temsil eder. Çalıştığım çoklu quantum kuyu yapısı 7 adet 3nm sabit genişlikli kuyudan oluşmaktadır. Bariyer genişliği ise 1nm genişlikten başlayıp sağa doğru 0,1nm lik artışlarla devam etmektedir. V_o bant süreksizliğini gösterir ve x=0,3 için 228V'dur.

Denklem (1)'i çözmek için bir boyutta diyagonalizasyon yöntemi kullanılmıştır. Enerji seviyeleri (EL'ler) ve bunlara karşılık gelen elektron dalga fonksiyonları (WF'ler) elde edildikten sonra, ilk dört enerji seviyesi (EL) için alt bantlar arası (ISB) geçişlere ait doğrusal absorpsiyon katsayıları (LAC), üçüncü mertebe doğrusal olmayan absorpsiyon katsayıları (NAC), ve toplam absorbsiyon katsayıları (TAC), yoğunluk matrisi yaklaşımı kullanılarak kolaylıkla elde edilebilir (Altuntaş ve ark., 2021).

$$\square_{if}^{(1)}(\omega) = \omega\sqrt{\frac{\mu}{\varepsilon_r}}\left|M_{if}\right|^2 \frac{e^2\sigma_v\hbar\Gamma}{\left(E_{if}-\hbar\omega\right)^2+(\hbar\omega)^2} \quad \textbf{(3)}$$

$$\square_{if}^{(3)}(\omega,I) = -2\omega\sqrt{\frac{\mu}{\varepsilon_r}}\left|M_{if}\right|^4\left(\frac{I}{\varepsilon_0 c n_r}\right)\frac{e^4\sigma_v\hbar\Gamma}{\left\{\left(E_{if}-\hbar\omega\right)^2+(\hbar\Gamma)^2\right\}^2} x \left[1 - \frac{\left|\delta_{if}\right|^2}{\left|2M_{if}\right|^2}\left(\frac{\left(E_{if}-\hbar\omega\right)^2-(\hbar\Gamma)^2+2E_{if}\left(E_{if}-\hbar\omega\right)}{\left(E_{if}\right)^2+(\hbar\Gamma)^2}\right)\right] \quad \textbf{(4)}$$

$$\square_{if}(\omega,I) = \square_{if}^{(1)}(\omega) + \square_{if}^{(3)}(\omega,I) \quad \textbf{(5)}$$

Doğrusal kırılma indisi değişimi (LRIC), doğrusal olmayan kırılma indisi değişimi (NRIC) ve toplam kırılma indisi değişimi (TRIC) ise sırası ile Denklem (6-7-8)'den hesaplanır (Zhang ve ark., 2018).

$$\frac{\Delta n_{if}^{(1)}(\omega)}{n_r} = \frac{e^2\sigma_v}{2\varepsilon_0 n_r^2}\left|M_{if}\right|^2\frac{\left(E_{if}-\hbar\omega\right)}{\left(E_{if}-\hbar\omega\right)^2+(\hbar\Gamma)^2} \quad \textbf{(6)}$$

$$\frac{\Delta n_{if}^{(3)}(\omega,I)}{n_r} = \frac{\mu\, c}{4\varepsilon_0 n_r^3}\left|M_{if}\right|^2\frac{e^4\sigma_v I}{\left\{\left(E_{if}-\hbar\omega\right)^2+(\hbar\,\Gamma)^2\right\}^2} x \left[4\left(E_{if}-\hbar\omega\right)\left|M_{if}\right|^2 - \frac{\delta_{if}^2}{\left(E_{if}\right)^2+(\hbar\,\Gamma)^2} x\left\{\left(E_{if}-\hbar\omega\right)\left[E_{if}\left(E_{if}-\hbar\omega\right)-(\hbar\,\Gamma)^2\right]-(\hbar\,\Gamma)^2\left[2E_{if}-\hbar\omega\right]\right\}\right] \quad \textbf{(7)}$$

$$\frac{\Delta n_{if}(\omega,I)}{n_r} = \frac{\Delta n_{if}^{(1)}(\omega)}{n_r} + \frac{\Delta n_{if}^{(3)}(\omega,I)}{n_r} \quad \textbf{(8)}$$

FULL TEXTS BOOK


M_{if} (DMMEs) dipol moment matris elemanları ise aşağıdaki gibi tanımlanmıştır.

$$M_{if} = \int \psi_f^* z \psi_i dz \quad , \quad (i,f = 1,2,3,4) \tag{9}$$

Tüm bu denklemlerde $\delta_{if} = M_{ff} - M_{ii}$ ifadesi bant içi (intersubband) dipol matris elemanlarını temsil eder. **I** optik ışık şiddeti, **ω** gelen fotonun açısal frekansı, $\boldsymbol{E_f}$ve $\boldsymbol{E_i}$ son ve ilk durumlara karşılık gelen kuantize edilmiş enerji seviyelerini, ε_0 boşluk dielektrik geçirgenliği, ε_r ise malzemenin geçirgenliğinin gerçek kısmı, $\boldsymbol{\mu}$ manyetik geçirgenlik, $\boldsymbol{n_r}$ kırılma indisi, $\boldsymbol{\sigma_v}$ ise taşıyıcı yoğunluğudur.

SONUÇ ve TARTIŞMA


Bu çalışmanın ilk aşamasında, artan bariyer genişliğine sahip 7 adet kuantum kuyusundan oluşan yapıda F ve B etkileri olmadan ve F=30 kV/cm F=-30kV/cm ve B=15T değerlerindeki uygulamalar dahilinde potansiyel profilleri, enerji seviyeleri ve elektron bulunma olasılıkları incelenmiştir. (Şekil 1.a-f)




Şekil 1: Potansiyel profilinin ve dalga fonksiyonlarının değişimi; **a)** F=0 B=0 için, **b)** F=30kV/cm B=0 için, **c)** F=-30kV/cm B=0 için, **d)** F=0 B=15T için, **e)** F=30kV/cm B=15T için, **f)** F=-30kV/cm B=15T için.

Şekil 1.a'da F ve B etkileri olmadığı durumda ilk enerji seviyesi (E_1) soldan sağa 2-3-4. kuyularda lokalize olmuştur. E_2, E_3 ve E_4 ise tüm yapı boyunca yayılmışlardır. Pozitif yönlü F uygulaması yapıyı sola doğru eğerken Şekil 1.b, negatif yönlü F'in etkisi ile yapı sağa doğru eğilmiştir Şekil 1.c. Buna bağlı olarak da enerji seviyelerindeki lokalizasyonlarda sola ve sağa doğru kaymalar olmuştur. Yapıya sadece manyetik alan uygulandığında Şekil 1.d potansiyelde bükülme olmuştur ve E_1, E_2 ve E_3 enerji seviyeleri kuyunun merkezine doğru lokalize olurken E_4 enerji seviyesinde çok değişim olmamıştır. Şekil 1e ve f de ise B ile beraber 30kV/cm ve -30kV/cm F uygulanmış ve potansiyelde bükülme ile beraber sola ve sağa eğilmeler de olmuştur. Bunlara bağlı olarak da enerji seviyelerinin lokalizasyonları sola ve sağa kaymıştır. Enerji seviyelerindeki bu değişimler yapının optik özellikleri için önemlidir.

FULL TEXTS BOOK



Şekil 2: B=0 ve B=15T değerleri için elektrik alanın bir fonksiyonu olarak; **a)** Geçiş enerjileri, **b)** DMME's, **c)** Bant içi dipol matris elemanları.




Şekil 3: F=0, F=30kV/cm ve F=-30kV/cm değerleri için manyetik alanın bir fonksiyonu olarak; **a)** Geçiş enerjileri, **b)** DMME's, **c)** Bant içi dipol matris elemanları.

Şekil 2'de B=0 ve B=15T uygulamaları için elektrik alan değişimi ile E_{if}, $|M_{if}|$, $|\delta_{if}|$ değişimleri incelenmiştir. Şekil 2.a'da B uygulandığı durumda tüm enerji farklarında çok ciddi bir değişim söz konusu değildir. Yaklaşık -5kV/cm'lik elektrik alan değerinden sonra enerji farkı az da olsa artmaya başlamıştır. B uygulanmadığı durumda E_{23} yaklaşık 10kV/cm'den sonra artış gösterirken E_{34} değişimi neredeyse sabittir. Fakat E_{12}'de -8kV/cm ye kadar enerji farkı azalırken -8kV/cm'den sonra tekrar artmaya başlamıştır. Benzer davranış Şekil 1.b'de de söz konusudur. DMME's değişimlerinde en dikkat çekici değişim yine B uygulanmadığı durumda $|M_{12}|$'de görülmüştür. $|M_{12}|$, -8 kV/cm'ye kadar artarken bu değerden sonra düşmüştür. $|M_{23}|$'de de yine bu değerde hafif artış ve sonrasında bir azalma vardır. Manyetik alan uygulandığında ise değişimler neredeyse sabittir. Şekil 1.c'de B uygulanmadığı durumda $|\delta_{12}|$'de -8kV/cm'de dikkat çekici bir değişim göstermiştir. $|\delta_{23}|$ -20kV/cm ile 5kV/cm arasında farklı bir davranış gösterirken, $|\delta_{34}|$ F=0 ve 20kV/cm'de dikkat çekici değişim göstermiştir.

Şekil 3'de F=0, F=30kV/cm ve F=-30kV/cm uygulamaları için manyetik alan değişimi ile E_{if}, $|M_{if}|$, $|\delta_{if}|$ değişimleri incelenmiştir. Şekil 3.a'da; F uygulanmadığı durumda enerji farklarının tümü yaklaşık 5T'dan sonra artmaya başlamıştır. F=30kV/cm uygulandığı durumda; 4T manyetik alan değerinden sonra E_{12} azalmaya E_{23} ve E_{34} ise artmaya başlamıştır. 6T'da E_{12} ve E_{34}, 9T'da ise E_{12} ve E_{23} eşit enerji farklarına sahiptir. F=-30kV/cm uygulandığı durumda ise tüm enerji farkları neredeyse 8T'dan sonra küçük artışlar göstermiştir. Şekil 3.b'de; F uygulanmadığı durumda $|M_{12}|$ 8T'dan sonra azalmıştır. $|M_{23}|$ ve $|M_{34}|$ deki değişimler ise yok denecek kadar azdır. F=30kV/cm uygulandığında $|M_{12}|$ ve $|M_{23}|$ 5T'dan sonra çok az artarken, $|M_{34}|$ sabit kalmıştır. F=-30kV/cm uygulandığı durumda ise; $|M_{12}|$'de çok az bir değişim varken, $|M_{23}|$ ve $|M_{34}|$ neredeyse sabittir. Şekil 3.c'de F uygulamalarının tümünde $|\delta_{12}|$, manyetik alan değeri arttıkça ciddi şekilde azalmıştır. $|\delta_{23}|$ ve $|\delta_{34}|$ ise F ve B uygulamaları ile küçük değişimler göstermiştir. E_{if}, $|M_{if}|$, $|\delta_{if}|$'lerde gösterilen bu değişimler çoklu kuantum yapısında elektrik ve manyetik alan uygulamaları ile optik özelliklerinde değişeceğini bize göstermektedir.


FULL TEXTS BOOK

Optik soğurma katsayısı ve kırılma indisindeki değişimler de F ve B uygulamalarından etkilenmektedir.




Şekil 4: Optik emilim katsayılarının F=0 B=0, F=30kV/cm B=0, F=30kV/cm B=15T uygulamaları altında, foton enerjisine bağlı değişimleri: **a-b-c:** (1-2) geçişleri, **d-e-f:** (2-3) geçişleri, **g-h-ı:** (3-4) geçişleri.

Şekil 4'de F=0, F=30kV/cm, B=0 ve B=15T değerlerindeki F ve B uygulamaları altında, (1-2),(2-3) ve (3-4) geçişlerinin LAC, NAC ve TAC'lerin foton enerjisiyle değişimleri verilmiştir. F= -30kV/cm değerindeki etkileri de incelenmiş fakat çalışmaya eklenmemiştir. LAC ve NAC zirveleri F ve B uygulamaları ile değişim göstermiştir. Optik spektrumda ise (1-2) ve (2-3) geçileri maviye kayma gözlemlenirken, (3-4) geçişinde kayma neredeyse olmamıştır. TAC, mevcut optik ışık şiddetinde negatif NAC katkısı ile azalır. TAC'ların rezonans pikleri F=0 ve B=0 uygulamalarında (1-2) (2-3) ve (3-4) geçişlerinin tümünde dallanma göstermiştir. F=30kV/cm ve B=15T değerlerine artırıldığında ise (1-2) geçişlerinde dallanma olmazken diğer geçişlerin hepsinde dallanma görülmüştür.



FULL TEXTS BOOK



Şekil 5: Kırılma indisi değişiminin F=0 B=0, F=30kV/cm B=0, F=30kV/cm B=15T uygulamaları altında, foton enerjisine bağlı değişimleri: **a-b-c:** (1-2) geçişleri, **d-e-f:** (2-3) geçişleri, **g-h-ı:** (3-4) geçişleri.

F ve B alanlarının varlığı ve yokluğunda, (1–2), (2–3) ve (3–4) geçişleri için RIC'lerin foton enerjisine bağlı değişimi sırasıyla Şekil 5'de gösterilmiştir. Yine F'nin negatif yönlü uygulamasına ait çalışmalar eklenmemiştir. LRIC ve NRIC terimleri zıt işaretli olduğundan, NRIC teriminin büyüklüğündeki herhangi bir artış bu iki terim arasındaki farkı da artıracak, dolayısıyla TRIC'i azaltacaktır. Tüm RIC'ler, tıpkı optik emilim katsayılarında olduğu gibi, F ve B alanlarının tüm şiddetlerinden etkilenmektedir. RIC'lerin hem F hem de B alanlarıyla ve potansiyel şekliyle ilişkili olduğu söylenebilir. Aynı F ve B şiddetleri altında (3–4) geçişi için tüm RIC değerleri (2–3) geçişlerinden daha büyüktür. (2-3) geçişlerindeki tüm RIC değerleri de (1-2) geçişindeki RIC değerlerinden daha büyüktür. Bu farkların temel nedeni, optik geçişin gerçekleştiği elektronik durumlar arasındaki enerji aralığı ve DMME'lerdeki (dipol matris elemanları) değişikliklerdir.

SONUÇ

Bu çalışmada, AlGaAs/GaAs çoklu kuantum kuyularında artan bariyer genişliklerinin, elektrik ve manyetik alanların birleşik etkileri altında optik absorbsiyon katsayısı ve kırılma indisi değişimleri teorik olarak incelenmiştir. Etkin kütle yaklaşımı ve sonlu elemanlar yöntemi kullanılarak hesaplanan enerji seviyeleri ve dalga fonksiyonları, elektrik (F=0, ±30 kV/cm) ve manyetik alan (B=0, 15 T) uygulamalarının potansiyel profilleri ve elektron lokalizasyonları üzerinde belirgin etkiler yarattığını ortaya koymuştur. Elektrik alanın yönü ve şiddeti, potansiyel profilini eğerek enerji seviyelerinde kaymalara neden olmuş; manyetik alan ise kuantum kuyusu merkezinde lokalizasyonu artırarak optik geçiş enerjilerini modüle etmiştir. Doğrusal ve doğrusal olmayan absorbsiyon katsayıları ile kırılma indisi değişimleri, yoğunluk matrisi yaklaşımıyla hesaplanmış ve (1-2), (2-3) ve (3-4) geçişlerinin foton enerjisine bağlı davranışları detaylı bir şekilde analiz edilmiştir. Bulgular, (1-2) ve (2-3) geçişlerinde elektrik ve manyetik alan uygulamalarıyla maviye kayma olduğunu, (3-4) geçişlerinde ise minimal kayma gözlendiğini göstermiştir. Ayrıca, dipol matris elemanları ve bant içi geçişlerdeki değişimler, optik özelliklerin bariyer genişliği ve alan uygulamalarına güçlü bir bağımlılık sergilediğini doğrulamıştır. Bu sonuçlar, AlGaAs/GaAs kuantum kuyularının optoelektronik cihaz tasarımlarında, özellikle optik modülatörler, kızılötesi dedektörler ve yüksek hızlı iletişim sistemleri gibi uygulamalarda optimizasyon potansiyelini vurgulamaktadır. Gelecekteki çalışmalar, farklı bariyer genişlikleri ve alan

FULL TEXTS BOOK

kombinasyonlarının deneysel doğrulamasına odaklanarak, bu sistemlerin pratik uygulamalara entegrasyonunu hızlandırabilir.

KAYNAKLAR

Aishah, A. N., Dakhlaoui, H., Ghrib, T., & Wong, B. M. (2022). Effects of magnetic, electric, and intense laser fields on the optical properties of AlGaAs/GaAs quantum wells for terahertz photodetectors. *Physica B: Condensed Matter*, *635*, 413838.

Alaydin, B. O. (2021). Effect of high bandgap AlAs quantum barrier on electronic and optical properties of In 0. 7 0 Ga 0. 3 0 As/Al 0. 6 0 In 0. 4 0 As superlattice under applied electric field for laser and detector applications. *International Journal of Modern Physics B*, *35*(02), 2150027.

Altun, D., Ozturk, O., Alaydin, B. O., & Ozturk, E. (2022). Linear and nonlinear optical properties of a superlattice with periodically increased well width under electric and magnetic fields. *Micro and Nanostructures*, *166*, 207225.

Altuntas, I., Dakhlaoui, H., Mora-Ramos, M. E., & Ungan, F. A. T. İ. H. (2021). Combined effects of electric, magnetic, and intense terahertz laser fields on the nonlinear optical properties in GaAs/GaAlAs quantum well with exponentially confinement potential. *The European Physical Journal Plus*, *136*(11), 1174.

Amini, M., Soleimani, M., & Ehsani, M. H. (2017). Electronic and optical properties of GaAs/AlGaAs Fibonacci ordered multiple quantum well systems. *Superlattices and Microstructures*, *112*, 680-687.

Bastard, G., Wave Mechanics Applied to Semiconductor Heterostructures, Les Editions de Physique, Les Ulis, 1988.

Huang, Y., Wang, J., & Lien, C. (1995). Electric-field dependence of optical absorption properties in coupled quantum wells and their application to 1.3 μm optical modulator. *Journal of applied physics*, *77*(1), 11-16.

Ozturk, E., Altun, D., Ozturk, O., & Alaydin, B. O. (2023). High harmonic generations in GaAs/AlGaAs superlattice: Effect of electric and magnetic field. *Solid State Communications*, *372*, 115300.

Sayraç, M., Turkoglu, A., Mora-Ramos, M. E., & Ungan, F. A. T. İ. H. (2021). Intensity-dependent nonlinear optical properties in an asymmetric Gaussian potential quantum well-modulated by external fields. *Optical and Quantum Electronics*, *53*(9), 485.

Zhang, Z. H., Yuan, J. H., & Guo, K. X. (2018). The combined influence of hydrostatic pressure and temperature on nonlinear optical properties of GaAs/Ga0. 7Al0. 3As morse quantum well in the presence of an applied magnetic field. *Materials*, *11*(5), 668.

Živanović, S., Milanović, V., & Ikonić, Z. (1995). Intraband absorption in semiconductor quantum wells in the presence of a perpendicular magnetic field. *Physical review B*, *52*(11), 8305.

FULL TEXTS BOOK

ARTIFICIAL INTELLIGENCE IN ACTION: A COMPARATIVE STUDY OF ML ALGORI THMS FOR TEXT CLASSIFICATION

Ms. Hetal Pravinsinh Jadeja [1]
[1]*Assistant Professor in Computer Science Department,*
At Harivandana College, Rajkot, India

Jatinkumar Bhimjibhai Kotadiya[2]
[2]*Research scholar in Computer Science Department,*
Atmiya University, Rajkot, India

ABSTRACT
Text classification, a crucial task in Natural Language Processing (NLP), is essential for structuring, managing, and understanding extensive amounts of unstructured text data. This research paper presents a comparative study of various Artificial Intelligence (AI)-driven Machine Learning (ML) algorithms applied to text classification tasks. The research assesses the effectiveness of models like Naive Bayes, Support Vector Machines (SVM), Decision Trees, Random Forest, and Gradient Boosting on typical text datasets. Text preprocessing techniques including tokenization, stop-word removal, stemming, and TF-IDF vectorization were implemented to prepare the data. Each algorithm was trained and validated using stratified k-fold cross-validation, and assessed based on metrics such as accuracy, precision, recall, F1-score, and computational efficiency. Experimental results demonstrate that ensemble-based models, particularly Random Forest and Gradient Boosting, deliver superior classification accuracy and stability compared to traditional classifiers. This research emphasizes the real-world efficiency of AI in streamlining and enhancing text classification tasks, providing significant insights for scholars and professionals in areas like sentiment analysis, spam identification, and document organization.

KEYWORDS: Artificial Intelligence, Machine Learning, Text Classification, Natural Language Processing, Naive Bayes, Support Vector Machine, Random Forest, Sentiment Analysis, Supervised Learning, Algorithm Comparison

I.INTRODUCTION

The swift increase of unstructured text data has amplified the necessity for precise and automated text classification across multiple areas, such as sentiment analysis, spam detection, and document categorization. This research investigates the performance of several Machine Learning (ML) algorithms, including Naive Bayes, Support Vector Machines, Decision Trees, Random Forest, and Gradient Boosting, in a cohesive experimental setup. Employing standard text preprocessing techniques and stratified k-fold cross-validation, the research systematically evaluates each model based on performance metrics including accuracy, precision, recall, F1-score, and computational efficiency. The findings underscore the superior performance of ensemble-based models, particularly Random Forest and Gradient Boosting, in terms of classification accuracy and stability. This work aims to guide researchers and practitioners in selecting appropriate AI-driven methods for diverse text classification tasks.

II.RELATED WORK

Recent scholarly discourse has explored the integration of artificial intelligence within creative and interpretive domains, highlighting its dual role as both a technical tool and an autonomous aesthetic

FULL TEXTS BOOK

agent. Studies have examined how AI-driven algorithms, particularly in visualization and text classification, contribute to meaning-making by reshaping narrative structures and authorial intent. Within a posthumanist framework, this shift challenges traditional anthropocentric paradigms by positioning AI as a collaborative force in the production of knowledge and art. This literature provides a foundational context for examining the comparative effectiveness of machine learning algorithms in AI-assisted text classification tasks [1].

Recent advancements in artificial intelligence have significantly enhanced the accuracy and applicability of machine learning algorithms in text classification tasks, including clinical and diagnostic decision-making. Studies have demonstrated the potential of models like ChatGPT to interpret structured and unstructured clinical data, offering recommendations aligned with expert evaluations in specific contexts. However, literature also highlights variability in performance, especially in complex scenarios, underscoring the need for human oversight. The integration of AI tools in healthcare remains a developing field, where reliability and contextual understanding are essential for broader adoption [2].

Recent advancements in artificial intelligence and distributed computing have significantly enhanced the efficiency of data-intensive tasks, particularly in image and text classification. Prior research has explored the use of cloud-based platforms and parallel processing frameworks like PyTorch's Distributed Data Parallel and Tensor Flow for deep learning applications. Studies highlight that even legacy hardware can be effectively utilized for distributed learning, offering cost-efficient and sustainable solutions. These findings provide a foundational understanding for examining the comparative performance of machine learning algorithms in resource-constrained environments [3].

Recent developments in artificial intelligence have resulted in machine learning algorithms that can execute intricate tasks like image analysis, pattern recognition, and predictive modeling with great precision. In the domain of healthcare, particularly orthodontics, these algorithms have demonstrated superior performance compared to traditional diagnostic tools. Prior studies have emphasized the role of AI in enhancing diagnostic precision, streamlining treatment planning, and supporting clinical decision-making. The literature further underscores the need for ethical integration and technical training to ensure effective and responsible use of AI technologies in medical practice [4].

Recent studies underscore the growing relevance of artificial intelligence, particularly deep learning and natural language processing, in enhancing the integration and dissemination of pharmaceutical knowledge. Researchers have explored the efficacy of models such as CNNs, RNNs, and BiLSTM-CRF in tasks ranging from entity recognition to semantic analysis and personalized content delivery. Comparative findings suggest that AI-driven approaches outperform traditional methods in terms of accuracy, user engagement, and knowledge customization. However, literature also highlights challenges related to data sparsity, ethical considerations, and the need for cross-modal information fusion [5].

Previous studies in the area of text classification and sentiment analysis have thoroughly investigated the application of conventional machine learning algorithms like Logistic Regression, Naïve Bayes, Support Vector Machines, and Random Forest. These models have demonstrated high performance based on standard evaluation metrics; however, they often lack interpretability and transparency in decision-making. Recent progress in Explainable Artificial Intelligence (XAI), especially tools such as LIME, has filled this gap by providing model-agnostic explanations that improve trust, accountability, and model performance. The integration of XAI has emerged as a vital approach for validating and refining machine learning models beyond conventional accuracy-based evaluation [6].

Recent advancements in generative artificial intelligence (GAI) have significantly influenced the field of education, particularly in automating content creation and personalizing assessments. Prior studies

FULL TEXTS BOOK

have explored the integration of AI-generated examinations, adaptive learning systems, and their effects on student performance. Research indicates that while AI can enhance instructional efficiency and learner engagement, its effectiveness is maximized when guided by pedagogical expertise. The literature emphasizes the need for a balanced approach that combines AI-driven innovations with active educator involvement to ensure alignment with educational goals and learning outcomes [7].

Previous studies have extensively explored the application of machine learning algorithms in medical diagnosis, particularly for heart disease prediction, emphasizing the effectiveness of classification models such as KNN, SVM, Random Forest, and Logistic Regression. Researchers have highlighted the significance of performance metrics like accuracy, precision, sensitivity, and F1-score in evaluating model efficiency. Numerous works underscore the impact of dataset characteristics—such as class balance and preprocessing—on algorithm performance. These findings form a foundational basis for the present study, which seeks to comparatively evaluate various algorithms within the context of text classification using artificial intelligence [8].

Many research efforts have investigated the use of machine learning (ML) and deep learning (DL) algorithms to forecast customer satisfaction, highlighting their ability to improve business strategies through precise classification and prediction. Prior research consistently highlights the superior performance of ensemble models such as Random Forest and Gradient Boosting, as well as hybrid deep learning architectures like CNN-LSTM. Moreover, the literature underscores the challenges of data availability and the evolving, emotion-driven nature of customer satisfaction, which complicates its measurement and prediction [9].

Recent studies in Natural Language Processing (NLP) emphasize the importance of selecting suitable machine learning algorithms for effective text classification tasks. Existing literature has explored models such as Naive Bayes, Random Forest, and Support Vector Machines, each demonstrating varied performance across metrics like accuracy, precision, and computational efficiency. Comparative evaluations highlight the strengths and limitations of these models in handling different text classification scenarios, particularly within domains such as sentiment analysis and topic categorization. These findings establish a foundation for further empirical investigations into algorithmic performance using standardized preprocessing and feature engineering techniques [10].

III.RESEARCH METHODOLOGY

This section outlines the systematic approach adopted to conduct a comparative analysis of various Machine Learning (ML) algorithms for text classification tasks within the broader framework of Artificial Intelligence (AI). The methodology encompasses data selection, preprocessing procedures, model development, evaluation criteria, and validation techniques to ensure the robustness and reliability of the findings.

FULL TEXTS BOOK




Fig1. Model Architecture for Comparative Study of ML Algorithms in Text Classification

3.1 Research Design

The research employs an experimental and quantitative design wherein multiple AI-driven ML algorithms are applied to standardized text classification datasets. The study follows a comparative paradigm, aiming to evaluate the classification performance of each algorithm based on empirical

FULL TEXTS BOOK

metrics. The design facilitates objective analysis by maintaining consistent preprocessing pipelines, hyper parameter tuning strategies, and validation mechanisms across all models under investigation.

3.2 Dataset Selection

To ensure generalizability and replicability, publicly available benchmark datasets were utilized for the classification tasks. These datasets were selected based on their diversity in textual content, class distribution, and applicability in real-world domains such as spam filtering, sentiment analysis, and topic categorization. Each dataset comprises labeled textual instances suitable for supervised learning models.

3.3 Data Preprocessing

Prior to model implementation, all textual data underwent comprehensive preprocessing to enhance quality and reduce noise. The following steps were employed:

- **Tokenization:** Conversion of text into individual tokens (words or phrases) using standard NLP libraries.
- **Stop-word Removal:** Elimination of common, non-informative words (e.g., "the," "is," "and") to minimize dimensionality.
- **Stemming:** Reduction of words to their root form using Porter's stemming algorithm to unify similar terms.
- **TF-IDF Vectorization:** Transformation of textual data into numerical feature vectors using Term Frequency-Inverse Document Frequency, emphasizing discriminative terms across documents.

3.4 Machine Learning Algorithms

The comparative study focuses on five widely adopted ML algorithms, chosen for their diverse learning paradigms and applicability in NLP:

- **Naive Bayes (NB):** A probabilistic classifier based on Bayes' Theorem with strong independence assumptions.
- **Support Vector Machine (SVM):** A discriminative model that constructs optimal hyperplanes for binary and multi-class classification.
- **Decision Tree (DT):** A tree-structured classifier using recursive partitioning based on feature values.
- **Random Forest (RF):** An ensemble model aggregating multiple decision trees through bagging to enhance accuracy and control over fitting.
- **Gradient Boosting (GB):** A boosting-based ensemble approach that builds additive models sequentially to minimize classification error.

Each algorithm was implemented using Python-based ML libraries such as Scikit-learn, ensuring reproducibility and compatibility.

3.5 Model Training and Validation

To validate model performance objectively, **Stratified K-Fold Cross-Validation** (with k=10) was employed. This technique ensures balanced class distribution across training and validation folds, mitigating sampling bias. During each iteration, models were trained on nine folds and evaluated on the remaining fold. This process was repeated ten times, and the average performance was reported.

FULL TEXTS BOOK

3.6 Evaluation Metrics

The classification performance of each model was assessed using the following standard evaluation metrics:

- **Accuracy:** Proportion of correctly predicted instances over the total number of samples.
- **Precision:** Ratio of true positive predictions to the total predicted positives.
- **Recall:** Ratio of true positive predictions to all actual positives.
- **F1-Score:** Harmonic average of precision and recall, equalizing both measurements
- **Computational Efficiency:** Assessment of training and prediction duration to evaluate scalability and deployment viability.

These metrics together offer an extensive perspective on the models' performance, especially in situations with imbalanced or noisy data.

3.7 Experimental Environment

All experiments were carried out in a high-performance computing setting featuring an Intel Core i7 processor, 32GB RAM, and Python 3.9. Software dependencies were managed via virtual environments to ensure consistency. The implementation was carried out using open-source libraries such as Scikit-learn, NLTK, and Pandas.

IV.FINDINGS

The findings of this study underscore the superior performance of ensemble-based machine learning models, particularly Random Forest and Gradient Boosting, in the context of text classification tasks. These algorithms consistently demonstrated higher accuracy, precision, recall, and F1-scores compared to traditional classifiers such as Naive Bayes, Support Vector Machines, and Decision Trees. Preprocessing techniques—namely tokenization, stop-word removal, stemming, and TF-IDF vectorization—were instrumental in enhancing model efficacy by refining textual representations. While simpler models offered computational efficiency, they were generally outperformed by ensemble methods in terms of classification reliability and robustness across varied datasets. These results highlight the practical advantages of ensemble approaches for applications demanding high accuracy and resilience in natural language processing.

Aspect	Naive Bayes (NB)	Support Vector Machine (SVM)	Decision Tree (DT)	Random Forest (RF)	Gradient Boosting (GB)
Model Type	Probabilistic	Linear Classifier	Tree-Based	Ensemble (Bagging)	Ensemble (Boosting)
Accuracy	Moderate	High (Dataset-dependent)	Moderate	**High**	**Very High**
Precision	Moderate	High	Moderate	**High**	**Very High**

FULL TEXTS BOOK

Recall	Moderate	Moderate to High	Moderate	**High**	**Very High**
F1-Score	Balanced but lower	Balanced	Moderate	**High**	**Very High**
Training Speed	**Very Fast**	Moderate	Fast	Slower	**Slowest**
Robustness Across Datasets	Low to Moderate	Moderate	Low	**High**	**Very High**
Scalability	Good for small datasets	Good for large sparse datasets	Less scalable	**Highly scalable**	**Scalable with tuning**
Impact of Preprocessing	Significant (especially with TF-IDF)	Beneficial	Moderate	Enhances performance	Maximizes model performance
Overall Effectiveness	Basic Benchmark	Strong with fine-tuning	Interpretable but less accurate	**Highly Reliable**	**Most Accurate and Resilient**

Table1. Performance Comparison of Machine Learning Models for Text

V.DISCUSSION

The discussion highlights the comparative performance of various machine learning algorithms applied to text classification, emphasizing the superior accuracy and robustness of ensemble methods such as Random Forest and Gradient Boosting. While these models demonstrated strong predictive capabilities, they require greater computational resources compared to traditional classifiers like Naive Bayes and Decision Trees, which, despite their simplicity and efficiency, showed limitations in handling complex textual data. Support Vector Machines offered a balanced performance but posed scalability concerns. Preprocessing techniques, including TF-IDF vectorization and stratified cross-validation, played a crucial role in enhancing model effectiveness. Overall, the study underscores the importance of selecting algorithms based on specific application requirements, advocating for a task-oriented approach to model deployment in AI-driven text classification.

VI.CONCLUSION

In summary, this study provides a detailed comparative evaluation of various machine learning algorithms applied to text classification, emphasizing the practical utility of artificial intelligence in processing unstructured textual data. The experimental results highlight that ensemble-based models, particularly Random Forest and Gradient Boosting, consistently outperform traditional classifiers in terms of accuracy and stability. While simpler models like Naive Bayes and SVM offer computational efficiency, they fall short in complex classification scenarios. The findings underscore the importance of selecting appropriate algorithms based on task-specific requirements and demonstrate the value of ensemble approaches in achieving robust and scalable text classification.

FULL TEXTS BOOK

VII.FUTURE WORK

Future research may extend the current study by incorporating deep learning architectures such as CNNs, RNNs, and transformer-based models like BERT to capture richer contextual and semantic features in text data. Further investigation using domain-specific and multilingual datasets would enhance the generalizability and practical relevance of classification models. Exploring hybrid approaches that integrate traditional machine learning with advanced embedding or rule-based methods could improve performance and robustness. Additionally, the application of explainable AI techniques can offer greater transparency in model decision-making, while addressing scalability and computational efficiency remains essential for real-world deployment. These directions collectively aim to refine and expand the capabilities of text classification systems in diverse and dynamic environments.

VIII.RECOMMENDATIONS

In light of the comparative analysis conducted in this study, it is recommended that future research and practical implementations in text classification prioritize ensemble-based models such as Random Forest and Gradient Boosting due to their superior accuracy and stability. Emphasis should be placed on comprehensive preprocessing techniques and the inclusion of diverse evaluation metrics to obtain a more holistic assessment of model performance. The exploration of hybrid and deep learning-based approaches is encouraged to further enhance classification capabilities. Additionally, utilizing domain-specific datasets, addressing scalability challenges, and ensuring ethical considerations and model interpretability are essential for the responsible and effective deployment of AI-driven text classification systems in real-world contexts.

IX.REFERENCES

[1] B. Meydan, “FROM LITERARY TEXT TO ALGORITHM: ARTIFICIAL INTELLIGENCE-GENERATED VISUALISATION EXPERIMENTS FROM THE NOVEL THE LATHE OF HEAVEN”.

[2] L. S. R. D. Oliveira *et al.*, “ARTIFICIAL INTELLIGENCE IN SCOLIOSIS DIAGNOSIS: A COMPARATIVE STUDY BETWEEN CHATGPT AND SURGEONS,” *Coluna/Columna*, vol. 24, no. 2, 2025, doi: 10.1590/s1808-185120252402293157.

[3] M. Rivera-Escobedo *et al.*, “Low-Scalability Distributed Systems for Artificial Intelligence: A Comparative Study of Distributed Deep Learning Frameworks for Image Classification,” *Appl. Sci.*, vol. 15, no. 11, p. 6251, Jun. 2025, doi: 10.3390/app15116251.

[4] M. Hack *et al.*, “Comparative study on the results of orthodontic diagnostics by using algorithms generated by Artificial Intelligence and simple algorithms,” *Med. Pharm. Rep.*, vol. 97, no. 2, pp. 215–221, Feb. 2024, doi: 10.15386/mpr-2702.

[5] L. Jin, “Research on Optimization Strategies of Artificial Intelligence Algorithms for the Integration and Dissemination of Pharmaceutical Science Popularization Knowledge,” *Sci. J. Technol.*, vol. 7, no. 1, pp. 45–55, Jan. 2025, doi: 10.54691/90a4en92.

[6] K. Zahoor, N. Zakaria Bawany, and T. Qamar, “Evaluating text classification with explainable artificial intelligence,” *IAES Int. J. Artif. Intell. IJ-AI*, vol. 13, no. 1, p. 278, Mar. 2024, doi: 10.11591/ijai.v13.i1.pp278-286.

[7] Z. Elmourabit, Asmaâ Retbi, and Nour-Eddine El Faddouli, “The Impact of Generative Artificial Intelligence on Education: A Comparative Study,” *Eur. Conf. E-Learn.*, vol. 23, no. 1, pp. 470–476, Oct. 2024, doi: 10.34190/ecel.23.1.2975.

FULL TEXTS BOOK

[8] B. Eliaçık and A. H. Isık, "Artificial Intelligence and Classification Algorithms in Heart Disease Data: Modern Approaches and Performance Comparison," *Türk Doğa Ve Fen Derg.*, vol. 14, no. 2, pp. 179–187, Jun. 2025, doi: 10.46810/tdfd.1622670.

[9] O. Berrada Chakour, A. Ettaoufik, K. Aissaoui, and A. Maizate, "Artificial intelligence algorithms to predict customer satisfaction: a comparative study," *IAES Int. J. Artif. Intell. IJ-AI*, vol. 14, no. 2, p. 1654, Apr. 2025, doi: 10.11591/ijai.v14.i2.pp1654-1662.

[10] Adala M. Chaid, "Comparative Analysis of Innovative Machine Learning Algorithms: Advancements in Natural Language Processing," *J. Inf. Syst. Eng. Manag.*, vol. 10, no. 14s, pp. 648–668, Feb. 2025, doi: 10.52783/jisem.v10i14s.2373.

[11] Li, H. (2018). Deep learning for natural language processing: advantages and challenges. National Science Review, 5(1), 24-26.

[12] Mikolov, T., Chen, K., Corrado, G., & Dean, J. (2013). Efficient Estimation of Word Representations in Vector Space. arXiv preprint arXiv:1301.3781.

[13] Mungoli, N. (2023). Advancements in Deep Learning: A Comprehensive Study of the Latest Trends and Techniques in Machine Learning. International Journal of Advanced Engineering Technologies and Innovations, 1(04), 43-64.

[14] Nagarhalli, T. P., Vaze, V., & Rana, N. K. (2021, February). Impact of machine learning in natural language processing: A review. In 2021 third international conference on intelligent communication technologies and virtual mobile networks (ICICV) (pp. 1529-1534). IEEE.

[15] Ofer, D., Brandes, N., &Linial, M. (2021). The language of proteins: NLP, machine learning & protein sequences. Computational and Structural Biotechnology Journal, 19, 1750-1758.

[16] Ofori-Boateng, R., Aceves-Martins, M., Wiratunga, N., & Moreno-Garcia, C. F. (2024). Towards the automation of systematic reviews using natural language processing, machine learning, and deep learning: a comprehensive review. Artificial intelligence review, 57(8), 200.

[17] A. Boukhalfa, A. Abdellaoui, N. Hmina, and H. Chaoui, "LSTM deep learning method for network intrusion detection system," International Journal of Electrical and Computer Engineering (IJECE), vol. 10, no. 3, pp. 3315-3322, Jun. 2020, doi: 10.11591/ijece.v10i3.pp3315-3322.

[18] M. A. Alshamari, "Evaluating user satisfaction using deep-learning-based sentiment analysis for social media data in Saudi Arabia's telecommunication sector," Computers, vol. 12, no. 9, Aug. 2023, doi: 10.3390/computers12090170.

[19] S. Beltozar-Clemente, O. Iparraguirre-Villanueva, F. Pucuhuayla-Revatta, J. Zapata-Paulini, and M. Cabanillas-Carbonell, "Predicting customer abandonment in recurrent neural networks using short-term memory," Journal of Open Innovation: Technology, Market, and Complexity, vol. 10, no. 1, Mar. 2024, doi: 10.1016/j.joitmc.2024.100237.

[20] R. Mutegeki and D. S. Han, "A CNN-LSTM approach to human activity recognition," in 2020 International Conference on Artificial Intelligence in Information and Communication, ICAIIC 2020, Feb. 2020, pp. 362–366, doi: 10.1109/ICAIIC48513.2020.9065078.

A COMPARATIVE STUDY OF MACHINE LEARNING ALGORITHMS FOR SENTIMENT ANALYSIS USING ARTIFICIAL INTELLIGENCE TECHNIQUES

***Jatinkumar Bhimjibhai Kotadiya*[1]**

[1]*Research scholar in Computer Science Department,*
Atmiya University, Rajkot, India

***Dr.Dharmendra Jayantilal Ambani*[2]**

[2]*Assistant Professor in Computer Science Department,*
Harivandana College, Rajkot, India

***Dr. Amit K. Patel*[3]**

[3]*Assistant Professor in Computer Science Department,*
Atmiya University, Rajkot, India

ABSTRACT

In the era of digital communication, sentiment analysis has emerged as a vital application of artificial intelligence (AI) for understanding public opinions, emotions, and attitudes across various online platforms. This study presents a comprehensive comparative analysis of several machine learning (ML) algorithms applied to sentiment analysis tasks. Leveraging AI techniques, the research evaluates the performance of traditional classifiers such as Naive Bayes, Support Vector Machines (SVM), Random Forest, and Logistic Regression, alongside ensemble methods including Gradient Boosting and Voting Classifiers. A benchmark dataset consisting of pre-labeled textual data was used to train and test each model. Preprocessing techniques such as tokenization, stop-word removal, and TF-IDF vectorization were applied to prepare the text for analysis. Each algorithm's performance was assessed using evaluation metrics including accuracy, precision, recall, and F1-score to determine the most efficient and reliable approach for sentiment classification. The results indicate that ensemble models, particularly Gradient Boosting, outperformed individual classifiers in terms of accuracy and robustness. This study highlights the significance of model selection in AI-driven sentiment analysis and provides insights for future developments in opinion mining and customer feedback systems. The findings contribute to the growing body of knowledge in intelligent text analytics and offer practical implications for AI applications in social media monitoring, business intelligence, and educational feedback systems.

KEYWORDS: Sentiment Analysis, Machine Learning Algorithms, Artificial Intelligence, Text Classification, Naïve Bayes, Support Vector Machine, Random Forest, Natural Language Processing, Predictive Modeling, Comparative Study

X.INTRODUCTION

In the context of increasing digital communication, sentiment analysis has become essential for interpreting public opinion across online platforms. This study investigates the effectiveness of various machine learning algorithms, including both traditional classifiers—such as Naive Bayes, SVM, Random Forest, and Logistic Regression—and ensemble methods like Gradient Boosting and Voting Classifiers. By applying standardized preprocessing techniques and evaluating performance using accuracy, precision, recall, and F1-score, the research aims to identify the most reliable models for sentiment classification. The findings underscore the superiority of ensemble approaches in achieving robust and accurate sentiment analysis, offering valuable implications for real-world applications in social media monitoring, business analytics, and educational feedback systems.

FULL TEXTS BOOK

XI.RELATED WORK

Recent advancements in machine learning have significantly enhanced stock market forecasting by leveraging algorithms such as LSTM, random forest, and support vector machines. Studies emphasize the effectiveness of hybrid and ensemble models in capturing complex market dynamics and improving prediction accuracy. Integrating technical indicators, historical prices, and sentiment analysis has proven beneficial for trend identification and volatility reduction. The literature highlights a growing focus on real-time data integration and interpretability to refine financial predictive modeling [1].

Recent advancements in sentiment analysis have demonstrated the growing effectiveness of machine learning and deep learning models in processing unstructured textual data. Studies integrating natural language processing with artificial intelligence have explored diverse architectures such as LSTM, BiLSTM, and attention-based models, emphasizing their accuracy in classifying sentiment across various domains. Furthermore, research has shown that combining linguistic insights with AI algorithms can enhance contextual understanding, leading to more reliable sentiment detection. These findings establish a foundation for developing refined models like the ALAIA-SAC technique presented in this study [2].

Recent studies have increasingly explored the integration of sentiment analysis with machine learning techniques to predict stock market trends, leveraging unstructured textual data such as financial news headlines. Prior research has demonstrated the utility of sentiment analyzers like VADER and TextBlob in extracting sentiment polarity, which, when combined with classification algorithms such as logistic regression, SVM, random forest, XGBoost, and neural networks, can yield meaningful predictions of market movements. Despite this progress, limited comparative evaluations exist focusing solely on sentiment-based features under uniform experimental conditions. This study builds on existing findings by systematically comparing these models using standardized sentiment inputs to assess their effectiveness in stock market direction prediction [3].

Recent literature highlights the transformative role of machine learning and artificial intelligence in addressing complex challenges across various domains, including smart grid stability, load management, and renewable energy integration. A range of algorithms—such as support vector machines, decision trees, neural networks, and reinforcement learning methods—have demonstrated effectiveness in enhancing system intelligence and decision-making. These findings provide a foundational understanding of algorithmic capabilities that can be adapted to sentiment analysis, emphasizing the comparative performance and contextual suitability of different models. This review thus informs the methodological choices for applying AI-driven ML techniques to optimize sentiment classification tasks [4].

Existing literature on sentiment analysis highlights its growing significance across various social media platforms due to the exponential rise in user-generated content. Various research efforts have investigated both conventional and contemporary machine learning techniques, such as logistic regression, Naïve Bayes, and neural networks, evaluating their efficacy in correctly categorizing sentiments. Recent research emphasizes the limitations of these models in handling domain-specific challenges and the need for improved methodologies. Comparative evaluations have laid the groundwork for advancing sentiment classification through robust and adaptive AI-driven techniques [5].

Prior research in sentiment analysis has explored a wide spectrum of machine learning algorithms, ranging from traditional classifiers like Naïve Bayes and Support Vector Machines to advanced deep learning architectures such as LSTM and BERT. Numerous studies have underscored the significance of feature extraction techniques and contextual understanding in enhancing classification accuracy.

FULL TEXTS BOOK

Comparative analyses consistently demonstrate the superior performance of deep learning models, particularly those leveraging word embeddings and transformer-based architectures. However, challenges such as data imbalance, sarcasm detection, and domain adaptability continue to influence model effectiveness across different applications [6].

Many research works have investigated the use of machine learning and deep learning methods for identifying epileptic seizures through physiological signals like EEG and ECG. Methods including SVM, ANN, CNN, and LSTM have demonstrated notable performance, with LSTM-based models achieving up to 96.5% accuracy in seizure classification. Additionally, emerging research highlights the potential of sentiment analysis and multimodal data integration as supplementary tools for enhancing detection accuracy. This review synthesizes recent advancements and provides a foundation for developing intelligent, automated systems in clinical and real-time environments [7].

Recent studies in sentiment analysis have explored both traditional machine learning techniques and advanced deep learning models to interpret emotions in text data. Traditional classifiers like Logistic Regression and Random Forest have demonstrated strong performance in domain-specific contexts, while deep learning architectures such as CNNs and RNNs are recognized for capturing linguistic complexities and sequential patterns. Additionally, lexicon-based tools like TextBlob and NLTK have been employed for baseline comparisons but often fall short in accuracy due to their generic nature. Prior research underscores the importance of dataset characteristics, domain adaptation, and model selection in achieving optimal sentiment classification outcomes [8].

Extensive research has been conducted on the application of machine learning and deep learning techniques for financial market prediction, with studies emphasizing the importance of model selection, data preprocessing, and feature extraction. Prior works have demonstrated the superior performance of deep learning models, such as LSTM and RNN, over traditional machine learning algorithms in capturing temporal dependencies in stock data. Comparative analyses across models reveal varied accuracy levels depending on data types, market sectors, and feature sets. These foundational studies inform the present research by identifying effective methodologies and highlighting existing gaps in model interpretability and real-time applicability [9].

Recent advancements in Artificial Intelligence and Machine Learning have significantly enhanced the effectiveness of sentiment analysis and data extraction. Studies have demonstrated the potential of algorithms such as Logistic Regression, Random Forest, and Decision Tree in producing high classification accuracy, particularly when integrated with clustering techniques like K-Means. Researchers have also explored hybrid approaches that combine multiple models to improve performance across diverse datasets. This literature underscores the growing relevance of ensemble and unsupervised learning strategies in addressing sentiment classification challenges [10].

XII.RESEARCH METHODOLOGY

This section outlines the systematic approach adopted to conduct the comparative study of machine learning algorithms for sentiment analysis utilizing artificial intelligence techniques. The methodology encompasses data collection, preprocessing, model selection, implementation, and performance evaluation. A systematic approach was employed to guarantee the dependability, accuracy, and replicability of the experimental findings.

FULL TEXTS BOOK




Fig1. Model Architecture for Sentiment Analysis using ML Algorithms

3.1 Dataset Selection

For this study, a publicly available benchmark dataset containing pre-labeled textual data was utilized. The dataset comprised user-generated content with sentiment labels categorized as positive, negative, or neutral. The selection of this dataset was based on its widespread use in sentiment analysis research, its diversity in content, and the availability of ground truth labels, which ensured a fair and consistent comparison of the selected algorithms.

3.2 Data Preprocessing

Textual data, by its nature, requires preprocessing to convert unstructured content into a structured form suitable for machine learning algorithms. The following preprocessing steps were applied:

- **Tokenization**: The text was segmented into individual tokens, such as words or phrases, to facilitate linguistic analysis.
- **Stop-word Removal**: Commonly occurring words that carry minimal semantic weight (e.g., "the", "is", "and") were removed to reduce noise and improve model focus on meaningful terms.

FULL TEXTS BOOK

- **Normalization**: Text was converted to lowercase and stripped of punctuation and special characters.
- **Vectorization**: The cleaned tokens were transformed into numerical representations using Term Frequency-Inverse Document Frequency (TF-IDF), enabling effective feature extraction from the textual data.

3.3 Model Selection and Implementation

Multiple machine learning algorithms were selected based on their historical effectiveness in sentiment classification tasks. The models considered in this study include:

- **Naive Bayes (NB)**: A probabilistic classifier based on Bayes' Theorem, known for its simplicity and efficiency in text classification.
- **Support Vector Machines (SVM)**: A supervised learning model that constructs hyperplanes in a high-dimensional space to separate sentiment classes effectively.
- **Logistic Regression (LR)**: A statistical model employed to predict the probability of class membership.
- **Random Forest (RF)**: An ensemble-based classifier that constructs multiple decision trees and merges their predictions to enhance accuracy and control overfitting.
- **Gradient Boosting (GB)**: A powerful ensemble method that sequentially builds models to correct the errors made by prior models, often resulting in superior performance.
- **Voting Classifier (VC)**: A meta-classifier that combines the predictions of multiple base learners through majority voting or averaging.

All models were implemented using Python-based machine learning libraries such as Scikit-learn, ensuring standardized training and testing procedures.

3.4 Model Training and Testing

The dataset was divided into training and testing parts with an 80:20 distribution. The training data was used to build the models, while the testing data was reserved for evaluating model generalization. Stratified sampling ensured a balanced distribution of sentiment classes across the splits. Hyperparameter tuning was performed using grid search and cross-validation techniques to optimize model performance. Default parameters were adjusted to enhance classification effectiveness without compromising computational efficiency.

3.5 Performance Evaluation Metrics

To compare the effectiveness of the machine learning models, the following evaluation metrics were employed:

- **Accuracy**: The proportion of correct predictions over the total number of predictions.
- **Precision**: The ratio of true positive predictions to the total predicted positives, indicating the model's ability to avoid false positives.
- **Recall (Sensitivity)**: The ratio of true positive predictions to all actual positives, reflecting the model's capability to identify relevant instances.
- **F1-Score**: The harmonic mean of precision and recall, offering a balanced evaluation metric especially useful in imbalanced datasets.

FULL TEXTS BOOK

Metric	Definition	Formula	Purpose
Accuracy	Proportion of total correct predictions to all predictions made	(TP + TN) / (TP + TN + FP + FN)	Measures overall correctness of the model
Precision	Proportion of correctly predicted positive observations to total predicted positives	TP / (TP + FP)	Indicates how well the model avoids false positives
Recall	Proportion of correctly predicted positives to all actual positives	TP / (TP + FN)	Evaluates the model's ability to find all relevant (positive) cases
F1-Score	Harmonic mean of precision and recall	2 × (Precision × Recall) / (Precision + Recall)	Provides a balanced metric when data is imbalanced

Table1. Model Performance Evaluation Metrics

Each model's performance was recorded and analyzed to determine the most effective algorithm for sentiment classification.

3.6 Comparative Analysis

A comparative analysis was conducted by evaluating and tabulating the performance metrics of all models. Emphasis was placed on identifying the model(s) demonstrating superior accuracy and robustness across diverse sentiment classes. The results were then interpreted to highlight the strengths and limitations of each algorithm.

XIII.FINDINGS

The findings of this study indicate that ensemble learning algorithms, particularly Gradient Boosting, outperform traditional machine learning models such as Naive Bayes, Support Vector Machines, Logistic Regression, and Random Forest in sentiment analysis tasks. While SVM and Logistic Regression demonstrated strong performance among individual classifiers, ensemble methods delivered superior accuracy, precision, recall, and F1-score, highlighting their robustness and adaptability in handling complex textual data. The consistent use of preprocessing techniques tokenization, stop-word removal, and TF-IDF vectorization—was found to significantly enhance model performance. These results underscore the critical role of model selection and preprocessing in optimizing AI-driven sentiment analysis, with practical implications for applications in social media analytics, customer feedback interpretation, and educational assessment systems.

Aspect	Description
Top Performing Model	**Gradient Boosting (Ensemble Learning)**
Other Ensemble Models	Random Forest

FULL TEXTS BOOK

Best Individual Classifiers	Support Vector Machines (SVM), Logistic Regression
Underperforming Model	Naive Bayes
Performance Metrics Considered	Accuracy, Precision, Recall, F1-Score
Preprocessing Techniques	Tokenization, Stop-word Removal, TF-IDF Vectorization
Impact of Preprocessing	Significantly improved performance across all models
Strengths of Ensemble Methods	Robustness, high accuracy, adaptability to complex text data
Application Domains	- Social Media Analytics- Customer Feedback Interpretation- Educational Assessment Systems
Key Implication	Appropriate model selection and consistent text preprocessing are critical for achieving optimal sentiment classification in AI applications.

Table2. Comparative Performance of Sentiment Analysis Models

XIV.DISCUSSION

The discussion of this study highlights the comparative performance of various machine learning algorithms in sentiment analysis tasks, emphasizing the superior effectiveness of ensemble methods, particularly Gradient Boosting, over traditional classifiers such as Naive Bayes and Logistic Regression. While conventional models offer computational efficiency, ensemble techniques demonstrated greater accuracy and robustness by integrating multiple learners and reducing classification errors. The application of preprocessing techniques, including tokenization and TF-IDF vectorization, significantly enhanced model performance by refining input data. The analysis also underscores the influence of dataset characteristics on algorithmic outcomes and suggests that future work may benefit from incorporating advanced models or hybrid approaches. Overall, the findings affirm the critical role of algorithm selection and preprocessing in the development of accurate and reliable sentiment analysis systems.

XV.CONCLUSION

This research undertook a detailed comparative study of various machine learning algorithms employed for sentiment analysis, emphasizing the integration of artificial intelligence techniques in understanding textual emotions and opinions. By systematically evaluating traditional classifiers—such as Naive Bayes, Support Vector Machines, Random Forest, and Logistic Regression—alongside ensemble models like Gradient Boosting and Voting Classifiers, the study aimed to identify the most effective and reliable methods for sentiment classification tasks. Comprehensive preprocessing steps, including tokenization, stop-word removal, and TF-IDF vectorization, were applied to ensure that the input data was suitably structured for analysis. The models were trained and tested on a benchmark dataset, with

FULL TEXTS BOOK

performance measured through accuracy, precision, recall, and F1-score. The findings revealed that ensemble approaches, particularly Gradient Boosting, consistently outperformed individual classifiers, demonstrating superior accuracy and robustness across evaluation metrics.

The comparative insights gained from this investigation underscore the importance of model selection in AI-driven sentiment analysis applications. Ensemble models, by leveraging the strengths of multiple algorithms, offered improved generalizability and predictive performance. This suggests their suitability for practical deployment in real-world systems, including social media monitoring, business intelligence platforms, and educational feedback mechanisms. In conclusion, this study contributes to the growing body of literature in the domain of intelligent text analytics by offering empirical evidence on the performance variations among commonly used machine learning techniques. Future work may focus on expanding this research by incorporating deep learning architectures, domain-specific datasets, and real-time analysis frameworks to further enhance the accuracy and adaptability of sentiment classification systems.

XVI.FUTURE WORK

Future research in sentiment analysis should focus on enhancing model generalizability through the use of multilingual and domain-specific datasets, and exploring advanced deep learning architectures such as LSTM and transformer-based models for improved contextual understanding. The integration of hybrid approaches combining lexicon-based and machine learning techniques, as well as the development of real-time adaptive systems, presents promising opportunities. Additionally, addressing ethical considerations including bias mitigation, data privacy, and algorithmic transparency will be essential. Expanding sentiment analysis to incorporate emotion detection and sarcasm recognition could further enrich analytical capabilities in diverse application domains.

XVII.RECOMMENDATIONS

Based on the comparative analysis conducted in this study, it is recommended that future sentiment analysis research and applications prioritize the use of ensemble learning techniques, particularly Gradient Boosting, due to their superior performance and robustness. Emphasis should be placed on careful dataset selection and preparation, including standardized text preprocessing methods such as tokenization and TF-IDF vectorization. Multiple evaluation metrics should be utilized to provide a holistic assessment of model effectiveness, especially in imbalanced data scenarios. Moreover, researchers are encouraged to explore hybrid and deep learning models for enhanced classification accuracy. Real-world implementation and continuous model updating are also essential to maintain relevance and ensure adaptability to evolving linguistic trends and user-generated content across digital platforms.

XVIII.REFERENCES

[1] “Prashant Awasthi, "International Journal of Engineering Technology Research & Management FORECASTING STOCK MARKET INDICES THROUGH THE INTEGRATION OF MACHINE LEARNING TECHNIQUES", Volume-07 Issue 02, February -2023”

[2] A. A. Alhashmi, A. Alshammari, M. K. Saeed, M. S. A. Alzaidi, F. Alanazi, and A. Sayed, “APPLIED LINGUISTICS-DRIVEN ARTIFICIAL INTELLIGENCE APPROACH FOR SENTIMENT ANALYSIS AND CLASSIFICATION ON SOCIAL MEDIA,” *Fractals*, vol. 33, no. 02, Jan. 2025, doi: 10.1142/s0218348x25400158.

FULL TEXTS BOOK

[3] K. R. Dahal, A. Gupta, and N. R. Pokhrel, “Predicting the Direction of NEPSE Index Movement with News Headlines Using Machine Learning,” *Econometrics*, vol. 12, no. 2, p. 16, Jun. 2024, doi: 10.3390/econometrics12020016.

[4] A. Fathollahi, “Machine Learning and Artificial Intelligence Techniques in Smart Grids Stability Analysis: A Review,” *Energies*, vol. 18, no. 13, p. 3431, Jun. 2025, doi: 10.3390/en18133431.

[5] N. Singh and U. C. Jaiswal, “Sentiment Analysis Using Machine Learning: A Comparative Study,” *ADCAIJ Adv. Distrib. Comput. Artif. Intell. J.*, vol. 12, p. e26785, Dec. 2023, doi: 10.14201/adcaij.26785.

[6] R. Shad, K. Potter, and A. Gracias, “Natural Language Processing (NLP) for Sentiment Analysis: A Comparative Study of Machine Learning Algorithms,” Oct. 30, 2024, *MDPI AG*. doi: 10.20944/preprints202410.2338.v1.

[7] D. P. Dash, M. Kolekar, C. Chakraborty, and M. R. Khosravi, “Review of Machine and Deep Learning Techniques in Epileptic Seizure Detection using Physiological Signals and Sentiment Analysis,” *ACM Trans. Asian Low-Resour. Lang. Inf. Process.*, vol. 23, no. 1, pp. 1–29, Jan. 2024, doi: 10.1145/3552512.

[8] L. Ashbaugh and Y. Zhang, “A Comparative Study of Sentiment Analysis on Customer Reviews Using Machine Learning and Deep Learning,” *Computers*, vol. 13, no. 12, p. 340, Dec. 2024, doi: 10.3390/computers13120340.

[9] D. A. Balaji, S. Nikhil, Y. V. R. Kiran, T. Jayaram, and V. Sivaiah, “Comparative Analysis of Machine Learning and Deep Learning Models for Stock Market Prediction Using Continuous and Binary Data,” vol. 9, no. 3, 2025.

[10] F. Hassan *et al.*, “Performance evolution for sentiment classification using machine learning algorithm,” *J. Appl. Res. Technol. Eng.*, vol. 4, no. 2, pp. 97–110, May 2023, doi: 10.4995/jarte.2023.19306.

[11] M. Usmani, S. H. Adil, K. Raza and S. S. A. Ali, "Stock market prediction using machine learning techniques," 2016 3rd International Conference on Computer and Information Sciences (ICCOINS), Kuala Lumpur, Malaysia, 2016, pp. 322-327, doi: 10.1109/ICCOINS.2016.7783235.

[12] Rouf, N.; Malik, M.B.; Arif, T.; Sharma, S.; Singh, S.; Aich, S.; Kim, H.-C. Stock Market Prediction Using Machine Learning Techniques: A Decade Survey on Methodologies, Recent Developments, and Future Directions. Electronics 2021, 10, 2717, doi:10.3390/electronics10212717.

[13] Strader, Troy J.; Rozycki, John J.; ROOT, THOMAS H.; and Huang, Yu-Hsiang John (2020) "Machine Learning Stock Market Prediction Studies: Review and Research Directions," Journal of International Technology and Information Management: Vol. 28: Iss. 4, Article 3, doi:10.58729/1941-6679.1435.

[14] Kumar, G., Jain, S. & Singh, U.P. Stock Market Forecasting Using Computational Intelligence: A Survey. Arch Computat Methods Eng 28, 1069–1101 (2021), doi:10.1007/s11831-020-09413-5.

[15] Ayala, J., García-Torres, M., Noguera, J. L. V., Gómez-Vela, F., & Divina, F. (2021). Technical analysis strategy optimization using a machine learning approach in stock market indices. Knowledge-Based Systems, 225, 107119,doi: 10.1016/j.knosys.2021.107119.

FULL TEXTS BOOK

[16] Marzband, M.; Parhizi, N.; Savaghebi, M.; Guerrero, J.M. Distributed smart decision-making for a multimicrogrid system basedon a hierarchical interactive architecture. IEEE Trans. Energy Convers. 2015, 31, 637–648.

[17] Liao, Z.; Kally, J.; Ru, S. Probabilistic modeling of renewable energy sources in smart grids: A stochastic optimization perspective. Sustain. Cities Soc. 2024, 109, 105522.

[18] Guo, Y.; Ten, C.-W.; Hu, S.; Weaver, W.W. Preventive maintenance for advanced metering infrastructure against malware propagation. IEEE Trans. Smart Grid 2015, 7, 1314–1328.

[19] Das, S.; Ohba, Y.; Kanda, M.; Famolari, D.; Das, S.K. A key management framework for AMI networks in smart grid. IEEE Commun. Mag. 2012, 50, 30–37.

[20] Usama, M.; Qadir, J.; Raza, A.; Arif, H.; Yau, K.l.A.; Elkhatib, Y. Unsupervised Machine Learning for Networking: Techniques, Applications and Research Challenges. IEEE Access 2019, 7, 65579–65615.

BUILDING A GLOBAL CAREER WHILE PURSUING AN ENGINEERING DEGREE: STRATEGIES FOR STUDENTS WORLDWIDE

Dr. Sameer Suresh Nanivadekar

A P SHAH INSTITUE OF TECHNOLOGY ,THANE ,MAHARSHTRA .INDIA

ABSTRACT

In an increasingly interconnected and competitive global job market, engineering students must adopt a proactive approach to career development well before graduation. This paper explores effective strategies for building a successful career while still enrolled in engineering programs, regardless of geographical location. It examines how academic performance, practical experiences, soft skills, global certifications, and digital presence contribute to employability. By analyzing international trends, case studies, and institutional support systems, the paper offers actionable insights for students to bridge the gap between learning and livelihood across global contexts.

Keywords: Career Development, Global Employability, Engineering Education, Internships, Skill Acquisition, Soft Skills, Professional Networking, Entrepreneurship, Remote Work, Student Leadership, Career Planning

1. Introduction

The traditional model of completing an engineering degree before considering career options is no longer viable in today's fast-paced, digital-first economy. The fourth industrial revolution, digital transformation, and post-pandemic employment shifts have redefined job-readiness for engineering students. Across regions—from North America to Africa, Europe to Southeast Asia—employers increasingly expect graduates to possess real-world experience, digital fluency, and cross-cultural communication abilities even before completing their studies.

This paper addresses the essential question: *How can engineering students worldwide actively build their careers while still studying?*

2. The Case for Early Career Preparation

Numerous studies by international organizations like the World Economic Forum (WEF), OECD, and UNESCO emphasize the urgency for future engineers to:

- Start career planning by their second year.
- Develop both domain expertise and global competencies.
- Engage in practical, project-based learning.
- Demonstrate adaptability in uncertain and hybrid job markets.

For example, the WEF Future of Jobs Report 2023 suggests that **over 44% of workers will need significant skill reskilling** within five years, even within STEM domains.

FULL TEXTS BOOK

3. Foundational Elements for Global Career Building

3.1 Academic Rigor and Curriculum Integration

Students must not only aim for good grades but focus on **learning outcomes** that align with global job roles—e.g., data analysis, AI fundamentals, sustainability engineering, or control systems.

- Choose electives strategically (AI, IoT, renewable energy, blockchain).
- Participate in international MOOC platforms (edX, Coursera, FutureLearn) to gain exposure to cross-border curriculum standards.

3.2 In-Demand Skills for the Global Market

Engineering students should actively develop the following:

- **Technical Skills**: Coding, system design, simulation tools (e.g., MATLAB, SolidWorks, AutoCAD), cloud platforms (AWS, Azure).
- **Soft Skills**: Cross-cultural communication, collaboration, time management, leadership.
- **Digital Presence**: LinkedIn profile optimization, GitHub repositories, personal portfolio websites.

3.3 International Internships and Exchange Programs

Securing global internships—virtual or physical—can significantly enhance one's professional profile:

- Apply through platforms like Internshala (Asia), WayUp (US), Erasmus Intern (EU), and IAESTE (global).
- Leverage university exchange programs and credit-transfer options to participate in international labs and companies.

3.4 Participation in Global Competitions

Technical competitions like:

- **Microsoft Imagine Cup**
- **Google Summer of Code**
- **Formula Student**
- **Shell Eco-marathon**

offer exposure to international standards and networking.

Many of these initiatives are now hybrid or remote, reducing the entry barrier for students in developing countries.

4. Diverse Career Pathways: Think Globally

4.1 Pathway 1: Global Corporate Employment

- Engage in **campus ambassador programs** of MNCs.
- Participate in **global hackathons** and coding sprints.
- Understand **ATS-optimized resumes**, global application platforms (Handshake, Glassdoor, Indeed).

FULL TEXTS BOOK

4.2 Pathway 2: Research and Academia

- Assist professors in internationally funded projects (e.g., Horizon Europe, NSF, DST).
- Publish research in journals indexed by Scopus, IEEE, Elsevier.
- Target international postgraduate education (MITACS, DAAD, Erasmus Mundus, Fulbright).

4.3 Pathway 3: Entrepreneurship and Startups

- Join startup incubators in your region or apply for international startup accelerators (Y Combinator, Seedstars, Techstars).
- Explore crowdfunding and MVP validation via platforms like Kickstarter, IndieGoGo.

4.4 Pathway 4: Freelancing and Remote Tech Careers

- Use Upwork, Toptal, Fiverr to work for clients globally.
- Build a niche portfolio (e.g., game development, WordPress, 3D modeling, data visualization).

5. Role of Global Mentorship and Professional Networking

Career growth is accelerated by relationships and reputation:

- **LinkedIn**: Follow industry leaders, contribute to discussions, publish short tech articles.
- **Global professional bodies**: IEEE, ASME, ACM, PMI, IFAC.
- **Alumni Connects**: Reach out to seniors working abroad or at leading companies.

Global mentorship platforms like **ADPList** and **Ten Thousand Coffees** offer structured mentoring for free.

6. Overcoming Regional and Economic Barriers

Challenge	**Global Solution**
Limited access to labs	Use open-source simulators and remote access labs
Language barriers	Practice English via Toastmasters or language exchanges
Financial constraints	Apply for scholarships, grants, and paid virtual internships
Visa restrictions	Opt for remote work or research collaborations
Lack of awareness	Follow international career counselors and join online communities

FULL TEXTS BOOK

7. Global Case Studies

Case Study A: Lucia from Brazil

Lucia, a mechanical engineering student, completed a remote internship with a German automotive company through IAESTE and published a paper in an international journal. This helped her secure a DAAD-funded master's degree in Munich.

Case Study B: Fahad from Pakistan

Fahad built a GitHub portfolio showcasing his Python projects and freelanced for European clients on Upwork. He was later hired remotely by a Canadian SaaS startup and now leads a distributed engineering team.

8. Institutional Responsibilities and Recommendations

To promote career-readiness, institutions across the globe should:

- Introduce **career credit systems** (for internships, freelancing, certifications).
- Offer **career bootcamps** on resume writing, interview skills, global job search platforms.
- Establish **international industry collaborations**.
- Enable **credit-linked international internships** and **virtual exchange programs**.

9. Conclusion

Career development is no longer confined to the post-graduation phase. It must be nurtured alongside academic education, regardless of geographical or economic barriers. By leveraging online tools, global opportunities, soft skill training, and remote work, engineering students from all regions can position themselves competitively on the world stage. The future belongs to those who learn early, adapt continuously, and connect globally.

10. REFERENCES

1. World Economic Forum (2023). *The Future of Jobs Report*.
2. OECD (2022). *Skills Outlook 2022: The Role of Education in Building Skills for Work*.
3. UNESCO-UNEVOC (2021). *Greening Technical and Vocational Education and Training*.
4. LinkedIn (2023). *Global Talent Trends Report*.
5. Coursera (2022). *Global Skills Report*.

FULL TEXTS BOOK

Mrs.E.Elavarasi,Yuvaraj AR,J.Jayadurka,Dr.R.Srinivasan

Faculty of Pharmacy, Bharath Institute of Higher Education and Research,Chennai-73

ABSTRACT

Pharmaceutical goods therapeutic efficacy, safety, and patient compliance are all improved by effective drug formulation and delivery methods. The exact blending of excipients and active pharmaceutical ingredients to produce a dosage form that guarantees medication stability, bioavailability, and ideal release patterns is known as formulation science. Delivery systems, on the other hand, are made to deliver medications to the intended location in the body as efficiently and as safely as possible. Recent developments in transdermal patches, liposomes, microspheres, nanotechnology, and controlled-release devices have completely changed how drugs are delivered. Site-specific administration, fewer doses, and better pharmacokinetics and pharmacodynamics are all made possible by these technologies. For instance, delivery systems based on nanoparticles can pass through biological barriers and offer prolonged medication release, which is especially advantageous for treating cancer and illnesses of the central nervous system. Likewise, buccal systems and oral thin films provide quick start of action and simplicity of administration, particularly for elderly and pediatrics patients. Customizing drug delivery systems to each patient's unique genetic profile and clinical condition, personalized medicine is also having an impact on formulation tactics. In order to guarantee safety and compliance, regulatory concerns and biocompatibility evaluations are still crucial when developing innovative delivery methods. More accuracy and personalization in drug distribution are promised by the combination of 3D printing and artificial intelligence in formulation creation. In conclusion, pharmacotherapy is changing as a result of advancements in drug formulation and delivery, which make treatment more patient-centered, effective, and focused.

Keywords: controlled release, nanotechnology, bioavailability, drug formulation, delivery methods, patient compliance.

INTRODUCTION

Over the past 100 years, the field of pharmaceutical sciences has made huge strides in how drugs are found, made, and given to patients. Drug formulation and delivery systems are one of the most important parts of this evolution. The main goal of drug formulation is to make sure that a drug is stable, bioavailable, effective, and acceptable to patients. The delivery system's main goal is to get the active pharmaceutical ingredient (API) to the site of action in the most efficient and controlled way possible. These systems are very important for turning the pharmacological potential of a substance into real-world therapeutic effects. Drug formulation is the process of mixing the active pharmaceutical ingredient (API) with other substances to make a final medicine in a certain dosage form, like tablets, capsules, injections, or creams that are put on the skin. It involves carefully mixing, processing, and stabilising the ingredients to make sure they are all the same, can be reproduced, and meet all regulatory standards. Formulation science looks at the drug's solubility, pKa, partition coefficient, and stability, among other physicochemical properties, to see how they affect the dosage form's performance. Drug delivery systems (DDS) are the methods and technologies used to move a drug through the body to get the desired therapeutic effect. These systems can be simple, like taking medicine by mouth or through an injection, or more advanced, like transdermal patches, nanoparticles, liposomes, or targeted drug

FULL TEXTS BOOK

delivery systems. A well-designed DDS makes sure that the drug gets to the right place at the right concentration and for the right amount of time, while reducing side effects throughout the body. Even when powerful new drug molecules are discovered, their clinical utility is frequently limited by inadequate formulation and delivery. Enzymatic breakdown, pH fluctuation, first-pass metabolism, and cellular membranes are some of the physiological barriers that a drug must overcome on its way from the site of administration to its site of action. Drug delivery systems are purposefully created to get around these obstacles and raise the drug's therapeutic index. For example, targeted delivery can concentrate the medication at a disease site, lowering off-target toxicity, while sustained-release formulations can decrease the frequency of dosing and enhance patient compliance. Additionally, formulation techniques are essential for repurposing outdated medications or enhancing the efficacy of currently used treatments. Novel excipients, delivery methods, or release profiles can all be used to increase the efficacy or acceptability of the same drug molecule. For instance, better oral bioavailability and clinical results have been achieved by reformulating poorly water-soluble medications using solid dispersions or nanoemulsions. Drug formulation has a long history that began with the preparation of crude herbal extracts using simple methods like infusion and decoction in ancient civilisations. More complex formulations like tablets, capsules, and injections were created as pharmacy became a recognised scientific field in the 19th and 20th centuries. Rapid developments in nanotechnology, biotechnology, and polymer science during the second half of the 20th century greatly increased the potential for drug delivery. Novel systems like transdermal patches, osmotic pumps, controlled-release matrices, and targeted delivery carriers were created as a result of this evolution. Pharmacokinetics absorption, distribution, metabolism, and excretion and pharmacodynamics drug action and effect principles are now frequently used to guide modern drug delivery systems. The demand for sophisticated delivery systems with precision targeting capabilities has increased in tandem with the rise of personalised medicine and biotechnology-based treatments such as gene therapy, siRNA, and monoclonal antibodies. A drug formulation is more than just a concoction of substances. Numerous excipients that serve as solubilizers, stabilisers, taste masks, or enhance absorption must be carefully chosen and optimised. These excipients consist of preservatives, lubricants, diluents, binders, disintegrants, and surfactants. Regulatory bodies like the FDA and EMA rigorously assess formulations for efficacy, safety, and quality, highlighting the significance of analytical validation, stability testing, and pre-formulation studies. A formulation's design also needs to take into account the intended onset and duration of action, the target patient population (e.g., paediatric or geriatric), the route of administration, and the physicochemical compatibility of its constituent parts. Scaling up procedures for commercial manufacturing is another aspect of formulation science that adds complexity, including choosing the right equipment, managing the process, and adhering to Good Manufacturing Practices.

Delivery System	Route of delivery		Advantages and benefits	Examples of drug
Oral	Oral		non-invasive, patient compliance	Paracetamol Omeprazole
Parenteral	IV, IM, SC		Rapid onset, 100% bioavailability	Insulin Ceftriaxone
Transdermal	Skin		liver metabolism, sustained release	Nicotine Fentanyl

FULL TEXTS BOOK

Pulmonary	Lungs		Fast absorption, systemic effect	Salbutamol Budesonide
Nasal	Nose		avoids first-pass metabolism	Oxymetazoline Flu vaccine
Rectal	Rectum		Useful in unconscious patient	Paracetamol Bisacodyl
Vaginal	Vagina		Local treatment, hormone therapy	Clotrimazole
Sustained Release	Various		Reduced dosing frequency, consistent plasma levels	Propranolol
Buccal	Mouth mucosa		Rapid onset, bypass liver metabolism	Nitroglycerin Fentanyl
Smart Systems	Various		personalized therapy	Thermo-sensitive gels
Molecular	Systemic/Cellular		Enables mRNA,	AAV gene therapies
3D-Printed Formulations	Oral/Custom		Personalized dosing, multilayer drugs	Levetiracetam
Mucoadhesive Systems	Mucosal surfaces		Prolonged retention at target site	Vaginal muco adhesive gels
Floating Drug Systems	Oral		Prolonged gastric retention, enhanced bioavailability	Floating Metformin

Table 1: The route of delivery system with drug advantages

Types of drug delivery system:

Because they control the way, time, and location of a drug's release within the body, drug delivery systems (DDS) are essential to contemporary therapeutics. In order to maximise the pharmacokinetic and pharmacodynamic characteristics of medications, these systems are purposefully made to increase effectiveness, decrease adverse effects, and boost patient compliance. Drug delivery systems can be categorised according to their targeting mechanism, drug release profile, and route of administration. The oral delivery system is one of the most popular and generally recognised drug delivery techniques. It is preferred because of its high patient compliance, non-invasiveness, and convenience. Oral medication delivery can be targeted to particular areas of the gastrointestinal tract, be immediate-release, or be modified-release (including delayed and sustained release). Nevertheless, issues like first-pass metabolism and enzymatic degradation could compromise the bioavailability of medications taken orally. Formulation scientists use strategies like enteric coatings, solid dispersions, and complexation with cyclodextrins to get around these restrictions. Parenteral delivery methods avoid the gastrointestinal system and are employed when a medication has a poor oral absorption rate or when a quick onset of action is needed. These comprise injections administered intravenously (IV), intramuscularly (IM),

FULL TEXTS BOOK

subcutaneously (SC), and intradermally. When exact dosage is required or in an emergency, intravenous delivery is utilised because it offers 100% bioavailability. Depot formulations, liposomal injectables, and polymer-based microspheres are examples of innovative injectable systems that enable controlled drug release over prolonged periods of time. Another non-invasive method for delivering drugs through the skin and into the bloodstream is through transdermal drug delivery systems (TDDS). Transdermal patches, gels, and ointments can minimise gastrointestinal side effects, circumvent hepatic metabolism, and deliver controlled and sustained drug release. Fentanyl patches for chronic pain, nicotine patches for quitting smoking, and hormonal patches for contraception are a few examples. Enhancers like microneedles, iontophoresis, or ultrasound are frequently used to get around the stratum corneum, which is the main barrier to transdermal absorption. Particularly useful for the local treatment of respiratory diseases like asthma and chronic obstructive pulmonary disease (COPD), pulmonary and nasal delivery systems are also becoming more and more popular for the systemic administration of proteins, peptides, and vaccines. Devices such as nebulisers, dry powder inhalers, and metered-dose inhalers (MDIs) guarantee that the medication reaches the alveoli, where absorption occurs quickly and widely. Since the olfactory pathway bypasses the blood-brain barrier and offers direct access to the brain, nasal sprays are being researched for drug delivery to the central nervous system (CNS). By getting past anatomical barriers like the cornea and conjunctiva, ocular delivery systems seek to treat eye conditions. Because of their poor corneal permeability and rapid drainage, conventional eye drops are not very effective. In situ gels, drug-eluting contact lenses, ocular inserts, and nanoparticles that enhance residence time and drug absorption are examples of advanced ocular DDS. Localised treatment for ailments like haemorrhoids, infections, or hormone replacement therapy is available through vaginal and rectal delivery systems. In addition to being formulated for systemic absorption in patients who are unconscious, vomiting, or incapable of taking oral medications, these include suppositories, pessaries, and gels. Targeted drug delivery, which involves delivering medications to particular tissues, cells, or even intracellular compartments in order to minimise systemic toxicity and maximise therapeutic effect, is a very complex and developing class of DDS. This is especially crucial in cancer treatment because traditional chemotherapy damages both healthy and cancerous cells. Liposomes, which are phospholipid vesicles that contain medications and can have their surfaces altered with ligands for cell-specific targeting, are examples of targeted delivery systems. Niosomes, which are composed of non-ionic surfactants and share structural similarities with liposomes, provide improved stability and biocompatibility. For drug targeting, dendrimers, highly branched polymeric nanocarriers, and polymeric nanoparticles offer regulated release, improved permeability, and customised surface chemistry. Lipid-based nanocarriers known as solid lipid nanoparticles (SLNs) and nanostructured lipid carriers (NLCs) combine the benefits of polymeric and liposome nanoparticles, particularly for hydrophobic medications. For formulations requiring controlled release, microspheres and microcapsules are frequently utilised. Made of biodegradable polymers like PLGA (poly-lactic-co-glycolic acid), these spherical particles can be made to release medications gradually over the course of days or months, making them perfect for depot injections or chronic illnesses. Long-term release is provided by implanted drug delivery systems, which insert a drug-containing device into a bodily cavity or beneath the skin. Insulin pumps, macular degeneration eye implants, and contraceptive implants are a few examples. These systems enhance adherence and lessen the need for frequent dosing, particularly in patients with long-term conditions. The purpose of controlled-release and sustained-release systems is to keep the drug concentration in plasma steady for an extended amount of time. These include gastroretentive devices that extend the duration of gastric residence, hydrophilic matrix systems, and osmotic pump tablets. By minimising peak-trough effects and lowering drug level fluctuations, controlled-release formulations enhance therapeutic outcomes. Drugs can be released by smart drug delivery systems, also referred to as stimuli-responsive systems, in response to particular biological signals or environmental triggers like pH, temperature, redox conditions, or enzymes. Because the

FULL TEXTS BOOK

tumour microenvironment can be used to release drugs at specific sites, these systems are at the forefront of precision medicine, particularly in diseases like cancer. Examples include glucose-responsive insulin systems for diabetes treatment and pH-sensitive nanoparticles for delivery to specific tumours. Delivery methods for genes and nucleic acids are a developing field, particularly in the age of mRNA vaccines and gene therapy. To transfer nucleic acids like siRNA, DNA, or mRNA into cells, lipid nanoparticles (LNPs), viral vectors (such as AAV or lentivirus), and polymeric carriers are employed. The effectiveness of lipid-based gene delivery platforms in conventional therapies has been highlighted by the recent success of mRNA COVID-19 vaccines. Additionally, drug absorption through the oral mucosa is made possible by buccal and sublingual delivery systems, which circumvent the hepatic first-pass effect. These routes provide a quick onset and are especially helpful for hormones, emergency medications, and cardiovascular drugs (like nitroglycerin). One kind of gastroretentive device called a floating drug delivery system (FDDS) is made to stay afloat in the stomach for long periods of time, which improves the absorption of medications with limited absorption windows. Similarly, mucoadhesive systems improve absorption and therapeutic efficacy by lengthening the duration that medications remain at mucosal sites. Lastly, the idea of personalised medicine is being revolutionised by 3D-printed drug delivery systems, which give precise control over the dosage, geometry, and patterns of drug release. Multiple APIs with complex release profiles that are suited to the needs of each patient can be accommodated by this technology in a single dosage form. Ultimately, the drug's nature, the ailment being treated, the patient's characteristics, and the intended therapeutic results all influence the drug delivery system selection. In order to optimise treatment, minimise side effects, and guarantee better patient outcomes, sophisticated and intelligent drug delivery systems are becoming more and more important as drug molecules become more complex and personalised therapies gain traction.

Recent innovations in formulation and delivery science

More accurate, effective, and patient-centered treatments are now possible thanks to recent advancements in formulation and delivery science, which have drastically changed the pharmaceutical development landscape. The increasing demand for targeted and personalised medicine, advances in material science, and a better understanding of how drugs behave in biological systems are the main forces behind these developments. The creation of drug delivery systems based on nanotechnology, such as solid lipid nanoparticles (SLNs), nanoparticles, nanomicelles, and nanostructured lipid carriers (NLCs), is among the most significant advancements. These nanoscale carriers allow for targeted delivery and controlled release while improving the solubility, stability, and bioavailability of medications that are poorly soluble in water. For instance, peptides, antibiotics, and anticancer medications are frequently encapsulated in polymeric nanoparticles derived from biodegradable polymers like PLGA or chitosan. The application of lipid-based drug delivery systems, especially liposomes and lipid nanoparticles (LNPs), represents yet another significant advancement. Both hydrophilic and lipophilic medications can be encapsulated in liposomes, which are spherical vesicles with phospholipid bilayers that improve delivery to particular tissues and lessen toxicity. By making it easier to transfer otherwise unstable genetic material into cells, lipid nanoparticles—particularly those employed in the delivery of mRNA vaccines, as in COVID-19 vaccines—have completely transformed nucleic acid-based treatments. In gene therapy and vaccine development, mRNA-based drug delivery has become a cutting-edge strategy that enables quick production and individualised treatment for genetic disorders, cancer, and infectious diseases. LNPs are essential for the effectiveness of contemporary vaccine platforms because they shield mRNA from degradation, enhance cellular uptake, and encourage endosomal escape.Additionally, by creating drug formulations that can be customised, 3D printing technology has ushered in a new era of personalised medicine. Pharmaceutical firms can create dosage forms with intricate geometries, multiple drug combinations, and customised release profiles using 3D printing. Dose strength, surface area, and disintegration time can all be precisely

FULL TEXTS BOOK

controlled with this technology. The potential of this strategy is demonstrated by SPRITAM® (levetiracetam), the first FDA-approved 3D-printed medication, which provides quick disintegration for patients who have trouble swallowing. Smart drug delivery or stimuli-responsive systems are becoming more popular in addition to 3D printing. Certain physiological or external stimuli, such as pH variations, temperature changes, redox conditions, enzymes, or even magnetic fields, cause these systems to release medications. For example, in the acidic tumour microenvironment, pH-sensitive nanoparticles can release cytotoxic drugs selectively while protecting healthy tissues in cancer therapy. Drugs can also be released on demand at inflammatory or feverish sites using temperature-sensitive hydrogels. These clever systems have a lot of potential for controlling the timing and location of a drug's action, which would increase its effectiveness and reduce its side effects. Mucoadhesive drug delivery is another cutting-edge field that enhances drug absorption through mucosal surfaces like the vaginal mucosa, gastrointestinal tract, nasal passages, and oral cavity. Mucoadhesive films, gels, and tablets improve the formulation's residence time at the absorption site, improving bioavailability and lowering the frequency of doses. These systems provide important benefits in conditions like vaginal infections or oral ulcers where localised therapy is desired. Another innovation in formulation is floating drug delivery systems (FDDS), which increase the absorption of medications with limited absorption windows in the upper gastrointestinal tract by extending the gastric residence time. Usually low-density formulations, these systems float on stomach fluid and release the medication gradually, enhancing the effectiveness of medications like ciprofloxacin or metformin. Another noteworthy development in formulation science is the use of biodegradable polymers. These polymers—like polylactic acid (PLA), polyglycolic acid (PGA), and their copolymers—are utilised to make injectable depots, implants, and microspheres. Over time, they break down into non-toxic byproducts that release the medication in a regulated way. This technology is used, for instance, in depot injections for the treatment of schizophrenia and long-acting contraceptive implants to guarantee steady drug release over several months. By improving the solubility, stability, and circulation time of therapeutic proteins and peptides, PEGylation—the attachment of polyethylene glycol chains to drugs—has enhanced their pharmacokinetics. Peginterferon and pegfilgrastim are examples of PEGylated medications that exhibit improved therapeutic efficacy and decreased dosage frequency. Numerous drawbacks of traditional drug delivery have been addressed by the creation of transdermal systems and microneedle patches. Transdermal patches offer steady-state plasma concentrations, enhanced patient adherence, and a non-invasive method of systemic medication delivery. The delivery of vaccines, insulin, and other biologics straight into the dermis or systemic circulation is made possible by microneedles, which form tiny channels in the skin without causing discomfort or bleeding. They are especially helpful for patients with needle phobia and during immunisation campaigns. The development of formulations that stay liquid during administration but gel when exposed to physiological conditions has been made possible by advancements in in situ gelling systems. This method encourages prolonged drug retention and sustained release, which makes it especially advantageous in ocular, rectal, nasal, and injectable formulations. Antibody-drug conjugates (ADCs) are a significant advancement in targeted delivery in the oncology field. These systems combine the strength of cytotoxic medications with the selectivity of monoclonal antibodies. The medication can be internalised and released inside the tumour while protecting healthy cells because the antibody binds to particular antigens on cancer cells. ADCs that treat HER2-positive breast cancer, such as trastuzumab emtansine (Kadcyla®), have demonstrated significant efficacy. In a similar vein, highly branched synthetic polymers called dendrimers have become multipurpose nanocarriers that can simultaneously deliver medications, genes, and imaging agents. They are perfect candidates for theranostic applications because of their structure, which permits precise control over size, shape, and surface functionality. Fast-dissolving films (FDFs) and oral disintegrating tablets (ODTs) are patient-friendly innovations, especially for psychiatric, paediatric, or elderly populations that have trouble swallowing traditional tablets. These dosage forms increase convenience and adherence by dissolving quickly in the mouth

FULL TEXTS BOOK

without the need for water. Supercritical fluid technology is also being investigated to enhance particle engineering and drug solubility. For medications with low aqueous solubility, this solvent-free and environmentally friendly process makes it possible to create nano- or microparticles with better dissolution profiles. A revolutionary change in pharmaceutical sciences is represented by the incorporation of machine learning (ML) and artificial intelligence (AI) into formulation development. These technologies are used to speed up the drug development process, screen excipients, simulate pharmacokinetic profiles, and predict the ideal formulation parameters. AI-powered modelling can shorten time-to-market for new products, increase scalability, and optimise nanocarrier design. Furthermore, the implementation of digital health technologies such as ingestible sensors, smart pills, and connected delivery devices enables real-time monitoring of therapeutic efficacy, physiological responses, and drug adherence, opening the door for closed-loop drug delivery systems. The development, administration, and patient experience of medications are being completely transformed by recent advances in formulation and drug delivery science. These developments, which range from smart hydrogels and 3D-printed dosage forms to gene delivery vectors and nanocarriers, are intended to improve patient outcomes, raise the therapeutic index, and aid in the transition to precision and personalised medicine. Future drug formulation and delivery will be more focused, effective, and individualised thanks to the ongoing integration of multidisciplinary approaches, such as nanotechnology, material science, biotechnology, and artificial intelligence.

Oral drug delivery:

The most popular and practical method of administering therapeutic agents is oral drug delivery. Pharmaceutical formulations are consumed orally, and the gastrointestinal (GI) tract is the primary site of drug absorption. Because of its cost-effectiveness, patient compliance, ease of administration, and non-invasiveness, this drug delivery method is widely used.A large variety of dosage forms, such as tablets, capsules, syrups, suspensions, powders, and emulsions, are included in the oral route. Because of their stability, portability, and ability to provide precise dosage, tablets and capsules are the most popular among these. The medication dissolves in the gastrointestinal fluids after administration, and then the intestinal epithelium absorbs it into the bloodstream. Oral drug delivery does, however, come with a number of difficulties. One of the main drawbacks is first-pass metabolism, which significantly lowers the drug's bioavailability after it has been absorbed and passes through the liver before entering the bloodstream. Efficacy may also be jeopardised by drug degradation brought on by the stomach's acidic pH or intestinal enzymatic activity. Drug performance is also impacted by food-drug interactions, low permeability, and poor solubility.Formulation scientists have created a number of sophisticated oral delivery techniques to get around these problems. Nanoparticle-based carriers to improve solubility and absorption, controlled-release and sustained-release systems for extended drug action, and enteric-coated formulations to stop gastric breakdown are a few examples. The efficacy of treatment is further enhanced by technologies like mucoadhesive polymers, self-emulsifying drug delivery systems (SEDDS), and solid lipid nanoparticles. Drug solubility, molecular size, pKa, GI motility, food presence, and physiological factors like gastric pH and enzyme activity are all factors that affect oral drug absorption. Addressing these factors is essential when developing a formulation.The possibilities of this approach are being expanded by recent developments in oral biologics delivery, oral dosage form 3D printing, and personalised medicine. Previously thought to be inappropriate for this route, oral insulin,




Figure 1: Oral route

FULL TEXTS BOOK

vaccines, and peptide medications are currently being actively studied to improve patient compliance. Because oral drug delivery is safe, easy to use, and well-received by patients, it continues to be a fundamental component of pharmaceutical therapy. Notwithstanding its drawbacks, ongoing advancements in drug delivery and formulation technologies have greatly increased the effectiveness and dependability of oral medications. Oral drug delivery has a bright and revolutionary future thanks to new developments like smart drug delivery and formulation design aided by artificial intelligence.




Figure 2: mechanism of oral drug delivery system

Parenteral Drug Delivery Systems:

The term "parenteral drug delivery" describes the administration of medications through methods other than the mouth or digestive system, most frequently by injection. This approach is essential in clinical settings where precise dosage, quick onset of action, and avoiding the first-pass metabolism are crucial. The Greek word "parenteral" literally means "administration beside the intestine." Although more specialised routes like intra-arterial, intrathecal, intra-articular, and intraperitoneal are also used under certain medical conditions, the primary parenteral administration routes are intravenous (IV), intramuscular (IM), subcutaneous (SC), and intradermal (ID). The intravenous route is the recommended approach for fluid/electrolyte therapy, chemotherapy, anaesthesia, and emergency treatments because it delivers the medication straight into the bloodstream, guaranteeing an immediate pharmacological effect and full bioavailability. Compared to intravenous administration, intramuscular administration—usually into the deltoid or gluteal muscle—allows for slower absorption, which is advantageous for vaccines and depot formulations. As demonstrated in insulin therapy, subcutaneous injections, which are administered beneath the skin, are preferred due to their ease of use and prolonged drug release. Vaccinations like BCG and allergy testing are the two main uses for intradermal injections, which are administered into the dermis. Parenteral formulations must meet strict manufacturing and storage requirements because they are usually sterile, pyrogen-free, and isotonic. Aqueous or oily solutions, suspensions, emulsions, and dry powders for reconstitution are examples of common forms. In order to prevent contamination and guarantee safety and effectiveness, parenteral product preparation requires the use of cutting-edge technologies and adherence to good manufacturing practices (GMP). To maintain formulation stability and patient comfort, tonicity adjusters, buffering agents, stabilisers, and preservatives are added. Syringes, ampoules, vials, prefilled syringes, infusion pumps, and autoinjectors are among the delivery devices used; there is a growing trend towards disposable and user-friendly systems that lower the risk of infections linked to healthcare. For medications like peptides, proteins, monoclonal antibodies, and vaccines that are poorly absorbed, unstable, or broken down in the gastrointestinal tract, parenteral delivery offers a significant benefit. Additionally, parenteral

FULL TEXTS BOOK

administration improves therapeutic compliance in chronic diseases by enabling controlled and sustained release of medications via depot or implantable formulations. Nevertheless, the system has disadvantages, such as injection site pain or discomfort, infection risk, tissue damage locally, and high sterile production costs. For targeted and regulated delivery, especially in oncology and disorders of the central nervous system, nanoparticles, liposomes, micelles, and dendrimers are being investigated. These carriers can lessen systemic toxicity while enhancing pharmacokinetics, stability, and solubility. The creation of long-acting injectables has been made possible by biodegradable polymers and PEGylation techniques. Furthermore, on-demand drug release is being studied using intelligent delivery systems that react to pH, temperature, or enzymes. Parenteral product regulations are strictly regulated, with regulations from organisations such as the FDA, EMA, and CDSCO defining strict standards for particulate matter limits, pyrogen testing, sterility, and container-closure integrity. Pharmacovigilance systems monitor adverse reactions after they occur, and pharmacopoeial standards guarantee that injectable medications fulfil predetermined quality standards. The field of parenteral drug delivery is constantly growing thanks to developments in formulation science, biotechnology, and device engineering. Injectable biosimilars that increase accessibility and affordability, self-administration tools, and customised injectables are the trends of the future. Combination products, in which the medication and its delivery system are combined, are increasingly being developed, particularly for chronic diseases like cancer, rheumatoid arthritis, and diabetes. Additionally, for conditions requiring long-term medication with low dosing frequency, depot injections and microsphere-based systems are being actively investigated. Since the majority of vaccines, monoclonal antibodies, and critical care medications were given by injection during the COVID-19 pandemic, the importance of parenteral delivery has been brought to light. This has underscored the necessity for stable, scalable, and sterile parenteral platforms. To sum up, parenteral drug delivery systems are essential to contemporary medicine because they provide quick, dependable, and efficient ways to administer medications. Their ability to deliver complex biologics and achieve precise pharmacokinetic profiles makes them indispensable, despite the challenges of invasiveness, cost, and sterility. As materials science, nanotechnology, and personalised medicine come together, parenteral delivery has a bright future ahead of it, with the potential to improve patient outcomes for a variety of illnesses.

Novel Drug Delivery Systems:

Novel Drug Delivery Systems (NDDS) are a cutting-edge method of creating and delivering medications in a way that maximises therapeutic effectiveness, increases patient compliance, and reduces side effects. With the help of cutting-edge technologies, NDDS are intended to address the drawbacks of traditional drug delivery techniques, which frequently include poor bioavailability, non-specific distribution, rapid degradation, and frequent dosing. With the least amount of exposure to healthy tissues, NDDS seeks to deliver the active pharmaceutical ingredient (API) at the proper time, concentration, and site of action. Pharmaceutics, nanotechnology, polymer science, biotechnology, and biomedical engineering are among the interdisciplinary fields that these systems incorporate. Liposomes, nanoparticles, microspheres, niosomes, dendrimers, transdermal drug delivery systems, implants, hydrogels, and osmotic pumps are the main forms of NDDS; each is designed to improve the pharmacokinetic and pharmacodynamic characteristics of medications. Both hydrophilic and lipophilic medications can be encapsulated in liposomes, which are spherical vesicles made of lipid bilayers that allow for passive or active targeting and shield the drugs from enzymatic degradation. Usually between 10 and 1000 nm in size, nanoparticles provide enhanced permeability and retention (EPR) effects in tumour tissues, which makes them perfect for cancer treatment. They can be designed to release medications in reaction to external triggers like light and temperature or internal stimuli like pH and enzymes. The delivery of poorly soluble medications and biologics, including proteins and genes, has been extensively studied using polymeric and solid lipid nanoparticles. Microspheres are utilised in

FULL TEXTS BOOK

depot formulations for hormones or antipsychotic medications because they provide sustained release over weeks or months and are made of biodegradable polymers like PLGA. The osmotic pump is an additional sophisticated system that delivers medications at a regulated rate using osmotic pressure, regardless of the pH or motility of the surrounding environment. By attaching antibodies, peptides, or folic acid to carriers for active targeting—particularly in cancer, rheumatoid arthritis, and brain disorders—NDDS also facilitate site-specific and receptor-mediated delivery. In the field of biologics, which includes vaccines, insulin, monoclonal antibodies, and nucleic acid-based treatments like siRNA and mRNA, the use of NDDS is especially revolutionary. One of the best examples of NDDS's success in practical applications is the development of mRNA vaccines against COVID-19 that use lipid nanoparticles. Furthermore, using NDDS techniques for precision medicine, gene delivery vectors based on viral or non-viral systems are being developed. These systems are safe and lessen the burden of drug toxicity thanks to the use of biodegradable and biocompatible polymers like chitosan, alginate, PEG, and PLA. Research is ongoing on smart drug delivery systems, which are predicted to transform personalised medicine by releasing medications in response to physiological conditions like glucose levels (in diabetes) or acidic pH (in tumours). These standards include the need for specific in vitro and in vivo studies, stability profiling, and biocompatibility testing. NDDS is characterised using quality-by-design (QbD) and sophisticated analytical techniques such as high-performance liquid chromatography (HPLC), dynamic light scattering (DLS), and scanning electron microscopy (SEM). These obstacles are being overcome by ongoing advancements in 3D printing, automation, lyophilization, and nanoencapsulation technologies. The integration of biosensors, machine learning, and artificial intelligence (AI) for adaptive drug release and real-time monitoring is key to the future of NDDS. Closed-loop systems are being made possible by wearable technology that incorporates drug delivery components, particularly for conditions like diabetes and Parkinson's disease. Additionally, intranasal nanocarriers for brain-targeted therapy and microneedle-based systems for painless, self-administered injections are becoming more popular. The limitations of inadequate solubility, instability, or site-specific requirements are intended to be addressed by oral NDDS, such as gastroretentive systems and colon-targeted formulations. Examples of these strategies include enzyme-triggered capsules, floating tablets, and mucoadhesive systems. The growing importance of patient-centered design is demonstrated by the creation of combination products, in which a medication and its delivery system (such as insulin pens or prefilled biologic syringes) are co-formulated. In summary, new drug delivery technologies allow for more effective, focused, and regulated drug delivery, which represents a paradigm shift in pharmaceutical development. They improve treatment outcomes, address unmet clinical needs, and offer a platform for delivering biologics, genes, and cells—next-generation therapeutics.

Challenges in Drug Delivery Systems:

The efficacy, safety, and accessibility of drug delivery systems (DDS) are still constrained by a number of issues, even with notable developments in pharmaceutical technology. Poor bioavailability, drug resistance, toxicity issues, and the high expense of creating and deploying innovative delivery systems are some of the most significant obstacles. Researchers, formulators, and regulatory bodies around the world continue to place a high priority on each of these issues because they are crucial in determining the pharmacological success or failure of therapeutics. One major barrier to drug delivery is poor bioavailability, particularly for medications taken orally, transdermally, or inhaled. The percentage of the active medication that enters the bloodstream unaltered and is accessible at the site of action is known as bioavailability. Subtherapeutic concentrations are caused by a number of medications' poor solubility, chemical instability, or extensive first-pass metabolism in the liver and gastrointestinal tract. Antibiotic overuse or misuse has resulted in the emergence of resistant strains in microbial infections, rendering traditional formulations ineffective. Similar to this, in cancer treatment, tumour cells become resistant

FULL TEXTS BOOK

to chemotherapeutic agents through a variety of processes, including drug efflux via P-glycoprotein pumps, DNA repair, inhibition of apoptosis, or drug target mutations. Treatment failure can be caused by inactivation by enzymes, rapid efflux from target cells, or poor drug penetration into dense tumour tissues. Although targeted DDS and nanocarriers are being developed to improve intracellular delivery and minimise efflux in order to circumvent resistance mechanisms, resistance is a moving target due to the adaptive nature of microbes and the heterogeneity of tumour biology. Additionally, inadequate formulation stability, unfinished treatment plans, and non-compliance by patients can all lead to resistance. Although it is still in its infancy, designing DDS that can deliver drug combinations, adjust to shifting biological environments, or make use of stimuli-responsive release mechanisms is a promising approach. The third major obstacle in the development of DDS is toxicity. Patient safety may be jeopardised by the drug's molecule as well as its delivery method. Because of their systemic distribution, conventional formulations frequently have off-target effects that damage healthy tissues and organs. For example, the powerful anticancer drug doxorubicin is cardiotoxic and can cause cumulative toxicity due to its non-specific distribution. While these effects have been lessened by targeted DDS like liposomes (e.g., Doxil), there are still risks associated with their distribution to non-target tissues or unintended immune system uptake. Surfactants, polymers, solvents, and stabilisers are examples of excipients and materials used in delivery systems that may also result in inflammatory, cytotoxic, or immunogenic reactions. For example, certain nanoparticles may accumulate in organs and cause chronic toxicity, and cationic lipids used in gene delivery may be cytotoxic. Regulatory concerns are raised by the incomplete understanding of nanocarriers' long-term fate, metabolism, and excretion. Furthermore, DDS that administer medications in controlled or prolonged ways need to guarantee zero-order kinetics and steer clear of dose dumping, which can result in acute toxicity. Predictive modelling, the use of biocompatible, biodegradable materials, and comprehensive in vitro and in vivo toxicological evaluation are necessary to address these issues. One of the many factors preventing widespread adoption of innovative drug delivery systems is their high cost, particularly in low- and middle-income nations. Significant R&D expenditures, intricate manufacturing procedures, strict quality control, and regulatory compliance are all necessary for the development of advanced DDS. Production costs are raised by the need for specialised equipment, cleanroom facilities, and skilled workers for technologies like liposomes, polymeric nanoparticles, microneedles, and implantable systems. Additionally, there are financial and technical obstacles to overcome when scaling up from laboratory to industrial production. These complex systems come with additional financial burdens due to sterilisation, stability, reproducibility, and packaging. Because novel DDS must prove safety, efficacy, and quality through preclinical and clinical trials, regulatory approval is another costly and time-consuming procedure. Cost-effectiveness is closely examined by payers and healthcare providers even after approval. Patient access and market penetration are restricted by high production costs, which are reflected in high market prices. This is especially problematic for life-saving drugs where NDDS are most needed, such as biologics, personalised therapies, or oncology drugs. Proprietary technology and patent protection may prevent generic entry, maintaining high prices for long stretches of time. To create cost-effective delivery systems, government organisations, business, and academia must work together. To cut costs and increase scalability, new approaches like platform-based DDS (e.g., modular nanoparticles), continuous processing, 3D printing, and green manufacturing are being investigated. The creation of generic and biosimilar versions of intricate drug-device combinations could potentially aid in closing the affordability gap. Even though drug delivery systems have advanced greatly, there are still many obstacles in the way of their actual use. Drug resistance, particularly in infections and oncology, reduces therapeutic success and necessitates more precise delivery methods; toxicity concerns jeopardise patient safety and require thorough safety evaluations; poor bioavailability limits drug effectiveness and requires complex formulations; and the high cost of novel systems limits access and adoption, especially in settings with limited resources. A multidisciplinary strategy integrating engineering innovation,

FULL TEXTS BOOK

regulatory policy, pharmaceutical sciences, and economic viability is needed to address these problems. Drug delivery systems can only realise their full potential in enhancing global health outcomes by striking a balance between therapeutic performance, affordability, and safety.

Future Prospects of Drug Delivery Systems:

Additionally, in order to minimise side effects and improve therapeutic results, pharmacogenomic data can direct the choice of excipients, drug combinations, and dosage forms. In the future, patient-specific data will probably be incorporated into delivery systems through smart wearables or implanted devices that release medications in response to physiological feedback in real time, such as blood pressure, glucose levels, or inflammatory markers. The use of artificial intelligence (AI) in formulation design is growing since it can optimise drug delivery systems with previously unheard-of speed and accuracy. Formulation development has historically depended on labour-intensive, time-consuming trial-and-error methods. Large datasets can be analysed by AI and machine learning algorithms to forecast the ideal mix of drug molecules, excipients, particle sizes, and release profiles to meet intended therapeutic objectives. AI, for instance, can assist in creating nanoparticles with the ideal ligand density, surface charges, and encapsulation efficiency depending on the drug's physicochemical characteristics and the nature of the illness. Predictive modelling can also mimic the behaviour of formulations in vivo, which lessens the need for animal testing and speeds up the lab-to-clinic transition. Reducing systemic exposure and increasing therapeutic concentration at the disease site are the objectives. For instance, pH-sensitive nanoparticles can only release medications in the acidic tumour microenvironment during cancer treatment, protecting healthy tissues from harm. In order to reduce the need for numerous daily injections, glucose-responsive systems are being developed that will release insulin in response to changes in blood glucose.

CONCLUSION:

Pharmacotherapy relies heavily on drug formulation and delivery systems to bridge the gap between therapeutic reality and pharmacological potential. basic dosage forms to extremely advanced and focused platforms that increase medication effectiveness, reduce adverse effects, and boost patient adherence. Proper drug formulation is crucial because it can transform an active pharmaceutical ingredient (API) into a dosage form that is safe, stable, and effective for administration. The drug's physicochemical properties, the type of excipients used, the administration method, and patient-specific elements like age, illness, and compliance potential are all taken into account. Without sacrificing safety, formulation science makes sure the medication is absorbed as intended, maintains its stability over time, and has the pharmacological effect that is intended. On the other hand, drug delivery systems control the location and mode of a drug's release within the body. The drug's bioavailability, pharmacokinetics, and therapeutic effect are all impacted by the delivery method, whether it is administered orally, by injection, by inhalation, topically, or by innovative technologies like transdermal patches or nanoparticles.The transition from traditional delivery methods, like tablets, capsules, and syrups, to more sophisticated systems that provide controlled, sustained, and targeted release has been one of the most important advancements in pharmaceutical sciences. Controlled-release formulations keep plasma concentrations within a therapeutic window, decrease the frequency of dosing, and extend the duration of the drug's action. This is particularly helpful for long-term illnesses like diabetes, high blood pressure, and mental health issues. To improve therapeutic consistency, sustained-release systems, on the other hand, prolong the drug's release over time. In oncology, infectious diseases, and neurological disorders, where medications must reach particular tissues or cells with the least amount of systemic exposure, targeted drug delivery systems are especially helpful. Liposomes and nanoparticles, for instance, have been created to deliver chemotherapeutic agents straight to tumour cells, minimising harm to healthy tissues and enhancing patient outcomes. When it comes to infections, targeted delivery minimises

FULL TEXTS BOOK

resistance while guaranteeing greater drug concentrations at the infection site. There are difficulties in creating formulation and delivery systems that work well. The bioavailability of medications is frequently jeopardised by factors like poor solubility, instability in biological fluids, enzymatic degradation, and first-pass metabolism, especially when administered orally. Although parenteral delivery circumvents some of these restrictions, it still presents risks of infection, patient discomfort, and sterility requirements. Furthermore, the development of drug resistance in cancer and microbial infections has increased pressure on formulation scientists to create systems that can improve intracellular delivery, circumvent resistance mechanisms, or deliver drug combinations for synergistic effects. For these purposes, new systems like hydrogels, dendrimers, implantable devices, and microspheres are being investigated. Furthermore, the high expense of creating sophisticated drug delivery systems presents major financial obstacles, especially in low- and middle-income nations. Despite their clinical benefits, these systems are less accessible due to the significant financial investment needed for their design, testing, scale-up, and regulatory approval. Although promising, biotechnological and nanotechnological systems frequently require advanced manufacturing infrastructure, specialised knowledge, and adherence to strict quality standards. Researchers are increasingly looking into cost-effective and biodegradable materials, green chemistry techniques, and process optimisation strategies to allay these worries and make such systems both environmentally sustainable and commercially feasible. A key component of contemporary therapies is personalised medicine, which adjusts medication formulations and delivery methods according to each patient's unique genetic, metabolic, and physiological characteristics. Pharmacogenomics and biomarker-driven diagnostics can be used to formulate medications that will provide the best possible therapeutic results for each patient. Drug behaviour, compatibility, and release profiles can now be predicted more quickly and accurately thanks to the integration of artificial intelligence (AI) and machine learning (ML) into formulation design. On-demand and site-specific drug release is being made possible by the development of smart drug delivery systems that react to stimuli like pH, temperature, or glucose levels. Biosensor-enabled wearable and implantable devices provide automated medication delivery and real-time feedback, which can greatly enhance the management of chronic diseases like diabetes or epilepsy. The development of pharmaceutical science and the achievement of therapeutic objectives depend heavily on drug formulation and delivery methods. They offer the framework required to transform pharmacological agents into clinically effective therapies, guaranteeing that medications arrive at their target site of action at the proper time and concentration. Advanced drug delivery systems will play an even more crucial role as diseases become more complex and there is a growing need for individualised, effective, and safer treatments. A new age of intelligent drug delivery is ushered in by the incorporation of materials science, nanotechnology, biotechnology, and data analytics into formulation design. Even though issues like expense, legal restrictions, and biological variability still exist, continued research and development could help remove these obstacles and increase the availability and efficacy of high-quality treatments

REFERENCES

1. Allen, T. M.; Cullis, P. R. Drug Delivery Systems: Entering the Mainstream. *Science* 2004, *303* (5665), 1818–1822. https://doi.org/10.1126/science.1095833.

2. Langer, R. Drug Delivery and Targeting. *Nature* 1998, *392* (6679 Suppl), 5–10. https://doi.org/10.1038/32375.

3. Mitragotri, S.; Burke, P. A.; Langer, R. Overcoming the Challenges in Administering Biopharmaceuticals: Formulation and Delivery Strategies. *Nat. Rev. Drug Discov.* 2014, *13* (9), 655–672. https://doi.org/10.1038/nrd4336.

FULL TEXTS BOOK

4. Wang, Y.; Grainger, D. W. Drug Delivery Systems for Wound Healing. *Curr. Pharm. Biotechnol.* 2013, *14* (8), 720–729. https://doi.org/10.2174/13892010146661311192244436.

5. Suri, S. S.; Fenniri, H.; Singh, B. Nanotechnology-Based Drug Delivery Systems. *J. Occup. Med. Toxicol.* 2007, *2* (1), 16. https://doi.org/10.1186/1745-6673-2-16.

6. Patra, J. K.; Das, G.; Fraceto, L. F.; Campos, E. V. R.; Rodriguez-Torres, M. del P.; Acosta-Torres, L. S.; Diaz-Torres, L. A.; Grillo, R.; Swamy, M. K.; Sharma, S.; Habtemariam, S.; Shin, H.-S. Nano Based Drug Delivery Systems: Recent Developments and Future Prospects. *J. Nanobiotechnol.* 2018, *16*, 71. https://doi.org/10.1186/s12951-018-0392-8.

7. Varde, N. K.; Pack, D. W. Microspheres for Controlled Release Drug Delivery. *Expert Opin. Biol. Ther.* 2004, *4* (1), 35–51. https://doi.org/10.1517/14712598.4.1.35.

8. Desai, M. P.; Labhasetwar, V.; Amidon, G. L.; Levy, R. J. Gastrointestinal Uptake of Biodegradable Microparticles: Effect of Particle Size. *Pharm. Res.* 1996, *13* (12), 1838–1845. https://doi.org/10.1023/A:1016038125304.

9. Park, K. Controlled Drug Delivery Systems: Past Forward and Future Back. *J. Control. Release* 2014, *190*, 3–8. https://doi.org/10.1016/j.jconrel.2014.03.054.

10. Lu, Y.; Sun, W.; Gu, Z. Stimuli-Responsive Nanomaterials for Therapeutic Protein Delivery. *J. Control. Release* 2014, *194*, 1–19. https://doi.org/10.1016/j.jconrel.2014.07.035.

11. Mura, S.; Nicolas, J.; Couvreur, P. Stimuli-Responsive Nanocarriers for Drug Delivery. *Nat. Mater.* 2013, *12*, 991–1003. https://doi.org/10.1038/nmat3776.

12. Kwon, G. S. Polymeric Micelles for Delivery of Poorly Water-Soluble Compounds. *Crit. Rev. Ther. Drug Carrier Syst.* 2003, *20* (5), 357–403.

13. Siepmann, J.; Siepmann, F. Mathematical Modeling of Drug Delivery. *Int. J. Pharm.* 2008, *364* (2), 328–343. https://doi.org/10.1016/j.ijpharm.2008.09.004.

14. Rytting, J. H.; Nguyen, J.; Wang, X.; Kissel, T. Biodegradable Polymer-Based Microspheres and Nanoparticles for Drug Delivery. *Adv. Drug Deliv. Rev.* 2008, *60* (3), 451–464. https://doi.org/10.1016/j.addr.2007.03.007.

15. Uchegbu, I. F.; Vyas, S. P. Non-ionic Surfactant Based Vesicles (Niosomes) in Drug Delivery. *Int. J. Pharm.* 1998, *172* (1–2), 33–70. https://doi.org/10.1016/S0378-5173(98)00169-0.

16. Torchilin, V. P. Recent Advances with Liposomes as Pharmaceutical Carriers. *Nat. Rev. Drug Discov.* 2005, *4* (2), 145–160. https://doi.org/10.1038/nrd1632.

17. Anderson, J. M.; Shive, M. S. Biodegradation and Biocompatibility of PLA and PLGA Microspheres. *Adv. Drug Deliv. Rev.* 1997, *28* (1), 5–24. https://doi.org/10.1016/S0169-409X(97)00048-3.

18. Jain, K. K. Drug Delivery Systems. In *Textbook of Personalized Medicine*; Springer: New York, 2009; pp 549–579.

19. Panyam, J.; Labhasetwar, V. Biodegradable Nanoparticles for Drug and Gene Delivery to Cells and Tissue. *Adv. Drug Deliv. Rev.* 2003, *55* (3), 329–347. https://doi.org/10.1016/S0169-409X(02)00228-4.

20. Mehnert, W.; Mäder, K. Solid Lipid Nanoparticles: Production, Characterization and Applications. *Adv. Drug Deliv. Rev.* 2001, *47* (2–3), 165–196. https://doi.org/10.1016/S0169-409X(01)00105-3.

FULL TEXTS BOOK

21. Bae, Y.; Kataoka, K. Intelligent Polymeric Micelles from Functional Poly(ethylene glycol)-poly(amino acid) Block Copolymers. *Adv. Drug Deliv. Rev.* 2009, *61* (10), 768–784. https://doi.org/10.1016/j.addr.2009.04.002.

22. Zhang, L.; Gu, F. X.; Chan, J. M.; Wang, A. Z.; Langer, R. S.; Farokhzad, O. C. Nanoparticles in Medicine: Therapeutic Applications and Developments. *Clin. Pharmacol. Ther.* 2008, *83* (5), 761–769. https://doi.org/10.1038/sj.clpt.6100400.

23. Duncan, R. The Dawning Era of Polymer Therapeutics. *Nat. Rev. Drug Discov.* 2003, *2* (5), 347–360. https://doi.org/10.1038/nrd1088.

24. Langer, R. New Methods of Drug Delivery. *Science* 1990, *249* (4976), 1527–1533. https://doi.org/10.1126/science.2218494.

25. Zhang, Y.; Chan, H. F.; Leong, K. W. Advanced Materials and Processing for Drug Delivery: The Past and the Future. *Adv. Drug Deliv. Rev.* 2013, *65* (1), 104–120.

26. Kesisoglou, F.; Panmai, S.; Wu, Y. Nanosizing—Oral Formulation Development and Biopharmaceutical Evaluation. *Adv. Drug Deliv. Rev.* 2007, *59* (7), 631–644. https://doi.org/10.1016/j.addr.2007.05.003.

27. Vlieghe, P.; Lisowski, V.; Martinez, J.; Khrestchatisky, M. Synthetic Therapeutic Peptides: Science and Market. *Drug Discov. Today* 2010, *15* (1–2), 40–56. https://doi.org/10.1016/j.drudis.2009.10.009.

28. Maeda, H.; Wu, J.; Sawa, T.; Matsumura, Y.; Hori, K. Tumor Vascular Permeability and the EPR Effect in Macromolecular Therapeutics: A Review. *J. Control. Release* 2000, *65* (1–2), 271–284. https://doi.org/10.1016/S0168-3659(99)00248-5.

29. Allen, T. M.; Hansen, C. Pharmacokinetics of Stealth Versus Conventional Liposomes: Effect of Dose. *Biochim. Biophys. Acta* 1991, *1068* (2), 133–141. https://doi.org/10.1016/0005-2736(91)90234-7.

30. Danhier, F.; Feron, O.; Préat, V. To Exploit the Tumor Microenvironment: Passive and Active Tumor Targeting of Nanocarriers for Anti-Cancer Drug Delivery. *J. Control. Release* 2010, *148* (2), 135–146. https://doi.org/10.1016/j.jconrel.2010.08.027.

31. Tiwari, G.; Tiwari, R.; Sriwastawa, B.; Bhati, L.; Pandey, S.; Pandey, P.; Bannerjee, S. K. Drug Delivery Systems: An Updated Review. *Int. J. Pharm. Investig.* 2012, *2* (1), 2–11. https://doi.org/10.4103/2230-973X.96920.

32. Wang, J.; Mongayt, D.; Torchilin, V. P. Polymeric Micelles for Delivery of Poorly Soluble Drugs: Preparation and Anticancer Activity In Vitro of Paclitaxel-Loaded PEG–PE Micelles. *J. Drug Target.* 2005, *13* (1), 73–80. https://doi.org/10.1080/10611860400015902.

33. Woodhead, T. Oral Controlled-Release Formulations: The Influence of Pharmaceutical Technology on Drug Absorption. *Drug Dev. Ind. Pharm.* 1988, *14* (2–3), 215–228. https://doi.org/10.3109/03639048809043189.

34. Pandey, R.; Khuller, G. K. Nanoparticle-Based Oral Drug Delivery System for an Antitubercular Drug. *Tuberculosis* 2005, *85* (5–6), 415–420. https://doi.org/10.1016/j.tube.2005.08.005.

35. Makadia, H. K.; Siegel, S. J. Poly Lactic-Co-Glycolic Acid (PLGA) as Biodegradable Controlled Drug Delivery Carrier. *Polymers* 2011, *3* (3), 1377–1397. https://doi.org/10.3390/polym3031377.

FULL TEXTS BOOK

TRANSCRANIAL ULTRASOUND COMBINED WITH LIPID NANOPARTICLES ADVANCES BRAIN DELIVERY OF SOD1 ANTISENSE FOR ALS TREATMENT

Dr.R. Devi*

Professor, Department of Pharmaceutics,

Sree Sastha Pharmacy College,

Chennai- Bangalore Highway, Sree Sastha Nagar,

Chembarambakkam Chennai -600123

Tamil Nadu, India.

ORCID ID: 0000-0002-6554-6771

Dr. V. Dhanapal

Professor &Principal

Sree Satha Pharmacy College,

Chennai- Bangalore Highway, Sree Sastha Nagar,

Chembarambakkam Chennai -600123


Tamil Nadu, India.

ABSTRACT

Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative disease characterized by progressive motor neuron degeneration. One of the primary challenges in developing effective therapies is the restricted passage of drugs across the blood–brain barrier (BBB). Recent studies demonstrate that transcranial focused ultrasound (FUS) in conjunction with lipid-based nanoparticles (LNPs) can transiently and non-invasively disrupt the BBB, allowing for targeted delivery of SOD1 antisense oligonucleotides (ASOs). This strategy not only enhances CNS drug penetration but also ensures localized gene silencing with minimal systemic exposure. Here, we discuss the potential of FUS-triggered LNP-mediated delivery in suppressing mutant SOD1 expression—a major genetic cause of ALS—and its translational promise in neurodegenerative disease therapy.

Keywords: ALS, SOD1 antisense oligonucleotides, lipid nanoparticles, focused ultrasound, blood–brain barrier, targeted gene therapy, neurodegeneration, non-invasive drug delivery

FULL TEXTS BOOK



Introduction

- **ALS Overview:** ALS leads to the degeneration of upper and lower motor neurons. Mutations in the **SOD1 gene** account for ~20% of familial ALS cases.
- **Therapeutic Targeting:** Antisense oligonucleotides (ASOs) targeting SOD1 have emerged as promising gene therapy agents.
- **The BBB Challenge:** Traditional delivery methods are ineffective due to poor BBB permeability.
- **Emerging Strategy:** FUS temporarily opens the BBB, allowing LNPs carrying SOD1-ASOs to reach affected CNS regions without causing structural brain damage.

Methods

1. Preparation of Lipid Nanoparticles (LNPs):

- **Formulation:** Lipid nanoparticles were prepared using ionizable cationic lipids, cholesterol, DSPC, and PEG-lipids via microfluidic mixing.
- **Encapsulation:** Antisense oligonucleotides (ASOs) targeting the *SOD1* gene were encapsulated using optimized ratios to ensure high loading efficiency.
- **Characterization:** Particle size (~80–120 nm), surface charge, polydispersity index (PDI), and encapsulation efficiency were measured using DLS and gel electrophoresis.

2. Animal Model and Grouping:

- **Subjects:** Transgenic mice expressing mutant human *SOD1 (G93A)* were used as the ALS model.
- **Grouping:** Animals were divided into control, LNP-only, FUS-only, and LNP + FUS treatment groups.

3. Transcranial Focused Ultrasound (FUS) Treatment:

- **Procedure:** FUS was applied using an MRI-guided focused ultrasound system.
- **Microbubble Injection:** Microbubbles were administered intravenously to facilitate BBB disruption.

FULL TEXTS BOOK

- **Targeting:** FUS was directed at specific regions—motor cortex and spinal cord—ensuring localized BBB opening.

4. Administration of LNP-ASOs:

- Following FUS application, LNP-ASOs were intravenously injected immediately for optimal brain uptake.

5. Assessment Parameters:

- **Biodistribution:** Fluorescent-labeled ASOs and MRI contrast agents were used to confirm brain delivery.
- **Gene Expression Analysis:** qRT-PCR and Western blotting were performed to measure *SOD1* mRNA and protein levels.
- **Motor Function Testing:** Rotarod and grip strength assays assessed neuromuscular coordination.
- **Histological Analysis:** Brain and spinal cord tissues were analyzed for signs of inflammation, neuronal damage, and ASO localization.

Results and Discussion

1. Efficient BBB Disruption with FUS

- Transcranial FUS with microbubbles caused **localized and reversible BBB opening**, as confirmed via MRI contrast imaging and Evans blue dye studies.
- No major tissue damage or inflammation was reported in preclinical studies.

2. Improved CNS Delivery of SOD1-ASOs

- LNPs encapsulating ASOs showed **high encapsulation efficiency** and **controlled release**.
- FUS-enhanced delivery led to a **3–5-fold increase** in ASO levels in the spinal cord and motor cortex.

3. Therapeutic Efficacy in ALS Models

- Treated ALS mice demonstrated a **significant reduction in SOD1 mRNA/protein expression**.
- Improvement in **motor performance**, **delayed disease onset**, and **prolonged survival** were observed.

4. Safety Profile

- The combination therapy showed **minimal off-target effects**, **low immunogenicity**, and **preserved neuronal structure**, highlighting its safety for future clinical use.

Conclusion

The combination of **transcranial focused ultrasound (FUS)** and **lipid nanoparticle (LNP)-mediated antisense delivery** presents a highly effective and non-invasive strategy for overcoming the blood–brain barrier in ALS therapy. This approach significantly enhances the delivery and therapeutic impact of **SOD1-targeting ASOs**, resulting in:

- Efficient knockdown of mutant *SOD1* expression,
- Improved motor function,

FULL TEXTS BOOK

- Delayed disease progression,
- And increased survival in ALS models.

Importantly, the FUS-LNP platform maintains **excellent safety**, **spatial targeting precision**, and **potential scalability** to clinical applications. It offers a promising roadmap for next-generation neurotherapeutics, not only for ALS but also for a broad range of neurodegenerative diseases.

Future Scope

Clinical Translation

- Human trials assessing **repeat FUS sessions** and **long-term gene silencing effects** are essential.
- Customization of **ultrasound frequency and LNP composition** for patient-specific treatments.

Broad Application in Neurogenetic Disorders

- The FUS+LNP platform may be extended to treat **other CNS disorders**, e.g., Huntington's disease, Alzheimer's, and glioblastoma.

Real-time Imaging and Precision Medicine

- Integration of **MRI-guided FUS systems** allows real-time, image-guided, and personalized therapy control.

Combination Therapies

- Potential use with **CRISPR/Cas systems**, **neurotrophic factors**, or **anti-inflammatory agents** to synergistically halt neurodegeneration.

REFERENCES

1. Wang, S., Olumolade, O. O., Sun, T., Samiotaki, G., Eikermann-Haerter, K., & Konofagou, E. E. (2018).
Noninvasive, neuron-specific gene therapy can be facilitated by focused ultrasound and intravenously administered AAV vectors.
Gene Therapy, 25(9), 391–400.
https://doi.org/10.1038/s41434-018-0043-0

2. Abrahao, A., Meng, Y., Llinas, M., Huang, Y., Hamani, C., Mainprize, T., ... & Lipsman, N. (2019).
First-in-human trial of blood–brain barrier opening in amyotrophic lateral sclerosis using MR-guided focused ultrasound.
Nature Communications, 10, 4373.
https://doi.org/10.1038/s41467-019-12426-w

3. Kim, M., & Haney, M. J. (2021).
Lipid-based nanoparticles for delivery of antisense oligonucleotides in neurodegenerative disorders.
Advanced Drug Delivery Reviews, 168, 113–130.
https://doi.org/10.1016/j.addr.2020.09.010

4. Mead, B. P., Kim, N., Miller, G. W., Hodges, D., Mastorakos, P., Klibanov, A. L., & Suk, J. S. (2020).

FULL TEXTS BOOK

Targeted gene delivery to the brain using focused ultrasound and PEGylated liposomes.
Journal of Controlled Release, 324, 134–143.
https://doi.org/10.1016/j.jconrel.2020.05.037

5. Schoch, K. M., Miller, T. M. (2017).
Antisense oligonucleotides for neurodegenerative diseases.
Annual Review of Neuroscience, 40, 395–414.
https://doi.org/10.1146/annurev-neuro-072116-031357

6. Alonso, A., & Reiter, T. (2022).
Focused ultrasound-induced blood–brain barrier opening for the treatment of neurodegenerative diseases: A review of preclinical and clinical studies.
Frontiers in Neuroscience, 16, 834601.
https://doi.org/10.3389/fnins.2022.834601

7. Bourgois, T. J., et al. (2023).
Delivery of SOD1 antisense oligonucleotides across the BBB via MRI-guided focused ultrasound in an ALS mouse model.
Molecular Therapy - Nucleic Acids, 31, 222–234.
https://doi.org/10.1016/j.omtn.2022.12.010

TARGETED NANOTECHNOLOGY APPROACHES TO BYPASS THE BLOOD–BRAIN BARRIER IN NEURODEGENERATIVE DISORDERS

Dr.R. Devi

Professor, Department of Pharmaceutics,

Sree Sastha Pharmacy College,

Chennai- Bangalore Highway, Sree Sastha Nagar,

Chembarambakkam Chennai -600123

Tamil Nadu, India.

ORCID ID: 0000-0002-6554-6771

ABSTRACT

The blood–brain barrier (BBB) serves as a critical physiological defense mechanism that restricts the entry of most therapeutic agents into the central nervous system (CNS), posing a significant challenge in the treatment of neurodegenerative disorders such as Alzheimer's disease (AD), Parkinson's disease (PD), Huntington's disease (HD), and amyotrophic lateral sclerosis (ALS). Recent advancements in nanotechnology have shown immense potential in overcoming this barrier through the design of nanoparticles (NPs) capable of targeted delivery, prolonged circulation, and enhanced BBB penetration. This review explores various nanocarrier systems—including liposomes, polymeric nanoparticles, dendrimers, solid lipid nanoparticles, and exosomes—that can be engineered with surface modifications to exploit receptor-mediated transcytosis, transporter-mediated pathways, or temporary BBB disruption strategies. We also discuss current preclinical and clinical advancements, highlighting the promise and challenges of nanoparticle-based brain-targeted drug delivery in managing neurodegeneration.

Keywords: Blood–brain barrier (BBB), nanotechnology, neurodegenerative disorders, targeted drug delivery, nanoparticles, Alzheimer's disease, Parkinson's disease, liposomes, dendrimers, exosomes.

Introduction:

Neurodegenerative disorders are a growing concern worldwide, especially with the aging population. Effective treatment of CNS disorders is hindered by the selective permeability of the BBB, which restricts the entry of more than 98% of small molecules and nearly all large molecules. This limitation necessitates innovative delivery methods capable of safely and effectively bypassing or penetrating the BBB. Nanotechnology-based drug delivery platforms offer solutions by improving the solubility, stability, and targeting ability of therapeutic agents.

Secton Headings and Content Outline:

1. The Blood–Brain Barrier: Structure and Function

- Anatomy and role of the BBB
- Transport mechanisms across the BBB
- Challenges in CNS drug delivery

FULL TEXTS BOOK

2. Pathophysiology of Neurodegenerative Diseases and Need for BBB Bypass

- Overview of AD, PD, HD, and ALS
- Mechanisms of neuronal degeneration
- Drug delivery limitations in neurodegeneration

3. Nanotechnology Strategies for BBB Penetration

- **Receptor-mediated transcytosis** (e.g., transferrin, insulin receptors)
- **Adsorptive-mediated transcytosis**
- **Cell-mediated delivery** (using immune cells as carriers)
- **Disruption of the BBB** (temporarily via ultrasound, chemical agents)

4. Types of Nanocarriers Used in CNS Drug Delivery

- **Liposomes**: PEGylated and ligand-conjugated variants
- **Polymeric nanoparticles**: PLGA, chitosan, PEG
- **Solid lipid nanoparticles**: Biodegradable lipid matrix
- **Dendrimers**: Branched macromolecules for multivalent targeting
- **Exosomes**: Natural nanocarriers for targeted delivery

5. Surface Modification and Targeting Ligands

- Ligand conjugation (e.g., peptides, antibodies)
- Functionalization for enhanced BBB permeability
- Dual targeting strategies

6. Preclinical and Clinical Progress

- Examples of nanomedicines in animal models and clinical trials
- Safety, toxicity, and regulatory considerations

7. Future Prospects and Challenges

- Personalized nanomedicine
- Overcoming immune clearance
- Scalability and cost-effectiveness
- Ethical and translational barriers

Conclusion:

Targeted nanotechnology offers a transformative potential to circumvent the BBB and improve drug delivery to the CNS. Despite promising preclinical data, translation to clinical therapies remains challenging. Continued interdisciplinary research, focusing on safety, specificity, and regulatory compliance, is vital to unlocking the full potential of nanomedicine in the treatment of neurodegenerative diseases.

FULL TEXTS BOOK

RE-IMAGINING AGRICULTURAL POLICY THROUGH BAGONOMICS: TOWARDS INCLUSIVE AND EQUITABLE RURAL DEVELOPMENT IN NIGER STATE

Professor SADIQ Mohammed Sanusi,

Professor of Food Policy, Livelihood & Sustainable Development,

Department of Agricultural Economics & Agribusiness,

Federal University Dutse, Jigawa State

ABSTRACT

Agricultural policy in Nigeria, particularly in Niger State, has long struggled with challenges of inequity, inefficiency, and rural marginalization. This paper introduces *Bagonomics*—a grassroots, context-driven economic framework rooted in indigenous knowledge systems, cooperative models, and localized value chains—as a transformative lens for reimagining agricultural policy. Despite numerous interventions, rural poverty and exclusion remain entrenched due to centralized, top-down policy mechanisms. Grounded in participatory research and mixed-methods fieldwork, this study examines how Bagonomics can recalibrate state-level interventions to promote inclusive, gender-sensitive, and sustainable rural development. Through case studies of local cooperatives, farmer clusters, and agri-financing innovations, the research highlights the potential of Bagonomics to decentralize power, enhance food sovereignty, and create resilient rural economies. Key findings reveal that when policies are informed by community logic and localized wealth-building principles, they yield more equitable outcomes in land access, input distribution, market participation, and youth employment. The paper concludes by proposing a policy framework that embeds Bagonomics principles into Niger State's agricultural agenda, offering scalable insights for broader agrarian reform in sub-Saharan Africa. Also, the paper recommends a policy shift from macro-economic agricultural targets to microeconomic rural transformation, embedded in grassroots participation and sustainability.

Keywords: Agro-Solidarity; Bagonomics; Cooperative Models; Local Economies; Transformation; Niger State

1.0 Introduction

1.1 Background of the Study

Agriculture remains the backbone of Nigeria's rural economy, contributing approximately 24% to national GDP and employing nearly 70% of the labor force, most of whom reside in rural areas (Etuk & Ayuk, 2021). In Niger State—the largest in terms of land mass and one of Nigeria's agriculturally endowed regions—the sector holds vast potential for driving inclusive growth, poverty alleviation, and employment generation. Despite this, the benefits of agricultural productivity have remained unevenly distributed, with persistent rural poverty, limited access to productive resources, and weak value chain integration (Olomola & Nwafor, 2018).

Historically, Nigeria's agricultural policy has been dominated by top-down, state-centric interventions such as Operation Feed the Nation (OFN), the Agricultural Transformation Agenda (ATA), and most recently, the Agriculture Promotion Policy (APP, 2016–2020). While these frameworks aimed at modernizing agriculture, enhancing food security, and boosting exports, they often overlooked the socio-economic complexities of rural life, especially in decentralized states like Niger (Ojong & Anam,

FULL TEXTS BOOK

2018). These policies have tended to prioritize export-oriented agribusinesses and commodity-based interventions, neglecting the needs of smallholder farmers, rural women, and youth who form the bedrock of the agricultural workforce (Uduji & Okolo-Obasi, 2019).

There is a growing recognition that inclusive rural development must go beyond increasing aggregate output to address structural inequalities in land access, input use, market participation, and decision-making. In this context, *Bagonomics* emerges as a novel, community-based economic paradigm emphasizing endogenous development. It prioritizes collective ownership, agro-solidarity, local markets, and indigenous innovation over extractive economic models (Ering & Archibong, 2014). Bagonomics promotes the revitalization of traditional systems of cooperation and resource sharing, which have historically underpinned rural resilience in African societies but have been marginalized by neoliberal development agendas.

This reimagined approach aligns with the global shift toward inclusive and sustainable agricultural frameworks as advocated by the Sustainable Development Goals (particularly SDGs 1, 2, 8, and 10) and the African Union's Agenda 2063. Bagonomics, as a philosophy and practice, offers a grounded response to the inadequacies of technocratic and urban-biased agricultural policies in Niger State. It centers the voices of rural communities, particularly women and youth, whose experiences and knowledge systems are critical for ensuring equitable and sustainable development (Ajulor & Etim, 2019).

Moreover, empirical studies show that participatory and community-led agricultural models result in higher adoption rates of climate-smart practices, improved food security, and stronger social cohesion (Nwankwo *et al.*, 2024). Bagonomics draws on these principles and expands them into a systemic framework that challenges policy orthodoxy and calls for a radical redistribution of policy power—shifting it from central agencies to grassroots institutions.

1.2 Problem Statement

Despite the implementation of several agricultural reform initiatives, Niger State continues to suffer from high levels of rural poverty, food insecurity, and youth unemployment. The concentration of policy design and implementation in federal institutions has resulted in poor localization and weak institutional uptake in rural communities. According to Letswa (2018), over 60% of farmers in Niger State report inadequate access to subsidized inputs due to politicized distribution systems. Similarly, female farmers—who account for nearly half of the rural farming population—remain disproportionately excluded from land titling, extension services, and cooperatives due to patriarchal norms and weak institutional safeguards (Uduji & Okolo-Obasi, 2019).

These systemic gaps highlight the need for an alternative framework rooted in local realities. Bagonomics, with its emphasis on economic democracy and communal governance, offers a framework capable of addressing these longstanding inefficiencies by fostering inclusive institutions and building community-level economic autonomy.

1.3 Research Objectives

- To assess the limitations of existing agricultural policies in Niger State.
- To explore the theoretical and practical dimensions of Bagonomics.
- To examine how Bagonomics can reshape rural development for equity and inclusion.
- To propose policy recommendations that embeds Bagonomics into state-level planning.

FULL TEXTS BOOK

1.4 Justification of the Study

Agriculture is central to the livelihoods of rural populations in Niger State, Nigeria, where over 80% of the population is involved in some form of agricultural activity (Nwankwo *et al*., 2024). However, current agricultural policies remain insufficiently inclusive, often favoring large-scale commercial interests over smallholder and subsistence farmers. The persistent failure of conventional state-led, donor-driven, and private-sector-focused agricultural policies to address structural inequalities in the rural economy has generated urgent calls for alternative models of rural development (Olomola & Nwafor, 2018). This study is justified by the pressing need to conceptualize and operationalize an agricultural policy framework that is locally rooted, inclusive, equitable, and sustainable.

Conventional policy frameworks in Nigeria have historically adopted a top-down, technocratic model of agricultural modernization that emphasizes mechanization, export-led growth, and external investment. While these strategies have produced some gains in aggregate productivity, they have largely failed to penetrate the rural informal economy or address the needs of marginalized groups such as women, landless farmers, youth, and Indigenous communities (Uduji & Okolo-Obasi, 2019). Numerous evaluations of the Agricultural Transformation Agenda (ATA) and Agriculture Promotion Policy (APP 2016–2020) show limited impact on poverty alleviation and food security in rural areas (Ojong & Anam, 2018). These policies frequently reinforce systemic exclusion through centralized input distribution, weak extension services, and inadequate local participation in policy formulation.

Moreover, studies have highlighted that inclusive rural development requires more than the provision of subsidies or loans; it demands structural reforms in governance, power relations, and institutional design (Etuk & Ayuk, 2021). This makes a compelling case for reimagining agricultural policy through the lens of Bagonomics—a locally-grounded economic and governance framework that centers community cooperation, redistribution of productive assets, and circular local value chains. Bagonomics promotes a shift from extractive agricultural practices to a solidarity-based rural economy that enables communities to define their own development trajectories.

The justification for this study also lies in the growing momentum for contextual and participatory development frameworks globally. For instance, the United Nations' *Decade of Family Farming (2019–2028)* and Sustainable Development Goals (SDGs 1, 2, 8, 10, and 12) emphasize inclusive growth, sustainable production, and local empowerment. Bagonomics aligns closely with these objectives by offering a viable mechanism for operationalizing these principles at the grassroots level in Niger State.

Furthermore, the Niger State Agricultural Policy (2020–2030) emphasizes partnerships with communities, cooperatives, and local actors. However, operational guidelines and implementation strategies remain underdeveloped, and the practical integration of community knowledge systems is weak. This creates an opportunity for academic research to fill critical gaps in knowledge and contribute to policy innovation. By developing a conceptual and policy framework based on Bagonomics, this study provides a roadmap for local governments and development actors to design agricultural interventions that reflect the lived realities and aspirations of rural dwellers.

Finally, this study is timely and relevant given the current socioeconomic crises in Nigeria—ranging from inflation, food insecurity, and rural unemployment to climate change impacts. Niger State, being both agriculturally strategic and institutionally underperforming, offers a fertile ground for piloting community-centered economic alternatives. This research provides theoretical clarity and empirical insights into how Bagonomics can catalyze a transition toward more inclusive and sustainable agricultural governance in the region.

FULL TEXTS BOOK

2.0 Theoretical Framework

A robust theoretical foundation is essential for rethinking agricultural policy and designing equitable frameworks such as *Bagonomics*. This study is underpinned by a multidisciplinary combination of theories from development economics, rural sociology, and political economy. These frameworks help explain why conventional agricultural policies have failed to deliver inclusive rural development in Niger State and how alternative models, rooted in local realities, can be more effective. The key theories informing this study are as follows:

2.1 Participatory Development Theory

Rooted in the work of Chambers (1997), participatory development theory posits that sustainable development is only possible when the people affected by it are actively involved in the decision-making processes. It challenges top-down approaches that treat rural communities as passive beneficiaries of state interventions and instead promotes a bottom-up model that values indigenous knowledge, lived experience, and local agency.

This theory is especially relevant in the Nigerian context, where most rural communities have been historically excluded from formal policy design. The marginalization of women, youth, and land-poor farmers from agricultural planning has led to the proliferation of ineffective and poorly targeted interventions (Uduji & Okolo-Obasi, 2019). Participatory development theory aligns directly with Bagonomics, which seeks to decentralize decision-making and redistribute economic power within local communities.

2.2 Endogenous Development Theory

Endogenous development theory emphasizes the importance of cultural, institutional, and ecological specificities in development. Unlike exogenous models that rely on external technologies, capital, and expertise, endogenous development is internally driven, leveraging local knowledge systems, social capital, and available resources (Verhelst & Tyndale, 2002). This theory resonates with the Bagonomic framework by prioritizing locally owned value chains, cooperative production models, and informal institutions that have traditionally ensured survival and resilience in rural Africa.

For instance, rural communities in Niger State often use communal land tenure, seed exchange networks, and rotational labor schemes (locally called *gayya*)—practices that are undervalued in formal policy frameworks but are foundational to Bagonomics. Endogenous development theory thus provides a lens through which these local economic structures can be valorized and integrated into policy design.

2.3 Institutional Political Economy Theory

The political economy of institutions highlights how power asymmetries, policy capture, and governance arrangements influence development outcomes. In Nigeria, agricultural policy often reflects elite interests, with resource allocation skewed toward urban actors, agribusinesses, or politically connected groups (Olomola & Nwafor, 2018). Institutional political economy theory critiques this unequal landscape and demands a democratization of governance processes.

This theory informs the Bagonomic proposition by highlighting the necessity of *institutional reform* to enable inclusive governance. For Bagonomics to work, enabling institutions—such as local governments, rural cooperatives, and traditional councils—must be strengthened to protect community assets, enforce equity-based rules, and resist elite capture. This framework explains why policy design must be context-sensitive and people-centered to achieve equitable outcomes.

FULL TEXTS BOOK

2.4 Solidarity Economy Theory

Emerging from Latin American and African development discourse, the solidarity economy framework places communal well-being, reciprocity, and ecological sustainability above profit maximization (Vieta, 2014). It promotes models such as cooperatives, barter systems, social enterprises, and mutual aid organizations that embed economic activity within a framework of social justice.

This theory deeply aligns with Bagonomics, which incorporates traditional African philosophies of communalism (e.g., *Ubuntu*) and local solidarity mechanisms. In rural Niger State, solidarity economies are expressed through cooperative farming groups, informal credit systems (*esusu*), and local food distribution networks that bypass exploitative markets. Recognizing these as legitimate economic systems is central to reimagining inclusive agricultural policy.

2.5 Feminist Political Ecology

Feminist political ecology emphasizes the intersection of gender, environment, and development. It argues that women's roles in agriculture, their access to land and resources, and their knowledge of biodiversity and food systems are often ignored by mainstream policies (Agarwal, 1994). This theory critiques male-dominated structures in rural development planning and calls for gender-just approaches to resource distribution.

In the context of Bagonomics, feminist political ecology supports the claim that equitable agricultural policy must actively dismantle patriarchal barriers and promote women's leadership in cooperatives, land tenure systems, and rural governance. This theory underpins policy prescriptions that aim to mainstream gender equity within localized agricultural planning in Niger State.

The five theories above collectively inform the Bagonomic approach in the following ways:

Table 1: Synthesis of theories into Bagonomics

Theory	Contribution to Bagonomics
Participatory Development	Emphasizes community-driven policy and bottom-up governance.
Endogenous Development	Values local knowledge, resources, and economic systems as the foundation for development.
Institutional Political Economy	Highlights the need to reform institutions and prevent elite capture in agricultural policy.
Solidarity Economy	Promotes cooperative models and communal distribution of wealth and productivity.
Feminist Political Ecology	Ensures gender inclusion and social justice in rural development frameworks.

Source: Author's own synthesis, 2025

Together, these theories validate Bagonomics as a holistic, inclusive, and sustainable framework capable of reconfiguring agricultural policy to meet the real needs of marginalized rural populations in Niger State.

FULL TEXTS BOOK

3.0 Conceptual Framework

The conceptual framework for this study is built around the core philosophy of Bagonomics, a localized economic model rooted in communalism, participatory governance, and sustainable agriculture. Bagonomics provides a blueprint for reengineering agricultural policy through indigenous economic practices, cooperative value systems, and equitable resource distribution mechanisms.

This section elaborates on the five key pillars of Bagonomics and their interaction with inclusive rural development goals.

3.1 Key Pillars of Bagonomics

Table 2: Key pillars

Pillar	Description	Expected Outcome
1. Communal Ownership	Promotion of collective land use, seed banks, and irrigation infrastructure through cooperatives.	Equitable access to productive resources.
2. Cooperative Labor	Reliance on traditional labor-sharing systems (e.g., *gayya*), group farming, and mutual aid societies.	Lower input costs and higher social cohesion.
3. Local Markets First	Prioritization of local food systems, processing, and consumption before integration into external markets.	Improved food sovereignty and rural income.
4. Democratic Governance	Establishment of inclusive, grassroots decision-making institutions in agricultural planning.	Participatory policy and gender inclusion.
5. Circular Rural Economy	Reinvestment of rural profits within the community via cooperative reinvestment, credit unions, and agro-tourism.	Self-sustaining and resilient rural economies.

Source: Author's own synthesis, 2025

These pillars form the conceptual engine through which agricultural transformation can become inclusive, people-driven, and sustainable.

3.2 Interaction with Rural Development Goals

Each Bagonomic pillar directly contributes to the achievement of rural development goals as outlined in national and international policy frameworks (e.g., SDG 1—No Poverty, SDG 2—Zero Hunger, SDG 10—Reduced Inequalities, and the African Union's Agenda 2063).

Table 3: Interaction with rural development goals

Bagonomic Element	Rural Development Goal Aligned	Mechanism of Influence
Communal Ownership	Land equity and reduced conflict	Land cooperatives, gender-inclusive land redistribution

FULL TEXTS BOOK

Cooperative Labor	Employment and productivity	Job creation, youth skill development, agro-labor mobilization
Local Markets First	Food security and rural livelihoods	Local value chains, community-supported agriculture, food affordability
Democratic Governance	Inclusion and local empowerment	Village-level planning councils, farmers' unions, participatory budgeting
Circular Rural Economy	Economic resilience and sustainability	Recycled rural surpluses, localized investment, community banking

Source: Author's own synthesis, 2025

This integrative model provides an operational structure to track and evaluate the transformation process under a Bagonomic lens.

3.3 Diagram of Conceptual Framework

Here is a visual representation of how the components of Bagonomics interact with the drivers of inclusive rural development:




Figure 1: Conceptual framework

Source: Phyton software, 2025

FULL TEXTS BOOK

4.0 Research Methodology

This study adopts a qualitative research methodology, enriched with systematic literature review techniques and informed by participatory rural appraisal (PRA) principles. Given that *Bagonomics* is a conceptual and community-driven framework not yet institutionalized in policy literature, a flexible and interpretive methodological approach was necessary to understand its relevance, map its dimensions, and evaluate its applicability to agricultural policy reform in Niger State.

4.1 Research Design

The research follows a multi-phase, exploratory review design, combining:

- **Desk-based literature analysis**: to identify and synthesize existing evidence on agricultural policy, rural exclusion, cooperative economics, and grassroots innovations in Nigeria and sub-Saharan Africa.
- **Conceptual synthesis**: of development models relevant to endogenous growth, solidarity economies, and participatory governance.
- **Theoretical triangulation**: applying multiple theoretical lenses—Participatory Development Theory, Endogenous Development, Feminist Political Ecology, etc.—to interpret findings from the literature.

This approach allows for a holistic analysis of both empirical evidence and normative arguments underpinning the Bagonomics framework.

4.2 Data Sources

The study utilized secondary data, obtained from the following categories:

- **Academic Literature**: Peer-reviewed journals accessed via Scopus, JSTOR, PubMed, and ScienceDirect databases from 2010–2024.
- **Policy Documents**: Nigerian government agricultural policies, particularly the Agriculture Promotion Policy (APP), National Agricultural Technology and Innovation Policy (NATIP, 2021–2025), and Niger State Agricultural Blueprint (2020–2030).
- **NGO and Development Reports**: Reports from UNDP, FAO, IFAD, and the Nigerian Bureau of Statistics (NBS) relevant to rural poverty, gender equity, and agricultural financing.
- **Grey Literature**: Think tank papers, grassroots innovation reports, and case studies from institutions such as IITA, ActionAid Nigeria, and the Nigeria Economic Summit Group.

Each document was reviewed using a content analysis framework that categorized findings by key themes: equity, inclusion, cooperative models, policy structure, and rural agency.

4.3 Inclusion and Exclusion Criteria

To ensure the relevance and credibility of sources:

- **Inclusion criteria**:
  - Studies published between 2010 and 2024.
  - Nigeria-focused or sub-Saharan African case studies with explicit analysis of agricultural policy, cooperative farming, rural governance, or gender dynamics.
  - Peer-reviewed journal articles, government white papers, and verified NGO publications.

FULL TEXTS BOOK

- **Exclusion criteria**:

  - Articles not focused on Africa or with purely econometric models without sociopolitical context.
  - Opinion pieces without empirical or theoretical grounding.

A total of **53 documents** were selected after screening 120 initial sources.

4.4 Analytical Framework

Data from the literature were coded and analyzed using a thematic analysis approach, guided by five pre-identified constructs from the Bagonomic framework:

1. Communal ownership of productive resources
2. Cooperative labor and knowledge-sharing
3. Local-first market models
4. Participatory governance structures
5. Circular and solidarity-based rural economies

Key concepts were coded manually using NVivo-style matrix development (though software was not used explicitly), and recurring patterns, gaps, contradictions, and innovations were identified and mapped against the rural development indicators relevant to Niger State.

The insights were then aligned with relevant SDG targets, gender equality indicators, and local policy frameworks to assess policy compatibility and innovation potential.

4.5 Participatory Rural Appraisal (PRA) Integration

Though no primary fieldwork was conducted, PRA principles were applied through secondary analysis of documented community-led pilot programs and cooperative development initiatives in Niger State, Benue, and Northern Kaduna. These included:

- Rotational farming cooperatives (*gayya* systems)
- Women-led agro-credit unions
- Indigenous irrigation schemes (e.g., *fadama* practices)
- Rural market cooperatives focused on local produce distribution

By synthesizing these real-world models, the study constructed plausible policy pathways for Bagonomics to be scaled into Niger State's agricultural framework.

4.6 Limitations of the Methodology

- Lack of primary data collection limited direct engagement with rural actors, which could have enriched the PRA dimension.
- Variability in data quality across grey literature sources posed challenges in comparative evaluation.
- Absence of disaggregated state-level data (e.g., gendered access to inputs in Niger State) constrained empirical depth.

Despite these limitations, triangulating findings across academic, policy, and community sources ensured the reliability and depth of the conceptual analysis.

FULL TEXTS BOOK

4.7 Ethical Considerations

As this study relied solely on secondary data, no formal ethical approval was required. However, care was taken to:

- Appropriately attribute all sources;
- Represent rural communities' experiences faithfully;
- Avoid over-generalizing localized practices to national contexts.

5.0 Results and Discussion

The review of recent academic and policy literature reveals five dominant thematic outcomes that affirm the potential of the Bagonomic framework to transform agricultural policy and rural development in Niger State. These are discussed below with supporting evidence.

5.1. Strengthened Agricultural Cooperatives Enhance Equity

One of the strongest findings across the reviewed literature is that agricultural cooperatives and community-based farming groups significantly enhance access to land, inputs, and markets for smallholder farmers—especially marginalized groups like women and youth.

A study by Ubandoma *et al.* (2025) in Mokwa Local Government Area of Niger State found that farmers who were members of rural cooperatives reported a 34% reduction in input costs and a 22% increase in access to extension services compared to non-members. These cooperatives functioned as social safety nets, enabling group-based access to fertilizers, irrigation, and even informal credit. The authors argue that "cooperative systems in rural Niger State are a viable entry point for agricultural equity if formally institutionalized" (Ubandoma *et al*., 2025).

This aligns with Bagonomics' principle of communal ownership and cooperative labor, offering a framework that not only supports group productivity but fosters economic democracy.

5.2. Community-Led Farming Enhances Sustainability and Agency

Community-driven agricultural initiatives—rooted in local governance and participatory decision-making—have proven to be more sustainable, adaptive, and resilient to climate stress.

Naadzenga & Palmiotto (2023) document a food cooperative model in Benue State that used participatory food system mapping and local seed preservation techniques. Farmers reported increased yield stability and greater trust in agro-ecological practices. The study emphasizes that "community-led cooperatives increased farmer agency, countering dependency on external inputs and state subsidies" (Naadzenga & Palmiotto, 2023).

This reinforces Bagonomics' commitment to local markets first and democratic governance, allowing rural communities to design agro-ecological practices that match their lived realities.

5.3. Gender Inclusion Remains an Incomplete Agenda

Although cooperatives have increased access to resources for many farmers, gender disparity persists in land access, decision-making, and cooperative leadership roles. According to Oseni (2023), while over 40% of agricultural workers in Southwest Nigeria are women, only 11% hold leadership roles in farming cooperatives, and fewer than 20% have titled land.

This gap is echoed in Niger State, where patriarchal structures continue to marginalize rural women from agri-financing and land titling schemes. Bagonomics, guided by feminist political ecology, calls for deliberate gender-transformative approaches—including quotas in cooperative governance and gender-sensitive land reforms.

FULL TEXTS BOOK

5.4. Policy-Community Disconnect Undermines Program Impact

A recurring issue in rural development interventions is the lack of alignment between policy goals and community needs. Hart (2024) found that donor-driven rural development projects in Nigeria often collapse due to weak local engagement and lack of contextual adaptation.

For example, in Niger State, several Anchor Borrowers Program clusters were reported to have failed due to poorly understood loan conditions and inflexible repayment structures. Bagonomics challenges this model by insisting that rural governance must emerge from the ground up, driven by local logic, seasonal realities, and customary practices.

5.5. Bagonomics Aligns with Proven Development Models

Finally, a global comparative review by Allen *et al*. (2022) supports the idea that community-led development frameworks outperform state-centric models in terms of inclusivity, resilience, and long-term sustainability. Their report shows that investments in locally owned seed systems, community irrigation schemes, and decentralized food networks led to a 30–50% increase in rural incomes in cases where rural agency was prioritized.

This aligns directly with Bagonomics' core principles and provides strong global validation for its application in the Niger State context.

Table 5: Synthesis of findings

Theme	Findings	Bagonomics Link
Agricultural Cooperatives	Improve input access, reduce costs, enhance extension access	Communal Ownership; Cooperative Labor
Community-Led Farming	Builds local trust, increases yield, and supports agro-ecology	Local Markets First
Gender Gaps in Rural Policy	Women underrepresented in land and cooperative leadership	Feminist Political Ecology
Policy-Community Disconnect	Projects fail due to lack of contextual fit and consultation	Democratic Governance
Global Support for Community Logic	Community-led development proven effective in Africa, Asia, Latin America	Validation of Bagonomics globally

Source: Author's own synthesis, 2025

6.0 Conclusion

This research review has demonstrated that conventional agricultural policies in Niger State, and Nigeria more broadly, have failed to achieve inclusive and equitable rural development due to systemic exclusion, policy centralization, and weak community engagement. While national frameworks such as the Agriculture Promotion Policy (APP) and the National Agricultural Technology and Innovation Plan (NATIP 2021–2025) aim for productivity gains, they largely remain extractive and top-down in nature, bypassing the socio-cultural realities and informal economies that dominate rural Nigeria.

FULL TEXTS BOOK

The conceptual framework of Bagonomics—with its pillars of communal ownership, cooperative labor, localized market logic, democratic governance, and circular rural economies—offers a transformative alternative. The findings reviewed here indicate that agricultural cooperatives, community-led farming initiatives, and participatory rural governance not only improve agricultural productivity and food security but also deepen economic inclusion and resilience, particularly for women, youth, and marginalized farmers.

Bagonomics represents more than a development strategy; it is a political and ethical commitment to justice, community agency, and sustainability. Its application in Niger State has the potential to recalibrate rural development from a model of dependency to one of empowerment and endogenous growth.

7.0 Recommendations

Based on the evidence gathered and the gaps identified in current policy practices, the following recommendations are offered:

7.1 Institutional Reforms

- Legalize and formalize rural cooperatives in Niger State through enabling legislation that guarantees access to land, credit, and capacity-building.
- Create local agricultural planning councils at ward and LGA levels to allow farmers and traditional authorities to co-design development priorities.

7.2 Gender-Responsive Interventions

- Implement affirmative action policies requiring 40% minimum representation of women in cooperative leadership and land allocation boards.
- Develop targeted rural women's innovation funds to support women-led food processing, seed systems, and extension networks.

7.3 Support for Agro-Solidarity Economies

- Establish community-owned seed banks and localized input systems that prioritize traditional crops, ecological methods, and community resilience.
- Incentivize internal local markets through tax breaks, public procurement from cooperatives, and infrastructure for village-level storage and transport.

7.4 Capacity Building and Knowledge Sharing

- Launch Bagonomics training hubs across Niger State to integrate youth into cooperative labor systems and agro-entrepreneurship.
- Support peer-to-peer knowledge exchanges between communities in Niger, Benue, and northern Kaduna to foster south-north solidarity learning.

8.0 Policy Implications

8.1 Reorientation of Development Metrics

- Shift policy success indicators from macro-output (tonnes of rice) to micro-impact (number of women landholders, cooperative retention rate, local food security scores).

8.2 Decentralization of Governance

- Empower local government councils with fiscal and legal authority to manage agriculture-led rural development in coordination with communities.

FULL TEXTS BOOK

8.3 Integration with National Frameworks

- Embed Bagonomic indicators within the Niger State Agricultural Policy (2020–2030) and align them with SDGs 1, 2, 5, 8, and 10 for development funding eligibility.

8.4 Donor and NGO Alignment

- Advocate for donors and multilateral agencies (e.g., IFAD, UNDP) to redirect technical assistance toward community-led policy formulation and institutional strengthening.

REFERENCES

Agarwal, B. (1994). *A field of one's own: Gender and land rights in South Asia* (No. 58). Cambridge University Press.

Ajulor, O. V., & Etim, E. (2019). Innovative agricultural policy and rural poverty reduction in Nigeria. *LASU Journal of Engineering, Science and Technology*, *1*(2), 39-43. Retrieved from https://www.researchgate.net/publication/343737538

Allen, B., Coonrod, J., Ashby, J., & Njiru, W. (2022). The case for inclusive agricultural development. *Image*.https://globalaffairs.org/sites/default/files/2022-06/IncDevRep%20FINAL%20DRAFT%204.pdf

Chambers, R. (1997). *Whose reality counts?: putting the first last* (pp. xx+-297). Intermediate Technology Publications.

Ering, S. O., Otu, J. E., & Archibong, E. P. (2014). Rural development policies in Nigeria: A critical appraisal. *International Journal of Education and Research*, *2*(9), 307-320.

Etuk, E. A., & Ayuk, J. O. (2021). Agricultural commercialisation, poverty reduction and pro-poor growth: evidence from commercial agricultural development project in Nigeria. *Heliyon*, *7*(5), e07997. https://doi.org/10.1016/j.heliyon.2021.e07997

Hart, A. O. (2024). Institutional framework for sustainable community engagement for development in Nigeria. *Journal of Humanities and Social Policy*, *10*(2), 14-37. https://www.iiardjournals.org/get/JHSP/VOL.%2010%20NO.%202%202024/Institutional%20Framework%20for%2014-37.pdf

Letswa, A. M. (2018). *Assessment of the National Fertilizer Policy Implementation on Farm Yields for Poverty Reduction among Farmers in Niger State, Nigeria* (Doctoral dissertation, Kwara State University (Nigeria). search.proquest.com

Naadzenga, A. I., & Palmiotto, P. A. (2023). Problem-solving with secondary climate refugees: identifying sustainable participatory processes through food security: a proposal for a food cooperative in Benue State–Nigeria. *Ethiopian Journal of Environmental Studies & Management*, *16*(2), 1–14. https://ejesm.org/wp-content/uploads/2023/06/ejesm.v16i2.1.pdf

Nwankwo, C. U., Ikehi, M. E., Ejiofor, T. E., & Ifeanyieze, F. O. (2024). Successive national agricultural policies/programmes, growth of gross domestic product (GDP) and expansion of agribusinesses in Nigeria. *Plos one*, *19*(2), e0291999. https://doi.org/10.1371/journal.pone.0291999

Ojong, F. E., & Anam, B. E. (2018). Agriculture promotion policy and rural development in Nigeria: Challenges and prospects. *Journal of Humanities and Social Science*, *23*(4), 31–40. https://staff.tiiame.uz/storage/users/43/articles/ufNH89h2b2thW6LIeDPazb2pGHGnxsS1CAyJ5aXS.pdf

FULL TEXTS BOOK

Olomola, A. S., & Nwafor, M. (2018). Nigeria agriculture sector performance review. *International Institute of Tropical Agriculture (IITA): Ibadan, Nigeria*, *3*. https://www.researchgate.net/publication/327666294

Oseni, A. O. (2023). *The impact of government policy on community enterprise in Nigeria* [Master's thesis, Estonian University of Life Sciences]. EMU DSpace. https://dspace.emu.ee/bitstream/10492/8530/1/Anuoluwapo_Oseni_2023MA_AFBM_fulltext.pdf

Ubandoma, G., Sule, B., Baba, K., Abubakar, S., & Yakubu, D. (2025). Assessment of rural cooperative societies on farming households'assistance in Mokwa Local Government Area of Niger State, Nigeria. *Journal of Renewable Agricultural Technology Research*.

Uduji, J. I., Okolo-Obasi, E. N., & Asongu, S. A. (2019). Corporate social responsibility and the role of rural women in sustainable agricultural development in sub-Saharan Africa: Evidence from the Niger Delta in Nigeria. *Sustainable Development*, *27*(4), 692-703. https://doi.org/10.1002/sd.1933

Verhelst, T., & Tyndale, W. R. (2002). Cultures, spirituality, and development. In *Development and culture: Selected essays from development in practice* (pp. 1-24).

Vieta, M. (2014). Social and solidarity economy: Towards inclusive and sustainable development. *Social and Solidarity Economy Academy, Campinas, Brazil*.

FULL TEXTS BOOK

REAL-TIME THERAPEUTIC DRUG MONITORING USING WEARABLE DEVICES

Puniparthi Sunitha

ORCID ID: 0009-0006-0930-4520

M.K. Vijiayalakshmi

Bharath institute of higher education and research

ABSTRACT

Therapeutic Drug Monitoring (TDM) plays a crucial role in optimizing drug efficacy and minimizing toxicity, particularly for medications with narrow therapeutic windows. Traditionally, TDM relies on intermittent blood sampling and laboratory assays, which can be invasive, time-consuming, and provide only snapshot information. In contrast, the integration of wearable technologies into modern healthcare introduces a paradigm shift, enabling real-time, continuous, and non-invasive monitoring of drug levels.

Wearable biosensors, including microneedle patches, smartwatches, and skin-interfaced electronics, can now detect drug concentrations in various biological fluids such as sweat, interstitial fluid, and saliva. These devices utilize electrochemical, optical, and microfluidic principles to measure drug levels with high sensitivity and accuracy. Combined with wireless communication and cloud-based data storage, wearable TDM systems offer patients and clinicians instant access to pharmacokinetic profiles, enhancing personalized drug dosing and improving therapeutic outcomes.

Applications of real-time TDM are particularly impactful in managing chronic conditions such as epilepsy, cardiovascular diseases, cancer, and post-transplant immunosuppression. These devices support proactive interventions by detecting early signs of subtherapeutic or toxic drug levels, thereby reducing hospitalizations and improving adherence.

However, challenges remain in sensor calibration, biological variability, data security, and regulatory approvals. Ethical concerns regarding patient privacy and the validation of wearable diagnostics must also be addressed. Despite these limitations, the future of TDM is promising with ongoing advances in artificial intelligence, predictive modeling, and digital health integration. Real-time TDM using wearable devices is emerging as a transformative tool in precision medicine, offering safer, smarter, and more responsive drug management strategies for diverse patient populations.

KEY WORDS: Therapeutic Drug Monitoring (TDM), Wearable Biosensors, Real-Time Drug Monitoring, Personalized Medicine, Smart Wearables in Healthcare, Pharmacokinetics

INTRODUCTION TO THERAPEUTIC DRUG MONITORING (TDM)

DEFINITION AND CLINICAL IMPORTANCE

Therapeutic Drug Monitoring (TDM) refers to the clinical practice of measuring specific drug concentrations in a patient's bloodstream to ensure that drug levels remain within the optimal therapeutic range. The purpose of TDM is to individualize dosing to maximize therapeutic effects while minimizing adverse reactions. This approach is especially critical for drugs with a narrow therapeutic index, where even small deviations in blood concentration can lead to toxicity or subtherapeutic effects.

TDM is not just about drug levels—it integrates pharmacokinetics, patient-specific variables (like renal or liver function), drug interactions, and disease state. This makes it a cornerstone of precision medicine,

FULL TEXTS BOOK

ensuring that patients receive the right drug at the right dose at the right time. The ultimate goal is improved clinical outcomes, fewer hospitalizations, and enhanced patient safety.

TRADITIONAL METHODS VS. REAL-TIME APPROACHES

Traditionally, TDM involves collecting blood samples at specific time points after drug administration. These samples are then sent to a laboratory for analysis using methods like high-performance liquid chromatography (HPLC) or mass spectrometry. Although accurate, these methods are invasive, time-consuming, and provide only periodic snapshots of drug concentration.

In contrast, real-time TDM uses wearable biosensors to continuously monitor drug levels in real-time. These devices can analyze interstitial fluid, sweat, or saliva non-invasively, sending live data to clinicians or apps for immediate interpretation. This technology offers several advantages: reduced need for venipuncture, real-time feedback, remote monitoring, and more precise dose titration.

By capturing the dynamic fluctuations of drug levels, real-time TDM enables personalized drug management in ways that traditional methods cannot.

DRUGS COMMONLY MONITORED (E.G., VANCOMYCIN, LITHIUM, ANTICONVULSANTS)

TDM is particularly useful for drugs that require tight control over blood concentrations. Common examples include:

- Vancomycin: A powerful antibiotic used to treat resistant infections. Inadequate levels may lead to treatment failure, while excess levels can cause nephrotoxicity.
- Lithium: Widely used in the management of bipolar disorder. Monitoring is essential due to its narrow therapeutic window and potential for toxicity.
- Anticonvulsants (e.g., phenytoin, valproic acid, carbamazepine): These medications must be carefully dosed to prevent seizures without causing sedation or toxicity.
- Immunosuppressants (e.g., tacrolimus, cyclosporine): In organ transplant patients, underdosing can lead to rejection, and overdosing can cause infection or organ damage.
- Theophylline and Digoxin: Older cardiovascular and respiratory drugs with known narrow therapeutic ranges.

Real-time wearable TDM systems hold the promise to transform the management of these drugs, especially in outpatient and home settings. By continuously monitoring levels, therapy can be adjusted proactively, avoiding complications before they arise.

WEARABLE DEVICES IN MODERN HEALTHCARE

TYPES OF WEARABLE BIOSENSORS

Wearable biosensors are compact, body-mounted devices designed to detect and measure physiological signals and biochemical markers continuously or periodically. These biosensors have revolutionized patient care by enabling real-time health monitoring outside clinical settings. Depending on the parameters they measure, wearable biosensors are categorized into several types:

- Electrochemical biosensors detect specific ions or molecules, like glucose or lactate, in body fluids such as sweat or interstitial fluid.
- Optical sensors use light to measure parameters like oxygen saturation, pulse, or blood flow (e.g., pulse oximeters).

FULL TEXTS BOOK

- Mechanical biosensors monitor physical movements, heart rate, or respiratory rate using pressure or stretch-sensitive materials.
- Microneedle-based sensors are minimally invasive and can sample interstitial fluid to analyze drug levels, offering an ideal tool for therapeutic drug monitoring (TDM).
- Smart patches and tattoos integrate sensors with skin-like flexibility to monitor hydration, pH, or drug metabolites.

These devices are typically connected to mobile phones or wearable hubs to allow continuous and personalized health tracking.

INTEGRATION WITH DIGITAL HEALTH AND IOT

Modern wearable biosensors are not standalone tools—they form part of a growing digital health ecosystem powered by the Internet of Things (IoT). Through Bluetooth or wireless connectivity, these devices transmit data in real time to smartphones, healthcare apps, or cloud-based platforms, enabling remote monitoring by clinicians.

This integration allows for:

- Telemedicine support, where clinicians can adjust medication remotely based on data.
- Smart alerts and notifications for patients when drug levels are too high or low.
- Data analytics powered by AI to detect patterns and predict future complications.

IoT-enabled wearables can be synced with Electronic Health Records (EHRs) to provide a more complete and dynamic patient profile. The convergence of biosensing, cloud computing, and real-time analytics enables a shift from reactive to predictive and personalized medicine.

ADVANTAGES OVER CONVENTIONAL MONITORING

Compared to traditional methods of monitoring, wearable biosensors offer numerous advantages that are reshaping how healthcare is delivered:

1. Non-Invasive or Minimally Invasive: Wearables eliminate the need for frequent blood draws or hospital visits, reducing patient discomfort and risk of infection.

2. Continuous Monitoring: Instead of relying on isolated data points, clinicians can observe drug levels and physiological trends throughout the day.

3. Remote Access: Healthcare providers can access patient data in real time, enabling early interventions and reducing hospital admissions.

4. Patient Empowerment: Patients become active participants in managing their health, as wearables promote greater self-awareness and adherence.

5. Cost-Effectiveness: By reducing the need for lab tests, hospital stays, and emergency care, wearables can lower overall healthcare costs.

Wearable biosensors are a game-changing innovation in modern medicine. Their ability to monitor health metrics continuously and transmit them wirelessly is redefining therapeutic monitoring, especially in chronic disease management and individualized treatment strategies.

PRINCIPLES OF REAL-TIME DRUG MONITORING

DETECTION MECHANISMS (ELECTROCHEMICAL, OPTICAL, ETC.)

FULL TEXTS BOOK

Real-time therapeutic drug monitoring (TDM) relies on highly sensitive detection mechanisms embedded within wearable biosensors. The two most commonly used technologies are electrochemical and optical detection.

- Electrochemical sensors measure changes in electrical properties (like current or voltage) when a specific drug or metabolite interacts with a sensor-coated electrode. These sensors are ideal for miniaturized, portable devices due to their low power requirements and rapid response times. They are especially useful for monitoring drugs like lithium, vancomycin, and theophylline in bodily fluids.

- Optical sensors, on the other hand, rely on changes in light absorption, fluorescence, or reflection. For example, some wearable devices shine light through the skin and measure the absorption spectrum to determine drug levels. Optical methods are non-invasive and useful for continuous, high-resolution monitoring.

Other emerging mechanisms include piezoelectric sensors (detecting mechanical changes) and colorimetric biosensors, which visually change color in the presence of a specific drug—useful for rapid at-home tests.

ANALYTES MEASURED (BLOOD, INTERSTITIAL FLUID, SWEAT)

For wearable real-time monitoring, accessing non-invasive or minimally invasive biological fluids is crucial. Traditional TDM uses blood samples, but real-time wearables have expanded to include:

- Interstitial Fluid (ISF): Found between cells, ISF reflects blood drug concentrations with only a slight delay. Microneedle-based sensors painlessly access ISF, making it ideal for long-term drug monitoring.

- Sweat: Easily accessible and non-invasive, sweat is used to detect several biomarkers and drug residues. However, drug levels in sweat may vary with hydration, temperature, and individual physiology.

- Saliva and Tears: Though less common, these fluids are also being explored for drug monitoring, particularly in neurological and psychiatric medications.

The challenge lies in ensuring the accuracy, correlation, and stability of drug levels in these alternative matrices compared to blood. Still, advances in sensor calibration are making these non-invasive fluids more reliable for real-time applications.

ROLE OF NANOTECHNOLOGY AND MICROFLUIDICS

The evolution of real-time TDM has been accelerated by the integration of nanotechnology and microfluidics, which enhance sensitivity, specificity, and user-friendliness.

- Nanotechnology enables the creation of nanoscale sensors and surfaces with high reactivity. Nanomaterials like graphene, gold nanoparticles, and carbon nanotubes are frequently used to coat sensors, significantly increasing their ability to detect small quantities of drugs even at low concentrations.

- Microfluidics involves manipulating tiny volumes of fluid (often less than a microliter) within miniaturized channels on a chip. This technology allows wearable devices to efficiently sample and process body fluids in real time. It also helps separate, filter, and guide analytes to sensor surfaces, increasing accuracy while minimizing sample contamination.

Together, nanotechnology and microfluidics create biosensors that are compact, highly sensitive, and well-suited for wearable platforms, allowing for seamless integration into daily life. Real-time drug monitoring through wearable devices relies on sophisticated detection principles and fluid sampling

FULL TEXTS BOOK

methods, made even more powerful by nanotech and microfluidic engineering. This integration allows patients to receive tailored, timely, and safer drug therapies, especially for chronic or high-risk medications.

TECHNOLOGIES ENABLING WEARABLE TDM

SMART PATCHES AND MICRONEEDLE SENSORS

Smart patches and microneedle sensors are among the most promising innovations in real-time therapeutic drug monitoring (TDM). These devices are designed to be non-invasive or minimally invasive and can continuously monitor drug levels in body fluids like interstitial fluid (ISF) or sweat.

Smart patches are flexible, skin-adherent devices embedded with biosensors, microelectronics, and often microheaters to enhance drug diffusion from the skin. These patches can monitor multiple parameters such as pH, temperature, and drug metabolites in real time. Some advanced patches integrate microneedle arrays to access ISF painlessly.

Microneedle sensors are especially significant for TDM. These are tiny projections, often smaller than a millimeter, that painlessly penetrate the skin to access ISF. They are typically coated with biosensing materials that react with specific drug molecules. The key advantage is their ability to provide highly accurate measurements while avoiding the discomfort and infection risks associated with traditional blood sampling.

Such technologies are currently being tested for monitoring drugs like vancomycin, lithium, and chemotherapeutic agents, with prototypes showing promising accuracy and patient comfort.

CONTINUOUS ANALYTE MONITORING CHIPS

Another major advancement is the development of continuous analyte monitoring chips that integrate multiple sensing and data-processing functions on a single, miniaturized platform. These chips, often referred to as lab-on-a-chip (LOC) systems, combine microfluidic channels with biosensors to detect and analyze drugs in a steady stream of biological fluid.

These chips are capable of real-time sampling, detection, signal processing, and result display—all within a few square millimeters. Using nanomaterials and precise engineering, the chips can detect drugs at low concentrations and even differentiate between structurally similar molecules.

A key advantage is their ability to perform multi-analyte monitoring, allowing the device to track not only the drug concentration but also relevant biomarkers such as creatinine (for kidney function) or liver enzymes. This helps clinicians interpret drug levels more accurately and tailor therapy more precisely.

WIRELESS DATA TRANSMISSION AND CLOUD STORAGE

Real-time TDM becomes truly powerful when it is integrated with wireless communication and cloud-based data management. Once a wearable sensor detects a drug concentration, the data must be transmitted securely and instantly to the relevant stakeholders—clinicians, pharmacists, or digital health platforms.

Most wearable devices use Bluetooth Low Energy (BLE), Wi-Fi, or Near Field Communication (NFC) to transmit data to a smartphone or wearable hub. From there, data is uploaded to cloud storage, where it can be accessed remotely by authorized healthcare providers.

This digital integration offers several key benefits:

- Remote monitoring and intervention, reducing hospital visits.

FULL TEXTS BOOK

- Automatic alerts when drug levels fall outside the therapeutic range.
- Long-term data tracking, enabling trend analysis and therapy optimization.
- AI-based analytics to predict dosing needs or flag potential side effects.

Moreover, with encrypted cloud systems and user authentication, patient data privacy is maintained while still allowing for collaborative care. The convergence of smart patches, microneedles, continuous monitoring chips, and wireless data systems is revolutionizing therapeutic drug monitoring. These technologies offer a future where drug levels are tracked continuously, accurately, and effortlessly—empowering both patients and providers with real-time insights.

APPLICATIONS IN CHRONIC DISEASE MANAGEMENT

MONITORING ANTIEPILEPTIC DRUGS

Patients with epilepsy often rely on antiepileptic drugs (AEDs) such as phenytoin, carbamazepine, valproic acid, and levetiracetam to prevent seizures. These medications have narrow therapeutic windows, meaning that even small fluctuations in blood levels can lead to either toxicity or inadequate seizure control. Moreover, AEDs often interact with other drugs, making dosage management complex.

Real-time wearable TDM offers a breakthrough in epilepsy care by continuously monitoring drug concentrations in sweat or interstitial fluid through non-invasive patches or microneedle sensors. This allows for personalized dosing adjustments, especially in patients with variable metabolism or during periods of stress, illness, or growth (as in children).

Additionally, these systems can send automated alerts to both patients and healthcare providers when drug levels drop below or rise above the safe range, helping prevent seizure episodes or drug-induced side effects like dizziness, fatigue, or liver toxicity. As a result, wearable TDM can significantly improve treatment adherence, seizure control, and quality of life for individuals with epilepsy.

CARDIOVASCULAR DRUGS (E.G., DIGOXIN)

Drugs like digoxin, used for managing heart failure and atrial fibrillation, require precise monitoring because of their high toxicity risk at elevated blood concentrations. Overdoses can lead to symptoms ranging from nausea and confusion to life-threatening arrhythmias.

Wearable biosensors designed for real-time digoxin monitoring can detect minute drug level changes in body fluids. For elderly patients or those with fluctuating renal function, these devices can help avoid adverse drug events by providing ongoing feedback on digoxin levels, prompting timely interventions such as dose changes or drug discontinuation.

Moreover, for patients on polypharmacy (multiple drugs), which is common in cardiovascular disease management, wearable TDM can reduce the chances of drug-drug interactions by tracking multiple drugs simultaneously. It helps clinicians balance therapeutic effectiveness and safety, especially in long-term outpatient care.

ONCOLOGY AND IMMUNOSUPPRESSIVE THERAPIES

In oncology, many chemotherapy agents (like methotrexate and cisplatin) and targeted therapies must be dosed precisely to avoid harming healthy tissue while effectively targeting cancer cells. However, traditional monitoring is reactive and often delayed.

Wearable TDM enables real-time tracking of chemotherapeutic drug levels, allowing oncologists to tailor doses based on patient-specific pharmacokinetics. This can reduce side effects like organ toxicity and enhance treatment response, especially in personalized cancer therapy protocols.

FULL TEXTS BOOK

Similarly, patients undergoing organ transplantation require lifelong use of immunosuppressive drugs (e.g., tacrolimus, cyclosporine). These medications must be carefully balanced to prevent rejection without suppressing the immune system too much. Real-time monitoring through smart patches and microneedles can detect subtherapeutic levels early—before rejection starts—or identify when doses are too high, putting the patient at risk for infection.

This kind of continuous, remote monitoring not only improves safety but also reduces hospital visits and enhances the overall management of transplant recipients, improving long-term outcomes. Wearable TDM technologies are poised to revolutionize chronic disease management by offering safer, more responsive, and highly individualized drug therapy for conditions like epilepsy, cardiovascular disease, cancer, and transplant care.

BENEFITS AND CLINICAL OUTCOMES

PERSONALIZED DOSAGE ADJUSTMENTS

One of the most impactful benefits of real-time therapeutic drug monitoring (TDM) using wearable devices is the ability to enable personalized dosage adjustments. Unlike conventional methods that rely on periodic blood tests and delayed results, wearable biosensors offer continuous, real-time data on drug levels in the body. This means healthcare providers can fine-tune medication doses based on a patient's current physiological status, not just population averages.

Each person metabolizes drugs differently due to genetic variations, age, organ function, and concurrent medications. Real-time TDM accounts for these differences by showing how a specific drug behaves in an individual's system over time. If drug levels are too low, the system can alert clinicians to increase the dose; if levels are too high, a reduction can be made immediately—preventing adverse effects before they occur.

This level of precision is especially valuable for managing drugs with narrow therapeutic windows, such as anticonvulsants, immunosuppressants, and chemotherapy agents. The result is a truly individualized treatment plan that evolves with the patient's needs.

IMPROVED ADHERENCE AND REDUCED TOXICITY

Medication adherence is a persistent challenge in chronic disease management. Many patients miss doses or take medications incorrectly due to forgetfulness, side effects, or lack of understanding. Wearable TDM helps address this issue by providing visual, real-time feedback on drug presence in the body. When patients see how drug levels fluctuate based on their behavior, they become more engaged and committed to following their regimen.

Moreover, by continuously monitoring drug concentrations, these devices can prevent drug accumulation and toxicity, which are common reasons for hospital admissions. For instance, patients taking lithium or digoxin are at high risk of toxic effects if blood levels rise even slightly above the therapeutic range. Wearable monitoring can detect this early and trigger alerts to modify the dose before harm occurs.

Clinicians, too, benefit from better insights into how well a patient is adhering to their treatment plan. If drug levels are consistently low, it may suggest missed doses or poor absorption. This allows for proactive interventions such as counseling or switching to alternative formulations, ultimately improving both safety and therapeutic effectiveness.

EARLY DETECTION OF UNDER- OR OVER-DOSING

Another critical clinical outcome of wearable TDM is the early detection of under- or over-dosing, which can have life-threatening consequences if not caught in time. With traditional monitoring, healthcare

FULL TEXTS BOOK

providers may only recognize dosing problems after symptoms develop or during routine lab tests—which may be too late.

Wearable biosensors continuously monitor the drug levels and can detect subtle trends or sudden changes. If levels are dropping too quickly, the system can flag potential under-dosing, prompting timely intervention before a relapse or flare-up occurs. Conversely, if drug levels spike unexpectedly, adjustments can be made before toxicity develops.

This level of vigilance ensures that treatment remains within the optimal therapeutic range at all times, minimizing fluctuations and enhancing clinical stability—especially in vulnerable populations like the elderly, transplant recipients, and oncology patients.

Wearable TDM empowers clinicians and patients alike through real-time insights, resulting in safer, more effective, and personalized care with fewer complications and better long-term outcomes.

REGULATORY AND ETHICAL CONSIDERATIONS

FDA AND EMA GUIDELINES FOR WEARABLE DIAGNOSTICS

As wearable devices for therapeutic drug monitoring (TDM) move from research to clinical practice, their regulation becomes crucial to ensure safety, accuracy, and reliability. Two major bodies that govern the approval of such medical devices are the U.S. Food and Drug Administration (FDA) and the European Medicines Agency (EMA).

In the United States, the FDA classifies wearable drug monitoring devices under Class II or Class III medical devices, depending on the level of risk involved. Devices that influence clinical decision-making, like those used in real-time TDM, often undergo Premarket Approval (PMA) or 510(k) clearance. Manufacturers must demonstrate that the wearable is substantially equivalent to existing approved technologies or prove its safety and effectiveness through clinical trials.

Similarly, the EMA requires compliance with the Medical Device Regulation (MDR) framework in the European Union. Wearables used in drug monitoring must go through Notified Body assessments, clinical evaluations, and post-market surveillance plans. The emphasis is on reliability of data, patient safety, and quality control. In both regions, regulatory frameworks are evolving to accommodate innovations like AI-powered biosensors and cloud-based diagnostics.

DATA PRIVACY AND PATIENT CONSENT

Ethical considerations are equally important, especially concerning data privacy and patient autonomy. Wearable TDM devices collect highly sensitive personal health data, often in real time and continuously. If this data is not properly protected, it can be misused, leading to discrimination, stigmatization, or breaches of confidentiality.

Hence, it is essential that patients give informed consent before using such technologies. This includes understanding what data is being collected, how it will be stored, who will access it, and how long it will be retained. Patients should also have the option to withdraw from monitoring without penalty.

Compliance with privacy laws such as the Health Insurance Portability and Accountability Act (HIPAA) in the U.S. and General Data Protection Regulation (GDPR) in the EU is mandatory. These laws require data encryption, anonymization, and secure cloud storage. Developers and clinicians must also ensure that wearable devices are not used to pressure patients into sharing more data than necessary.

In short, ethical TDM practices must strike a balance between clinical utility and patient rights.

CLINICAL VALIDATION AND APPROVAL HURDLES

FULL TEXTS BOOK

Before wearable drug monitoring devices can be widely adopted, they must undergo rigorous clinical validation to prove their accuracy, consistency, and real-world usability. Unlike traditional lab-based methods, wearables face unique challenges such as skin compatibility, sensor drift, motion artifacts, and sample variability in non-invasive fluids like sweat or interstitial fluid.

Validation involves comparing wearable readings with gold-standard lab results in diverse populations under various conditions. Furthermore, these devices must be tested across different age groups, ethnicities, and disease states to confirm broad applicability.

Another hurdle is ensuring interoperability with healthcare systems, including integration into electronic health records (EHRs). Without clinical proof of reliability and usability, regulatory approval becomes difficult, delaying patient access to this life-changing technology.

While wearable TDM devices offer great promise, their success depends on meeting strict regulatory standards, protecting patient data, and passing scientific validation to ensure trust, safety, and ethical deployment in healthcare.

LIMITATIONS AND CHALLENGES

SENSOR CALIBRATION AND STABILITY

One of the primary technical challenges in real-time therapeutic drug monitoring (TDM) using wearable devices is ensuring accurate sensor calibration and long-term stability. Unlike controlled laboratory environments, wearable sensors are exposed to variable conditions such as temperature, humidity, sweat, and skin movement. These factors can affect the precision of measurements over time.

Sensors must be regularly calibrated to maintain accuracy, especially when detecting drugs at very low concentrations. However, frequent manual calibration is not practical for at-home or ambulatory use. Therefore, ongoing research is focused on developing self-calibrating sensors or those that use built-in reference markers to adjust readings automatically.

Another issue is signal drift, where sensor readings become less reliable with extended use. Material fatigue, enzyme degradation (in biosensors), and mechanical wear can all contribute to reduced performance. This instability can lead to false readings, which, in the context of drug monitoring, could result in dangerous dosing decisions.

Improving the durability, responsiveness, and lifespan of wearable biosensors is essential to making them more viable for long-term clinical application.

BIOLOGICAL VARIABILITY IN SAMPLE MATRICES

Real-time TDM often relies on alternative biological fluids such as interstitial fluid (ISF), sweat, saliva, or tears rather than traditional blood samples. While these fluids offer a non-invasive route for monitoring, they introduce challenges related to biological variability.

For example, drug concentrations in sweat or ISF may not correlate perfectly with blood levels due to delay in diffusion, pH differences, or individual variations in skin properties. Factors like hydration, diet, skin health, and exercise can all influence analyte concentrations in these fluids.

In addition, variability between patients—or even within the same patient under different conditions—makes it difficult to establish universal calibration curves for all users. This lack of standardization poses a barrier to widespread clinical adoption, as clinicians must have confidence that readings from wearable devices are both consistent and representative of true therapeutic levels.

Addressing this issue requires refined algorithms, personalized baselines, and possibly the use of multi-analyte sensors that can cross-reference multiple biomarkers to enhance accuracy.

FULL TEXTS BOOK

COST AND ACCESSIBILITY

While wearable TDM devices offer numerous benefits, they currently remain financially inaccessible to many patients, particularly in low-resource settings. The cost includes not only the sensor hardware but also supporting technologies like smartphones, wireless connectivity, cloud-based storage, and subscription fees for medical monitoring platforms.

Moreover, frequent replacement of consumable parts (like sensor patches) and the need for specialized apps or software further increase the expense. For healthcare providers, integrating wearable data into clinical workflows also requires investment in infrastructure and training.

Beyond cost, there is a digital divide that limits access for populations with limited technological literacy or internet connectivity. Elderly patients, rural communities, and underserved regions may find it difficult to adopt these technologies despite their clinical potential.

To achieve equitable healthcare benefits, future developments must focus on cost-effective, scalable designs and inclusive technology education, ensuring wearable TDM becomes a tool for global health, not just for high-income groups. While wearable devices for drug monitoring hold enormous promise, challenges related to calibration, biological variability, and affordability must be addressed to ensure accurate, stable, and widespread clinical use.

FUTURE PERSPECTIVES AND RESEARCH DIRECTIONS

AI AND PREDICTIVE ANALYTICS IN TDM

Artificial intelligence (AI) is set to revolutionize therapeutic drug monitoring (TDM) by transforming how we interpret data collected from wearable devices. Wearable biosensors generate a vast amount of continuous, real-time data regarding drug levels, physiological responses, and lifestyle factors. AI algorithms, particularly those based on machine learning (ML) and deep learning (DL), can analyze this complex data to uncover patterns that are often too subtle or complex for human interpretation.

By integrating pharmacokinetic and pharmacodynamic models with patient-specific variables (like weight, organ function, and genetic markers), AI can predict optimal drug dosages, warn about potential adverse effects before they occur, and even forecast how a patient's response to a drug may evolve over time. These predictive models will shift TDM from a reactive to a proactive approach, minimizing trial-and-error and enhancing treatment precision.

In the near future, AI-powered TDM systems could automatically adjust dosages or suggest real-time therapeutic decisions through clinician alerts or patient apps—making precision medicine truly intelligent.

INTEGRATION WITH ELECTRONIC HEALTH RECORDS (EHRS)

Another promising area of research is the seamless integration of wearable TDM data with Electronic Health Records (EHRs). EHRs serve as the central hub for patient medical information, including history, medications, lab results, and physician notes. When wearable biosensors are connected to EHR platforms, healthcare providers can view real-time drug levels alongside other clinical data, improving decision-making.

This integration would support more coordinated, efficient care, especially in multidisciplinary teams where multiple providers must manage a patient's treatment. For example, an oncologist and pharmacist can review real-time drug metabolism data before adjusting chemotherapy doses. Additionally, automated alerts for out-of-range drug levels can be incorporated into EHR systems, ensuring that critical issues are not missed during routine care.

FULL TEXTS BOOK

Future research should focus on interoperability standards, data security protocols, and software development to allow widespread, safe, and effective EHR integration.

WEARABLES FOR PEDIATRIC AND ELDERLY POPULATIONS

Although wearable TDM holds great promise for all age groups, special attention is needed for pediatric and elderly populations. These groups often have unique challenges in drug metabolism, body fluid composition, and skin sensitivity—all of which impact the performance of wearable sensors.

In pediatrics, dosing is particularly sensitive due to rapid growth, developmental changes, and limited communication abilities. Wearables designed for children must be non-invasive, comfortable, and easy to use. Real-time monitoring could significantly reduce dosing errors and avoid side effects in neonates and children on long-term medications such as anticonvulsants or antibiotics.

For the elderly, age-related decline in kidney or liver function, comorbidities, and polypharmacy increase the risk of drug toxicity. Wearable TDM can offer continuous, at-home monitoring, reducing the need for frequent hospital visits. However, future devices must consider fragile skin, dexterity limitations, and ease of data interpretation for older adults and caregivers.

Developing age-specific, user-friendly wearable systems is a key area for ongoing and future research, the future of wearable TDM is bright, with AI, EHR integration, and population-specific innovations paving the way for smarter, safer, and more inclusive drug monitoring solutions.

CONCLUSION

Real-time therapeutic drug monitoring (TDM) using wearable devices represents a transformative step in modern, personalized healthcare. These innovative systems provide continuous, non-invasive tracking of drug levels through fluids such as interstitial fluid, sweat, or saliva—offering an effective alternative to traditional blood-based methods. With technologies like microneedle sensors, smart patches, and lab-on-a-chip platforms, wearables have begun bridging the gap between the clinic and the patient's daily environment.

By enabling personalized dosage adjustments, improving patient adherence, and detecting drug level abnormalities early, wearable TDM has demonstrated potential to improve outcomes in chronic diseases such as epilepsy, heart failure, cancer, and post-transplant care. The integration of AI and predictive analytics further strengthens this field by converting raw sensor data into meaningful clinical decisions, enhancing safety and responsiveness in treatment.

However, despite its promise, wearable TDM still faces challenges. Sensor calibration, biological variability in non-blood matrices, cost, and accessibility are key limitations that need further research and innovation. Additionally, regulatory clarity, data privacy, and clinical validation are crucial for widespread adoption and trust among healthcare professionals and patients.

Looking ahead, the success of wearable drug monitoring depends not only on technological advancements but also on interdisciplinary collaboration—involving engineers, clinicians, data scientists, regulatory authorities, and patients. With growing demand for remote and personalized care, wearable TDM systems are well-positioned to become central tools in digital health and precision medicine. Real-time wearable drug monitoring is not just a futuristic idea—it is a fast-approaching reality that promises to redefine how medications are managed, making therapies safer, more effective, and more tailored to individual needs.

FULL TEXTS BOOK

REFERANCE

1. Ates, I., Kaplan, M., & Yilmaz, N. (2020). Therapeutic drug monitoring: A review of current practice and future trends. *Frontiers in Pharmacology, 11*, 601427. https://doi.org/10.3389/fphar.2020.601427

2. Liu, Y., Luo, Y., Jin, Y., et al. (2023). Revolutionizing precision medicine: Exploring wearable sensors for therapeutic drug monitoring and personalized therapy. *Biosensors, 13*(8), 720. https://doi.org/10.3390/bios13080720

3. Conejo-Cuevas, G., Wang, Y., & Zhang, Y. (2023). Microneedles for continuous, minimally invasive drug monitoring. *Biosensors and Bioelectronics, 235*, 115341. https://doi.org/10.1016/j.bios.2023.115341

4. Turner, A. P., & Newman, J. D. (2022). Aptamer-based microneedle biosensors for drug monitoring. *Analytical Chemistry, 94*(11), 4560–4568. https://doi.org/10.1021/acs.analchem.2c00829

5. Kang, J. S., & Lee, M. H. (2009). Overview of therapeutic drug monitoring. *The Korean Journal of Internal Medicine, 24*(1), 1–10. https://doi.org/10.3904/kjim.2009.24.1.1

6. Clarke, W. (2016). Analytical issues in therapeutic drug monitoring. *Journal of Clinical Pharmacology, 56*(1), 7–14. https://doi.org/10.1002/jcph.597

7. Shalannanda, W., et al. (2023). Biosensors for therapeutic drug monitoring: A review. *F1000Research, 12*, 171. https://doi.org/10.12688/f1000research.128648.1

8. Emaminejad, S., et al. (2017). Autonomous sweat extraction and analysis applied to monitoring alcohol consumption. *PNAS, 114*(18), 4625–4630. https://doi.org/10.1073/pnas.1701740114

9. Yetisen, A. K., et al. (2018). Wearables in medicine. *Advanced Materials, 30*(33), 1706910. https://doi.org/10.1002/adma.201706910

10. Parlak, O., et al. (2018). Noninvasive monitoring of drug levels in sweat. *Science Advances, 4*(7), eaar2904. https://doi.org/10.1126/sciadv.aar2904

11. Heikenfeld, J., et al. (2019). Wearable sensors: Modalities, challenges, and prospects. *Lab on a Chip, 19*(2), 199–207. https://doi.org/10.1039/C8LC00917J

12. Kim, J., et al. (2019). Soft wearable systems for colorimetric analysis of body fluids. *Advanced Functional Materials, 29*(4), 1807310. https://doi.org/10.1002/adfm.201807310

13. Yang, Y., et al. (2021). Integrated wearable sensors for real-time sweat analysis. *Biosensors and Bioelectronics, 177*, 112989. https://doi.org/10.1016/j.bios.2020.112989

14. Bandodkar, A. J., & Wang, J. (2014). Non-invasive wearable electrochemical sensors. *TrAC Trends in Analytical Chemistry, 76*, 173–179. https://doi.org/10.1016/j.trac.2015.07.001

15. Gao, W., et al. (2016). Fully integrated wearable sensor arrays for multiplexed in situ perspiration analysis. *Nature, 529*(7587), 509–514. https://doi.org/10.1038/nature16521

16. Sempionatto, J. R., et al. (2020). Epidermal enzymatic biosensors for real-time monitoring of lactate and glucose. *Nature Biomedical Engineering, 4*(12), 1176–1186. https://doi.org/10.1038/s41551-020-00636-w

17. Ji, W., et al. (2023). Lab-on-a-chip platforms for real-time drug monitoring. *Microfluidics and Nanofluidics, 27*(3), 45. https://doi.org/10.1007/s10404-023-02473-3

FULL TEXTS BOOK

18. Imani, S., et al. (2016). A wearable chemical sensor for real-time sweat monitoring. *Nature Communications, 7*, 11650. https://doi.org/10.1038/ncomms11650

19. Lee, H., et al. (2021). Stretchable graphene–gold biosensor arrays for wearable health monitoring. *ACS Nano, 15*(3), 5440–5451. https://doi.org/10.1021/acsnano.0c08689

20. Ray, T. R., et al. (2019). Bio-integrated wearable systems. *Chemical Reviews, 119*(8), 5461–5533. https://doi.org/10.1021/acs.chemrev.8b00573

21. Samiei, N., et al. (2020). Personalized therapy using wearable sensors. *NPJ Digital Medicine, 3*, 20. https://doi.org/10.1038/s41746-020-0220-6

22. Alshahrani, S., et al. (2023). Microneedle biosensors for chronic disease monitoring. *Biosensors, 13*(2), 155. https://doi.org/10.3390/bios13020155

23. Oster, J., et al. (2016). Continuous monitoring of digoxin using wearable biosensors. *Journal of Biomedical Science and Engineering, 9*(6), 276–285. https://doi.org/10.4236/jbise.2016.96023

24. Mariani, S., & Nicolini, L. (2021). Wearable devices for cancer drug monitoring. *Cancers, 13*(12), 3046. https://doi.org/10.3390/cancers13123046

25. Vanderslott, S., et al. (2022). Wearables for transplant recipients. *Transplantation Reviews, 36*(4), 100675. https://doi.org/10.1016/j.trre.2022.100675

26. FDA. (2023). *Premarket Notification 510(k)*. https://www.fda.gov/medical-devices

27. EMA. (2023). *Medical Device Regulation (EU 2017/745)*. https://www.ema.europa.eu

28. U.S. Department of Health & Human Services. (2020). *Summary of the HIPAA Privacy Rule*. https://www.hhs.gov/hipaa

29. European Union. (2018). *General Data Protection Regulation (GDPR)*. https://gdpr.eu/

30. Mathews, A. W. (2020). Wearables and the ethical dilemma. *Journal of Ethics in Health Tech, 7*(1), 14–22. https://doi.org/10.1016/j.jeht.2020.05.003

31. Topol, E. (2019). High-performance medicine: The convergence of human and artificial intelligence. *Nature Medicine, 25*, 44–56. https://doi.org/10.1038/s41591-018-0300-7

32. Esteva, A., et al. (2021). A guide to deep learning in healthcare. *Nature Medicine, 27*(5), 750–765. https://doi.org/10.1038/s41591-021-01301-3

33. Rajpurkar, P., et al. (2022). Machine learning for medical diagnostics. *Annual Review of Biomedical Data Science, 5*, 307–331. https://doi.org/10.1146/annurev-biodatasci-030320-100218

34. Ahmed, M. U., & Penzel, T. (2023). Smart wearables for predictive healthcare. *Sensors, 23*(1), 86. https://doi.org/10.3390/s23010086

35. Chen, M., et al. (2020). Wearable 2.0: Enabling predictive and precision health. *IEEE Communications Magazine, 58*(1), 44–50. https://doi.org/10.1109/MCOM.001.1900422

36. Shi, Q., et al. (2021). Pediatric wearable biosensors: Design and implementation. *Pediatric Research, 90*, 214–220. https://doi.org/10.1038/s41390-020-01188-9

37. Chan, M., et al. (2020). Wearables for elderly care: A review. *Sensors, 20*(17), 4826. https://doi.org/10.3390/s20174826

38. Coorey, G. M., et al. (2022). Barriers to wearable tech in rural areas. *JMIR mHealth and uHealth, 10*(3), e28823. https://doi.org/10.2196/28823

FULL TEXTS BOOK

39. Ahmed, S., et al. (2021). Affordable wearable diagnostics: Bridging gaps in global health. *Global Health Action, 14*(1), 1878866. https://doi.org/10.1080/16549716.2021.1878866

40. Gupta, S., & Arora, S. (2023). Low-cost biosensing for underserved populations. *Biomedical Engineering Letters, 13*(2), 271–278. https://doi.org/10.1007/s13534-023-00291-2

41. Wu, W., et al. (2022). Challenges in wearable sensor calibration. *Sensors, 22*(12), 4439. https://doi.org/10.3390/s22124439

42. Nguyen, H. T., et al. (2023). Performance variability in wearable TDM. *Biomedical Signal Processing and Control, 81*, 104328. https://doi.org/10.1016/j.bspc.2023.104328

43. Ali, F., et al. (2021). Stability testing of wearable biosensors. *IEEE Sensors Journal, 21*(9), 10723–10731. https://doi.org/10.1109/JSEN.2021.3065614

44. Zhao, Y., et al. (2021). Real-time biofluid monitoring: Challenges and prospects. *ACS Sensors, 6*(9), 3299–3312. https://doi.org/10.1021/acssensors.1c00777

45. Luo, L., et al. (2022). Signal drift and correction strategies in wearable biosensors. *Sensors and Actuators B: Chemical, 360*, 131481. https://doi.org/10.1016/j.snb.2021.131481

46. Jones, B., et al. (2023). Real-time drug monitoring in clinical trials. *Contemporary Clinical Trials, 124*, 106962. https://doi.org/10.1016/j.cct.2023.106962

47. Williams, C., et al. (2021). Integration of wearable data into clinical workflow. *Journal of the American Medical Informatics Association, 28*(3), 554–560. https://doi.org/10.1093/jamia/ocaa304

48. Novak, J., et al. (2020). Clinical acceptability of wearable biosensors. *Digital Biomarkers, 4*(3), 125–134. https://doi.org/10.1159/000509651

49. Snyder, C. F., et al. (2022). Remote patient monitoring in cancer care. *JCO Clinical Cancer Informatics, 6*, e2200065. https://doi.org/10.1200/CCI.22.00065

50. Lindus Health. (2024). Enhancing clinical trial efficiency through wearable monitoring. Retrieved from https://www.lindushealth.com/blog

FULL TEXTS BOOK

THE ROLE OF GUT MICROBIOTA IN CHRONIC DISEASES

Puniparthi Sunitha

ORCID ID: 0009-0006-0930-4520

M.K. Vijiayalakshmi

Bharath institute of higher education and research

ABSTRACT

The human gut microbiota—a complex and dynamic community of trillions of microorganisms—plays a critical role in maintaining overall health. Beyond its traditional association with digestion, recent research has unveiled its deep-rooted influence on the development and progression of numerous chronic diseases. This article explores the multifaceted roles of the gut microbiome in major chronic conditions, highlighting its emerging importance in modern medicine.

First, we delve into the gut-brain axis, revealing how microbial metabolites and signaling molecules affect neurological function and are implicated in disorders such as depression, autism, and Parkinson's disease. We then examine its connection to metabolic conditions, including obesity and type 2 diabetes, where altered microbial composition contributes to systemic inflammation and insulin resistance. Cardiovascular diseases are also discussed, focusing on gut-derived metabolites like TMAO that promote atherosclerosis and cardiac dysfunction.

Additionally, the microbiota's ability to shape immune responses is explored in the context of autoimmune disorders such as inflammatory bowel disease and multiple sclerosis. The article also investigates how certain bacterial species and toxins influence cancer risk, particularly in colorectal and gastric cancers.

Diet, antibiotic use, and environmental factors are shown to significantly influence microbial balance, emphasizing the need for tailored dietary strategies and microbiota-based therapies. Promising interventions like fecal microbiota transplantation, probiotics, and microbiome-targeted drugs are presented as potential tools for restoring microbial harmony and improving clinical outcomes.

While research into gut microbiota is expanding rapidly, challenges remain in translating findings into consistent clinical practice. A deeper understanding of microbial interactions and host factors is essential for advancing personalized medicine and chronic disease prevention in the years ahead.

KEY WORDS: Gut Microbiota, Chronic Diseases, Dysbiosis, Gut-Brain Axis, Metabolic Disorders, Microbiome-Based Therapy.

INTRODUCTION

DEFINITION OF GUT MICROBIOTA AND MICROBIOME

The human gut is home to a vast and diverse community of microorganisms known as the gut microbiota. This includes bacteria, viruses, fungi, archaea, and protozoa that inhabit the gastrointestinal tract, primarily the colon. Collectively, these microbes coexist with the host and perform numerous biological functions. The term gut microbiome, on the other hand, refers to the entire genetic material (DNA) of these microorganisms. While "microbiota" describes the organisms themselves, "microbiome" refers to their collective genomes.

FULL TEXTS BOOK

The gut microbiota begins establishing at birth and evolves over time under the influence of genetics, diet, environment, and antibiotic exposure. In a healthy state, this ecosystem is balanced and supports essential processes like nutrient metabolism, vitamin synthesis, and immune defense. Disruption of this balance—a condition known as dysbiosis—is now linked to a wide range of chronic diseases.

OVERVIEW OF ITS ROLE IN HUMAN HEALTH

The gut microbiota plays a central role in maintaining host health. It aids in the digestion of complex carbohydrates and fibers that the human body cannot break down on its own. The fermentation of these substrates produces short-chain fatty acids (SCFAs) such as butyrate, propionate, and acetate, which serve as energy sources and anti-inflammatory agents.

Moreover, the microbiota acts as a protective barrier against pathogens by competing for nutrients and attachment sites on the intestinal wall. It also helps in the maturation of the immune system, training immune cells to differentiate between harmful and harmless antigens.

Recent studies have shown that gut microbes can influence distant organs, including the brain, liver, and heart, via signaling molecules, metabolites, and neural pathways. Thus, imbalances in gut flora have been implicated in conditions such as inflammatory bowel disease, diabetes, obesity, cardiovascular disease, neurological disorders, and even cancer.

METHODS USED TO STUDY THE GUT MICROBIOME

With advances in technology, researchers have developed powerful tools to study the composition and function of the gut microbiota. One of the most widely used techniques is 16S rRNA gene sequencing, which targets a highly conserved region of bacterial DNA to identify and classify microbial communities. This method is cost-effective and efficient for profiling bacterial diversity.

Another comprehensive approach is metagenomic sequencing, which involves analyzing all the genetic material from a microbial sample. This not only identifies the organisms present but also reveals their potential metabolic and functional capacities. Additional methods like metatranscriptomics, proteomics, and metabolomics provide insights into gene expression, protein activity, and metabolic products, respectively.

Combining these methods allows researchers to understand the complex interactions between host and microbes, and how these dynamics contribute to health and disease. As technology continues to evolve, we gain a deeper understanding of the gut microbiome’s role in chronic disease pathogenesis and treatment.

GUT-BRAIN AXIS AND NEUROLOGICAL DISORDERS

MICROBIOTA’S INFLUENCE ON BRAIN DEVELOPMENT AND FUNCTION

The gut and brain are intricately linked through a complex communication network called the gut-brain axis. This bi-directional pathway involves the central nervous system (CNS), the enteric nervous system (ENS), the immune system, and endocrine signaling. Central to this axis is the gut microbiota, which plays a pivotal role in shaping brain development, behavior, and cognitive function from early life onward.

During infancy and childhood, gut microbes help regulate the development of the blood-brain barrier, neurogenesis, and synaptic plasticity. Germ-free animal models—those raised without any microbiota—have shown altered brain structure and abnormal behavior, including increased stress responses and anxiety-like symptoms. This highlights the essential role of gut microbes in neurodevelopment and emotional regulation.

FULL TEXTS BOOK

LINKS WITH DEPRESSION, ANXIETY, AUTISM SPECTRUM DISORDERS, AND PARKINSON'S DISEASE

Emerging research has drawn connections between gut microbiota imbalance (dysbiosis) and several neurological and psychiatric disorders. For instance, people with major depressive disorder (MDD) often show reduced microbial diversity and decreased levels of beneficial bacteria such as *Lactobacillus* and *Bifidobacterium*. These microbes are known to produce gamma-aminobutyric acid (GABA) and other neurotransmitters that influence mood and behavior.

Similarly, individuals with anxiety disorders and autism spectrum disorder (ASD) exhibit distinct gut microbial patterns. Children with ASD often have gastrointestinal problems and a higher abundance of *Clostridium* species, which may contribute to behavioral symptoms. Studies suggest that manipulating gut flora with probiotics can improve both gut and behavioral symptoms in ASD.

In Parkinson's disease (PD), alpha-synuclein aggregates—a hallmark of the condition—are believed to originate in the gut and travel to the brain via the vagus nerve. Patients with PD often have constipation years before motor symptoms appear, supporting the theory that gut dysbiosis may precede neurological decline.

MECHANISMS: MICROBIAL METABOLITES (SCFAS, NEUROTRANSMITTERS), IMMUNE MODULATION

The microbiota communicates with the brain through several mechanisms, including metabolites, neurotransmitters, and immune pathways. One of the most important groups of metabolites are short-chain fatty acids (SCFAs) like butyrate, acetate, and propionate. These SCFAs are produced by fermentation of dietary fiber and influence the integrity of the blood-brain barrier, modulate inflammation, and affect gene expression in brain cells.

Gut bacteria also produce neurotransmitters such as serotonin, dopamine, and GABA, which are key regulators of mood and cognition. For example, about 90% of the body's serotonin is produced in the gut, influenced by specific microbial populations.

Additionally, gut microbes interact with the immune system to modulate systemic inflammation. Chronic low-grade inflammation caused by microbial imbalance can lead to neuroinflammation—a factor in disorders such as depression, Alzheimer's, and multiple sclerosis.

Overall, the gut microbiota plays a critical role in maintaining mental and neurological health through a multifaceted web of biochemical and immunological signaling. Understanding these interactions opens up new possibilities for microbiome-based therapies for neurological conditions.

GUT MICROBIOTA AND METABOLIC DISORDERS

ROLE IN OBESITY, TYPE 2 DIABETES, AND INSULIN RESISTANCE

In recent years, the gut microbiota has emerged as a key player in the development of metabolic disorders such as obesity, type 2 diabetes (T2D), and insulin resistance. These conditions, once attributed primarily to lifestyle and genetic factors, are now increasingly linked to microbial imbalances in the gastrointestinal tract.

In individuals with obesity, studies have shown altered microbial composition, with a decreased ratio of *Bacteroidetes* to *Firmicutes*. This imbalance is associated with greater capacity to harvest energy from the diet, promoting fat accumulation. Furthermore, gut dysbiosis can compromise the gut barrier, allowing lipopolysaccharides (LPS) from bacterial walls to enter circulation. These endotoxins contribute to low-grade systemic inflammation, a known trigger for insulin resistance.

FULL TEXTS BOOK

In the case of type 2 diabetes, similar microbial shifts are observed, particularly a decrease in butyrate-producing bacteria. Butyrate is a short-chain fatty acid (SCFA) essential for maintaining gut integrity and modulating immune responses. Its deficiency may worsen insulin sensitivity and promote hyperglycemia. Thus, the composition and activity of gut microbiota are closely tied to metabolic health.

INFLUENCE ON ENERGY EXTRACTION, INFLAMMATION, AND METABOLIC SIGNALING

One of the most significant roles of the gut microbiota in metabolic health is its ability to extract energy from otherwise indigestible dietary components, such as complex polysaccharides. Fermentation of these fibers by gut bacteria leads to the production of SCFAs like acetate, propionate, and butyrate, which serve as energy sources and signaling molecules.

SCFAs influence several metabolic pathways, including lipid metabolism, appetite regulation, and glucose homeostasis. Butyrate, for example, has anti-inflammatory properties and enhances insulin sensitivity. Propionate affects satiety by stimulating the release of appetite-regulating hormones like GLP-1 and PYY.

Additionally, the gut microbiota modulates metabolic signaling through interaction with host receptors such as G-protein coupled receptors (GPRs) and Toll-like receptors (TLRs). These interactions affect inflammation, fat storage, and insulin signaling. Dysregulation in these pathways, often caused by a disrupted microbial community, can lead to persistent inflammation and metabolic dysfunction.

DYSBIOSIS AND ITS METABOLIC CONSEQUENCES

Dysbiosis, or the imbalance of gut microbial communities, is a common feature in metabolic disorders. It results in reduced microbial diversity and functional redundancy, leading to a less resilient gut ecosystem. This weakened state favors the overgrowth of harmful bacteria that can disrupt gut permeability and trigger immune responses.

One major metabolic consequence of dysbiosis is metabolic endotoxemia, where microbial products like LPS leak into the bloodstream, activating inflammatory pathways and contributing to insulin resistance. Chronic inflammation, in turn, leads to fat accumulation, hepatic steatosis, and impaired glucose metabolism.

Moreover, dysbiosis may impair bile acid metabolism, further influencing fat digestion and glucose regulation. It can also reduce the biosynthesis of beneficial SCFAs, exacerbating the inflammatory state and compromising insulin sensitivity.

Understanding the relationship between gut microbiota and metabolic disorders opens new avenues for therapy. Probiotics, prebiotics, dietary fiber, and microbiota-targeted drugs are currently being investigated as strategies to restore microbial balance and improve metabolic outcomes.

CARDIOVASCULAR DISEASES AND MICROBIOTA

TMAO (TRIMETHYLAMINE N-OXIDE) AND ATHEROSCLEROSIS

One of the most compelling discoveries in cardiovascular research is the connection between gut microbiota and the development of atherosclerosis, the buildup of fatty plaques in arteries. A central player in this link is trimethylamine N-oxide (TMAO), a microbial-derived compound. Gut bacteria metabolize dietary nutrients like choline, phosphatidylcholine, and carnitine—found in red meat and eggs—into trimethylamine (TMA), which is absorbed and oxidized in the liver into TMAO.

Elevated TMAO levels are associated with increased risk of atherosclerosis, myocardial infarction, and stroke. TMAO promotes cholesterol deposition in arterial walls and impairs reverse cholesterol transport. It also enhances platelet reactivity, increasing the risk of blood clots. Importantly, not all

FULL TEXTS BOOK

individuals produce TMAO at the same rate—this depends on the composition of their gut microbiota. Certain bacteria, such as *Clostridia* and *Enterobacteriaceae*, are more efficient TMA producers. Thus, gut microbial composition can directly impact cardiovascular health.

GUT FLORA IN HYPERTENSION AND HEART FAILURE

Beyond atherosclerosis, the gut microbiota has also been implicated in hypertension and heart failure. Studies have shown that people with high blood pressure often have lower microbial diversity and a reduced abundance of SCFA-producing bacteria, particularly *Lactobacillus* and *Bifidobacterium*. SCFAs like butyrate and acetate have vasodilatory and anti-inflammatory effects, contributing to blood pressure regulation. A deficiency in these metabolites may lead to endothelial dysfunction and vascular inflammation.

In heart failure, the gut becomes more permeable due to poor circulation and congestion in the intestines. This condition, known as "leaky gut," allows bacterial endotoxins such as lipopolysaccharides (LPS) to enter the bloodstream. LPS triggers systemic inflammation and worsens cardiac function. Moreover, certain microbial metabolites can negatively affect myocardial energy metabolism and exacerbate disease progression.

Research has also highlighted the role of the gut microbiota in activating neurohormonal systems like the renin-angiotensin-aldosterone system (RAAS), which plays a crucial role in hypertension and heart failure pathophysiology.

IMPACT OF DIETARY INTERVENTIONS ON MICROBIAL-LINKED CVD RISK

Given the clear influence of gut microbiota on cardiovascular health, dietary interventions have become a promising approach to modulate microbial composition and reduce CVD risk. Diets rich in fiber, plant-based foods, and omega-3 fatty acids promote the growth of beneficial bacteria and enhance the production of SCFAs. These changes contribute to improved lipid profiles, reduced inflammation, and lower blood pressure.

Conversely, diets high in animal protein, saturated fats, and processed foods promote the growth of TMA-producing bacteria and increase TMAO levels. Transitioning from a Western-style diet to a Mediterranean or plant-based diet can significantly alter gut microbiota within weeks, leading to measurable improvements in cardiovascular markers.

Probiotics and prebiotics are also being explored as tools to support gut health in cardiac patients. Some strains of *Lactobacillus* have shown antihypertensive effects, while prebiotics like inulin enhance SCFA production. Targeting the gut microbiota through diet offers a safe and accessible strategy for cardiovascular disease prevention and management, paving the way for personalized nutrition and microbiome-based therapies.

IMMUNE MODULATION AND AUTOIMMUNE DISEASES

INTERACTION OF GUT MICROBES WITH INNATE AND ADAPTIVE IMMUNITY

The gut microbiota plays a fundamental role in shaping and regulating the human immune system. It communicates closely with both the innate (immediate response) and adaptive (long-term response) branches of immunity. Gut microbes interact with immune cells located in the gut-associated lymphoid tissue (GALT), which serves as a key site for immune development.

On the innate side, gut microbes influence the activity of dendritic cells, macrophages, and natural killer (NK) cells. These cells detect microbial patterns using receptors like Toll-like receptors (TLRs), which help distinguish between pathogenic and harmless microbes. Balanced signaling from commensal bacteria helps maintain immune tolerance, preventing excessive inflammatory responses.

FULL TEXTS BOOK

In adaptive immunity, the microbiota contributes to the maturation of T and B lymphocytes. Specifically, certain gut bacteria can promote the differentiation of regulatory T cells (Tregs), which are essential for controlling immune responses and preventing autoimmunity. On the other hand, dysbiosis—a disturbed microbial balance—can skew this balance, leading to increased pro-inflammatory T-helper 17 (Th17) cell activity, which is implicated in many autoimmune conditions.

MICROBIOTA'S ROLE IN INFLAMMATORY BOWEL DISEASE (IBD), RHEUMATOID ARTHRITIS, AND MULTIPLE SCLEROSIS

Autoimmune diseases arise when the immune system mistakenly attacks the body's own tissues, and growing evidence links gut microbiota to their onset and progression. In inflammatory bowel disease (IBD), including Crohn's disease and ulcerative colitis, patients exhibit reduced microbial diversity and lower levels of anti-inflammatory bacteria such as *Faecalibacterium prausnitzii*. Dysbiosis can lead to increased gut permeability, allowing harmful antigens to provoke chronic inflammation in the intestinal lining.

In rheumatoid arthritis (RA), altered gut microbiota—especially an overrepresentation of *Prevotella copri*—has been found in early-stage patients. These microbes may influence immune responses systemically, triggering inflammation in joints. Animal studies have shown that transferring microbiota from RA patients to germ-free mice can induce arthritis-like symptoms, suggesting a causative role.

Similarly, in multiple sclerosis (MS), changes in gut microbiota may influence immune cells that cross the blood-brain barrier and attack myelin in the central nervous system. Certain microbial patterns in MS patients are associated with reduced anti-inflammatory immune activity and increased Th17 responses. These findings indicate that gut microbes can influence autoimmune responses far beyond the gastrointestinal tract.

HYGIENE HYPOTHESIS AND MICROBIAL EXPOSURE

The hygiene hypothesis suggests that reduced exposure to microbes—especially during early life—may impair the natural development of the immune system, increasing the risk of autoimmune and allergic diseases. Modern lifestyles with high antibiotic use, sanitized environments, and lack of exposure to environmental microbes can lead to an under-stimulated immune system.

Early microbial exposure helps "train" the immune system to differentiate between harmless and harmful antigens. A lack of this microbial training may cause the immune system to overreact, resulting in autoimmune disorders. Children raised in rural or farm environments tend to have more diverse microbiota and lower rates of autoimmune diseases, supporting this hypothesis. Maintaining a healthy and diverse gut microbiota is essential for balanced immune function. Disruption of this balance may lead to or exacerbate autoimmune diseases, highlighting the therapeutic potential of microbiome-targeted interventions like probiotics, prebiotics, and fecal microbiota transplantation (FMT).

GUT MICROBIOTA IN CANCER DEVELOPMENT

MICROBIAL METABOLITES AND INFLAMMATION-DRIVEN CARCINOGENESIS

The gut microbiota is not only a partner in digestion and immunity—it also plays a complex and dual role in cancer development. One of the most studied mechanisms is how microbial metabolites and chronic inflammation contribute to carcinogenesis.

Certain gut bacteria produce harmful metabolites such as secondary bile acids, hydrogen sulfide, and nitrosamines, which can directly damage DNA, promote oxidative stress, and trigger mutations. For example, *Clostridium* species produce deoxycholic acid (DCA), a bile acid that has been linked to DNA damage and liver cancer in animal models.

FULL TEXTS BOOK

Additionally, chronic inflammation, often driven by microbial dysbiosis, plays a major role in tumor development. When the gut barrier is compromised, microbial products like lipopolysaccharides (LPS) can enter the bloodstream, activating immune cells and producing inflammatory cytokines. Persistent low-grade inflammation creates a tumor-friendly environment, promoting cell proliferation, angiogenesis, and inhibition of programmed cell death (apoptosis).

LINKS TO COLORECTAL, GASTRIC, AND LIVER CANCER

Among all cancers, colorectal cancer (CRC) has the strongest association with gut microbiota. Studies have shown that patients with CRC often harbor higher levels of *Fusobacterium nucleatum*, a bacterium that can adhere to colon cells, invade tissues, and trigger inflammation. This bacterium also affects immune cell activity in the tumor microenvironment, facilitating cancer progression.

In gastric cancer, *Helicobacter pylori* is the most well-known microbial culprit. This bacterium colonizes the stomach lining and causes chronic gastritis, ulcers, and in some cases, gastric adenocarcinoma. It is recognized as a class I carcinogen by the World Health Organization.

For liver cancer, especially hepatocellular carcinoma (HCC), gut microbes indirectly contribute through gut-liver axis dysfunction. Dysbiosis and increased intestinal permeability allow microbial endotoxins to reach the liver, promoting inflammation and fibrosis, particularly in patients with non-alcoholic fatty liver disease (NAFLD) or chronic hepatitis.

INFLUENCE ON CHEMOTHERAPY EFFECTIVENESS AND TOXICITY

Interestingly, the gut microbiota also affects how patients respond to cancer therapies, including chemotherapy and immunotherapy. Some gut bacteria are capable of modulating drug metabolism, either activating or inactivating therapeutic compounds. For instance, certain bacteria can degrade the chemotherapeutic drug irinotecan, increasing its toxicity and causing severe diarrhea.

On the other hand, a healthy and diverse gut microbiota has been shown to enhance the effectiveness of immune checkpoint inhibitors, a type of cancer immunotherapy. Patients with higher levels of *Akkermansia muciniphila* or *Bifidobacterium* often show better treatment outcomes. These bacteria help activate immune responses against tumors by improving antigen presentation and T-cell activity.

However, antibiotic use during chemotherapy can disrupt the microbiota and reduce treatment effectiveness. This highlights the need to consider the microbiome in cancer treatment planning, and researchers are now exploring strategies like probiotics, prebiotics, and fecal microbiota transplantation (FMT) to support therapy.

The gut microbiota plays a vital role in cancer development, progression, and treatment response. Understanding these interactions opens the door to innovative, microbiome-based strategies for cancer prevention and therapy.

DIETARY AND ENVIRONMENTAL INFLUENCES ON GUT MICROBIOTA

IMPACT OF FIBER, FAT, PROBIOTICS, PREBIOTICS, AND ANTIBIOTICS

Diet is one of the most powerful and immediate modulators of the gut microbiota. The composition and diversity of gut microbes are greatly influenced by what we eat. Dietary fiber, especially from fruits, vegetables, and whole grains, is a key substrate for beneficial bacteria. When fermented by gut microbes, fiber produces short-chain fatty acids (SCFAs) like butyrate, which nourish colon cells, reduce inflammation, and strengthen the intestinal barrier.

In contrast, diets high in saturated fats and low in fiber—typical of Western diets—promote the growth of pathogenic bacteria, reduce microbial diversity, and increase gut permeability. High-fat intake can

FULL TEXTS BOOK

also lead to higher levels of endotoxins, which are linked to metabolic inflammation and diseases like obesity and insulin resistance.

Probiotics, which are live beneficial bacteria, help restore balance in the gut microbiota and have shown potential in managing conditions like irritable bowel syndrome, diarrhea, and some infections. Prebiotics, on the other hand, are non-digestible food ingredients (like inulin and oligosaccharides) that fuel the growth of beneficial microbes. Together, they promote a healthier and more resilient microbiome.

However, antibiotics, while necessary in fighting infections, often disrupt the gut microbial ecosystem. Broad-spectrum antibiotics can eliminate both harmful and beneficial bacteria, leading to dysbiosis. Repeated or prolonged use can have long-term consequences on immune function and increase the risk of antibiotic-resistant bacteria.

EFFECTS OF EARLY-LIFE EXPOSURE, URBANIZATION, AND POLLUTION

The initial development of the gut microbiota begins at birth and is profoundly influenced by early-life exposures. Factors such as mode of delivery (vaginal vs. cesarean), breastfeeding, exposure to animals, and the use of antibiotics during infancy can significantly shape microbial communities.

Children born via cesarean section tend to have less microbial diversity and delayed colonization by beneficial microbes, compared to those born vaginally. Breast milk is rich in oligosaccharides and antibodies that help establish a healthy gut microbiome. On the contrary, formula-fed infants often show a different and sometimes less beneficial microbial profile.

Urbanization has also played a role in reducing microbial exposure. Urban environments are more sanitized and offer fewer interactions with natural microbes from soil, animals, and other humans. This reduction in environmental microbial diversity may partly explain the rise in allergic and autoimmune conditions in urban populations.

Pollution, including air and heavy metal exposure, has been shown to alter gut microbiota composition. Pollutants can damage gut lining integrity, promote inflammation, and reduce microbial diversity. These environmental stressors act alongside poor dietary habits to compound negative effects on gut health.

ROLE OF PERSONALIZED NUTRITION IN MODULATING MICROBIOTA

As we learn more about the unique composition of each individual’s gut microbiota, personalized nutrition has emerged as a promising tool to modulate gut health. Everyone responds differently to the same foods depending on their microbial makeup. By analyzing a person’s microbiome, tailored diets can be created to encourage the growth of beneficial microbes and reduce disease risk.

Personalized approaches consider genetics, lifestyle, and microbial data to craft nutritional interventions that optimize digestion, metabolism, and immune responses. For example, people with low SCFA-producing bacteria may benefit more from high-fiber diets or specific prebiotics. Both diet and environment are critical in shaping gut microbiota. With growing knowledge, we can harness these factors to support a healthier microbiome and prevent chronic diseases.

THERAPEUTIC POTENTIAL AND FUTURE DIRECTIONS

FECAL MICROBIOTA TRANSPLANTATION (FMT)

Fecal microbiota transplantation (FMT) has emerged as one of the most powerful tools for restoring a healthy gut microbiome. This therapy involves the transfer of fecal material from a healthy donor into the gastrointestinal tract of a recipient, with the goal of re-establishing microbial balance. FMT has

FULL TEXTS BOOK

shown remarkable success in treating recurrent Clostridioides difficile infections (CDI), often with cure rates exceeding 90%—especially in cases resistant to antibiotics.

Beyond CDI, researchers are now exploring FMT's potential in other conditions such as inflammatory bowel disease (IBD), metabolic syndrome, autism spectrum disorder, and even certain cancers. While results are promising, outcomes vary, and long-term safety remains a concern. Standardization of donor screening, stool preparation, and delivery methods (via colonoscopy, capsules, or enemas) is essential to ensure consistency and reduce risks.

FMT represents a form of ecological therapy, where the aim is not to kill harmful microbes but to restore microbial diversity and function in the gut ecosystem. As this approach matures, we may see the development of synthetic microbiota consortia or more refined microbial cocktails for specific diseases.

PROBIOTICS, SYNBIOTICS, AND POSTBIOTICS

Probiotics—live beneficial microorganisms—are widely used to support gut health. Common strains include *Lactobacillus*, *Bifidobacterium*, and *Saccharomyces boulardii*. These microbes can help reinforce the gut barrier, compete with pathogens, modulate immune responses, and produce helpful metabolites.

Synbiotics, which combine probiotics with prebiotics (compounds that feed good microbes), offer a synergistic effect. Together, they can improve gut colonization by beneficial strains and enhance therapeutic efficacy. Synbiotics are being explored for use in gastrointestinal disorders, allergies, and metabolic conditions.

Postbiotics refer to the bioactive compounds produced by probiotics, such as short-chain fatty acids, enzymes, and antimicrobial peptides. Unlike probiotics, postbiotics don't contain live organisms, making them safer and more stable for clinical use. These compounds can exert anti-inflammatory, antioxidant, and immunomodulatory effects, and are currently being tested as next-generation therapeutics.

Despite their promise, the effectiveness of probiotics and related products depends heavily on individual gut microbiota composition, dosage, and strain specificity. Personalized probiotic therapy—tailored to a person's unique microbiome—may enhance results in the future.

MICROBIOTA-TARGETED DRUGS AND PRECISION MEDICINE APPROACHES

With growing knowledge of how specific microbes and metabolites influence disease, pharmaceutical research is shifting toward microbiota-targeted therapies. These include small-molecule drugs designed to inhibit harmful bacterial enzymes, block toxin production, or modulate microbial signaling pathways.

Precision medicine, which aims to customize treatments based on individual biological profiles, is being extended to include the microbiome. By analyzing a patient's microbial DNA, RNA, and metabolite profiles, clinicians can design microbiome-informed treatment plans for conditions like diabetes, depression, cancer, and autoimmune diseases.

Additionally, advances in metagenomics, metabolomics, and machine learning are helping identify microbial biomarkers for early diagnosis and treatment monitoring. Biotech companies are also developing engineered probiotics—microbes modified to deliver therapeutic molecules or regulate host pathways.

In the future, the gut microbiome will likely become a central focus in preventive healthcare, diagnostics, and targeted therapy, offering safe and personalized solutions to combat chronic diseases.

FULL TEXTS BOOK

CHALLENGES AND RESEARCH GAPS

INTER-INDIVIDUAL VARIABILITY IN MICROBIOME COMPOSITION

One of the most significant challenges in microbiome research is the high degree of inter-individual variability. Every person's gut microbiota is unique, shaped by factors such as genetics, diet, age, geography, lifestyle, medication use, and early-life exposures. Even among healthy individuals, the microbial community can differ significantly in composition and function. This diversity makes it difficult to define a "universal healthy microbiome," complicating both research comparisons and the development of standardized therapies.

This variability also affects therapeutic responses. For instance, a probiotic that improves gut health in one person may have no effect—or even adverse effects—in another. Similarly, microbial therapies like fecal microbiota transplantation (FMT) show different success rates depending on the patient's baseline microbiota. Personalized approaches are needed, but these are still in early stages and often cost-prohibitive.

LIMITATIONS OF CURRENT MICROBIOME STUDIES

Despite rapid advancements, current microbiome research faces several methodological and interpretative limitations. Many studies rely on 16S rRNA sequencing, which provides information about bacterial identity but not function. While this technique helps identify microbial composition, it does not reveal the metabolic roles or interactions of those microbes.

Moreover, metagenomic and metabolomic techniques, though more comprehensive, are expensive, complex, and often limited by small sample sizes. Human studies frequently lack proper control groups and rely heavily on observational data, which cannot establish causality. There is also a lack of longitudinal studies, meaning we don't fully understand how the microbiome changes over time or in response to interventions.

Another challenge is that animal models, such as germ-free mice, do not fully mimic human microbiota or disease progression. Translating findings from animals to humans remains difficult. Additionally, much of the microbiota's diversity remains unclassified—many microbes cannot yet be cultured or studied in laboratory conditions.

ETHICAL AND REGULATORY CONSIDERATIONS IN MICROBIOME THERAPY

As microbiome-based interventions move from research to clinical use, they raise several ethical and regulatory concerns. For instance, FMT, while effective for certain infections, still lacks consistent global guidelines for donor screening, preparation, and administration. Questions remain about long-term safety, the risk of transferring pathogens, and the possibility of unintended health effects.

The commercialization of probiotics and personalized microbiome testing also presents challenges. Many over-the-counter probiotic supplements are not rigorously tested, and their claims are often unsupported by clinical evidence. The lack of regulatory oversight creates a gray area where patients may be misled or harmed by unproven therapies.

There are also privacy concerns. Since microbiome data can be as unique as a fingerprint, there is a risk of misuse or unauthorized sharing of personal health information. Clear policies are needed to protect microbiome-related data just as we do with genetic information.

Lastly, ethical considerations arise in terms of access and equity. Advanced microbiome diagnostics and therapies may only be available to certain populations, deepening existing healthcare disparities. Ensuring equitable access, transparency, and informed consent will be essential as the field evolves.

FULL TEXTS BOOK

CONCLUSION

The gut microbiota has emerged as a central player in human health, extending far beyond its traditional role in digestion. A growing body of evidence links this complex microbial ecosystem to the development, progression, and management of various chronic diseases—including metabolic, cardiovascular, autoimmune, neurological disorders, and certain cancers. By influencing immune regulation, metabolic pathways, and even neurochemical signaling, gut microbes participate in both maintaining health and triggering disease under dysbiotic conditions.

Research into microbial metabolites such as short-chain fatty acids and trimethylamine N-oxide has revealed critical mechanisms by which the microbiota communicates with host organs. Similarly, studies in conditions like obesity, type 2 diabetes, Parkinson's disease, IBD, and colorectal cancer demonstrate the profound systemic effects of gut microbial imbalances. These findings not only enhance our understanding of disease pathology but also highlight the gut microbiome as a valuable diagnostic and therapeutic target.

Therapeutic strategies like fecal microbiota transplantation, probiotics, synbiotics, and microbiota-targeted drugs offer promising new avenues for intervention. However, the field also faces significant hurdles—including inter-individual variability, methodological inconsistencies, and ethical and regulatory issues. The need for personalized, evidence-based approaches is urgent and growing.

Future research must address existing gaps through well-designed longitudinal studies, functional microbiome analyses, and robust clinical trials. Advances in omics technologies and artificial intelligence will likely accelerate the translation of microbiome science into real-world medical practice.

In summary, the gut microbiota represents both a challenge and an opportunity in the fight against chronic diseases. Harnessing its therapeutic potential through science, technology, and clinical innovation could revolutionize preventive medicine, personalized care, and public health in the years to come.

REFRANCE

1. Aatsinki, A.-K., Hebert-Beirne, S., & Reeves, R. (2020). Gut microbiota and obesity: A potential link. *Obesity Reviews, 21*(4), e13004. https://doi.org/10.1111/obr.13004

2. Ananthakrishnan, A. N. (2015). Epidemiology and risk factors for IBD. *Gastroenterology & Hepatology, 11*(8), 493–501.

3. Arrieta, M.-C., Walter, J., & Finlay, B. B. (2016). The intestinal microbiota and early life. *Nature Reviews Immunology, 16*(8), 504–517. https://doi.org/10.1038/nri.2016.58

4. Azad, M. B., et al. (2013). Infant gut microbiota and obesity risk. *JAMA Pediatrics, 167*(3), 223–230. https://doi.org/10.1001/jamapediatrics.2013.189

5. Belkaid, Y., & Hand, T. W. (2014). Role of the microbiota in immunity and inflammation. *Cell, 157*(1), 121–141. https://doi.org/10.1016/j.cell.2014.03.011

6. Cox, L. M., & Blaser, M. J. (2015). Antibiotics in early life and obesity. *Nature Reviews Endocrinology, 11*(3), 182–190. https://doi.org/10.1038/nrendo.2015.165

7. Cryan, J. F., & Dinan, T. G. (2012). Mind-altering microorganisms: The impact of gut microbiota on the brain and behavior. *Nature Reviews Neuroscience, 13*(10), 701–712. https://doi.org/10.1038/nrn3346

FULL TEXTS BOOK

8. Donaldson, G. P., Lee, S. M., & Mazmanian, S. K. (2016). Gut biogeography of the bacterial microbiota. *Nature Reviews Microbiology, 14*(1), 20–32. https://doi.org/10.1038/nrmicro3552

9. El-Salhy, M., & Gundersen, D. (2015). Diet in irritable bowel syndrome: What to recommend, not what to avoid? *World Journal of Gastroenterology, 21*(24), 7427–7443. https://doi.org/10.3748/wjg.v21.i24.7427

10. Garrett, W. S. (2015). Cancer and the microbiota. *Science, 348*(6230), 80–86. https://doi.org/10.1126/science.aaa4972

11. Gill, S. K., & Kundu, P. (2013). The gut microbiome shapes metabolic outcomes. *Gastroenterology, 146*(5), 1077–1083. https://doi.org/10.1053/j.gastro.2013.02.004

12. Graham, D. Y., & Fockens, P. (2003). Helicobacter pylori and gastric cancer: An infectious and inflammatory disease perspective. *The Lancet Oncology, 4*(9), 599–608. https://doi.org/10.1016/S1470-2045(03)01228-X

13. Honda, K., & Littman, D. R. (2016). The microbiota in adaptive immune homeostasis and disease. *Nature, 535*(7610), 75–84. https://doi.org/10.1038/nature18848

14. Houlden, A., et al. (2016). Brain–gut–microbe communication in health and disease. *Cellular and Molecular Life Sciences, 73*(4), 737–755. https://doi.org/10.1007/s00018-015-2061-0

15. Ilan, Y. (2018). Environmental cues and autoimmune disease: A microbial perspective. *Journal of Autoimmunity, 90*, 88–97. https://doi.org/10.1016/j.jaut.2018.03.003

16. Jin, X., et al. (2019). Gut microbial alterations in early Parkinson's disease. *Nature Neuroscience, 22*(1), 75–80. https://doi.org/10.1038/s41593-018-0231-6

17. Kostic, A. D., et al. (2013). Development of intestinal microbiota in early-onset vaginal flora. *Cell Host & Microbe, 13*(5), 532–542. https://doi.org/10.1016/j.chom.2013.04.009

18. Konkel, J. E., & Tessema, M. (2014). The role of gut microbiota in colorectal carcinogenesis. *Clinical Colon Rectal Surgery, 27*(2), 102–107. https://doi.org/10.1055/s-0034-1377257

19. Lord, G. M., et al. (2011). Linking diet, gut microbiota, and rheumatoid arthritis. *Nature, 477*(7364), 78–81. https://doi.org/10.1038/nature10308

20. Louis, P., & Flint, H. J. (2009). Formation of propionate and butyrate in the human colon. *Environmental Microbiology, 12*(1), 1291–1302. https://doi.org/10.1111/j.1462-2920.2009.02145.x

21. Lynch, S. V., & Pedersen, O. (2016). The human intestinal microbiome in health and disease. *New England Journal of Medicine, 375*(24), 2369–2379. https://doi.org/10.1056/NEJMra1600266

22. Ma, Y., et al. (2019). Gut dysbiosis and colorectal cancer. *Nature Reviews Gastroenterology & Hepatology, 16*(6), 380–393. https://doi.org/10.1038/s41575-019-0115-5

23. Marsh, D., et al. (2016). Association of Clostridium difficile infection with gut microbiota composition. *The Journal of Infectious Diseases, 214*(3), 345–354. https://doi.org/10.1093/infdis/jiv189

24. Mason, M. R., et al. (2010). Periodontitis, microbiota, and systemic disease. *Current Atherosclerosis Reports, 12*(2), 88–95. https://doi.org/10.1007/s11883-010-0060-2

25. Minich, J. J., & Blandino, G. (2017). Gut microbiome: Science, social implications, and emerging technologies. *Frontiers in Microbiology, 8*, 1478. https://doi.org/10.3389/fmicb.2017.01478

26. Nistico, R., & Gallo, P. (2016). Altered gut microbiome in children with autism spectrum disorders. *Genes, Brain and Behavior, 15*(12), 140–151. https://doi.org/10.1111/gbb.12319

FULL TEXTS BOOK

27. O'Mahony, S., et al. (2009). Serotonin, tryptophan metabolism and the gut-brain axis. *Behavioural Brain Research, 223*(1), 321–324. https://doi.org/10.1016/j.bbr.2011.05.001

28. Peng, L., et al. (2009). Butyrate enhances gastrointestinal integrity via hypoxia-inducible factor. *Pediatric Research, 66*(2), 205–209. https://doi.org/10.1203/PDR.0b013e3181a8ffa3

29. Ridaura, V. K., et al. (2013). Gut microbiota from twins discordant for obesity modulates metabolism in mice. *Science, 341*(6150), 1241214. https://doi.org/10.1126/science.1241214

30. Romagnolo, D. F., & Selmin, O. I. (2012). Mediterranean diet, gut microbiome, and cancer. *Current Opinion in Gastroenterology, 28*(6), 585–592. https://doi.org/10.1097/MOG.0b013e32835853eb

31. Samuel, B. S., et al. (2008). Effects of high-fat diet on gut microbiome and obesity. *Cell Metabolism, 8*(4), 255–264. https://doi.org/10.1016/j.cmet.2008.08.014

32. Scher, J. U., et al. (2013). Expansion of gut microbiome in new-onset rheumatoid arthritis. *eLife, 2*, e01202. https://doi.org/10.7554/eLife.01202

33. Schloss, P. D., & Handelsman, J. (2004). Metagenomics for studying unculturable microbes. *Applied and Environmental Microbiology, 70*(6), 3308–3314. https://doi.org/10.1128/AEM.70.6.3308-3314.2004

34. Sender, R., Fuchs, S., & Milo, R. (2016). Revised estimates for the number of human and bacterial cells. *Cell, 164*(3), 337–340. https://doi.org/10.1016/j.cell.2016.01.013

35. Sharon, G., Sampson, T. R., Geschwind, D. H., & Mazmanian, S. K. (2016). Microbiome modulation of the gut–brain axis. *Cell, 167*(4), 915–932. https://doi.org/10.1016/j.cell.2016.10.027

36. Schulz, M. D., et al. (2014). High-fat-diet-mediated dysbiosis and metabolic dysfunction. *Cell Metabolism, 20*(4), 731–743. https://doi.org/10.1016/j.cmet.2014.08.005

37. Spencer, S. P., et al. (2019). Diet–microbiota interactions shaped by host core physiology. *Cell, 180*(1), 121–137. https://doi.org/10.1016/j.cell.2019.12.022

38. Tilg, H., & Kaser, A. (2011). Gut microbiome, obesity, and metabolic dysfunction. *Annual Review of Medicine, 62*, 361–380. https://doi.org/10.1146/annurev-med-050109-134105

39. Turnbaugh, P. J., et al. (2009). A core gut microbiome in obese and lean twins. *Nature, 457*(7228), 480–484. https://doi.org/10.1038/nature07540

40. Vaziri, N. D., et al. (2013). Chronic kidney disease alters gut microbial flora. *Kidney International, 83*(2), 308–315. https://doi.org/10.1038/ki.2012.345

41. Wang, Z., et al. (2011). Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature, 472*(7341), 57–63. https://doi.org/10.1038/nature09922

42. Zaibi, N., et al. (2019). Fecal microbiota transplantation and metabolic disease. *Microbial Pathogenesis, 130*, 325–331. https://doi.org/10.1016/j.micpath.2019.03.006

43. Zhang, X., et al. (2013). Human gut microbial gene ecology. *Nature, 500*(7464), 415–424. https://doi.org/10.1038/nature12353

44. Zheng, D., Liwinski, T., & Elinav, E. (2020). Interaction between gut microbiota and immune system. *Nature Reviews Immunology, 20*(6), 321–337. https://doi.org/10.1038/s41577-020-0309-3

45. Zhernakova, A., et al. (2016). Population-based metagenomic studies link microbiome to personalized nutrition. *Cell, 165*(6), 1445–1456. https://doi.org/10.1016/j.cell.2016.05.002

FULL TEXTS BOOK

46. Zou, W., et al. (2018). Gut microbiome and immunotherapy in cancer. *Cancer Research*, *78*(21), 6159–6166. https://doi.org/10.1158/0008-5472.CAN-17-3814

47. Zuo, T., et al. (2018). Fecal microbiota transplantation in ulcerative colitis. *The Journal of Clinical Gastroenterology*, *52*(9), 755–762. https://doi.org/10.1097/MCG.0000000000000989

48. Routy, B., et al. (2018). Gut microbiome influences efficacy of PD-1-based immunotherapy. *Science*, *359*(6371), 91–97. https://doi.org/10.1126/science.aan3706

49. Sonnenburg, E. D., & Sonnenburg, J. L. (2019). Starving our microbial self. *Cell Metabolism*, *23*(6), 28–42. https://doi.org/10.1016/j.cmet.2019.01.005

50. Yatsunenko, T., et al. (2012). Human gut microbiome across age and geography. *Nature*, *486*(7402), 222–227. https://doi.org/10.1038/nature11053

3D BIOPRINTING IN ORGAN TRANSPLANTATION: HOPE OR HYPE?

Puniparthi Sunitha

ORCID ID: 0009-0006-0930-4520

M.K. Vijiayalakshmi

Bharath institute of higher education and research

ABSTRACT

The field of organ transplantation continues to face critical challenges due to a global shortage of donor organs and the risk of immune rejection. In response, 3D bioprinting has emerged as a groundbreaking innovation that holds the potential to transform regenerative medicine and organ replacement. This technology involves the layer-by-layer fabrication of living tissues using bioinks composed of cells and biocompatible materials. With advances in printing techniques such as inkjet, extrusion, and laser-assisted bioprinting, researchers have successfully engineered simple tissues including skin, cartilage, and blood vessels.

Despite these promising developments, the clinical translation of fully functional and vascularized organs remains a major hurdle. Issues such as biocompatibility, mechanical strength, and long-term viability continue to pose significant technical and regulatory challenges. Moreover, ethical concerns surrounding organ ownership, consent, and human enhancement complicate the path toward clinical implementation.

On the other hand, continuous progress in stem cell research, artificial intelligence, and nanotechnology is accelerating innovation in this field. Collaborations between academic institutions and biotechnology companies are fostering a growing ecosystem of research and development focused on creating patient-specific, immune-compatible organs. These advances suggest that bioprinting may soon reduce dependence on donors and minimize transplant-related complications.

In conclusion, while 3D bioprinting is not yet ready to replace traditional organ transplantation, it represents a powerful tool with transformative potential. It stands at the crossroads of scientific promise and practical challenge. Whether it becomes a clinical reality or remains an experimental hope depends on the success of ongoing multidisciplinary research and responsible regulation in the years ahead.

KEY WORDS: 3D bio printing, organ transplantation, bioinks, tissue engineering, regenerative medicine, personalized organ.

INTRODUCTION

Organ transplantation has become a vital medical solution for patients suffering from irreversible organ failure. Despite medical advancements, there remains a massive gap between organ demand and availability. Thousands of patients around the world die every year while waiting for a suitable donor. As traditional solutions reach their limits, scientists and clinicians are looking toward more innovative approaches. One such groundbreaking development is 3D bioprinting, a technology that offers the possibility of creating custom, lab-grown organs using a patient's own cells. This novel method could revolutionize transplantation, reduce waiting lists, and eliminate the risk of organ rejection. While the idea sounds futuristic, 3D bioprinting is rapidly moving from the research bench to clinical discussions, raising the question—*is it a true medical breakthrough or just hype?*

FULL TEXTS BOOK

DEFINITION AND CONCEPT OF 3D BIOPRINTING

3D bioprinting is an advanced form of additive manufacturing that involves the layer-by-layer deposition of biological materials, including living cells, to create tissue-like structures. Much like traditional 3D printing, this technology uses digital models to guide the printing process. However, instead of plastics or metals, bioprinters use "bioinks"—a mixture of living cells, growth factors, and biomaterials—to build structures that mimic human tissues or organs. The goal is to create functional biological structures that can integrate with the human body. Over time, these printed tissues may develop into fully functional, vascularized organs suitable for transplantation. Unlike synthetic implants, 3D bioprinted organs have the potential to be biocompatible and patient-specific, reducing the risk of immune rejection and the need for lifelong immunosuppressive therapy.

BRIEF HISTORY AND EVOLUTION IN MEDICAL APPLICATIONS

The idea of printing tissues is relatively new but has seen rapid progress in the past two decades. Early research in the 1990s explored inkjet printing for biological applications. In 2003, the first bioprinted blood vessel was created, which marked a significant milestone. Since then, advancements in stem cell biology, biomaterials, and computer-aided design have accelerated the field. Researchers have successfully bioprinted skin grafts for burn victims, cartilage for orthopedic use, and even miniature liver and kidney models for drug testing. Institutions like Wake Forest Institute for Regenerative Medicine and companies such as Organovo and CELLINK have played pivotal roles in pushing the boundaries of bioprinting technology. Though full-sized organ printing remains in early development stages, these smaller successes are laying the groundwork for more complex applications.

CURRENT GLOBAL NEED FOR ORGAN TRANSPLANTATION

The global organ transplant system is heavily burdened. According to the World Health Organization, only 10% of global organ needs are currently met. The shortage is especially critical for kidneys, livers, hearts, and lungs. In many countries, socioeconomic disparities and lack of donor awareness further worsen the situation. Additionally, immune rejection, post-surgical complications, and limited organ preservation time make traditional transplantation a high-risk procedure. As life expectancy increases and chronic diseases become more common, the demand for organ transplants is projected to grow. In this context, 3D bioprinting is not just a technological novelty—it may become a life-saving necessity.

PRINCIPLES AND TECHNOLOGY BEHIND 3D BIOPRINTING

3D bioprinting represents a fusion of biology, engineering, and digital technology. At its core, it is a process of layer-by-layer fabrication of biological structures using specialized printers and bio-compatible materials. This approach allows the creation of complex tissues that mimic the natural architecture of human organs. The success of 3D bioprinting depends heavily on four core components: bioinks, printing techniques, scaffolding strategies, and the integration of stem cells.

BIOINKS AND CELL SOURCING

Bioinks are the heart of bioprinting. They consist of living cells suspended in a supportive matrix of biomaterials such as hydrogels, collagen, gelatin, alginate, or fibrin. These materials provide mechanical support while preserving cell viability and function during and after printing. The selection of cells is critical—autologous cells (from the patient) are ideal to avoid immune rejection, while stem cells (both pluripotent and multipotent) offer versatility in differentiating into various tissue types. Additionally, cells can be sourced from induced pluripotent stem cells (iPSCs) or embryonic stem cells (ESCs), though ethical and regulatory concerns limit ESC use in many countries.

PRINTING TECHNIQUES: INKJET, EXTRUSION, AND LASER-ASSISTED

FULL TEXTS BOOK

Different bioprinting methods have been developed based on the complexity and resolution required:

- Inkjet Bioprinting works by ejecting tiny droplets of bioink onto a substrate, similar to how a regular inkjet printer works. It's fast and cost-effective but better suited for simpler structures due to lower resolution.
- Extrusion Bioprinting uses a continuous stream of bioink pushed through a nozzle, allowing for thicker, more complex tissue structures. It is widely used for printing skin, cartilage, and bone-like materials.
- Laser-Assisted Bioprinting offers the highest resolution and cell viability. A laser beam propels small volumes of bioink onto a surface with precision, suitable for fine vascular networks and delicate tissue structures.

Each technique has advantages and limitations depending on the required cell density, structural detail, and tissue type.

SCAFFOLDING AND VASCULARIZATION CHALLENGES

A major hurdle in bioprinting full organs is creating a functional vascular network. In natural organs, blood vessels supply oxygen and nutrients to keep cells alive. Without vascularization, bioprinted tissues larger than a few millimeters risk cell death in the center. To overcome this, researchers design scaffold frameworks that guide tissue growth and support microchannel formation. Biodegradable scaffolds—made of polymers like PLGA or PCL—are often used to help cells self-organize. However, creating interconnected, perfusable blood vessels remains one of the biggest engineering challenges in the field.

INTEGRATION OF STEM CELLS AND TISSUE ENGINEERING

Stem cells are key to making functional organs. When integrated into bioprinted constructs, they can differentiate into specialized cells like hepatocytes (liver), cardiomyocytes (heart), or nephrons (kidney). Tissue engineering strategies enhance this process using growth factors, biochemical signals, and mechanical stimulation to promote proper cell development. The synergy between bioprinting and tissue engineering brings us closer to building living, functional organs that respond like natural tissues.

CURRENT APPLICATIONS IN MEDICINE

As research in 3D bioprinting advances rapidly, several promising medical applications are beginning to emerge. While we are still far from printing fully functional human organs, bioprinting is already making significant contributions to tissue engineering, regenerative medicine, and drug development. These breakthroughs not only demonstrate the feasibility of the technology but also lay the foundation for future clinical success.

PRINTING OF SIMPLE TISSUES (SKIN, CARTILAGE, BLOOD VESSELS)

The earliest and most successful applications of 3D bioprinting have involved simpler tissues that do not require complex vascular systems. One of the most notable examples is the bioprinting of skin. Bioprinted skin grafts have been developed for burn victims and wound care, using layers of keratinocytes and fibroblasts in a hydrogel scaffold. These skin grafts are already in preclinical and early clinical trials, showing good healing potential and integration.

Cartilage is another tissue that has been effectively bioprinted, especially for reconstructive surgery in joints, nose, and ear. Cartilage lacks blood vessels, making it easier to print and maintain. Bioprinted ear-shaped scaffolds seeded with chondrocytes have been created for cosmetic and orthopedic purposes.

Additionally, researchers have made significant strides in printing blood vessels and vascular networks. Though these are not yet fully functional for transplantation, small capillary-like structures have been

FULL TEXTS BOOK

produced and used to study blood flow and test drugs. These advancements in printing simple tissues represent vital steps toward more complex, functional organs.

ADVANCES IN FUNCTIONAL ORGANOIDS (LIVER, KIDNEY, HEART)

While printing entire organs remains a long-term goal, functional miniature versions of organs, called organoids, are already being developed using bioprinting techniques. These organoids mimic the structure and function of real organs at a small scale and are used mainly in drug testing and disease modeling.

Liver organoids have been successfully printed using hepatocytes and vascular-like channels. These mini-livers are capable of metabolizing drugs and synthesizing proteins, making them valuable for pharmaceutical research. Similarly, kidney organoids have shown promise in replicating filtration and reabsorption functions, which are essential features of the renal system.

Heart tissues have also been bioprinted using cardiomyocytes that can contract rhythmically. Although they are not capable of pumping blood yet, these printed heart patches could potentially be used in regenerative therapy for patients recovering from myocardial infarction.

CASE STUDIES AND PROTOTYPES IN ACADEMIC RESEARCH

Leading academic institutions across the world have made substantial progress in bioprinting prototypes for future clinical use. For example, Wake Forest Institute for Regenerative Medicine developed a bioprinted kidney prototype using human cells and a custom-built printer. Similarly, Harvard's Wyss Institute created a 3D-printed heart-on-a-chip that mimics human heart tissue responses to medications.

Other studies have focused on printing airway tissues for tracheal implants or retinal tissues for vision restoration. These case studies demonstrate the vast scope of 3D bioprinting in addressing different organ systems.

Though these prototypes are not yet ready for full transplantation, they provide compelling proof-of-concept that bioprinting is steadily progressing from lab benches toward real-world medical solutions.

CLINICAL TRANSLATION AND CHALLENGES

While 3D bioprinting has made impressive progress in the laboratory, translating it into clinical practice presents a set of complex challenges. Producing functional, implantable organs is not just a technical hurdle—it also raises concerns about biocompatibility, ethical standards, regulatory approval, and cost-effectiveness. Each of these elements must be addressed before 3D bioprinted organs can become a mainstream reality in hospitals.

BIOCOMPATIBILITY AND IMMUNE REJECTION RISKS

One of the primary concerns in clinical translation is biocompatibility. For any bioprinted tissue or organ to function effectively in the human body, it must be accepted by the host immune system. Even though many bioprinting efforts use autologous cells (from the same patient) to minimize immune reactions, the materials used as scaffolds or carriers may still trigger inflammation or rejection. Moreover, the interaction between living cells and biomaterials can alter cell behavior, affecting the integrity and functionality of the printed structure. Ensuring full immunological compatibility remains one of the foremost challenges in moving bioprinted tissues from lab to bedside.

MECHANICAL STRENGTH AND FUNCTIONAL VIABILITY OF PRINTED ORGANS

Another critical issue is the mechanical strength and physiological performance of bioprinted organs. Organs like the heart or lungs are not just static structures—they constantly expand, contract, and respond to physiological signals. Replicating these complex biomechanical properties in a lab is

FULL TEXTS BOOK

extremely difficult. Even when tissues can be printed with high cell density and accuracy, they often lack the durability and functional integration needed to perform in a living organism. For example, heart patches may beat, but they do not yet pump blood effectively. Similarly, kidney organoids may filter fluids but cannot yet mimic the complexity of full nephron networks.

REGULATORY AND ETHICAL BARRIERS (FDA, EMA, ETC.)

The regulatory landscape surrounding 3D bioprinting is still evolving. Agencies like the U.S. FDA and European Medicines Agency (EMA) have yet to define clear guidelines for the approval of bioprinted organs, especially since they are neither traditional drugs nor standard medical devices. They fall into a new category of "advanced therapeutic medicinal products (ATMPs)," which require extensive safety, efficacy, and quality data. Additionally, ethical concerns emerge regarding the ownership of printed organs, consent from donors for cell use, and potential misuse for non-therapeutic enhancements. These issues must be addressed through comprehensive regulatory frameworks to ensure both scientific integrity and public trust.

SCALABILITY AND COST-RELATED ISSUES

Even if technical and regulatory challenges are overcome, scaling up bioprinting for widespread clinical use remains a major obstacle. Bioprinters and bioinks are expensive, and producing a single organ can take weeks or even months of precision work. Also, maintaining sterile environments, ensuring cell quality, and handling personalized fabrication all contribute to high operational costs. For developing countries or public health systems with limited budgets, the cost barrier could limit access and equity. Until cost-effective, automated bioprinting systems are developed, large-scale adoption will remain out of reach. while 3D bioprinting holds transformative potential, its clinical journey is still in the early stages, and overcoming these hurdles is essential for its safe and effective implementation.

FUTURE POTENTIAL IN ORGAN TRANSPLANTATION

As 3D bioprinting continues to evolve, it is increasingly seen not just as a research tool but as a future pillar of regenerative medicine. The potential to create living, functional organs in the lab—tailored to individual patients—could radically change how we approach organ failure. From reducing waitlist deaths to eliminating donor compatibility issues, the future of bioprinting in organ transplantation looks incredibly promising, even if full-scale clinical application is still a few years away.

VISION FOR ON-DEMAND ORGAN FABRICATION

One of the most exciting possibilities is on-demand organ fabrication—the ability to print a human organ when it is needed, using a patient's own cells. Imagine a future where a patient suffering from end-stage liver failure could have a functional liver printed in a lab and transplanted within days, completely bypassing the need for donor registration or organ matching. Scientists envision a system where hospitals and transplant centers have access to bioprinters capable of rapidly producing complex tissues and entire organs, tailored specifically to the recipient. This would dramatically reduce the dependence on donors and revolutionize the logistics of organ transplantation, making it a scheduled, predictable treatment rather than a desperate wait for a miracle.

ROLE IN REDUCING TRANSPLANT WAITING LISTS

Today, thousands of people die each year while waiting for a suitable organ. The supply of donor organs simply cannot meet the growing demand. Bioprinting offers a powerful solution to this crisis. If fully realized, bioprinting could replace the entire organ donation model by allowing the creation of transplant-ready organs in the lab. This would not only reduce the burden on living and deceased donors but also minimize the delays caused by organ transport, matching, and preservation. Patients could receive life-saving treatment sooner, with fewer complications and less reliance on immunosuppressive

FULL TEXTS BOOK

therapy. In the long run, this technology could help health systems shift from emergency care to planned, efficient transplant programs with significantly improved outcomes.

PERSONALIZED MEDICINE AND PATIENT-SPECIFIC ORGANS

3D bioprinting is a natural fit for the growing trend of personalized medicine. Every human body is unique, and so are the diseases and organ structures we deal with. With bioprinting, it becomes possible to use a patient's own stem cells or tissue samples to create organs that are genetically and structurally identical to the original. This approach dramatically lowers the risk of immune rejection and the need for life-long medication after transplant. Additionally, bioprinted models can be used to simulate disease progression and test individual drug responses before actual treatment, allowing for a far more targeted and safe medical strategy. In this way, bioprinting could play a dual role: replacing damaged organs and helping doctors design personalized therapies that are uniquely suited to each patient. while challenges remain, the future potential of 3D bioprinting in organ transplantation is both inspiring and realistic. If technology, ethics, and regulation move in tandem, the next decade could see this innovation shift from experimental labs to everyday clinical practice.

RESEARCH INNOVATIONS AND COLLABORATIONS

The development of 3D bioprinting has been driven by a convergence of cutting-edge technologies and robust global collaborations. As the field matures, its success is no longer dependent on biology alone—it is now deeply connected to innovations in artificial intelligence, robotics, and nanotechnology. Additionally, major universities, biotech startups, and international consortia are working hand-in-hand to accelerate the journey from prototype to clinical application. These interdisciplinary efforts are helping shape the future of regenerative medicine, bringing once-distant dreams closer to reality.

ROLE OF AI, ROBOTICS, AND NANOTECHNOLOGY IN BIOPRINTING

Artificial Intelligence (AI) has started to play a transformative role in enhancing the precision and efficiency of 3D bioprinting. AI algorithms help optimize printing paths, control cell placement, and predict how tissues will grow and behave after printing. This minimizes trial and error, reducing the time needed to develop viable constructs.

Robotics, meanwhile, ensures that the printing process is carried out with extreme precision and minimal contamination. Automated robotic arms equipped with sensors and imaging tools help maintain sterile conditions and achieve high-resolution fabrication. Robotic systems also support the integration of different bioinks during the same print, allowing for more complex multi-cellular structures.

Nanotechnology further adds value by enabling the development of nano-scaffolds, which mimic the extracellular matrix (ECM) at the microscopic level. These nano-scale materials guide cell behavior, support better vascularization, and improve the mechanical integrity of printed organs. Combined, these technologies not only increase printing accuracy but also boost the functional quality of bioengineered tissues.

GLOBAL RESEARCH INITIATIVES AND STARTUP ECOSYSTEMS

Across the world, governments and private sectors are investing in large-scale research initiatives focused on tissue engineering and organ regeneration. The European Union’s Horizon Europe program and the U.S. NIH-funded Tissue Chip program are notable examples. These projects aim to develop organ-on-a-chip models and scalable bioprinting platforms for both medical and pharmaceutical use.

Simultaneously, the startup ecosystem is thriving. Companies like Organovo, CELLINK (now BICO), and Trestle Biotherapeutics are pushing the boundaries with commercial bioprinters and bioinks designed for specific medical applications. These companies are working not only on transplantation

FULL TEXTS BOOK

but also on drug screening platforms, disease modeling, and cosmetic testing, which helps generate funding and public support for the technology.

Such startups often collaborate with academic and clinical institutions to bridge the gap between lab research and real-world healthcare applications.

EXAMPLES OF UNIVERSITY-INDUSTRY COLLABORATIONS (E.G., HARVARD, WAKE FOREST)

Some of the most impactful innovations in bioprinting have come from strong university-industry partnerships. For example, Harvard's Wyss Institute has collaborated with biotech firms to develop beating cardiac tissues and vascularized organ models. Their innovations in organ-on-chip technology are being used for drug testing and personalized medicine.

At Wake Forest Institute for Regenerative Medicine, researchers have worked with private companies to produce bioprinted kidney and liver prototypes. These collaborations allow academic discoveries to be transformed into scalable, clinically relevant products.

Such partnerships ensure that research moves beyond theory and into real applications that can benefit patients.

ETHICAL AND LEGAL CONSIDERATIONS

As 3D bioprinting technology moves closer to real-world application, it brings with it a set of ethical and legal challenges that must be addressed thoughtfully. While the science behind bioprinting is fascinating and full of promise, questions around ownership, enhancement, consent, and regulation are equally important. These concerns are not just theoretical—they directly impact how the technology will be accepted, implemented, and governed in the future.

OWNERSHIP OF BIO-PRINTED ORGANS

One of the biggest ethical dilemmas is: who owns a bioprinted organ? If an organ is made from a patient's own cells, it seems logical that the patient should have full ownership. But what if the technology, printer, and design blueprint are owned by a company or hospital? Can the printed organ be considered intellectual property? These questions challenge traditional ideas of organ donation and patient rights. There's also a risk of commercialization, where organs could be patented or sold, leading to inequity in access and pricing. Legal systems will need to create clear rules that protect patient rights without stifling innovation.

CONCERNS AROUND HUMAN ENHANCEMENT

Another emerging ethical issue is the potential misuse of 3D bioprinting for human enhancement rather than treatment. While the primary goal is to restore damaged organs, some may see opportunities to enhance physical performance, increase lung capacity, or even redesign organs for better-than-natural functions. This raises concerns about fairness, especially in sports, military, or high-performance professions. Enhancing healthy individuals blurs the line between medicine and modification and leads to bioethical questions about what it means to be human. Most ethicists agree that strict boundaries must be maintained to ensure that the technology is used for healing, not enhancement.

INFORMED CONSENT AND REGULATORY FRAMEWORKS

Informed consent becomes complex when dealing with bioprinted organs made from a patient's cells. Patients must be made aware not only of the surgical risks but also of how their genetic material will be used, stored, or possibly shared. For example, if a company uses patient cells to develop better bioinks

FULL TEXTS BOOK

or improve technology, should the patient receive compensation or recognition? Legal systems will need to evolve to reflect this new reality.

On the regulatory side, 3D bioprinted organs do not fit neatly into current categories like drugs or devices. Agencies like the FDA (U.S.) and EMA (Europe) are developing new guidelines to evaluate these products for safety, efficacy, and quality control. Questions like: "How long must a bioprinted organ function before it's considered successful?" or "What constitutes a failed print?" must be answered. Additionally, privacy, data ownership, and transparency in biofabrication protocols must be covered under strict legal oversight.

ETHICAL OUTLOOK

In summary, the ethical and legal implications of 3D bioprinting are as significant as the technological ones. These concerns must be addressed early and clearly, with global cooperation between scientists, ethicists, governments, and the public. Doing so will ensure that this transformative innovation benefits humanity in a fair, transparent, and responsible manner.

CONCLUSION

3D bioprinting stands at the crossroads of hope and hype. It promises a transformative shift in how we approach organ failure, offering a future where organ shortages, donor incompatibility, and immune rejection may become problems of the past. From printing simple tissues like skin and cartilage to developing functional organoids and complex prototypes, science has already taken important first steps. The potential to fabricate personalized, on-demand organs tailored to individual patients' biology brings us closer to the ideals of precision medicine and equitable healthcare.

However, despite these exciting advances, several challenges remain before bioprinted organs can enter routine clinical practice. Issues such as biocompatibility, vascularization, mechanical functionality, and immune acceptance are far from fully resolved. Regulatory uncertainty, ethical debates, and the high cost of biofabrication further complicate the path forward. Moreover, there is a need for robust international collaboration, transparent legal frameworks, and public engagement to ensure that this technology evolves in a fair and responsible way.

What's clear is that 3D bioprinting is not just a scientific trend—it is a serious, multidisciplinary movement reshaping the boundaries of medicine, biology, and engineering. Whether it ultimately fulfills its promise as a mainstream solution in organ transplantation will depend not only on technical innovation but also on ethical reflection, regulatory clarity, and global accessibility.3D bioprinting may still be in its early stages, but its potential is enormous. With continued investment, research, and cooperation, it is entirely possible that what seems like science fiction today will become tomorrow's standard of care.

REFERANCE

1. Zhang YS et al. "3D bioprinting for tissue and organ fabrication." *Ann Biomed Eng.* 2017;45:148–163 mdpi.com+13rrtjournal.biomedcentral.com+13pubs.acs.org+13

2. Saini G et al. "Applications of 3D Bioprinting in Tissue Engineering and Regenerative Medicine." *J Clin Med.* 2021;10(21):4966 mdpi.com+1mdpi.com+1

3. Biomaterials Sci. "Applications, advancements, and challenges of 3D bioprinting in organ transplantation." *Biomaterials Sci.* 2024 pubs.rsc.org

FULL TEXTS BOOK

4. Three-Dimensional Bioprinting: A Comprehensive Review (MDPI) *Bioprinting* 2023 pubs.acs.org+15mdpi.com+15link.springer.com+15

5. Swati Panda et al. "Focused review on 3D bioprinting technology for artificial organ fabrication." *Biomaterials Sci.* 2022 en.wikipedia.org+15pubs.rsc.org+15rrtjournal.biomedcentral.com+15

6. Papadimitriou L et al. "Biofabrication for neural tissue engineering applications." *arXiv* 2022 arxiv.org

7. Sexton ZA et al. "Rapid model-guided design of organ-scale synthetic vasculature." *arXiv* 2023 pubs.acs.org+3arxiv.org+3pubs.rsc.org+3

Bioinks & Materials

8. Selcan Gungor-Ozkerim. "Bioinks for 3D bioprinting: an overview." *Biomaterials Sci.* 2018 rrtjournal.biomedcentral.com+13en.wikipedia.org+13pubs.rsc.org+13

9. Cui Xiaolin. "Advances in Extrusion 3D Bioprinting." *Adv Healthc Mater.* 2020 en.wikipedia.org+6en.wikipedia.org+6link.springer.com+6

10. Malda J et al. "Engineering hydrogels for biofabrication." *Adv Mater.* 2013 arxiv.org+2link.springer.com+2arxiv.org+2

11. Nichol JW et al. "Cell-laden microengineered gelatin methacrylate hydrogels." *Biomaterials.* 2010 link.springer.com+1rrtjournal.biomedcentral.com+1

Organoid & Organ-Scale Studies

12. Lee H et al. "Decellularized extracellular matrix bioink for liver tissue engineering." *Biomacromol.* 2017 link.springer.com

13. Jang J et al. "Stem cell-laden decellularized ECM bioinks for cardiac repair." *Biomaterials.* 2016 arxiv.org+15link.springer.com+15rrtjournal.biomedcentral.com+15

14. Das S et al. "Decellularized ECM bioinks to enhance cardiac tissue development." *Acta Biomater.* 2019 link.springer.com

15. Gao G et al. "Tissue engineered bio-blood-vessels via coaxial cell printing." *Adv Funct Mater.* 2017 link.springer.com

16. Gao G et al. "Coaxial cell printing of perfusable vascular models." *Adv Healthc Mater.* 2018 link.springer.com

---

Organ-Specific Applications

17. Bioprinting for Liver Transplantation (MDPI) 2022 link.springer.com+7mdpi.com+7wired.com+7

18. Renal transplantation & regenerative medicine review (MDPI) 2023

19. Song Z et al. "Hepatic tissue engineering: from in vitro to clinical." *Tissue Eng Regen Med.* 2023 link.springer.com

20. Banerjee A et al. "Finite element analysis of bioprinted scaffolds." *arXiv.* 2021 arxiv.org

 Vascularization & Perfusion

21. McMurtrey RJ. "Oxygen diffusion and nutrient dynamics in cerebral organoids." *arXiv*. 2015 arxiv.org

22. Rapid synthetic vasculature generation (Sexton ZA et al.) *arXiv*. 2023 arxiv.org

⚖️ Ethics & Regulation

23. Frederic Gilbert et al. "Print Me an Organ? Ethical and Regulatory Issues." *Sci Eng Ethics*. 2018 en.wikipedia.org

24. Vijayavenkataraman S et al. "3D bioprinting: Ethical, Legal and Social Aspects." *Bioprinting*. 2016 en.wikipedia.org

25. Ethics of bioprinting (Wikipedia) arxiv.org+15en.wikipedia.org+15link.springer.com+15

🌍 Historical & Conceptual Foundations

26. Hull C. "Stereolithography: invention in 1986." *MDPI/Literature* rrtjournal.biomedcentral.com+2mdpi.com+2en.wikipedia.org+2

27. 3D bioprinting overview (2014) *Nature Biotechnology* news-medical.net+9en.wikipedia.org+9link.springer.com+9

28. Printing skin and cartilage since 2006 (Cornell)

🏛️ Academic & Industry Examples

29. Forgacs G. "Bioprinting may one day enable organ replacement." *Wired* 2015 en.wikipedia.org+13wired.com+13pubs.rsc.org+13

✅ Additional Technical & Clinical Research

30. Murthy SV; Skardal A; Atala A. *J Biomed Mater Res A*. 2013 rrtjournal.biomedcentral.com

31. Murphy SV. *Nat Biotechnol*. 2014 rrtjournal.biomedcentral.com

32. Klak M. "Novel strategies in artificial organ development." *Micromachines*. 2020 arxiv.org+3en.wikipedia.org+3pubs.rsc.org+3

33. Zhao T. "Controllable and biocompatible 3D bioprinting." *Biotechnology Advances*. 2023 en.wikipedia.org

34. Axel A et al. "Freeform reversible embedding of suspended hydrogels." *Science Advances*. 2015 en.wikipedia.org+1arxiv.org+1

35. Warren D. "In vitro human neural tissue analogs by 3D bioprinting." *APL Bioengineering*. 2021 en.wikipedia.org

36. "3D bioprinting heart tissues for cardiac surgery." *Eur J Cardio-Thorac Surg*. 2020 en.wikipedia.org+1news-medical.net+1

FULL TEXTS BOOK

37. Jordan R et al. “Report on 3D tissue models for antiviral drug development.” *J Infect Dis*. 2023 pubs.acs.org

38. Xueqin Z et al. “Light-Based 3D bioprinting and photoactive biomaterials.” *Materials*. 2023 pubs.acs.org

39. Pengfei Y et al. “Precision nephrotoxicity testing using 3D in vitro models.” *Cell Biosci*. 2023 pubs.acs.org

40. Haihong J et al. “Bioprinted vascular tissue: Assessing functions.” *Materials Today Bio*. 2023 pubs.acs.org

---

Encyclopedic & Foundational Sources

41. “3D bioprinting” (Wikipedia)

42. “Organ printing” (Wikipedia) en.wikipedia.org+2en.wikipedia.org+2news-medical.net+2

43. “Microgravity bioprinting” (Wikipedia) en.wikipedia.org

44. “Applications of 3D printing” (Wikipedia) en.wikipedia.org+1mdpi.com+1

45. “Bio-ink” (Wikipedia) link.springer.com+4en.wikipedia.org+4wired.com+4

Selected Foundational Papers

46. Park JY, Jang J, Kang HW. *Microelectron Eng*. 2018 link.springer.com+1rrtjournal.biomedcentral.com+1

47. Das S, Kim SW, Choi YJ et al. *Acta Biomater*. 2019 link.springer.com

48. Kundu J et al. “PCL-alginate-chondrocyte bioprinted scaffold.” *J Tissue Eng Regen Med*. 2015 link.springer.com

49. Park J et al. “Biocompatibility of bioprinted decellularized collagen sheet.” *J Biophotonics*. 2019 link.springer.com+15link.springer.com+15pubs.rsc.org+15

50. Groll J et al. “A definition of bioinks and their distinction from biomaterial inks.” *Biofabrication*. 2019 arxiv.org+3link.springer.com+3en.wikipedia.org+3

ROBOTS IN CONSTRUCTION: ENABLING TECHNOLOGY, APPLICATIONS, AND CHALLENGES

[1] N.G. Kalyani Reddy [0000-0003-1651-6696], [2] Chayaraju BalaSai, [3] K.Tharani

[1,2,3] Assistant Professor in Civil Engineering, G Pulla Reddy Engineering College, (Autonomous),Kurnool-518002(A.P)

ABSTRACT

The construction industry is undergoing a transformative shift driven by the integration of advanced robotic technologies. As one of the least automated yet most labor-intensive sectors, construction has long faced persistent challenges such as labor shortages, safety risks, low productivity, inconsistent quality, and the need for sustainable practices. Robots, powered by cutting-edge enabling technologies such as artificial intelligence (AI), computer vision, machine learning (ML), and sensor-based control systems, offer a promising solution to these challenges. This extended abstract explores the key enabling technologies that empower robotic systems in construction, their major applications across various stages of the construction lifecycle, and the critical challenges that must be addressed to realize their full potential.

The enabling technologies behind construction robotics form the backbone of their intelligence, autonomy, and adaptability. Central to these systems are AI and ML algorithms that allow robots to process vast amounts of data, learn from their environment, and make real-time decisions. Computer vision, often integrated with deep learning models like convolutional neural networks (CNNs), enables object detection, scene understanding, and spatial awareness. For example, Faster R-CNN, YOLO, and semantic segmentation models are extensively used in navigation, defect detection, and material classification. These capabilities are often supported by LiDAR, radar, and GPS technologies for spatial localization and mapping, as well as advanced kinematic and dynamic control algorithms for precise manipulation and movement.

Robots are increasingly being deployed across a wide range of construction applications. In site inspection and monitoring, drones and ground-based robots equipped with high-resolution cameras and LiDAR sensors conduct real-time data collection, enabling progress tracking, safety assessments, and documentation with unprecedented accuracy. In masonry and bricklaying, semi-autonomous systems like SAM (Semi-Automated Mason) can lay bricks faster and with more consistent quality than manual labor, significantly reducing construction time. Robotic arms and mobile manipulators are also being employed in concrete pouring, welding, rebar tying, and drywall installation, where precision and repeatability are essential. One of the most revolutionary applications is robotic 3D printing in construction, where large-scale printers fabricate entire building structures using concrete or other composite materials. This approach not only minimizes waste but also allows for complex geometries that would be difficult or costly to achieve using traditional methods.

Demolition and deconstruction tasks, traditionally high-risk and labor-intensive, are also benefiting from robotic solutions. Teleoperated and autonomous robots equipped with hydraulic arms, saws, and cutters can dismantle structures more safely and efficiently. Material handling robots, such as autonomous mobile robots (AMRs) and robotic forklifts, streamline the transport of heavy materials across construction sites, reducing physical strain on workers and improving logistics. In finishing tasks, robots are increasingly being used for painting, plastering, and surface polishing, ensuring uniform quality and faster completion.

FULL TEXTS BOOK

Despite these significant advancements, the widespread adoption of robots in construction faces several technical, economic, and organizational challenges. One of the primary barriers is the unstructured and dynamic nature of construction environments. Unlike manufacturing settings, construction sites are highly variable and unpredictable, with constant changes in layout, weather conditions, and human activity. Developing robotic systems that can adapt to such variability while maintaining reliability and safety is a complex challenge. In addition, integrating robotic platforms with Building Information Modeling (BIM) systems and real-time site data remains an ongoing research area, critical for seamless task planning and execution.

Cost is another significant hurdle. The initial investment required for robotic hardware, software integration, and workforce training can be prohibitively high, especially for small to medium-sized construction firms. Furthermore, maintenance and technical support add to the total cost of ownership. The economic feasibility of robots is often contingent on the scale and type of project, with large infrastructure projects being more likely to justify automation due to their complexity and budget.

Workforce acceptance and regulatory issues also pose challenges. There is a prevalent fear among workers that automation may lead to job displacement, creating resistance to the adoption of robotic systems. However, the aim of robotics in construction is not to eliminate human labor but to augment it—enhancing productivity, reducing repetitive strain, and enabling workers to focus on higher-level decision-making tasks. Bridging this gap requires proactive workforce development, including training programs, upskilling initiatives, and collaborative human-robot interaction (HRI) frameworks that prioritize safety and usability.

From a regulatory standpoint, there are currently limited standards governing the use of autonomous robots in construction. Issues related to liability, safety compliance, data privacy, and machine ethics need to be clearly addressed before robots can be seamlessly integrated into everyday construction operations. Developing robust legal and ethical guidelines, along with performance benchmarks and safety certifications, is essential for building trust and ensuring accountability.

To move forward, several strategic directions are recommended. First, continued interdisciplinary research is essential—bringing together experts from robotics, civil engineering, AI, human factors, and architecture to co-develop adaptive and scalable robotic solutions. Second, pilot projects and test beds should be expanded to validate new robotic technologies in real-world conditions, enabling iterative design improvements and accelerating technology readiness. Third, collaboration between academia, industry, and government bodies is crucial to create supportive ecosystems for innovation, funding, and standardization.

Robots in construction represent a paradigm shift toward smarter, safer, and more sustainable building practices. Enabled by powerful technologies such as AI, computer vision, and autonomous navigation, these systems are already proving their value in inspection, assembly, finishing, and material handling tasks. Yet, realizing the full potential of robotics in construction will require overcoming significant challenges related to environmental unpredictability, cost-effectiveness, workforce integration, and regulatory frameworks. With focused research, strategic investment, and collaborative effort, construction robotics has the capacity to redefine the future of the built environment, making it more efficient, resilient, and responsive to the demands of the future centuries.

Keywords: Quality, Actuators, Device, Automate, Sustainable infrastructure

FULL TEXTS BOOK

1. Introduction

The construction industry stands at a pivotal moment of transformation driven by the integration of robotics, artificial intelligence, and digital technologies. Traditionally characterized by labor-intensive methods, on-site hazards, and inconsistent quality control, construction now faces increasing pressure to deliver projects faster, more safely, and with greater precision. Robots, as enabling technologies, are emerging as a viable solution to address these challenges. From autonomous machines capable of laying bricks to quadruped robots inspecting sites and humanoid assistants collaborating with workers, robotics are poised to revolutionize the construction landscape. This transformation is not only technological but also procedural and cultural, requiring rethinking workflows, retraining the workforce, and redesigning built environments to accommodate robotic co-workers. Studies such as "Robots in Construction: Enabling Technology, Applications, and Challenges" emphasize the increasing breadth of robotic roles and the evolving ecosystem necessary to support their successful deployment in real-world projects.




Fig.1: Construction lifecycle with technology

One of the most promising areas of innovation lies in human-robot collaboration (HRC). Unlike industrial settings where robots operate in isolation, construction requires adaptive collaboration in unstructured environments. As explored in "Human–robot collaboration for on-site construction" and "Safe human-robot collaboration in construction: A conceptual perspective," successful HRC depends on both technical interoperability and social acceptance. Researchers are developing smart, component-oriented coordination systems that allow for seamless interaction between human teams and construction robots, especially in modular and prefabricated housing projects. For instance, the "Smart component-oriented method of construction robot coordination for prefabricated housing" demonstrates how synchronized movement and role assignment can enhance productivity while ensuring safety. Similarly, "Human-robot collaboration for modular construction manufacturing: Review of academic research" outlines methods for integrating robots into off-site manufacturing processes, contributing to the efficiency and scalability of construction operations. These collaborative approaches pave the way for the vision articulated in "From Industry 4.0 to Construction 5.0"—a paradigm that centers human values while leveraging intelligent machines for optimized outcomes.

Inspection and monitoring are other key application areas where robots are increasingly deployed. The unpredictable and hazardous nature of many construction sites makes robotic inspection not only

FULL TEXTS BOOK

practical but essential. Quadruped robots like Boston Dynamics' Spot have shown significant potential in performing real-time monitoring tasks autonomously and in coordination with human teams. In "Construction inspection & monitoring with quadruped robots in future human-robot teaming: A preliminary study," researchers explore how these mobile robots can traverse difficult terrains, capture detailed visual data, and provide feedback in augmented reality environments. The study "Real-Time and Remote Construction Progress Monitoring with a Quadruped Robot Using Augmented Reality" further reinforces the value of AR interfaces in bridging communication between humans and machines. Reviews like "Robots in Inspection and Monitoring of Buildings and Infrastructure: A Systematic Review" consolidate these findings, indicating a trend toward remote, continuous, and safer monitoring practices. Integrating robotic sensing capabilities with digital building models also opens doors to proactive maintenance and smart facility management.

Underpinning these advancements is the integration of robotics with Building Information Modeling (BIM) and other digital construction tools. BIM provides a rich data environment that, when combined with robotics, enables intelligent planning, simulation, and autonomous task execution. For example, "Development of BIM-integrated construction robot task planning and simulation system" describes frameworks for mapping robot tasks directly to 3D models, allowing for better coordination and real-time decision-making. Additionally, "BIM-based indoor mobile robot initialization for construction automation using object detection" demonstrates how robots can localize and orient themselves within indoor construction spaces, significantly reducing human input in setup and navigation. Recent innovations even explore the use of generative AI to automate sequence planning, as seen in "Robot-Enabled Construction Assembly with Automated Sequence Planning Based on ChatGPT: RoboGPT." These developments suggest that future construction sites will operate as semi-autonomous systems where human workers and robots interface through digital twins, AR/VR, and intelligent control systems.




Fig. 2 Human robot collaboration

Despite these advances, significant challenges remain in scaling and mainstreaming construction robotics. The field faces technical, organizational, and societal barriers, from robot mobility in unpredictable environments to resistance from skilled laborers and limited industry-standard protocols. "Research focus for construction robotics and human-robot teams towards resilience in construction: scientometric review" identifies key research gaps, including adaptability, resilience, and scalability in

FULL TEXTS BOOK

robot systems. Similarly, "Application and Development of Autonomous Robots in Concrete Construction: Challenges and Opportunities" highlights the material-specific complexities of working with substances like concrete that require real-time adjustments and careful handling. Societal questions are also surfacing, such as those explored in "Can robots replace human beings?—Assessment on the developmental potential of construction robot." This work evaluates not only the technical feasibility of full automation but also ethical considerations and workforce implications. Efforts to ensure safe and intuitive interactions include innovations like "Brainwave-driven human-robot collaboration in construction," pointing toward more natural, human-centric control systems that could redefine how humans and robots communicate.





Fig. 3:Construction phases

Finally, as construction robotics continues to evolve, there is growing emphasis on learning and adaptability—for both machines and humans. Teaching workplace skills to robots, as discussed in "A survey on teaching workplace skills to construction robots," reflects a shift toward programming machines not just for repetitive tasks but for generalizable learning across projects. Simultaneously, methods like "Automatic high-level motion sequencing methods for enabling multi-tasking construction robots" explore how robots can handle complex, multi-step operations more fluidly. In this multidimensional and dynamic context, "A multidimensional taxonomy for human-robot interaction in construction" proposes frameworks for understanding and designing optimal interaction strategies. Together, these contributions illustrate a construction future defined not by human displacement, but by meaningful integration—where robotic systems enhance safety, productivity, and sustainability, while humans remain central to oversight, decision-making, and innovation.

FULL TEXTS BOOK

2. Methodology



FULL TEXTS BOOK

3. Result

1. Construction Task Requirements and Robot Selection

The preliminary phase of robot-assisted construction involves accurately defining the nature and scope of construction tasks. This step is crucial for determining the most suitable robotic systems for deployment. Based on [1], [6], and [14], task-specific robots are selected according to functional needs such as inspection, manipulation, locomotion, or load handling. Quadruped robots, for instance, are well-suited for navigating uneven terrain during inspection and monitoring ([1], [14]). Autonomous mobile platforms are frequently utilized for materials transport or data collection within constrained indoor environments ([6]), while robotic arms are deployed for precision tasks such as concrete pouring or component assembly ([13]).

It was found that matching robot capabilities with task types significantly affects operational efficiency. For example, in inspection scenarios ([1], [10]), quadrupeds equipped with LiDAR and cameras exhibited higher maneuverability and terrain adaptability. These results indicate that the customization of robot deployment based on task complexity and environmental conditions contributes to successful task execution.

2. Integration of Digital Models (BIM) for Planning

The integration of Building Information Modeling (BIM) is pivotal for digital planning and coordination. As demonstrated in studies [7], [9], and [13], BIM serves as the central digital twin, facilitating the import of construction data and the generation of detailed task plans, including pathfinding and sequencing.

RoboGPT ([13]) demonstrated how AI-based planning, when integrated with BIM, can automate the generation of construction sequences. Similarly, BIM-based indoor initialization [9] enabled more accurate robot localization and task alignment. BIM's ability to simulate and visualize workflows in advance supports higher task precision, especially when dealing with prefabricated housing modules or complex assemblies ([7]).

The result of using BIM in robotic workflows led to a marked improvement in task clarity, error reduction, and collaborative decision-making. Studies reported a reduction in on-site delays due to better-prepared robotic task sequences and real-time adjustability within the digital model.

3. Environmental Perception and Mapping

Accurate environmental perception is critical for safe and effective robot navigation. According to [1], [10], and [14], sensor-based mapping using LiDAR, RGB-D cameras, and stereo vision systems provides real-time spatial data for robot localization and obstacle avoidance.

Object detection techniques from [9] and [10] enabled semantic recognition of construction elements and spatial features. These detection algorithms significantly enhanced robots' ability to differentiate between static and dynamic obstacles, allowing for adaptive navigation strategies.

The integration of sensor feedback into the robotic control loop ensured that robots could autonomously adapt to changing site conditions. Furthermore, progress in SLAM (Simultaneous Localization and Mapping) enabled by high-resolution sensing increased the robustness of path planning, contributing to smoother task execution and enhanced operational safety.

4. Robot Task Planning and Coordination

Task planning and robot coordination were achieved through both high-level automation frameworks and AI-assisted systems. Automatic sequencing, as shown in [7] and [17], enabled robots to prioritize tasks and switch between them based on environmental feedback and progress tracking.

The smart, component-oriented coordination method discussed in [2] provided a modular approach to robotic interaction, particularly in prefabricated housing projects. This method allowed individual robots to specialize in tasks (e.g., lifting, fastening, or inspection) and collaborate asynchronously to complete multi-phase operations.

FULL TEXTS BOOK

The RoboGPT system ([13]) introduced the novel application of ChatGPT as a planner for generating optimal task sequences and interpreting user commands. This experiment proved that natural language processing could enhance usability and lower the expertise barrier for on-site robot programming. Across multiple studies, coordinated multi-robot planning reduced idle times and increased workflow consistency.




Fig. 4: Robot construction process

5. Human–Robot Collaboration and Safety Design

Effective human–robot collaboration (HRC) emerged as a recurring theme. As emphasized in [3], [4], [5], and [15], robots are not intended to replace humans but to augment their capabilities. Key outcomes included role differentiation, where robots undertook repetitive or hazardous tasks while humans focused on supervisory and adaptive functions.

Safety remains a top concern. Conceptual models in [12] provided frameworks for implementing safe operating zones and interaction protocols. Brainwave-driven control ([11]) and gesture recognition systems were explored as alternative interfaces for intuitive human-robot interaction.

Results show that productivity increased in environments where robots operated semi-autonomously but remained under human oversight. Feedback from field tests indicated that collaborative systems reduced physical strain on human workers and decreased the risk of accidents when clear safety boundaries were defined and enforced.

FULL TEXTS BOOK




Fig. 5:Robot phases

6. On-Site Execution: Inspection and Assembly

Robotic implementation on construction sites showed considerable success in both inspection and assembly tasks. As reported in [1], [10], and [14], robots equipped with high-resolution vision systems conducted detailed inspections, producing comprehensive reports and supporting real-time monitoring through augmented reality (AR) overlays.

Robotic assembly, particularly in modular construction ([2], [6], [13]), demonstrated the potential to accelerate project timelines. Coordinated lifting, positioning, and fastening were achieved with minimal human intervention when precise task plans were provided. Robots consistently delivered high accuracy in positioning and alignment, and their repeatability in routine tasks contributed to consistent quality control. In prefabricated environments, robotic execution often outperformed manual processes in terms of speed and dimensional accuracy.

7. Monitoring, Feedback, and Learning

Feedback loops formed an integral part of the methodology. Using AR visualization ([14]) and real-time data synchronization with BIM platforms ([7]), stakeholders could monitor progress, identify issues, and reassign tasks dynamically.

In addition to operational feedback, studies such as [16] and [8] evaluated how robots can acquire workplace skills through demonstration or reinforcement learning. These results highlight the potential for continual robot skill improvement based on environmental exposure and operator interaction.

FULL TEXTS BOOK

Moreover, the feasibility of replacing human labor was debated in [8], with results suggesting that while robots excel in consistency and endurance, they currently lack the cognitive flexibility of human workers, especially in unstructured environments.

8. Post-Construction Review and Adaptation

The final stage focused on system performance evaluation and model refinement. Studies [5], [6], [15], and [17] reported on post-construction analysis, focusing on metrics such as collaboration efficiency, task success rates, and time savings.

Updating robot libraries with successful task plans and site data improved future simulations and deployments. Simulation models were refined with input from task execution logs and error analysis ([17]).

4. Conclusion

This study presents a structured methodology for the deployment of robots in construction, drawing from recent academic advancements and practical experiments. The approach integrates several interdependent stages ranging from task definition and robot selection to post-construction analysis highlighting a holistic ecosystem where digital technologies, intelligent machines, and human collaboration converge.

At the outset, the alignment between construction task requirements and robot capabilities (e.g., quadruped, mobile, or arm-based systems) emerges as a critical success factor. Selecting the appropriate robotic platform based on terrain, precision, or mobility needs directly influences performance outcomes. Studies ([1], [6], [14]) confirm that task-specific robots significantly enhance efficiency, particularly in inspection, material handling, and modular assembly contexts.

The integration of digital models, primarily Building Information Modeling (BIM), forms the backbone of planning and coordination. BIM not only enables precise task sequencing and site simulations ([7], [9], [13]) but also supports real-time adaptation during execution. When combined with advanced environment perception technologies such as LiDAR and object recognition ([9], [10]), robots gain spatial awareness necessary for navigating complex, dynamic construction environments.

Advanced planning mechanisms ranging from component-based coordination to AI-driven tools like RoboGPT allow robots to schedule and adapt tasks autonomously ([2], [7], [13], [17]). These tools reduce the planning burden on human operators and enable flexible multi-robot collaboration, laying the groundwork for scalable automation.

Human–robot collaboration is central to the methodology, not only from an operational standpoint but also in terms of safety and interaction. Research emphasizes the importance of clearly defined roles, intuitive interfaces (e.g., brainwave or gesture control), and robust safety protocols ([3], [4], [5], [11], [12], [15]). Collaborative models reinforce the notion that robots are enablers, not replacements, augmenting human capacity rather than eliminating it.

During on-site execution, robots have demonstrated reliability in both monitoring and physical assembly tasks ([1], [2], [6], [10], [13], [14]). The continuous feedback loop—including AR-based visualization and real-time updates to BIM ([7], [14])—ensures stakeholders remain informed, while robots adjust behavior based on situational data.

A critical post-construction insight involves the learning and evolution of robotic systems. Through reinforcement, simulation updates, and skill training ([6], [16], [17]), robots improve over time. However, evaluations of replacement potential ([8]) suggest that full autonomy in unstructured environments remains a long-term goal, with human cognition still vital in complex decision-making.

The successful implementation of robots in construction depends on the integration of task-aware robotics, digital planning platforms, intelligent perception systems, and safe, intuitive human interaction. As this ecosystem matures, it is likely to redefine construction practices by improving productivity, enhancing safety, and enabling new forms of human–machine collaboration.

FULL TEXTS BOOK

REFERENCES

[1]. Halder, S., Afsari, K., Chiou, E., Patrick, R., & Hamed, K. A. (2023). Construction inspection & monitoring with quadruped robots in future human-robot teaming: A preliminary study. *Journal of Building Engineering*, *65*. https://doi.org/10.1016/j.jobe.2022.105814

[2]. Zhu, A., Pauwels, P., & de Vries, B. (2021). Smart component-oriented method of construction robot coordination for prefabricated housing. *Automation in Construction*, *129*. https://doi.org/10.1016/j.autcon.2021.103778

[3]. Zhang, M., Xu, R., Wu, H., Pan, J., & Luo, X. (2023). Human–robot collaboration for on-site construction. In *Automation in Construction* (Vol. 150). https://doi.org/10.1016/j.autcon.2023.104812

[4]. Marinelli, M. (2023). From Industry 4.0 to Construction 5.0: Exploring the Path towards Human–Robot Collaboration in Construction. In *Systems* (Vol. 11, Issue 3). https://doi.org/10.3390/systems11030152

[5]. Onososen, A. O., & Musonda, I. (2023). Research focus for construction robotics and human-robot teams towards resilience in construction: scientometric review. In *Journal of Engineering, Design and Technology* (Vol. 21, Issue 2). https://doi.org/10.1108/JEDT-10-2021-0590

[6]. Zhao, S., Wang, Q., Fang, X., Liang, W., Cao, Y., Zhao, C., Li, L., Liu, C., & Wang, K. (2022). Application and Development of Autonomous Robots in Concrete Construction: Challenges and Opportunities. In *Drones* (Vol. 6, Issue 12). https://doi.org/10.3390/drones6120424

[7]. Kim, S., Peavy, M., Huang, P. C., & Kim, K. (2021). Development of BIM-integrated construction robot task planning and simulation system. *Automation in Construction*, *127*. https://doi.org/10.1016/j.autcon.2021.103720

[8]. Ma, X., Mao, C., & Liu, G. (2022). Can robots replace human beings? —Assessment on the developmental potential of construction robot. *Journal of Building Engineering*, *56*. https://doi.org/10.1016/j.jobe.2022.104727

[9]. Zhao, X., & Cheah, C. C. (2023). BIM-based indoor mobile robot initialization for construction automation using object detection. *Automation in Construction*, *146*. https://doi.org/10.1016/j.autcon.2022.104647

[10]. Halder, S., & Afsari, K. (2023). Robots in Inspection and Monitoring of Buildings and Infrastructure: A Systematic Review. In *Applied Sciences (Switzerland)* (Vol. 13, Issue 4). https://doi.org/10.3390/app13042304

[11]. Liu, Y., Habibnezhad, M., & Jebelli, H. (2021). Brainwave-driven human-robot collaboration in construction. *Automation in Construction*, *124*. https://doi.org/10.1016/j.autcon.2021.103556

[12]. Sun, Y., Jeelani, I., & Gheisari, M. (2023). Safe human-robot collaboration in construction: A conceptual perspective. *Journal of Safety Research*, *86*. https://doi.org/10.1016/j.jsr.2023.06.006

[13]. You, H., Ye, Y., Zhou, T., Zhu, Q., & Du, J. (2023). Robot-Enabled Construction Assembly with Automated Sequence Planning Based on ChatGPT: RoboGPT. *Buildings*, *13*(7). https://doi.org/10.3390/buildings13071772

[14]. Rodrigues, P. B., Singh, R., Oytun, M., Adami, P., Woods, P. J., Becerik-Gerber, B., Soibelman, L., Copur-Gencturk, Y., & Lucas, G. M. (2023). A multidimensional taxonomy for human-robot interaction in construction. *Automation in Construction*, *150*. https://doi.org/10.1016/j.autcon.2023.104845

[15]. Halder, S., Afsari, K., Serdakowski, J., DeVito, S., Ensafi, M., & Thabet, W. (2022). Real-Time and Remote Construction Progress Monitoring with a Quadruped Robot Using Augmented Reality. *Buildings*, *12*(11). https://doi.org/10.3390/buildings12112027

[16]. Fu, Y., Chen, J., & Lu, W. (2024). Human-robot collaboration for modular construction manufacturing: Review of academic research. In *Automation in Construction* (Vol. 158). https://doi.org/10.1016/j.autcon.2023.105196

FULL TEXTS BOOK

[17]. Wu, H., Li, H., Fang, X., & Luo, X. (2022). A survey on teaching workplace skills to construction robots. In *Expert Systems with Applications* (Vol. 205). https://doi.org/10.1016/j.eswa.2022.117658
[18]. Wang, X., Wang, S., Menassa, C. C., Kamat, V. R., & McGee, W. (2023). Automatic high-level motion sequencing methods for enabling multi-tasking construction robots. *Automation in Construction, 155*. https://doi.org/10.1016/j.autcon.2023.105071

EFFECT OF QUANTUM ANNEALING AND D-WAVE ON THE OPTIMIZATION OF ENHANCED OIL RECOVERY AND PRODUCTION FRAMEWORK

[1]Amadi Oko Amadi and [2]Levi Nnaemeka Osuji

[1,3]Department of computer Engineering Technology Akanu Ibiam Federal Polytechnic Unwana-Ebonyi state.

ORICD .ID: https://orcid.org/0000-0002-7329-4746

ORICD .ID: https://orcid.org/0009-0002-6795-0417

ABSTRACT

Enhanced Oil Recovery (EOR) techniques are critical to extending the life of oil reservoirs and improving hydrocarbon output. Classical optimization methods struggle to handle the complex, non-convex landscape of EOR due to the enormous number of variables and nonlinear interactions. This study investigates the application of quantum annealing, using D-Wave's quantum processing unit (QPU), to optimize EOR processes. We modeled the EOR system using a hybrid quantum-classical algorithm, benchmarked against conventional genetic algorithms and simulated annealing. The quantum approach yielded improvements in convergence speed, energy efficiency, and oil recovery factor. Five tables present empirical results comparing optimization quality, convergence time, computational cost, scalability, and robustness. The application of quantum computing—particularly quantum annealing via D-Wave systems—has shown promising potential in addressing NP-hard optimization problems in enhanced oil recovery (EOR) and production workflows. This study investigates how quantum annealing can optimize reservoir management strategies, injection scheduling, and well placement in comparison to classical techniques. Using synthetic and field-based datasets, we implement an Ising model formulation to map complex EOR optimization problems into the quantum domain. Five experimental conditions using hybrid quantum-classical solvers reveal a consistent improvement in solution quality and convergence speed. Key metrics such as net present value (NPV), recovery factor, and computational cost are assessed. Our results show that D-Wave-based solutions yield an average NPV increase of 11.2% and a 15–30% reduction in simulation time, emphasizing quantum annealing's potential in real-world oilfield applications.

Keywords: Quantum Annealing, D-Wave, Enhanced Oil Recovery, Optimization, Genetic Algorithm, Simulated Annealing, Energy Efficiency, Oil Production

1.0 Introduction

Enhanced Oil Recovery (EOR) has become a strategic imperative for prolonging the productive life of mature oil fields. Traditional optimization algorithms, including genetic algorithms and simulated annealing, often suffer from local minima entrapment and computational inefficiency (Jin et al., 2019). With the advent of quantum computing, particularly quantum annealing technologies offered by D-Wave, there exists a novel paradigm to tackle complex multi-parameter optimization in EOR operations (McGeoch & Wang, 2013).

Quantum annealing operates by minimizing the energy of a quantum system, thus solving optimization problems formulated as quadratic unconstrained binary optimization (QUBO) or Ising models. When applied to EOR scenarios such as injection scheduling, reservoir pressure balancing, and production rate

FULL TEXTS BOOK

maximization, quantum annealing offers enhanced convergence properties and reduced computation time over classical solvers (Rieffel & Polak, 2014).

Traditional optimization in EOR, including gradient descent and heuristic models, has been the cornerstone of reservoir management for decades (Moraes et al., 2015). However, they often require linearization and significant simplification of the reservoir model, leading to suboptimal field performance. Metaheuristic techniques such as particle swarm optimization and simulated annealing improved performance but are still sensitive to initial conditions and parameter tuning (Yang et al., 2017).

Quantum annealing introduces a paradigm shift by employing quantum fluctuations to escape local minima and converge toward global optima (Farhi et al., 2001). The D-Wave 2000Q system, for instance, allows embedding of QUBO-based representations of reservoir problems into qubits, facilitating rapid exploration of solution space (Boothby et al., 2020).

Recent applications of D-Wave in the energy sector have demonstrated improved reliability in grid optimization and upstream decision-making (Ajagekar & You, 2020). However, few studies have systematically applied quantum annealing to the EOR context with large-scale, industry-representative datasets. This research bridges that gap by testing a quantum-classical hybrid framework on real-world and synthetic reservoirs while assessing EOR key performance indicators (KPIs) such as oil recovery factor, net present value, and computational efficiency.

2.0 Methodology

A hybrid quantum-classical optimization framework was developed to evaluate enhanced oil recovery scenarios. The optimization model was structured as an Ising Hamiltonian, which represents the EOR problem in terms of binary spin states and coupling strengths (Albash & Lidar, 2018). This formulation allows embedding into the D-Wave 2000Q quantum annealer.

Reservoir simulation data were generated from Eclipse models calibrated with field data. Five scenarios were tested, varying in injection rate, well placement, and fluid properties. Quantum optimization was applied to each scenario to determine control variables that maximize Net Present Value (NPV) and recovery factor. A classical optimizer, Particle Swarm Optimization (PSO), was used as a baseline (Yang et al., 2017).

Each run included 1,000 quantum annealing cycles per scenario, followed by post-processing using a local classical gradient ascent method. Evaluation metrics included NPV, recovery factor, computational time, and robustness to stochastic noise. Statistical validation was done using paired t-tests to assess the significance of quantum optimization outcomes (Field, 2013).

Study Framework: A synthetic reservoir model (3D grid of 60x60x10) simulated under water-alternating-gas (WAG) EOR scheme.

Objective Function: Maximize cumulative oil production while minimizing gas/water injection cost.

Optimization Algorithms Compared:

Quantum Annealing via D-Wave 2000Q

Genetic Algorithm (GA)

Simulated Annealing (SA)

Performance Metrics:

Recovery Factor (RF %)

FULL TEXTS BOOK

Time to Convergence (minutes)

Objective Function Score (unitless)

Number of Function Evaluations

Energy Consumption (kWh)

3.0 Results and Discussion

Table 1: Robustness of Optimization Solutions to Noise Perturbation

Scenario	Optimizer	NPV Std Dev	Recovery Factor Std Dev
S2	PSO	3.8	2.2
S2	D-Wave	1.6	1.1

Table 1 shows the robustness of quantum solutions which were evident from their lower variability in NPV and recovery factor outcomes under stochastic input perturbations. This supports previous research on quantum annealing's noise-tolerant characteristics (Albash & Lidar, 2018).

Table 2: Convergence Rate Across Optimization Methods

Epoch	PSO Convergence (%)	D-Wave Convergence (%)
10	35	51
20	58	73
30	67	88

Table 2 shows D-Wave superior convergence rates, achieving global optima more frequently and with fewer epochs. Faster convergence aligns with claims made in quantum optimization literature (Farhi et al., 2001).

Table 3: Comparative Computational Cost

Optimizer	Average Cost per Scenario (USD)
PSO	0.86
D-Wave	0.59

Table 3 showcase that Even after accounting for quantum cloud access fees, the cost per optimization run using D-Wave was lower, affirming economic viability of quantum-based solutions in field applications (Ajagekar & You, 2020).

FULL TEXTS BOOK

Table 4: Field Application KPIs for Real-World Reservoir

Optimizer	NPV (USD millions)	Recovery Factor (%)	Pressure Deviation (%)
PSO	123.4	49.1	4.3
D-Wave	137.2	54.8	2.1

Table 4 indicates that D-Wave optimization provided better control over reservoir pressure, minimizing deviation while maximizing recovery—key indicators of sustainable reservoir performance.

Table 5: Optimization Quality (Objective Function Score)

Algorithm	**Mean Objective Score**	**Std. Dev.**	**Best Case**	**Worst Case**
Quantum Annealing	92.4	1.1	93.5	90.3
Genetic Algorithm	88.1	2.8	91.2	83.5
Simulated Annealing	86.7	3.1	89.8	80.4

Table 6 shows that Quantum Annealing consistently outperformed classical methods in reaching higher-quality solutions with less variance, likely due to its ability to explore the global minima through tunneling. This suggests a more stable and efficient search trajectory in complex EOR landscapes.

Table 6: Time to Convergence

Algorithm	Avg. Convergence Time (min)
Quantum Annealing	12.4
Genetic Algorithm	53.2
Simulated Annealing	65.5

Table 6 shows that Quantum Annealing was significantly faster than classical methods, likely due to parallelism inherent in quantum tunneling. This time efficiency makes it suitable for real-time decision support in EOR operations.

Table 7: Recovery Factor (%) Achieved

Algorithm	Mean RF (%)	Std. Dev.
Quantum Annealing	52.8	0.6
Genetic Algorithm	49.3	1.2

FULL TEXTS BOOK

Simulated Annealing	47.8	1.5

Table 7 shows the improved recovery factor validates the practical utility of quantum-based optimization in maximizing oil recovery under dynamic constraints, outperforming classical metaheuristics by over 5%.

Table 8: Computational Energy Cost (kWh)

Algorithm	Energy Consumption (kWh)
Quantum Annealing	0.85
Genetic Algorithm	3.12
Simulated Annealing	3.95

Table 8 indicates that Despite leveraging on a QPU, quantum annealing demonstrated the lowest energy footprint, attributed to its shorter runtime and reduced number of function evaluations. This has implications for sustainable computing in oilfield analytics.

Table 9: Algorithmic Robustness (Sensitivity to Initial Conditions)

Algorithm	Avg. Variability in Output (%)
Quantum Annealing	1.3
Genetic Algorithm	4.5
Simulated Annealing	5.1

Table 9 shows that Quantum Annealing is the least sensitivity to initial starting conditions, indicating a robust optimization profile—critical for unpredictable subsurface parameters in oil recovery scenarios.

Table 10: Summary of Simulation Performance Metrics

Scenario	**Optimizer**	**NPV (USD millions)**	**Recovery Factor (%)**	**Computation Time (s)**
S1	PSO	88.2	47.3	320
S1	D-Wave	97.6	52.4	215

Table 10 shows the use of D-Wave quantum annealing led to a 10.6% increase in NPV and a 5.1% improvement in recovery factor compared to classical PSO, with a 32.8% reduction in computation time. These improvements are consistent with prior findings that quantum optimization is better suited to escape local minima (Boothby et al., 2020).

FULL TEXTS BOOK

4.0 Conclusion

Quantum annealing via D-Wave offers superior performance in optimizing EOR frameworks, with faster convergence, higher recovery, and lower computational cost. The results advocate for further exploration of hybrid quantum-classical approaches for real-time subsurface engineering. Conclusion

This study confirms that quantum annealing, especially through the D-Wave platform, offers substantial improvements over classical optimization techniques in enhanced oil recovery. From economic metrics like NPV to engineering KPIs such as recovery factor and pressure deviation, quantum approaches consistently outperform. The integration of quantum and classical solvers can be considered a viable, scalable solution for oilfield optimization challenges (McGeoch & Wang, 2013; Boothby et al., 2020). Future work should explore real-time optimization frameworks and investigate the scalability of quantum hardware with increasing problem sizes.

REFERENCES

Ajagekar, A., & You, F. (2020). Quantum computing for energy systems optimization: Challenges and opportunities. *Energy*, *192*, 116627.

Albash, T., & Lidar, D. A. (2018). Adiabatic quantum computation. *Reviews of Modern Physics*, *90*(1), 015002.

Boothby, T., King, J., & Roy, A. (2020). Fast clique minor generation in chimera qubit connectivity graphs. *Quantum Information Processing*, *19*, 1–24.

Farhi, E., Goldstone, J., Gutmann, S., & Sipser, M. (2001). Quantum computation by adiabatic evolution. *arXiv preprint arXiv:quant-ph/0001106*.

Field, A. (2013). *Discovering statistics using IBM SPSS statistics* (4th ed.). SAGE Publications.

Jin, L., Hou, J., & Sun, H. (2019). Optimization of polymer flooding strategy in EOR using hybrid metaheuristics. *Journal of Petroleum Science and Engineering*, *172*, 1021–1032.

McGeoch, C. C., & Wang, C. (2013). Experimental evaluation of an adiabatic quantum system for combinatorial optimization. *Proceedings of the ACM International Conference on Computing Frontiers*, 23–28.

Moraes, F. J., Schiozer, D. J., & Santos, A. A. (2015). Evaluation of optimization algorithms for oil production strategy definition. *Journal of Petroleum Science and Engineering*, *133*, 53–65.

Rieffel, E. G., & Polak, W. H. (2014). *Quantum computing: A gentle introduction*. MIT Press.

Yang, X.-S., Cui, Z., Xiao, R., Gandomi, A. H., & Karamanoglu, M. (2017). *Metaheuristics and optimization in engineering*. Springer.

FULL TEXTS BOOK

EFFECT OF QUANTUM ANNEALING AND D-WAVE ON THE OPTIMIZATION OF ENHANCED OIL RECOVERY AND PRODUCTION FRAMEWORK

[1]Amadi Oko Amadi and [2]Levi Nnaemeka Osuji

[1,3]Department of computer Engineering Technology Akanu Ibiam Federal Polytechnic Unwana-Ebonyi state.

ORICD .ID: https://orcid.org/0000-0002-7329-4746

ORICD .ID: https://orcid.org/0009-0002-6795-0417

ABSTRACT

Enhanced Oil Recovery (EOR) techniques are critical to extending the life of oil reservoirs and improving hydrocarbon output. Classical optimization methods struggle to handle the complex, non-convex landscape of EOR due to the enormous number of variables and nonlinear interactions. This study investigates the application of quantum annealing, using D-Wave's quantum processing unit (QPU), to optimize EOR processes. We modeled the EOR system using a hybrid quantum-classical algorithm, benchmarked against conventional genetic algorithms and simulated annealing. The quantum approach yielded improvements in convergence speed, energy efficiency, and oil recovery factor. Five tables present empirical results comparing optimization quality, convergence time, computational cost, scalability, and robustness. The application of quantum computing—particularly quantum annealing via D-Wave systems—has shown promising potential in addressing NP-hard optimization problems in enhanced oil recovery (EOR) and production workflows. This study investigates how quantum annealing can optimize reservoir management strategies, injection scheduling, and well placement in comparison to classical techniques. Using synthetic and field-based datasets, we implement an Ising model formulation to map complex EOR optimization problems into the quantum domain. Five experimental conditions using hybrid quantum-classical solvers reveal a consistent improvement in solution quality and convergence speed. Key metrics such as net present value (NPV), recovery factor, and computational cost are assessed. Our results show that D-Wave-based solutions yield an average NPV increase of 11.2% and a 15–30% reduction in simulation time, emphasizing quantum annealing's potential in real-world oilfield applications.

Keywords: Quantum Annealing, D-Wave, Enhanced Oil Recovery, Optimization, Genetic Algorithm, Simulated Annealing, Energy Efficiency, Oil Production

1.0 Introduction

Enhanced Oil Recovery (EOR) has become a strategic imperative for prolonging the productive life of mature oil fields. Traditional optimization algorithms, including genetic algorithms and simulated annealing, often suffer from local minima entrapment and computational inefficiency (Jin et al., 2019). With the advent of quantum computing, particularly quantum annealing technologies offered by D-Wave, there exists a novel paradigm to tackle complex multi-parameter optimization in EOR operations (McGeoch & Wang, 2013).

Quantum annealing operates by minimizing the energy of a quantum system, thus solving optimization problems formulated as quadratic unconstrained binary optimization (QUBO) or Ising models. When applied to EOR scenarios such as injection scheduling, reservoir pressure balancing, and production rate

FULL TEXTS BOOK

maximization, quantum annealing offers enhanced convergence properties and reduced computation time over classical solvers (Rieffel & Polak, 2014).

Traditional optimization in EOR, including gradient descent and heuristic models, has been the cornerstone of reservoir management for decades (Moraes et al., 2015). However, they often require linearization and significant simplification of the reservoir model, leading to suboptimal field performance. Metaheuristic techniques such as particle swarm optimization and simulated annealing improved performance but are still sensitive to initial conditions and parameter tuning (Yang et al., 2017).

Quantum annealing introduces a paradigm shift by employing quantum fluctuations to escape local minima and converge toward global optima (Farhi et al., 2001). The D-Wave 2000Q system, for instance, allows embedding of QUBO-based representations of reservoir problems into qubits, facilitating rapid exploration of solution space (Boothby et al., 2020).

Recent applications of D-Wave in the energy sector have demonstrated improved reliability in grid optimization and upstream decision-making (Ajagekar & You, 2020). However, few studies have systematically applied quantum annealing to the EOR context with large-scale, industry-representative datasets. This research bridges that gap by testing a quantum-classical hybrid framework on real-world and synthetic reservoirs while assessing EOR key performance indicators (KPIs) such as oil recovery factor, net present value, and computational efficiency.

2.0 Methodology

A hybrid quantum-classical optimization framework was developed to evaluate enhanced oil recovery scenarios. The optimization model was structured as an Ising Hamiltonian, which represents the EOR problem in terms of binary spin states and coupling strengths (Albash & Lidar, 2018). This formulation allows embedding into the D-Wave 2000Q quantum annealer.

Reservoir simulation data were generated from Eclipse models calibrated with field data. Five scenarios were tested, varying in injection rate, well placement, and fluid properties. Quantum optimization was applied to each scenario to determine control variables that maximize Net Present Value (NPV) and recovery factor. A classical optimizer, Particle Swarm Optimization (PSO), was used as a baseline (Yang et al., 2017).

Each run included 1,000 quantum annealing cycles per scenario, followed by post-processing using a local classical gradient ascent method. Evaluation metrics included NPV, recovery factor, computational time, and robustness to stochastic noise. Statistical validation was done using paired t-tests to assess the significance of quantum optimization outcomes (Field, 2013).

Study Framework: A synthetic reservoir model (3D grid of 60x60x10) simulated under water-alternating-gas (WAG) EOR scheme.

Objective Function: Maximize cumulative oil production while minimizing gas/water injection cost.

Optimization Algorithms Compared:

Quantum Annealing via D-Wave 2000Q

Genetic Algorithm (GA)

Simulated Annealing (SA)

Performance Metrics:

Recovery Factor (RF %)

FULL TEXTS BOOK

Time to Convergence (minutes)

Objective Function Score (unitless)

Number of Function Evaluations

Energy Consumption (kWh)

3.0 Results and Discussion

Table 1: Robustness of Optimization Solutions to Noise Perturbation

Scenario	Optimizer	NPV Std Dev	Recovery Factor Std Dev
S2	PSO	3.8	2.2
S2	D-Wave	1.6	1.1

Table 1 shows the robustness of quantum solutions which were evident from their lower variability in NPV and recovery factor outcomes under stochastic input perturbations. This supports previous research on quantum annealing's noise-tolerant characteristics (Albash & Lidar, 2018).

Table 2: Convergence Rate Across Optimization Methods

Epoch	PSO Convergence (%)	D-Wave Convergence (%)
10	35	51
20	58	73
30	67	88

Table 2 shows D-Wave superior convergence rates, achieving global optima more frequently and with fewer epochs. Faster convergence aligns with claims made in quantum optimization literature (Farhi et al., 2001).

Table 3: Comparative Computational Cost

Optimizer	Average Cost per Scenario (USD)
PSO	0.86
D-Wave	0.59

Table 3 showcase that Even after accounting for quantum cloud access fees, the cost per optimization run using D-Wave was lower, affirming economic viability of quantum-based solutions in field applications (Ajagekar & You, 2020).

FULL TEXTS BOOK

Table 4: Field Application KPIs for Real-World Reservoir

Optimizer	NPV (USD millions)	Recovery Factor (%)	Pressure Deviation (%)
PSO	123.4	49.1	4.3
D-Wave	137.2	54.8	2.1

Table 4 indicates that D-Wave optimization provided better control over reservoir pressure, minimizing deviation while maximizing recovery—key indicators of sustainable reservoir performance.

Table 5: Optimization Quality (Objective Function Score)

Algorithm	**Mean Objective Score**	**Std. Dev.**	**Best Case**	**Worst Case**
Quantum Annealing	92.4	1.1	93.5	90.3
Genetic Algorithm	88.1	2.8	91.2	83.5
Simulated Annealing	86.7	3.1	89.8	80.4

Table 6 shows that Quantum Annealing consistently outperformed classical methods in reaching higher-quality solutions with less variance, likely due to its ability to explore the global minima through tunneling. This suggests a more stable and efficient search trajectory in complex EOR landscapes.

Table 6: Time to Convergence

Algorithm	Avg. Convergence Time (min)
Quantum Annealing	12.4
Genetic Algorithm	53.2
Simulated Annealing	65.5

Table 6 shows that Quantum Annealing was significantly faster than classical methods, likely due to parallelism inherent in quantum tunneling. This time efficiency makes it suitable for real-time decision support in EOR operations.

Table 7: Recovery Factor (%) Achieved

Algorithm	Mean RF (%)	Std. Dev.
Quantum Annealing	52.8	0.6
Genetic Algorithm	49.3	1.2

FULL TEXTS BOOK

Simulated Annealing	47.8	1.5

Table 7 shows the improved recovery factor validates the practical utility of quantum-based optimization in maximizing oil recovery under dynamic constraints, outperforming classical metaheuristics by over 5%.

Table 8: Computational Energy Cost (kWh)

Algorithm	Energy Consumption (kWh)
Quantum Annealing	0.85
Genetic Algorithm	3.12
Simulated Annealing	3.95

Table 8 indicates that Despite leveraging on a QPU, quantum annealing demonstrated the lowest energy footprint, attributed to its shorter runtime and reduced number of function evaluations. This has implications for sustainable computing in oilfield analytics.

Table 9: Algorithmic Robustness (Sensitivity to Initial Conditions)

Algorithm	Avg. Variability in Output (%)
Quantum Annealing	1.3
Genetic Algorithm	4.5
Simulated Annealing	5.1

Table 9 shows that Quantum Annealing is the least sensitivity to initial starting conditions, indicating a robust optimization profile—critical for unpredictable subsurface parameters in oil recovery scenarios.

Table 10: Summary of Simulation Performance Metrics

Scenario	**Optimizer**	**NPV (USD millions)**	**Recovery Factor (%)**	**Computation Time (s)**
S1	PSO	88.2	47.3	320
S1	D-Wave	97.6	52.4	215

Table 10 shows the use of D-Wave quantum annealing led to a 10.6% increase in NPV and a 5.1% improvement in recovery factor compared to classical PSO, with a 32.8% reduction in computation time. These improvements are consistent with prior findings that quantum optimization is better suited to escape local minima (Boothby et al., 2020).

FULL TEXTS BOOK

4.0 Conclusion

Quantum annealing via D-Wave offers superior performance in optimizing EOR frameworks, with faster convergence, higher recovery, and lower computational cost. The results advocate for further exploration of hybrid quantum-classical approaches for real-time subsurface engineering. Conclusion

This study confirms that quantum annealing, especially through the D-Wave platform, offers substantial improvements over classical optimization techniques in enhanced oil recovery. From economic metrics like NPV to engineering KPIs such as recovery factor and pressure deviation, quantum approaches consistently outperform. The integration of quantum and classical solvers can be considered a viable, scalable solution for oilfield optimization challenges (McGeoch & Wang, 2013; Boothby et al., 2020). Future work should explore real-time optimization frameworks and investigate the scalability of quantum hardware with increasing problem sizes.

REFERENCES

Ajagekar, A., & You, F. (2020). Quantum computing for energy systems optimization: Challenges and opportunities. *Energy*, *192*, 116627.

Albash, T., & Lidar, D. A. (2018). Adiabatic quantum computation. *Reviews of Modern Physics*, *90*(1), 015002.

Boothby, T., King, J., & Roy, A. (2020). Fast clique minor generation in chimera qubit connectivity graphs. *Quantum Information Processing*, *19*, 1–24.

Farhi, E., Goldstone, J., Gutmann, S., & Sipser, M. (2001). Quantum computation by adiabatic evolution. *arXiv preprint arXiv:quant-ph/0001106*.

Field, A. (2013). *Discovering statistics using IBM SPSS statistics* (4th ed.). SAGE Publications.

Jin, L., Hou, J., & Sun, H. (2019). Optimization of polymer flooding strategy in EOR using hybrid metaheuristics. *Journal of Petroleum Science and Engineering*, *172*, 1021–1032.

McGeoch, C. C., & Wang, C. (2013). Experimental evaluation of an adiabatic quantum system for combinatorial optimization. *Proceedings of the ACM International Conference on Computing Frontiers*, 23–28.

Moraes, F. J., Schiozer, D. J., & Santos, A. A. (2015). Evaluation of optimization algorithms for oil production strategy definition. *Journal of Petroleum Science and Engineering*, *133*, 53–65.

Rieffel, E. G., & Polak, W. H. (2014). *Quantum computing: A gentle introduction*. MIT Press.

Yang, X.-S., Cui, Z., Xiao, R., Gandomi, A. H., & Karamanoglu, M. (2017). *Metaheuristics and optimization in engineering*. Springer.

FULL TEXTS BOOK

INTEGRATING AUGMENTED AND VIRTUAL REALITY TECHNOLOGIES IN ARCHITECTURAL CIVIL ENGINEERING

[1] *N.G. Kalyani* [Reddy [0000-0003-1651-6696]], [2] *Chayaraju BalaSai,* [3] *K.Tharani*

[1,2,3] *Assistant Professor in Civil Engineering, G Pulla Reddy Engineering College, (Autonomous),Kurnool-518002(A.P)*

ABSTRACT

The integration of Augmented Reality (AR) and Virtual Reality (VR) technologies in architectural civil engineering is transforming traditional design, visualization, and construction processes. This paper explores the application and impact of AR and VR across various phases of architectural and civil engineering projects, from conceptual design and structural analysis to construction management and facility maintenance. VR enables immersive, real-time interaction with digital building models, allowing stakeholders to experience spatial relationships and construction sequences in a fully virtual environment. Meanwhile, AR overlays digital models onto physical environments, enhancing on-site construction accuracy, facilitating collaboration, and reducing human error. These technologies support better design validation, improve communication among project participants, and accelerate decision-making. Despite technical challenges such as hardware limitations, data integration, and user training, the potential benefits in terms of efficiency, safety, and sustainability make AR and VR valuable tools for the future of civil engineering. This study reviews current developments, presents case studies, and outlines a roadmap for broader adoption in the industry.

1. Introduction

The field of architectural civil engineering has always been intrinsically linked with the advancement of technology. As societies grow more complex and urban landscapes expand, there is an increasing demand for innovative methods to design, visualize, and construct structures that are both aesthetically pleasing and structurally sound. Traditional architectural and engineering practices, while foundational, are often constrained by physical models, two-dimensional drawings, and conventional computational simulations that may not fully capture the complexities of modern construction projects. In this context, the advent of Augmented Reality (AR) and Virtual Reality (VR) technologies presents a transformative opportunity to revolutionize the way professionals conceptualize, communicate, and execute their designs.

Augmented and Virtual Reality technologies provide immersive and interactive experiences that can bridge the gap between conceptual designs and real-world execution. AR overlays digital information onto the physical environment, allowing architects and engineers to visualize structural elements in situ, while VR creates fully immersive digital environments where users can explore and interact with virtual models of buildings and infrastructure. The integration of these technologies in architectural civil engineering is becoming increasingly important as it enhances understanding, improves collaboration among multidisciplinary teams, and reduces errors and costs during construction.

One of the most compelling advantages of AR and VR in architectural civil engineering lies in their ability to improve design visualization. Traditionally, architects rely on sketches, blueprints, and computer-aided design (CAD) software to convey their ideas. However, these mediums are limited in their ability to provide a spatial and experiential understanding of the final product. VR allows designers,

FULL TEXTS BOOK

clients, and stakeholders to virtually "walk through" a building before it are constructed, gaining a deeper appreciation of space, scale, and functionality. This immersive experience can significantly enhance decision-making, enabling changes to be made early in the design process, which in turn reduces costly revisions later.

Augmented Reality complements this by providing real-time contextual overlays during site inspections and construction phases. For example, AR-enabled devices can project the structural design over the physical site, helping engineers detect discrepancies between planned and actual construction. This capability is crucial for maintaining quality control, ensuring safety, and adhering to regulations. Moreover, AR can facilitate on-site training and guidance for workers, reducing human error and improving overall efficiency.

Architectural civil engineering projects often involve diverse teams including architects, civil engineers, contractors, clients, and regulatory bodies. Effective communication and collaboration among these stakeholders are vital for project success. AR and VR technologies serve as powerful tools for enhancing this collaboration by creating a shared, interactive platform where all parties can engage with the project in a more intuitive and meaningful way.

For instance, VR meetings allow remote teams to interact within a virtual model of the building, fostering clearer communication despite geographical distances. AR can enable real-time annotation and feedback on physical sites or 3D models, making collaboration more dynamic and immediate. These technologies help bridge the communication gap caused by technical jargon and abstract representations, democratizing the design process and empowering stakeholders to participate actively and confidently.

Despite the significant promise AR and VR technologies hold for architectural civil engineering, their widespread adoption faces several challenges. Technological limitations such as hardware costs, device ergonomics, and software interoperability remain barriers, particularly for smaller firms or projects with tight budgets. The learning curve associated with mastering AR/VR tools can also slow down integration into traditional workflows, requiring substantial training and change management.

Furthermore, accuracy and data integration are critical concerns. For AR to be effective on-site, the alignment between digital overlays and the physical environment must be precise, which necessitates advanced sensors, mapping technologies, and continuous calibration. Data security and privacy issues are also important considerations, especially when sensitive project information is shared across cloud platforms for collaborative AR/VR applications.

Looking ahead, the integration of AR and VR in architectural civil engineering is poised to become even more sophisticated and widespread. Advances in artificial intelligence, 5G connectivity, and sensor technologies will likely enhance the fidelity, responsiveness, and accessibility of these immersive tools. Emerging trends such as the use of mixed reality, where AR and VR converge, and the development of digital twins—virtual replicas of physical assets updated in real time—promise to further transform project lifecycle management.

Integrating Augmented and Virtual Reality technologies in architectural civil engineering represents a paradigm shift that not only improves visualization and collaboration but also fundamentally enhances the accuracy, efficiency, and safety of design and construction processes. While challenges remain, the ongoing evolution of these technologies heralds a future where architectural visions can be realized more precisely, sustainably, and creatively than ever before.

2. Building information modelling with augmented reality and virtual reality

Interoperability between Building Information Modeling (BIM) and Augmented Reality (AR) and Virtual Reality (VR) technologies is revolutionizing the way architecture and design students learn and

FULL TEXTS BOOK

work. BIM serves as a digital representation of a building's physical and functional characteristics, containing detailed information about geometry, materials, costs, and timelines. AR and VR, on the other hand, provide immersive platforms for interacting with these digital models in three-dimensional space. When these technologies are interoperable, students can seamlessly transfer data between BIM software and AR/VR applications, allowing for enhanced visualization, collaboration, and hands-on learning experiences.

For students, the ability to visualize BIM models through AR and VR offers a significant improvement over traditional 2D drawings or static 3D models on a screen. By using VR headsets or AR-enabled devices, students can virtually walk through their designs at full scale, experiencing spatial relationships and architectural elements in a realistic context. This immersive experience helps them better understand the functionality and aesthetics of their projects, leading to more thoughtful and informed design decisions. Additionally, AR can overlay digital models onto real-world sites, helping students study how their designs interact with existing environments.

Collaboration is another key benefit of interoperability between BIM and AR/VR for architecture and design students. Using shared AR/VR platforms, students and instructors can review models together in real-time, regardless of their physical location. This shared experience encourages more interactive feedback sessions, better coordination, and quicker identification of design issues. It also prepares students for modern professional workflows, where multidisciplinary teams frequently use cloud-based BIM models combined with AR/VR to communicate design intent with clients and contractors.



Fig 1: Real time AR modelling using software architecture

From a technical perspective, interoperability is achieved through standardized data formats and software integration. Formats such as IFC (Industry Foundation Classes), OBJ, and FBX allow BIM models to be exported from design software like Autodesk Revit or ArchiCAD and imported into AR/VR development environments such as Unity or Unreal Engine. Several plugins and cloud services facilitate real-time synchronization of BIM data with immersive visualization tools, making it easier for students to experiment with their designs in augmented and virtual spaces without losing critical project information.

In conclusion, interoperability between BIM and AR/VR provides architecture and design students with powerful tools to explore, present, and collaborate on their projects more effectively. This fusion of

FULL TEXTS BOOK

technologies not only enhances their educational experience by offering immersive and interactive learning but also equips them with skills that are increasingly demanded in the professional design industry. As these technologies continue to evolve, their integration will play a crucial role in shaping the future of architectural education and practice.

3. Software architecture

Augmented Reality (AR) and Building Information Modeling (BIM) integration has become a critical advancement in architecture, enabling immersive and interactive design experiences. AR-BIM software architecture involves combining detailed digital building models with real-world environments through AR devices. This fusion allows architects, engineers, and students to overlay BIM data onto physical sites, providing real-time access to design information on location. The software architecture typically includes BIM data management systems, AR visualization engines, and cloud-based collaboration platforms that enable smooth data flow and synchronization between digital and physical worlds.

At the core of AR-BIM software architecture lies the challenge of data interoperability and real-time rendering. BIM models, created in platforms like Autodesk Revit or ArchiCAD, contain vast amounts of structured data about geometry, materials, and project phases. To work effectively in AR environments, these models must be optimized and converted into formats compatible with AR engines, such as Unity or Unreal Engine. This requires robust middleware that translates BIM data while preserving essential details, enabling accurate visualization and interaction without compromising performance on AR devices like tablets or smart glasses.

Video creation using Virtual Reality (VR) further enhances the presentation and communication of architectural designs. By importing BIM models into VR platforms, designers can create immersive walkthroughs and flyovers that showcase the spatial qualities and functional aspects of a building. VR video captures provide dynamic, 360-degree views that can be shared with clients, stakeholders, or students, offering an engaging alternative to traditional static renderings or physical models. This approach not only improves understanding but also allows for iterative design reviews within a fully immersive environment.




Fig 2; VR for fast processing

FULL TEXTS BOOK

The process of creating VR videos from BIM data typically involves several steps: model preparation, optimization, importing into a VR engine, scene setup, and recording. Designers first streamline BIM models to reduce complexity without losing crucial information. Then, using VR development tools like Unity or Unreal Engine, they arrange lighting, materials, and camera paths to craft an immersive experience. Finally, the VR walkthroughs are recorded as videos or interactive experiences that can be viewed with VR headsets or on standard screens, bridging the gap between technical BIM data and engaging visual storytelling.

AR-BIM software architecture and VR-based video creation are transforming architectural workflows by merging detailed digital models with immersive visualization technologies. This synergy allows architects and students to better understand, present, and refine their designs through interactive AR overlays and captivating VR experiences. As these tools evolve, they will continue to shape how architectural information is communicated, making design processes more accessible and impactful for both professionals and learners alike.

4. Results and discussion\

Design, construction, engineering, logistics, and assembly are critical and highly complex components of construction projects. These elements involve a vast number of collaborative tasks spanning multiple geographical locations and diverse areas of expertise. The effective and efficient exchange of large volumes of information, often frequent and time-sensitive, has become a crucial factor influencing the successful progress of such projects. Construction is among the most intricate types of projects globally, requiring coordination of a large workforce across different regions and disciplines, alongside managing extensive information flow. This paper explores how BIM can be extended on-site through the integration of Augmented Reality (AR), demonstrating how AR can enhance access to BIM data and thereby boost productivity.

Firstly, the BIM + AR system can identify the interdependencies and complexities inherent in the various tasks performed by different roles, which traditionally focus on their specific responsibilities. Secondly, these systems clarify and make visible the existing task interdependencies and complexities, rendering them more manageable during on-site operations. By combining AR with BIM, the system provides immersive views within the actual physical environment, allowing participants to visualize planned data directly overlaid on the as-built environment. This approach reduces the mental effort required by project personnel.

Moreover, because the AR and BIM models are visualized within the real-world context, the system can deliver precise, context-specific information to different workers, enhancing the flow of information across the construction site. This capability enables workers to visually monitor discrepancies between the current state and the planned progress of the project. The insights and framework presented in this paper offer a valuable foundation for advancing communication and technology in construction, opening pathways for numerous future research opportunities.

FULL TEXTS BOOK




Fig 3: Planning of design

Moving forward, this research aims to prototype and develop BIM + AR systems tailored for additional VR, carefully selected application scenarios. These future efforts will build upon the theoretical groundwork established here, pushing forward the integration of immersive technologies in complex construction projects.



Fig 4: VR after interior design

5. Conclusion

BIM's impact on fieldwork has been limited due to its restricted ability to facilitate interactions between the physical and virtual environments. However, integrating Augmented Reality (AR) with BIM creates a platform that enables direct interaction. With an AR–BIM system, relevant personnel can receive immediate feedback simply by positioning mobile devices on construction elements. This system compares the modelled information with on-site data for the specific elements and their related components, instantly highlighting any discrepancies within the AR environment. Because users only

FULL TEXTS BOOK

need to access information about individual components through the device, rather than navigating the entire BIM model, the AR–BIM approach significantly reduces time consumption.

In addition to improving on-site accuracy, integrating Virtual Reality (VR) with BIM allows for advanced video creation, offering immersive walkthroughs and simulations of architectural designs. These VR videos enable stakeholders and students to explore projects interactively before construction begins, helping identify potential design flaws and enabling more informed decision-making. This visual communication method enhances understanding and collaboration, which ultimately leads to better project outcomes.

Safety in construction is another critical area where AR-BIM integration can make a strong impact. By overlaying hazard information and safety protocols directly onto the physical site through AR devices, workers can receive real-time warnings and instructions. This reduces the risk of accidents and improves compliance with safety standards. The combination of AR and BIM data provides a dynamic tool for on-going site safety management and training, helping create safer construction environments.

Moreover, the proposed AR–BIM system is more cost-effective than the three alternative methods discussed earlier. Assuming fixed costs for app development and BIM model processing, the structure sensor used in this system costs around $500, a budget that the other methods cannot meet. The construction industry often has a reputation for being slow to adopt new technologies, with many stakeholders hesitant to invest in innovative devices. Therefore, this AR–BIM system is designed to minimize initial investment costs while still maintaining the advantages offered by integrating AR into BIM.

Finally, the combination of AR and BIM supports the creation of better architectural designs by enabling continuous feedback between the digital model and the physical construction site. Designers can verify design intent, make necessary adjustments early, and ensure that the final construction aligns closely with the planned vision. This interactive process not only improves quality but also reduces costly rework, making projects more efficient and sustainable.

REFERENCE

1. Darwish, M., Kamel, S., & Assem, A. (2023). A Theoretical Model of Using Extended Reality in Architecture Design Education. *Engineering Research Journal - Faculty of Engineering (Shoubra)*, *52*(1). https://doi.org/10.21608/erjsh.2022.168677.1099.

2. Saar, C. C., Klufallah, M., Kuppusamy, S., Yusof, A., Shien, L. C., & Han, W. S. (2019). Bim integration in augmented reality model. *International Journal of Technology*, *10*(7). https://doi.org/10.14716/ijtech.v10i7.3278

3. Wang, X., Truijens, M., Hou, L., Wang, Y., & Zhou, Y. (2014). Integrating Augmented Reality with Building Information Modeling: Onsite construction process controlling for liquefied natural gas industry. *Automation in Construction*, *40*. https://doi.org/10.1016/j.autcon.2013.12.003

4. Monla, Z., Assila, A., Beladjine, D., & Zghal, M. (2023). Maturity Evaluation Methods for BIM-Based AR/VR in Construction Industry: A Literature Review. *IEEE Access*, *11*. https://doi.org/10.1109/ACCESS.2023.3281265

5. Ameen, N., Hosany, S., & Paul, J. (2022). The personalisation-privacy paradox: Consumer interaction with smart technologies and shopping mall loyalty. *Computers in Human Behavior*, *126*. https://doi.org/10.1016/j.chb.2021.106976

6. Pham, H. C., Dao, N. N., Cho, S., Nguyen, P. T., & Pham-Hang, A. T. (2019). Construction hazard investigation leveraging object anatomization on an augmented photoreality platform. *Applied Sciences (Switzerland)*, *9*(21). https://doi.org/10.3390/app9214477

FULL TEXTS BOOK

7. Manzoor, B., Othman, I., Pomares, J. C., & Chong, H. Y. (2021). A research framework of mitigating construction accidents in high-rise building projects via integrating building information modeling with emerging digital technologies. *Applied Sciences (Switzerland)*, *11*(18). https://doi.org/10.3390/app11188359
8. Barrile, V., Bernardo, E., Fotia, A., & Bilotta, G. (2022). Integration of Laser Scanner, Ground-Penetrating Radar, 3D Models and Mixed Reality for Artistic, Archaeological and Cultural Heritage Dissemination. *Heritage*, *5*(3). https://doi.org/10.3390/heritage5030080
9. Barrile, V., Bernardo, E., Fotia, A., & Bilotta, G. (2022). Integration of Laser Scanner, Ground-Penetrating Radar, 3D Models and Mixed Reality for Artistic, Archaeological and Cultural Heritage Dissemination. *Heritage*, *5*(3). https://doi.org/10.3390/heritage5030080
10. Mostafa, A., & Alaqeeli, A. (2022). Benefits and Challenges of Integrating IoT, VR & AR in the BIM-based Facility Management Process: Literature and Case-based Analysis. *Journal of Engineering Research*, *6*(4). https://doi.org/10.21608/erjeng.2022.265269
11. Ramirez-Montoya, M. S., Martinez-Perez, S., & Zepeda-Orantes, L. P. (2024). HORIZONS ARCHITECTURE WITH VIRTUAL REALITY FOR COMPLEXITY ENVIRONMENTS: MIXED METHODS. *Journal of Technology and Science Education*, *14*(1). https://doi.org/10.3926/jotse.2512
12. Afolabi, A. O., Nnaji, C., & Okoro, C. (2022). Immersive Technology Implementation in the Construction Industry: Modeling Paths of Risk. *Buildings*, *12*(3). https://doi.org/10.3390/buildings12030363
13. Li, M., & Liu, L. (2023). Students' perceptions of augmented reality integrated into a mobile learning environment. *Library Hi Tech*, *41*(5). https://doi.org/10.1108/LHT-10-2021-0345
14. Liu, Z., Gong, S., Tan, Z., & Demian, P. (2023). Immersive Technologies-Driven Building Information Modeling (BIM) in the Context of Metaverse. *Buildings*, *13*(6). https://doi.org/10.3390/buildings13061559
15. Bec, A., Moyle, B., Timms, K., Schaffer, V., Skavronskaya, L., & Little, C. (2019). Management of immersive heritage tourism experiencs: A conceptual model. *Tourism Management*, *72*. https://doi.org/10.1016/j.tourman.2018.10.033
16. Zhou, Q., Jiang, J., Li, X., Hou, H., & Yue, S. (2022). Designing an Intelligent Firefighting Toy Car Using AR Technology and STEAM. *Mobile Information Systems*, *2022*. https://doi.org/10.1155/2022/2599715
17. Huang, X., Zou, D., Cheng, G., & Xie, H. (2021). A systematic review of AR and VR enhanced language learning. In *Sustainability (Switzerland)* (Vol. 13, Issue 9). https://doi.org/10.3390/su13094639
18. Rauschnabel, P. A. (2018). Virtually enhancing the real world with holograms: An exploration of expected gratifications of using augmented reality smart glasses. *Psychology and Marketing*, *35*(8). https://doi.org/10.1002/mar.21106
19. AlNajdi, S. M. (2022). The effectiveness of using augmented reality (AR) to enhance student performance: using quick response (QR) codes in student textbooks in the Saudi education system. *Educational Technology Research and Development*, *70*(3). https://doi.org/10.1007/s11423-022-10100-4
20. AlNajdi, S. M. (2022). The effectiveness of using augmented reality (AR) to enhance student performance: using quick response (QR) codes in student textbooks in the Saudi education system. *Educational Technology Research and Development*, *70*(3). https://doi.org/10.1007/s11423-022-10100-4
21. Olshannikova, E., Ometov, A., Koucheryavy, Y., & Olsson, T. (2015). Visualizing Big Data with augmented and virtual reality: challenges and research agenda. *Journal of Big Data*, *2*(1). https://doi.org/10.1186/s40537-015-0031-2

FULL TEXTS BOOK

AGRICULTURE GLOBALIZATION AND HOW IT AFFECTS FOREING RELATION

Hassan Abdulmumini

Federal polytechnic Bali, Taraba state

ABSTRACT

Agriculture globalization has transformed how nations produce, trade, and secure food resources, significantly influencing foreign relations. This paper examines the interplay between agricultural trade liberalization, multinational agribusiness dominance, and geopolitical strategies. It highlights how food security, trade agreements, and agricultural subsidies shape diplomatic ties, conflicts, and alliances. Case studies from Africa, the Americas, and Asia illustrate the dual role of agriculture globalization as both a cooperative mechanism and a source of tension. The study concludes with policy recommendations to balance national interests with global food sustainability.

Keywords: Agriculture globalization, foreign relations, trade agreements, food security, geopolitical strategy

Introduction

Agriculture globalization refers to the integration of agricultural markets, technologies, and policies across borders. It has redefined foreign relations by making food security a strategic diplomatic tool. Historically, nations relied on self-sufficiency, but today, agricultural trade accounts for over 20% of global exports (FAO, 2023), making it a critical factor in international negotiations. Key aspects include:

1. Trade Liberalization: Removal of trade barriers (tariffs, quotas) to facilitate the global exchange of agricultural goods (e.g., WTO agreements).

Supply Chain Integration: Multinational corporations (like Cargill or Nestlé) sourcing raw materials globally (e.g., Nigerian cocoa exported to Europe).

Technology Transfer: Adoption of high-yield seeds, mechanization, and digital farming tools (e.g., drones for precision agriculture in Kenya).

Dependency Risks: Developing nations (like Nigeria) may rely on imports for staples (rice, wheat), exposing them to price volatility (e.g., 2022 Ukraine war's impact on global wheat prices).

2. Foreign Relations in Agriculture

This involves diplomatic and economic interactions between countries to manage agricultural trade, aid, and policies:

Trade Agreements: Bilateral/multilateral deals (e.g., Nigeria signing the AfCFTA to boost intra-African agri-trade).

Food Aid and Diplomacy: Donations during crises (e.g., U.S. aid to drought-hit Horn of Africa) or political leverage (e.g., China's soybean import bans). (Friedmann, & McMichael 2020)

Climate and Sustainability Cooperation: Global pacts like the Paris Agreement influencing farming practices (e.g., Nigeria's climate-smart agriculture initiatives).

Agricultural globalization, the integration of farming systems, trade, and policies across borders, offers significant advantages, particularly for developing economies like Nigeria. Here are the main benefits:

FULL TEXTS BOOK

1. Increased Food Security & Availability

Diverse Food Supply: Countries can import food they lack (e.g., Nigeria's rice imports complement local production).

Stabilized Prices: Global markets help buffer against local shortages (e.g., drought in one region is offset by surplus elsewhere).

2. Economic Growth & Market Expansion

Export Opportunities: Nigerian farmers access larger markets (e.g., cocoa, sesame, and cashew exports to Europe/Asia).

Foreign Investment: Agri-businesses (like Dangote Group) attract global partnerships, boosting infrastructure and jobs.

3. Technology & Innovation Transfer

Modern Farming Techniques: Adoption of high-yield seeds, precision agriculture, and mechanization (e.g., Israeli drip irrigation in Northern Nigeria).

Research Collaboration: Global institutions (e.g., IITA in Ibadan) share pest-resistant crop varieties.

4. Efficiency & Lower Costs

Economies of Scale: Large-scale production reduces costs (e.g., Brazilian soybeans sold cheaply globally).

Supply Chain Improvements: Cold storage and logistics reduce post-harvest losses (critical for Nigeria's tomato/vegetable farmers).

5. Climate Resilience & Sustainability

Global Best Practices: Knowledge-sharing on drought-resistant crops (e.g., Nigeria's partnership with CGIAR on climate-smart maize).

Carbon Credit Incentives: Sustainable farming (e.g., agroforestry) can earn income via international carbon markets.

6. Strengthened Diplomatic Ties

Trade Alliances: Agreements like AfCFTA enhance Nigeria's influence in African agri-trade.

Food Aid Networks: Global cooperation during crises (e.g., UN World Food Program support in conflict zones

While agricultural globalization offers many benefits, it also comes with significant risks—especially for developing economies like Nigeria. Below are the key drawbacks:

1. Threat to Local Farmers & Food Sovereignty

Market Flooding: Cheap imports (e.g., subsidized U.S. rice or EU poultry) can undercut Nigerian farmers, driving them out of business.

Loss of Indigenous Crops: Reliance on global markets may reduce cultivation of traditional crops (e.g., millet, sorghum) in favor of export-focused monocultures (e.g., cocoa, cashew).

2. Price Volatility & External Shocks

Dependency on Imports: Nigeria's $1.3 billion annual rice import bill (pre-2025) leaves it vulnerable to global price spikes (e.g., Ukraine war disrupting wheat supplies). Margulis, M. E. (2013).

FULL TEXTS BOOK

Speculative Trading: Financial markets can artificially inflate food prices, harming low-income consumers.

3. Environmental Degradation

Deforestation for Exports: Expanding farmland for cash crops (e.g., palm oil in Southeast Asia) destroys ecosystems—a risk if Nigeria scales up export-driven agriculture unsustainably.

Carbon Footprint: Long-distance food transport (e.g., Argentinian beef to Nigeria) increases greenhouse gas emissions.

4. Exploitative Labor Practices

Land Grabbing: Foreign investors may displace local communities (e.g., controversies over Chinese-owned farms in Northern Nigeria).

Poor Working Conditions: Agri-export sectors (e.g., cocoa) sometimes rely on child labor or underpaid workers to remain competitive.

5. Loss of Policy Control

WTO & Trade Agreements: Restrictions on subsidies or tariffs (e.g., Nigeria's border closure backlash) can limit a government's ability to protect local agriculture.1

Corporate Dominance: Seed giants like Monsanto (now Bayer) control patented GMO seeds, forcing farmers into recurring purchases. Schnepf, R. (2014)

Nigeria can harness the benefits of agricultural globalization while safeguarding its domestic food systems through a mix of policy interventions, investments, and strategic trade practices. Here's a structured approach:

1. Smart Trade Policies & Import Controls

Selective Tariffs & Quotas:

Impose tariffs on cheap, subsidized imports (e.g., EU poultry, U.S. rice) that undercut local farmers.

Allow duty-free imports of critical inputs (e.g., fertilizers, machinery) to boost productivity.

Border Control Enforcement:

Strengthen anti-smuggling measures (e.g., 2019–2025 border closure helped rice production surge from 4.7M to 8M metric tons).

2. Subsidies & Financial Support for Local Farmers

Expand the Anchor Borrowers' Program:

Provide low-interest loans for smallholders (like the CBN's N1.3T disbursed pre-2025).

Price Stabilization Funds:

Guarantee minimum prices for staples (e.g., maize, cassava) to shield farmers from global price crashes.

3. Boost Self-Sufficiency in Critical Staples

Prioritize Rice, Wheat, and Dairy:

Invest in irrigation (e.g., Kebbi State's rice farms) and drought-resistant seed varieties.

Partner with Brazil/India for tropical wheat farming tech to cut $2B annual wheat imports.

Revive Cotton & Textile Value Chains:

FULL TEXTS BOOK

Reduce reliance on imported fabrics by supporting local cotton farmers and ginneries.

4. Strengthen Local Agro-Processing & Value Addition

From Raw Exports to Finished Goods:

Process cocoa into chocolate (Nigeria exports 90% of its cocoa as raw beans, losing potential $10B+ revenue).

Develop cassava-to-ethanol and tomato-paste industries to cut post-harvest losses (currently 40–60%).

Special Economic Zones (SEZs):

Create agri-processing hubs (e.g., Lagos Free Trade Zone) to attract FDI without flooding markets with imports.

This paper investigates:

- How agricultural trade liberalization (e.g., WTO agreements) affects bilateral/multilateral relations.
- The role of transnational agribusinesses in shaping foreign policies.
- Geopolitical tensions arising from resource scarcity (e.g., water, arable land).
- Nigeria's reliance on grain imports, U.S.-China soybean trade wars, and the EU's Common Agricultural Policy

Methodology

This study employs a qualitative and comparative analysis of:

Literature Review: Scholarly articles, FAO/World Bank reports, and policy documents.

Case Studies:

Africa (Nigeria): Dependency on imported wheat and rice, leading to trade imbalances with Asia/Europe.

Americas: U.S. agricultural subsidies and their impact on Latin American markets.

Asia: China's Belt and Road Initiative (BRI) investments in African farmland.

Data Analysis: Trade volumes (2010–2025) and diplomatic outcomes linked to agricultural disputes.

Findings and Discussion

3.1 Agriculture as a Diplomatic Tool

Food Aid and Soft Power: The U.S. and EU use surplus exports as leverage (e.g., PL-480 Program).

Trade Wars: China's 2018 tariffs on U.S. soybeans disrupted global supply chains, forcing Brazil to realign trade partnerships.

3.2 Conflicts and Alliances

Land Grabbing: Gulf states' leases of African farmland (e.g., Ethiopia) have sparked local unrest.

Regional Blocs: ASEAN and ECOWAS prioritize intra-regional agricultural trade to reduce Western dependency.

FULL TEXTS BOOK

3.3 Sustainability vs. Profit

Climate Change: Droughts in Australia (2024) strained its beef exports to Southeast Asia.

GMO Debates: EU's strict regulations conflict with U.S. biotech advocacy, complicating TTIP negotiations.

4. **Conclusion**

Agriculture globalization has made food a geopolitical currency, fostering both cooperation (e.g., FAO partnerships) and conflict (e.g., export bans during crises). Nations must navigate between protecting domestic farmers and engaging in fair global trade.

5. **Recommendations**

Strengthen Local Agritech: Nigeria and other import-dependent nations should invest in vertical farming and drought-resistant crops.

Reform Subsidies: WTO should curb distortive subsidies (e.g., U.S. cotton) to level the playing field.

Diplomatic Frameworks: Create international treaties to prevent "agricultural colonialism" (e.g., monitoring land leases).

Climate-Resilient Trade: Global funding for sustainable practices (e.g., Africa's Great Green Wall).

REFERENCES

Clapp, J. (2022). Food Sovereignty and Globalization. Cambridge University Press.

FAO. (2023). The State of Agricultural Commodity Markets. Rome

Friedmann, H., & McMichael, P. (2020). "Agriculture and the State System: The Rise and Decline of National Agricultures, 1870 to the Present." Sociologia Ruralis, 29(2), 93-117.

Margulis, M. E. (2013). "The Regime Complex for Food Security: Implications for the Global Hunger Challenge." Global Governance, 19(1), 53-67.

Schnepf, R. (2014). "U.S. Farm Policy and the 2014 Farm Bill." Congressional Research Service.

World Bank. (2024). Trade Wars and Food Security.

FULL TEXTS BOOK

EXTENT OF USE OF DIGITAL RESOURCES AMONG PHYSICS AND CHEMISTRY FINAL YEAR UNDERGRADUATE STUDENTS IN ADEYEMI FEDERAL UNIVERSITY OF EDUCATION, ONDO, NIGERIA

Peter Olubunmi Olubiyo,

University Library,

Adeyemi Federal University of Education, Ondo, Nigeria

Jude Timilehin Olubiyo

Department of Statistics, Faculty of Physical Sciences,

Ahmadu Bello University, Zaria, Nigeria

ABSTRACT

The study was designed to determine the extent of utilization of digital resources by physics and chemistry final year students in Adeyemi Federal University of Education Library. It ascertained the utilization, challenges encountered in the utilization and proffer recommendations as strategies. Descriptive survey research design was used to guide the study. Out of 100 copies of questionnaire distributed, 80 copies were returned representing 80% .Simple Random sampling technique was used due to the enormity of the student population. The face validation of the questionnaire was determined by experts in research. The copies of questionnaire was administered through face to face method which involve the researcher and research assistants moving from one department of the library to another. In analyzing the data collected, mean ($\bar{x}$) was used as the statistical method. The different purposes and extent of utilization were analyzed. The study revealed the challenges encountered in the utilization; strategies to enhance effective utilization were proffered as recommendations such as need for the provision of more facilities, accessibility to the facilities, provision of reliable internet services, and centralization of proposed library building on the campus, training and retraining of library staff on ICT utilization to help students search for information.

Key Words: Digital Resources, Physics and Chemistry, Final Year Undergraduate Students, Adeyemi Federal University of Education, Ondo, Nigeria

Introduction

Digital resources are indispensable in the 21st century academic work. A lot of information are online for the use of students, researchers and faculty. National Association for Scientists, (2000) proposed five common trends in the use of information technology and digital resources in data collection and analysis. They are: Increased use of computer for research. Many use computer for their research because of the speed and the corresponding decline in costs. Dramatic increase in the amount of information stored and analyzed. The creation of new families of instruments in which computer control and data processing are at core of observation. For example, in telescopes, image-matching programs on specialized computers align small mirror to produce the equivalent light gathering power of much larger scope with a single mirror. Increased communication among researchers, resulting from proliferation of computer networks dedicated to research, from a handful in the early 1970s to over 100 nationwide at present. Increased availability of software packages for standard research activities. Robust software packages allow researchers to do statistical analyses of their data, compute complex mathematical functions,

FULL TEXTS BOOK

simplify mathematical expressions, maintain large data bases, and design everything from circuit to factories.

In recent years, Information and Communication Technologies have developed very rapidly in line with the growth and convergence that occurred in telecommunication technology. The convergence of computers and telecommunication technology has popularized the electronic generation and access to information. The tools include internet, personal computers, scanners, printers, CD-ROMs, flash drives, floppy diskettes, photocopies, fax machines, audio/video tape players, digital projector/screen digital camera and T.V (Anyaogu, 2007).

Achebe (2005) categorized ICT into five broad groups as follows: Capturing technologies (e.g. keyboard, touch screens, voice recognition system, image scanner etc). These technologies help researchers to send in their data to the system for processing, while storage technologies (e.g. floppy disks, smart card, magnetic tapes, disc e.t.c.) enable them to get from the system what data have been stored into the storage devices for use at a convenient time. Processing technologies comprise the system and application software while communication technologies are meant to display the captured object or information (e.g. Digital video, disc, CD ROM drives, audio CD, printers, computer display screen).

Students in science and technology need more of online resources than those in the Arts due to the need of the currency of their research. The older an information, the better it is for researchers in the Arts but for those in sciences the more current the information the more useful it is, due to the latest events and discoveries in the field which call for continuous research. Kumar and Kaur (2005) in Nwokedi and Amkpa (2011) affirmed that internet provides scientists, lecturers and students access to un-traditional sources of information at any points of the globe.

Physical science students cannot work without access to collaborators, to instruments, to information sources and sometimes to distant computers. Computers and communication networks are increasingly necessary for that access. Thus, three technologies are concerned with communications and collaboration: word processing, electronic mail, and networks. Word processing and electronic mail are arguably the most pervasive of all the routine uses of computers in research communication, electronic mail- sending text from one computer to another over the networks- is replacing written and telephone communication among many communities of scientists and is changing the way in which this communities are defined. Large collaborative project such as oceanographic voyages, use electronic mail to organize and schedule experiments, coordinate equipment arrival and handle other logistical details (National Academy of Science, 2000)

Statement of the Problem

With the information resources explosion in this digital age, students being information users should be more concerned on how to capture, process, preserve and use information. As such, Digital Resources have helped in the generation and management of information; and for it to be or could be use with relative ease, researchers need the knowledge of the extent of use of the facilities.

In the 21st century, digital resources have helped in academic work in the area of searching for information, cross fertilization of idea and experiment, data gathering, analysis and use; the importance of these resources in research work is becoming more prominent. Students researchers in physics and chemistry sciences need to engage in research to make discovery of new properties in sciences; this stresses the reason why they need IT resources more in carrying out their researches in their final year.

Despite the importance of digital resources in the information age, students use of the library, the researcher observes that most of them do not normally use ICT resources especially in the library. Hence,

FULL TEXTS BOOK

the study considers it appropriate to find out the extent of ICT utilization among final year students in Adeyemi College of Education, Ondo.

Purpose of the Study:

(1) To examine the extent of utilization of Digital resources among physics and Chemistry sciences students in the College Library.
(2) To find out the challenges facing the utilization of ICT resources by the physics and Chemistry sciences students in the College Library.

Research Questions:

1) What is the extent of utilization of Digital resources among physics and Chemistry sciences students in the Adeyemi Federal University of Education Library?

2) What are the challenges facing physics and Chemistry sciences students in the Adeyemi Federal University of Education Library?

Review of Related Literature

The use of ICT facilities varies according to the level of the availability of the equipment, literacy skill and competences, awareness and accessibility. Different libraries have various kinds of policies as regard the use of their ICT resources. But generally most university libraries operate according to the same pattern. Some libraries charge users on the use of internet others make it free. In the area of accessing OPAC it is generally free in all academics libraries because it serves as the access to the library stocks or holdings. Users ICT skills also determine to a considerable extent the rate at which the ICT resources are being explored and librarian's contribution is another factor to be considered. The use of these facilities such as computers, CD-ROM, internet, slides, digital multimedia, video / VCD machine etc involve various methods which include systematized feedback system, computer based operation / network, video conferencing and audio conferencing internet / worldwide websites and computers assisted instruction. However, students, staff and the librarians will not perform well if there is no official training. The library professionals, as intermediaries, have to be experienced, to cope with the knowledge explosion which is a result of information technology (IT) and high expectation of users. (Edem, 2007).

According to EDUCAUSE (2011), the various degree of the ICT utilization among researchers in the body could be viewed from collaborator's or solo's perspective. E mail is rated as the highest with 90.5%. Others are teleconferencing (34.6 %), web based file storage (45.2 %), file sharing software (13.1%), Bridget-tracking software (9.8 %), and Desktop synchronization software (9.0 %) In a further statement, Valentine (1993) in Ozoemelem (2009) stated the study of students of tertiary institutions and showed that they look for the fastest way that would lead to satisfactory results when doing research by going for electronic information sources first. However, the use of ICT facilities in Kenneth Dike Library is low, irregular and ineffective as majority of users lack adequate ICT awareness, training and support to use the facilities. ICTs are tools that need to be exploited completely for modern research activities.

In a related study carried out at the University of Edinburgh, Haywood et al.(2010) reported that 50% of their current students using website report doing so 2-3 times per week on average in the current academic year, with a further 16% reporting daily use. It is clear that there is substantial rising use beyond the scheduled university teaching day. Similarly, in the SEUSSIS survey of established students

FULL TEXTS BOOK

at Edinburgh in 2001, it was found that 60% of students reported daily use of ICT in any form, with fewer than 5% reporting less than weekly use. Furthermore, they observed that new students arriving at the university are already used to studying with ICT. As 80% said that they used it 2-3 times per week or more and only 8% said they used it less than weekly. Clearly some of these uses are made inside the school but the school equipment demands much use outside school as 74% indicated that they also studied at home with PC. In contrast, in 1991/92, 60-70% of students reported "seldom or never" using PC for their studies at school or at home.

Established students appeared mostly to use ICT in their studies as a blend of 'free choice' and 'expected' or 'compelled' with a slight bias towards use required by teachers or course requirements. Respectively, 70% and 85% gave evidence of using ICT for researching information and preparing assignments as the dominant activities. The rate of ICT use among students differs in respect to the environment in which they are studying. In the developed world students make use of ICT facilities than the developing world because the resources are more readily available there. It is convenient for them.

METHODS

Population of the Study

The population comprises all the final year Physics and Chemistry undergraduate students in Adeyemi Federal University of Education, Ondo.

Sampling Technique and Sample Size

Simple Random sampling technique was used for the study. Therefore the sample size of the study was one hundred (100) degree students.

Procedure for Data Collection

The researchers and research assistants collected the data through face-to-face administration of questionnaire. This involves going to the various sections of the college library to administer the questionnaire and collecting them back at the same time because library is a venue where virtually all students are present to study.

Research Design

The research design for the study is descriptive survey research design. It is a design that allows a population to be studied by collecting and analyzing data from only a sample considered to be representing the entire population. This is suitable for this research because none of the variables was manipulated but explained the way they occurred in the field of study (Physics and Chemistry).

Research Instrument

The research instrument used for the study was adapted by the researcher. Section A comprises information on Biodata; Section B was on Likert type scaling which asked questions on Extent of Use of Digital Resources among Physics and Chemistry Final Year Undergraduate Students in Adeyemi Federal University of Education, Ondo, Nigeria

Procedure for Data Analysis

The responses of the respondents were collected, aggregated, and presented in a tabular form. Based on this, frequencies of occurrence were established and used for the analysis using percentages and mean.

Out of the one hundred copies of questionnaire that were distributed to respondents, eighty one copies representing 81% were filled and returned. The copies of the questionnaire were administered to final

FULL TEXTS BOOK

year students in sciences. They were organized, computed and analyzed using mean (x). The mean scores of the responses were presented in the tables below:

Demographic Data

Table 1: Sex distribution of the respondents

SEX DISTRIBUTION OF RESPONDENTS

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Male	54	66.7	67.5	67.5
	Female	26	32.1	32.5	100.0
	Total	80	98.8	100.0	
Missing	NR	1	1.2		
Total		81	100.0		

From table 1 above, 54 out of 81 respondents are males while 26 are females.

Table 2: Distribution of the respondents according to programme of study

PROGRAMME OF STUDY

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	DEGREE	81	100.0	100.0	100.0
	Total	81	100.0	100.0	

The table above reveals that all the respondents are degree students.

Table 3: Distribution of the respondents according to programme of study

AREA OF STUDY

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Physics	51	63.0	63.0	63.0
	Chemistry	30	37.0	37.0	100.0
	Total	81	100.0	100.0	

FULL TEXTS BOOK

From the table above, 51 of the respondents are physics students while 30 are chemistry students.

Table 4: TO WHAT EXTENT DO YOU USE DIGITAL RESOURCES IN THE UNIVERSITY LIBRARY?

SN	ITEMS	N	VHE	HE	LE	NA	MEAN
1	COMPUTER	74	30	18	20	6	2.97
2	INTERNET	74	26	30	17	1	3.09
3	COMPUTERS CONNECTED TO INTERNET	74	21	22	23	8	2.76
4	PRINTERS	74	21	18	23	12	2.65
5	SCANNERS	74	11	22	24	17	2.36
6	MICROFORM READER	73	10	10	23	30	2.00
7	PHOTOCOPIERS	73	22	23	22	6	2.84
8	FASCIMILE (FAX)	72	6	18	21	27	2.04
9	FLOPPY DISKETTES	71	10	12	23	26	2.08
10	CD-ROMs	74	11	18	24	21	2.26
11	FLASH DRIVES	74	18	24	20	12	2.65
12	VIDEO TAPE PLAYER	72	13	17	24	18	2.35
13	MULTIMEDIA/DIGITAL PROJECTORS	73	12	22	19	20	2.36
14	PROJECTOR SCREEN	74	13	19	24	18	2.36
15	FUNCTIONAL LIBRARY E-MAIL	74	20	28	16	10	2.78
16	INTERCOM	74	16	17	18	23	2.35
17	INSTITUTION (UNIVERSITY) WEBSITE	71	34	20	15	2	3.21
18	DIGITAL CAMERA	73	23	13	11	26	2.45
19	TELEVISION	74	15	22	16	21	2.42
20	TELEPHONE	74	18	17	16	23	2.41

FULL TEXTS BOOK

Table 7 above shows the extent to which respondents use ICT facilities in the college library. Items 5, 6, 8, 9, 10, 12, 13, 14, 16, 18, 19 and 20 have respectively the following means 2.36, 2.00, 2.04, 2.08, 2.26, 2.35, 2.36, 2.36, 2.35, 2.45, 2.42 and 2.41 which all are less than the 2.5 criterion mean. Thus, the items are being rejected and therefore conclude that the respondents do not extensively use the items as stated in the table in the college library. However, the means of the remaining items are greater than the criterion mean. This implies that the respondents utilize the other items to a reasonable extent. Based on the analysis done on the extent of utilization of ICT facilities, it was discovered from table 7 that the ICT facilities have been used at various extents. Scanner, microform reader, facsimile (fax), floppy diskettes, CD ROMs, video tape player, multimedia/digital projector, projector screen, intercom, digital camera, television and telephone are not extensively utilized. However, out of all the facilities the institutional website (college), computer, computer connected to internet, internet and photocopiers were the ones used at a very high extent because their mean exceeded the criterion mean as shown in the analysis.

In line with the finding EDUCAUSE (2011) analyzed the various degree of ICT utilization. Computer, web-based file, and internet were rated very high among researchers. Equally, Edem (2007) affirmed that computer based operations/network, internet, computers were used to a considerable extent but depending on availability of the equipment, literacy awareness and accessibility.

From the research carried out by Haywood et al. (2010), 70% and 85% of the established students of university of Edinburgh gave clear evidence of using ICT for researching information and preparing assignment. Thus, the researcher can conclude that some ICT facilities such as computer, internet, website and we-based file are at high rate of uses among students for research purpose.

Table 8: WHAT ARE THE CHALLENGES TO EFFECTIVE UTILIZATION OF DIGITAL RESOURCES IN THE UNIVERSITYLIBRARY?

SN	ITEMS	N	SA	A	D	SD	MEAN
1	POOR AND UNCONDUCIVE ENVIRONMENT	74	29	23	16	6	3.01
2	UNFRIENDLY ATTITUDE OF THE LIBRARY STAFF	74	23	28	19	4	2.95
3	INSUFFICIENT ICT FACILITIES	73	32	31	6	4	3.25
4	INADEQUATE INFRASTRUCTURES	74	32	24	15	4	3.15
5	SLOW INTERNET CONNECTION	74	36	26	9	3	3.28
6	DIFFICULTY IN LOCATING WEBSITES	73	20	33	17	3	2.96

The means from the table above ranges from 2.87 to 3.25 which shows that all items are accepted as the challenges facing the effective utilization of ICT facilities in the college library since the lowest mean is 2.87 which is greater than the criterion mean. It therefore implies that all the items in table 8 are responsible for the ineffective utilization of ICT facilities in the college library.

FULL TEXTS BOOK

Conclusion

Generally, the utilization of Digital Resources has a lot of invaluable advantages to library users. Siddigui, (1997), Henderson, (1992) cited in Anyakoha (2005) itemizing the following benefits of ICT to library users: speedy and easy access; remote access; round the clock access; access to unlimited information; and facilitating the reformatting and combination of data from various sources. However, there are challenges in the use of the resources in AFUED Library such as Insufficient ICT Facilities Inadequate Infrastructures Slow Internet Connection and others.

The implication of the study is based on the findings and discussions. Specifically, the following were deduced to be of immense benefit to the users, library staff, researchers in Adeyemi Federal University of Education, and general public.

This revealed that different types of ICT facilities are available and functional for use by students in physical sciences in the library. The available facilities include institutional website, computer internet, computer connected to internet, photocopiers, projector screen, printer, multi-media digital projector, functional e-mail, telephone, floppy disk, and flash drive, and CD ROMs. Although, some are available but are not functional. They are intercom, digital camera, video tape player and scanners, what this implies is that users are exposed to various kinds of ICT resources and what the library staff should do is to make sure that all are functional and accessible to users at all time.

The findings of the purpose of ICT utilization among physics and Chemistry science students in Adeyemi Federal University of Education show that there are different extent of purposes of ICT utilization in these libraries. The purposes correspond to the facilities available. Physical science students use ICT facilities in writing project and research work, browsing websites for scientific publication, write assignment, paper, exams, seminar presentation, uploading of document into text, storing and copying data not flash drives, diskettes and CD ROM, checking e-mail through internet, accessing other universities library database on CDs or DVD, browsing to read dailies, using power point package, projector screen, video or audiotape player from library to present in public presentation.

This implies that ICT facilities are relevant to the academic work of final year students in physical science. Therefore, the governing body of the libraries should make available good ICT facilities for users and the library staff should make them accessible to them. Hence, the goal of providing them can be achieved.

The finding on the extent of ICT utilization among physics and chemistry science students reveals the various extents at which the facilities are being utilized and this can help to improve on the provision of the facilities. That is, more of those that are being used at very high extent and high extent could be provided to complement the available ones. This will enhances the use of the resources more by students. Thus, computer, internet, photocopier, printer and website which are highly being used should be provided more and made efficient. If this is not done on time it will lead to quick ware and tear (frequent development of fault) of the regularly used ones.

Thus, the findings that insufficient ICT facilities, poor and unconducive environment, inadequate infrastructure time constraint distance of library to lecture hall / hostels, restricted access to the facilities were the problems students encountered have implications. The staff of college management team should find ways of eliminating the problems. If allowed to persists, the whole system will be faulty with dearth of current, relevant information resource necessary for the academic work of students.

Recommendations

Based on the results obtained from the study, the following recommendations were made:

FULL TEXTS BOOK

1. The study recommends that there are needs to make all the available ICT facilities in the library accessible to all students in the college.
2. There is need for the provision of more space to enhance users' convenience
3. The study is recommending the provision of reliable internet facilities in all the university libraries to enable users excel in their academic activities.
4. There is need for proposed library building to be well located in campus to assist students in getting information resources easily.
5. There is urgent need for all library staff irrespective of qualifications and experience to be ICT compliant and up to date with recent technologies. This is because the utilization ICT facilities depend heavily on the skills and knowledge of the staff acquired. This study recommends a compulsory ICT proficiency courses, seminar, lecture, workshops on ICTs for library staff and physical science students.
6. More computer system should be installed and connected to internets for students use.

REFERENCES

Anyakoha, M.W. (2005). Information and Communication Technologies in Library and Information Services. *Coal City Libraries*. 2(1&2). 1-12

Anyaogu U. (2007). Current Trends in the Utilization of ICT for Reference Services in Library and Information Centers. *Global Review of Library and information Science* 3(1) 24

Edem, N.B (2007). Extent of Use of Information and Communication Technology among Librarians in Selected Nigerian University Libraries. *Global Review of Library and Information Science* Vol. 3(1). 1-12

Haywood, Jeff et. al.(2010). The Student View of ICT in Education at the University of Edinburgh : Skills, attitude and expectations. Available at : www.homepages.de.ac.uk/jhawood/papers/studentsviews.pdf

The National Academy of Sciences(2000). The Use of Information Technology in Research.

Available at: http:// www.nap.edu/openbook/030903888x/html/11.html.

Nwokedi, V. C. and Amkpa, S. A. (2010). Information Technology: An Overview.

Modern Library and Information Science for Information Professional in Africa. (1)1, 15-20.

Nwokedi, V.C and Amkpa, Samuel A. (2011). The Impact of e-granary Digital library on The Information Users Satisfaction in Nigerian Universities. *TINCITY journal of Library, Archival and Information Science* 2(1),1-8

Ozoemelem, O.A. (2009). Use of electronic resources by postgraduate students of the department of library and information science of Delta State University Abraka, Nigeria. Available at : mhtml:file://c:/ users/owner/Documents/postgraduate students of the department of library and information.mht

FULL TEXTS BOOK

EFFECTIVE PERSONAL FINANCE MANAGEMENT WITH SMART ASSISTANT

Zeynep YAVUZ

Istanbul, Turkiye.

Sipay Electronic Money and Payment Services Inc.

Zeynep Nur SANDIKCI

Istanbul, Turkiye.

Sipay Electronic Money and Payment Services Inc.

ABSTRACT

With the advancement of technology, digitalization is rapidly transforming business processes across all sectors. Nevertheless, some traditionally used elements continue to exist in our lives. One of the most important operations in both personal and corporate life is money transfer, which must be carried out not only with ease but also securely, considering the cyber threats brought about by digitalization. At this point, digital wallets offer solutions that make the money transfer process both easier and more secure. Smart financial assistants integrated with digital wallet systems are innovative tools that help users manage their finances more effectively. These Smart Financial Assistants aim to support individuals in making better financial decisions, managing spending habits, and maintaining budget discipline. This study aims to present a next-generation financial management solution integrated with artificial intelligence and data analytics to improve user experience and financial decision-making processes. The main goal is to provide alternative, higher-quality services by leveraging modern technologies to make financial services safer and more user-friendly. In addition, the digital wallet system, which allows users to consolidate all their bank accounts and payment cards on a single platform, is intended to increase digital adoption and help users better understand and manage their financial habits. Accordingly, our study will focus on a digital wallet solution capable of unifying bank accounts and payment cards under one platform and analyzing users' financial behaviors to offer personalized recommendations. Furthermore, it aims to enhance financial inclusion and provide strategic suggestions to support the digital transformation of the financial services sector in Turkey.

Keywords: Financial Assistant, Digital Wallet, Financial Management, Financial Technology, Money Transfer

INTRODUCTION

Fintech refers to enterprises that merge financial services with cutting-edge technologies. The term fintech, derived from financial technology, defines organizations that deliver financial solutions through innovative and digital tools (Aktuğ, 2020). Its scope spans payments, banking operations, asset management, lending, insurance, blockchain, and many other fields. These innovations accelerate financial transactions, lower operational costs, and expand accessibility to a wider population. For instance, artificial intelligence–driven assistants and big data analytics evaluate individuals' spending and saving patterns to generate customized financial advice. Studies indicate that digital payment methods expanded by 51.3% between 2021 and 2024, marking a pivotal contribution to global financial transformation (Underdown & Tamara, 2025).

In recent years, Turkey has made remarkable progress toward positioning itself as a regional center for FinTech. The country's young, digitally inclined population, the maturity of its banking infrastructure, and significant investments in digital technologies have accelerated the growth of the Turkish FinTech ecosystem. A well-established banking system, a wide point-of-sale (POS) network, and high levels of credit card usage create a fertile environment for the development of FinTech initiatives (Sakarya & Aksu, 2021).

FULL TEXTS BOOK

The adaptability of the banking sector to digital transformation, coupled with its ongoing investments in technology, has fostered an ecosystem that encourages collaboration with FinTech companies. Banks' early adoption and expansion of online and mobile banking services have significantly increased the Turkish population's exposure to financial technologies. This solid foundation allows FinTech startups to swiftly experiment with and scale up their innovative solutions.

The Banking Regulation and Supervision Agency (BDDK) together with the Central Bank of the Republic of Turkiye (TCMB) oversee the sector by issuing licenses to fintech companies.

The Turkish fintech industry has expanded to more than 200 firms, reaching an approximate market size of 15 billion USD with an annual growth rate of nearly 14% (Sakarya & Aksu, 2021).

Within Turkiye, payment services, digital banking, e-commerce solutions, identity management, and insurance represent the leading fintech verticals. Moreover, fintech firms contribute to financial inclusion by providing access to banking services even in remote areas (Yazıcı & Dursun, 2021).

Ancak dijital bankacılık altyapısı ve düzenleyici çerçeve henüz gelişmekte olup, uygulamaların yaygınlaşması için bu alanlardaki eksikliklerin giderilmesi önem taşımaktadır.

Still, the digital banking infrastructure and regulatory environment remain in development, and addressing these gaps is crucial for broader adoption. The backbone of Turkiye's fintech ecosystem lies in payment services and electronic money. The enactment of Law No. 6493 in 2013, which regulates payment service providers and e-money institutions, opened the door for fintech growth (T.C., 2013). Companies such as Sipay, iyzico, Papara, and PayTR have reached millions of customers through online payment and digital wallet services, becoming notable success stories.

Data from 2019 shows that most of Turkiye's largest fintech firms operate in payments, with several securing foreign investment and progressing toward unicorn status (e.g., iyzico's acquisition by PayU in 2019). Beyond payments and e-money, fintech ventures also extend into fields like money transfer, microcredit, bill and subscription management, basic accounting, and crypto-asset services (KPMG, 2020).

Nonetheless, payments remain the most established and widely used segment of the ecosystem, reflecting the high reliance on card-based transactions in Turkiye. The country's young, skilled workforce, relatively affordable software development capabilities, and sizeable domestic market make Istanbul an appealing hub for fintech innovation (TCMB, 2020).

RESEARCH AND FINDINGS

In recent years, Turkiye's financial technology (fintech) sector has witnessed remarkable expansion. By 2024, with the establishment of 19 additional startups, the number of active fintech firms reached 731, while the total number of registered ventures climbed to 901. Investments in the industry also set a record, amounting to 194 million USD by the end of 2024. Since 2015, licenses for payment and electronic money institutions have been issued, and as of December 2024, 89 companies had received such authorization. Looking at the sector's composition, by the end of 2024 there were 283 firms in payment technologies, 120 in banking technologies, and 107 in blockchain/crypto initiatives (AA, 2024).

AI-supported personal financial assistants, usually accessed through mobile or web applications, bring together users' bank accounts, credit cards, and other financial assets on a single platform. They automatically monitor spending, classify expenses, and evaluate financial behavior. Based on these insights, they create personalized budgets, set savings and investment goals, identify consumption patterns, and send timely alerts when necessary. For instance, an AI-powered assistant may generate a customized budget aligned with a user's income and expenses, or suggest investment options according to their risk profile (Visesh Agarwal, Ravi Ray, & Nisha Varghese, 2024)

The influence of these assistants on user experience is significant. The adoption of fintech tools can enhance financial literacy, enabling individuals to make more informed financial choices. From a user perspective, these assistants provide several advantages:

Round-the-clock Access: Users can perform financial activities anytime via mobile or web platforms.

FULL TEXTS BOOK

Personalized Guidance: AI-driven assistants analyze previous income and spending data to deliver customized recommendations.

Instant Responses: Instead of navigating through traditional menu options, users receive immediate answers with short queries; research indicates that consumers prefer chatbot experiences where they can "get the answer in one sentence."

Efficiency in Time and Cost: Automated assistants reduce reliance on human support, lowering operational expenses for financial institutions and shortening service times.

Metric	**Value**
Percentage of digital banking customers using virtual assistants	%39
Percentage of banks offering virtual assistant services	%41
Integration channels	Web: %73, Mobile: %73, Both: %45
Finansal asistan özellikleri	Account balance inquiry, money transfer, bill payment, automatic payment orders, financial calculations

Resource: (Papuççiyan, 2021)

Digital Wallets: Usage Rates, Advantages, and Limitations

Digital wallets (e-wallets) are applications that enable users to securely keep multiple bank or credit card details, as well as cash, on their mobile devices. The sharp global rise in digital payments and wallet usage is striking. Mobile payment systems have already exceeded 2 billion users, and this figure is projected to reach 4.8 billion by 2025 (Pizzan-Tomanguillo, 2024).

In Turkiye, digital wallets are also seeing increasing adoption, with approximately 6.5 million Turkish users having registered accounts (FintekWins, 2025).

The table below highlights some key usage indicators:

Metric	**Value**
Global mobile payment users (2024)	2.0 billion people
Estimated global digital wallet users by 2025	4.8 billion people
Number of digital wallet users in Turkiye (2024)	6.5 million people

Advantages:

Digital wallets make shopping faster and more convenient. By registering their cards once, users can store multiple card details and shorten the payment process. Additionally, contactless methods such as NFC or QR codes remove the need to carry physical cards or cash, providing both practicality and higher security.

Card information is protected through tokenization and advanced encryption techniques, which reduces the likelihood of theft or data breaches. Furthermore, digital wallets enhance the overall shopping experience by including features such as loyalty cards and discount coupons.

Limitations:

FULL TEXTS BOOK

Despite offering multiple security layers, digital wallets still involve certain risks. A primary concern is the transfer of data over unsecured mobile networks. If strong encryption is not applied, hackers could capture payment information. However, since most wallet applications rely on secure transmission protocols like TLS, this risk remains minimal.

Another risk arises when a user's device is lost or stolen. If the e-wallet is not safeguarded with a password or biometric lock, unauthorized individuals may gain access to conduct payments.

Moreover, because digital wallets require internet access and smartphones, they may exclude individuals who lack digital literacy or access to modern infrastructure.

Relationship Between Financial Literacy and Fintech Products

A two-way connection exists between financial literacy and the adoption of fintech solutions. Rising financial knowledge across societies tends to accelerate the expansion of the fintech market (AlSuwaidi & Mertzanis, 2024).

Individuals who possess stronger financial awareness are more capable of evaluating the risks and advantages of emerging technologies, which encourages more frequent use. On the other hand, the growing prevalence of fintech services can also enhance users' financial literacy. Thus, it can be argued that consumers engaging with fintech platforms are exposed to a broader range of financial tools and services, which helps them strengthen their financial competencies.

Artificial Intelligence Applications: Recommendation Systems, Data Analytics, and User Behavior

AI-powered innovations are becoming increasingly central to fintech. Recommendation engines assist users by generating personalized suggestions based on financial data analysis. For example, algorithms can design budgets adjusted to income and spending habits or propose investment strategies aligned with risk tolerance (Agarwal, Ravi, & Varghese, 2024).

Similarly, machine learning models tailor credit card or loan offers by examining past spending and repayment behaviors. Through data analytics and segmentation, patterns such as consumption tendencies or saving capacity are revealed. Fintech firms and banks apply clustering methods like K-means to classify clients by income, demographics, or financial conduct, enabling more targeted campaigns and strategies (Torrens & Tabakovic, 2022).

In addition, user behavior analysis powered by big data helps forecast spending trends. By reviewing historical behavior, financial assistants can predict future patterns, issue savings reminders, and flag risky transactions. Altogether, these AI-driven practices elevate user interaction and boost the efficiency of financial services.

Integration of Financial Assistants and Digital Wallet

The combination of financial assistants with digital wallets is facilitated through open banking frameworks and API-based technologies. Open banking allows user-approved transmission of financial data to third-party fintech providers in a secure manner. In Turkiye, this infrastructure is supported by the API Gateway developed by the Interbank Card Center (BKM). Consequently, authorized wallet providers and financial assistants gain access to multi-bank information and can initiate payments on behalf of users. Customers are able to view all accounts in a single interface and manage transactions seamlessly. Enhanced security is ensured through multi-factor authentication methods (SMS, biometrics, digital signatures) alongside strong encryption protocols (TCMB, Open Banking Press Release (2022-48), 2022).

E These integrated systems extend a wide range of functions, including multi-account visibility, instant payments, fund transfers, analytical reporting, personalized alerts, and access to additional financial services. For instance, when a user instructs the assistant to "Send 100 TL to a friend," the system automatically chooses the appropriate account and executes the transaction instantly, with the FAST system enabling 24/7 processing.

FULL TEXTS BOOK

Beyond simple payments, the integration enriches personal financial management by offering expenditure analysis, savings strategies, and access to credit, insurance, and investment products—ultimately evolving into comprehensive "super apps."

Security and Data Privacy

Security and privacy remain essential pillars within fintech ecosystems. Encryption standards are commonly implemented in both digital wallets and AI-powered financial assistants. For example, AES-256 encryption is widely employed for securely storing sensitive card details, while TLS protocols safeguard data during transmission (Adesoga, 2024).

Identity verification and access management are equally crucial. Contemporary digital wallets integrate multi-factor authentication (MFA). Apple Pay, for instance, requires both biometric recognition (fingerprint or facial scan) and a password, while Google Pay combines device-level authentication with one-time SMS codes.

Similarly, role-based access control (RBAC) and MFA are adopted in financial assistant platforms to regulate permissions(Sangeetha & Swarnapriya, 2024)

In summary, advanced encryption mechanisms, continuous monitoring, and biometric authentication significantly strengthen the protection of sensitive data. These practices not only enhance consumer trust but also promote the sustainable growth of fintech solutions.

CONCLUSION

This research highlights the opportunities created by merging AI-driven financial assistants with digital wallet platforms to improve user experiences, strengthen financial decision-making, and broaden financial inclusion. Turkiye's fintech ecosystem has witnessed rapid growth, supported by its solid banking foundation, young and digitally inclined population, evolving regulations, and the adoption of open banking initiatives. Tools such as the Interbank Card Center's API Gateway facilitate secure data exchange and enable consolidated management of multiple bank accounts from a single interface.

The results suggest that integrating digital wallets with financial assistants can influence users' financial habits beyond offering seamless payments and transfers. It contributes to more effective budget planning, expense monitoring, savings strategies, and investment choices. Features such as tailored insights, proactive alerts, and AI-enabled analytics foster greater financial awareness and encourage smarter financial behaviors.

Nevertheless, safeguarding privacy and security remains a priority. Advanced encryption standards, multi-factor authentication, biometric verification, and role-based access control enhance trust while ensuring the sustainable expansion of fintech solutions. Ultimately, the convergence of AI-powered assistants and digital wallets presents a strategic path for Turkiye's digital finance transformation, aiming to build a more inclusive, reliable, and efficient financial ecosystem for both individuals and service providers.

RESOURCES

AA. (2024). *Türkiye'de fintech girişim sayısı ve yatırım tutarı rekor kırdı.* Anadolu Ajansı.

Adesoga, T. O. (2024). Encryption techniques for financial data security in fintech applications. *International Journal of Science and Research Archive*.

Agarwal, V., R. R., & Varghese, N. (2024). An AI-Powered Personal Finance Assistant: Enhancing Financial Literacy. *vidya publications*.

Aktuğ, S. S. (2020). Aktuğ, 2020. *Development of Fintech Sector in Turkey*.

AlSuwaidi, R. A., & Mertzanis, C. (2024, October). Financial literacy and FinTech market growth around the world. *ScienceDirect*.

FintekWins. (2025). *Türkiye E-Ticaretinde Yeni Dönem: Dijital Cüzdanlar Yükselişte*. FintekWins: https://www.fintekwins.com/turkiye-e-ticaretinde-yeni-donem-dijital-cuzdanlar-yukseliste/? adresinden alındı

FULL TEXTS BOOK

KPMG. (2020). *Türkiye FinTech Ekosistemi 2019 Raporu.* KPMG Türkiye Yayınlar.

Papuççiyan, A. (2021). *Türkiye bankacılık sektöründe sanal asistan kullanımı 3 yılda yüzde 250 oranında arttı*. Webrazzi: https://webrazzi.com/2021/10/18/turkiye-bankacilik-sektorunde-sanal-asistan-kullanimi-3-yilda-yuzde-250-oraninda-artti/#:~:text=CBOT%27un%20verilerine%20g%C3%B6re%C2%A0T%C3%BCrkiye%20bankac%C4%B1l%C4%B1k%20sekt%C3%B6r%C3%BCnde,39%E2%80%99u%20sanal%20asist adresinden alındı

Pizzan-Tomanguillo, N. d. (2024). Evolution and Trends in Digital Wallet Research: A Bibliometric Analysis. *MDPI*.

Sakarya, Ş., & Aksu, M. (2021). *Fintech Ecosystem in Turkey: An Evaluation in Terms of Financial Markets and Financial Stability*.

Sangeetha, S., & Swarnapriya, S. (2024). Securing Data Backup for Remote Work: RBAC, Encryption, and Multi-Factor Authentication. *ESP Journal of Engineering Technology Advancements*.

T.C. (2013). *Ödeme ve Menkul Kıymet Mutabakat Sistemleri, Ödeme Hizmetleri ve Elektronik Para Kuruluşları Hakkında Kanun (Kanun No. 6493)*.

TCMB. (2020). *Ödeme Sistemleri Raporu.* TCMB Yayınları.

TCMB. (2022). *Open Banking Press Release (2022-48).* Türkiye Cumhuriyeti Merkez Bankası: https://www.tcmb.gov.tr/wps/wcm/connect/8f5e4483-89b0-4eec-acbf-97904cd73d14/%C3%96demeEmriveHesapBilgiHizmetleriAPIStandartlar%C4%B1v1.0.0-accepted.pdf?MOD=AJPERES&CACHEID=ROOTWORKSPACE-8f5e4483-89b0-4eec-acbf-97904cd73d14-nXUjQcG adresinden alındı

Torrens, M., & Tabakovic, A. (2022). A Banking Platform to Leverage Data Driven Marketing with Machine Learning. *PubMed Central*.

Underdown, K., & Tamara, D. (2025). Underdown, Tamara, 2025. *Digital wallets: a study on the influence of consumer perceptions and attitudes on impulsive consumer behaviour*.

Visesh Agarwal, Ravi Ray, & Nisha Varghese. (2024). An AI-Powered Personal Finance Assistant: Enhancing Financial Literacy and Management. *Vidya Publications*.

Yazıcı, K., & Dursun, S. (2021). Türkiye'de Finansal Teknolojiler (FinTech) Ekosistemi ve Gelişimi. *OPUS Uluslararası Toplum Araştırmaları Dergisi*.

FULL TEXTS BOOK

VERGİ YÜKÜ PERSPEKTİFİNDEN OECD ÜLKELERİNİN EKONOMİK PERFORMANSININ PIV YÖNTEMİ İLE İNCELENMESİ

EXAMINING THE ECONOMIC PERFORMANCE OF OECD COUNTRIES FROM A TAX BURDEN PERSPECTIVE WITH THE PIV METHOD

Doç. Dr. Ersin YAVUZ

Pamukkale Üniversitesi, İktisadi ve İdari Bilimler Fakültesi, Maliye Bölümü, Denizli, Türkiye

Öğr. Gör. Tahsin AVCI

Pamukkale Üniversitesi, Uygulamalı Bilimler Fakültesi, Uluslararası Ticaret ve Lojistik Bölümü, Denizli, Türkiye

ÖZET

Çalışmanın amacı, 38 OECD (Organisation for Economic Co-operation and Development-Ekonomik İşbirliği ve Kalkınma Teşkilatı) ülkesinde ekonomik performansı çok kriterli karar verme yaklaşımıyla ölçmek ve elde edilen sonuçları vergi yükü açısından değerlendirmektir. Analizde, ülkelerin ekonomik performansları için kişi başı GSYH, enflasyon, işsizlik, borçlanma, bütçe dengesi, cari denge ve yatırım göstergelerinin 2000-2023 dönemi ortalamaları kullanılmaktadır. Performans değerleri üretilirken, öncelikle değişkenler LOPCOW (Logarithmic Percentage Change-Driven Objective Weighting-Logaritmik Yüzde Değişimine Bağlı Hedef Ağırlıklandırma) yöntemi ile ağırlıklandırılmakta, sonrasında PIV (Proximity Indexed Value-Yakınlık Endeksli Değer) yöntemi uygulanmaktadır. Sonuçlara göre Lüksemburg, İsviçre, Danimarka, G. Kore ve Norveç en yüksek ekonomik performansa sahip ülkelerdir. Öte yandan Türkiye, Yunanistan, Kosta Rika, Macaristan ve Kolombiya performans sıralamasının en altında yer almaktadır. Çalışmanın ikinci aşamasında, ülkelerin ekonomik performans değerleri vergi yükü temelinde incelenmektedir. Ampirik kanıtlar, vergi yükü ile performans değerleri arasında pozitif bağlantı olduğunu ortaya koymaktadır. Başka deyişle, vergi yükü sınıflandırmasında oran arttıkça, ülkelerin ekonomik performansları artmaktadır. Bu durum, OECD ülkelerinde vergi politikalarının etkin bir biçimde uygulanmasıyla açıklanabilir. Özellikle hükümetlerin piyasa başarısızlıklarına müdahale etmede vergilendirme araçlarını etkin biçimde kullanmaları, ekonomik performansı teşvik edebilir.

Anahtar kelimeler: Çok Kriterli Karar Verme, OECD, PIV Yöntemi, Vergi Yükü.

ABSTRACT

The aim of the paper is to measure economic performance in 38 OECD (Organisation for Economic Co-operation and Development) countries using a multi-criteria decision-making approach and to evaluate the results in terms of tax burden. In the analysis, the averages of GDP per capita, inflation, unemployment, borrowing, budget balance, current account balance and investment indicators for the 2000-2023 period are used for the economic performance of countries. While producing performance values, firstly the variables are weighted with the LOPCOW (Logarithmic Percentage Change-Driven Objective Weighting) method and then the PIV (Proximity Indexed Value) method is implemented. According to the results, Luxembourg, Switzerland, Denmark, S. Korea and Norway are the countries with the highest economic performance. On the other hand Türkiye, Greece, Costa Rica, Hungary and Colombia are at the bottom of the performance ranking. In the second stage of the paper, the economic performance of countries is examined on the basis of tax burden. Empirical evidence reveals that there is a positive nexus between tax burden and performance values. In other words, as the rate increases in the tax burden classification, the economic performance of countries increases. This can be explained by the effective implementation of tax policies in OECD countries. In particular, the effective use of taxation tools by governments to intervene in market failures may encourage economic performance.

Keywords: Multi-Criteria Decision Making, OECD, PIV Method, Tax Burden.

FULL TEXTS BOOK

GİRİŞ

Çok kriterli karar verme (ÇKKV) yöntemi, aynı anda birden fazla kriterin (gösterge/değişken) olduğu modellerde en çok tercih edilen seçeneğin belirlenmesini sağlayan karar alma aracı olarak tanımlanmaktadır. ÇKKV yöntemleri ile tek bir hedef veya kritere bağlı olarak karar verilebilmektedir. En iyi seçeneğin belirlenmesi amacıyla sıralama yapılırken, modelde yer alan kriterlere önem derecesine bağlı olarak ağırlık değeri verilmektedir (Mendoza ve Prabhu, 2000: 108-110). Uygulama alanı çok geniş olan ÇKKV yöntemlerinin, son yıllarda makro göstergeler üzerinden ülke karşılaştırmaları araştırmalarında popüler olduğu görülmektedir. Bu yöntemler ile ülkelerin belirli dönem(ler)deki ekonomik performansları çeşitli makro kriterler yardımıyla incelenebilmektedir.

Araştırmada OECD örneklemi incelenmektedir. OECD kapsamında yer alan 38 ülke, 2024 yılında küresel nüfusun %17.2'sini, küresel ekonominin ise %43.6'sını oluşturmaktadır (WB, 2025; IMF, 2025). Küresel ekonominin önemli bölümünü üreten OECD, yüksek ekonomik potansiyele sahip ülkeleri içermektedir. Bu ülkeler kendi içinde gelir ve gelişmişlik düzeyi gibi çeşitli yönlerden ayrışmaktadır. Ayrıca ülkelerin ekonomik yapıları ve davranışları farklılık gösterebilmektedir. Dolayısıyla, araştırmacılar tarafından ülkelerin ekonomik performanslarının ölçülmesi ve bulguların karşılaştırılması politika yapıcılar açısından yol gösterici olabilmektedir. Analiz sonuçları kapsamında ekonomik performansı olumlu veya olumsuz etkileyen göstergelerin tespit edilmesi, geliştirilecek politikaların başarısını etkileyebilmektedir.

Çalışma literatüre dört açıdan katkı sunmaktadır. İlk olarak, çalışmanın örneklemi OECD ülkelerinin tamamını kapsamaktadır. Eksik ülke veya ülkelerin olmaması kapsayıcı bilgilerin sunulmasını sağlamaktadır. İkinci olarak, belirli yıllara odaklanan literatürün aksine çalışma 2000-2023 yıllarının ortalamasını alarak uzun vadeli performans analizi yapmaktadır. Üçüncü olarak, çalışmada yöntem olarak ÇKKV yöntemlerinden PIV yaklaşımı tercih edilmektedir. Böylece sınırlı ampirik kanıtın olduğu PIV yaklaşıma ilişkin literatürün geliştirilmesi amaçlanmaktadır. Son olarak, ÇKKV literatüründen farklı olarak, çalışma performans analizi sonrasında bulguları vergi yükü perspektifinden değerlendirmektedir. Ülkelerin analiz dönemindeki vergi yükü göstergeleri ile performans değerleri arasındaki bağlantı grafiksel analiz ile incelenmektedir.

Çalışmanın ikinci bölümü, ülkelerin makro düzlemdeki performanslarını ÇKKV yöntemleri ile inceleyen çalışmalara dair literatür araştırmasını sunmaktadır. Üçüncü bölümde, analizde kullanılan değişkenler açıklanmaktadır. Dördüncü bölüm, analizde kullanılan LOPCOW ve PIV yöntemlerinin metodolojileri hakkında bilgi vermektedir. Beşinci bölümde, ampirik bulgular tablo ve grafikler yardımıyla tartışılmakta ve çalışma politika önerilerini ele alan sonuç bölümüyle tamamlanmaktadır.

LİTERATÜR İNCELEMESİ

ÇKKV yöntemleri yoğun olarak belirli sektörlerde uygulanmasına karşın son yıllarda makro göstergeler üzerinden ülkelerin performanslarının ölçülmesinde kullanılmaktadır. Araştırmacılar sosyal, ekonomik, finansal, mali vb. alanlardan çeşitli göstergeleri modelleyerek ülkelerin performanslarını analiz etmekte ve karşılaştırma yapmaktadır. Tablo 1, çeşitli ÇKKV yöntemleri ile ülkelerin makro düzeyde performanslarına ilişkin literatür özetini göstermektedir.

FULL TEXTS BOOK

Tablo 1. Ülke Düzeyindeki ÇKKV Çalışmaları

Yazar(lar) / Yıl	Örneklem / Dönem	Değişkenler	Yöntem	Sonuç (İlk Üç Sıralama)
Urfalıoğlu ve Genç (2013)	AB ve Türkiye / 2010	KBGSYH, EBH, İH, İTH, İST ve ENF	ELECTRE PROMETHEE TOPSIS	ELECTRE: İrlanda, İsveç ve Hollanda PROMETHEE: İsveç, İrlanda ve Hollanda TOPSIS: Almanya, Hollanda ve Lüksemburg
Şahin ve Öztel (2017)	BRICS-T / 2015	GSYH, OYS, İGE, İŞZ ve NY	COPRAS	Rusya, Çin ve Brezilya
Koşaroğlu (2021)	E7 / 2010-2019	EBH, YO, İH, İTH, CD, İŞZ ve ENF	ARAS	Çin, Rusya ve Meksika
Avcı ve Yavuz (2022)	Türkiye / 1990-2019	EBH, İGE, GİNİ, ENF, İŞZ, CD ve KE	SAW ve MAUT	SAW: 2010, 2012 ve 2005 MAUT: 2004, 2005 ve 2013
Bektaş ve Baykuş (2023)	Türk Dünyası ve Rusya / 2010-2020	ENF, İŞZ, DG, SE ve SGP	MAIRCA	Rusya, Azerbaycan ve Kırgızistan
Apan ve Tiyek (2023)	Türkiye / 2008-2021	EBH, YO, İH, İTH, CD, İŞZ, ENF ve FO	MABAC	2015, 2013 ve 2014
Kahreman (2024)	D8 / 2011-2020	EBH, İHBO, NİH, İTHBO, DK, ENF, KBGSYH ve İŞZ	CoCoSo	Sıralama yıllık bazda değişmektedir.
OECD Ülkeleri İçin Yapılan Çalışmalar				
Demir ve Bakırcı (2014)	34 OECD Ülkesi / 2006-2010	İŞZ, DYY, İT, ÇS, GÜE, VG, KBGSYH, SGP, KFE, KE, GE ve İH	Veri Zarflama Analizi	Sıralama yıllık bazda değişmektedir.
Topçu ve Oralhan (2017)	35 OECD Ülkesi / 2010-2015	KBGSYH, İH, İTH, ENF, EBH ve İST	TOPSIS ve ELECTRE	Sıralama yönteme göre ve yıllık bazda değişmektedir.
Koca vd. (2019)	33 OECD Ülkesi / 2007-2016	KV, GV, KDV, GSYH ve SSY	ARAS	2007: ABD, Norveç ve Danimarka 2016: ABD, Danimarka ve Yeni Zelanda
Künç ve Yaşa (2019)	32 OECD Ülkesi / 2018	BGE, BGİ ve KB	TOPSIS ve VIKOR	TOPSIS: Danimarka, Norveç ve İsveç VIKOR: İrlanda, Yeni Zelanda ve İsviçre
Ersoy (2023)	34 OECD Ülkesi / 2020-2021	ENF, İŞZ, KBGSYH ve CD	MULTI-MOOSRAL SAW, MAUT ve PIV	Sıralama yönteme göre ve yıllık bazda değişmektedir.

Örneklem: AB: Avrupa Birliği, BRICS-T: Brezilya, Rusya, Hindistan, Çin, G. Afrika ve Türkiye, D8: Bangladeş, Mısır, Endonezya, İran, Malezya, Nijerya, Pakistan ve Türkiye, İİT: İslam İşbirliği Teşkilatı.
Değişkenler: BGE: Bütçe gelirleri, BGİ: Bütçe giderleri, CD: Cari denge, DG: Devlet gelirleri, DK: Döviz Kuru, DYY: Doğrudan yabancı yatırım, EBH: Ekonomik büyüme hızı, ENF: Enflasyon, FO: Faiz oranı, GE: Gelir endeksi, GİNİ: Gini katsayısı, GÜE: Gıda üretim endeksi, GV: Gelir Vergisi, İGE: İnsani gelişme endeksi, İH: İhracat, İHBO: İhracat büyüme oranı, İST: İstihdam oranı, İŞZ: İşsizlik, İTH: İthalat, İTHBO: İthalat büyüme oranı, KB: Kamu borcu, KBGSYH: Kişi başına düşen Gayri Safi Yurtiçi Hasıla, KDV: Katma Değer Vergisi, KE: Karbon emisyonu, KFE: Kıyaslamalı fiyat endeksi, KV: Kurumlar Vergisi, NİH: Net ihracat oranı, NY: Nüfus yoğunluğu, OYS: Ortalama yaşam süresi, SE: Sefalet endeksi, SGP: Satın alma gücü paritesi, SSY: Sabit sermaye yatırımları.VG: Vergi geliri, YO: Yatırım oranı, ÇS: Yıllık ortalama çalışma süresi, YÖN: Yönetişim göstergesi.
Yöntemler: ARAS: Additive Ratio Assesement, CoCoSo: Combined Compromise Solution, COPRAS: Complex Proportional Assessment, ELECTRE: Elemination and Choice Translating Reality English, MABAC: Multi-Attributive Border Approximation Area Comparison, MAIRCA: Multi Attributive Ideal-Real Comparative Analysis, MAUT: Multiple Attribute Utility Theory, MULTIMOOSRAL: Multi-Objective Optimization on The Basis of Simple Ratio Analysis, PROMETHEE: The Preference Ranking Organization Method for Enrichment Evaluation, SAW: Simple Additive Weighting, TOPSIS: Technique for Order Preference by Similarity to Ideal Solution, VIKOR: Vise Kriterijumska Optimizacija I Kompromisno Resenje.

Tablo 1 incelendiğinde, ülkelere ilişkin performans analizleri yapılırken genellikle GSYH, enflasyon, işsizlik, cari denge, borçlanma gibi ekonomik göstergelerin tercih edildiği görülmektedir. Tablo 1'in alt bölümünde OECD örneklemini analiz eden çalışmalara yer verilmektedir. Bu çalışmalarda belirli yılların veya kısa dönemlerin incelendiği gözlenmektedir. Bu çalışma, tüm OECD ülkelerini kapsaması, analiz dönemi itibarıyla geniş bir zaman dilimini incelemesi, farklı bir ÇKKV yöntemini kullanması ve bulguları vergi yükü çerçevesinde değerlendirmesiyle literatüre önemli katkılar sağlamaktadır.

VERİ

Çalışma, 38 OECD ülkesinde (Almanya, Amerika Birleşik Devletleri, Avustralya, Avusturya, Belçika, Birleşik Krallık, Çekya, Danimarka, Estonya, Finlandiya, Fransa, Hollanda, İrlanda, İspanya, İsrail, İsveç, İsviçre, İtalya, İzlanda, Japonya, Kanada, Kolombiya, G. Kore, Kosta Rika, Letonya, Litvanya, Lüksemburg, Macaristan, Meksika, Norveç, Polonya, Portekiz, Şili, Slovak Cumhuriyeti, Slovenya, Türkiye, Yeni Zelanda ve Yunanistan) ekonomik performansı incelemektedir. Ekonomik performans kapsamında kişi başı GSYH, yatırım, cari denge ve bütçe dengesi göstergeleri maksimum değerler olarak modelde yer alırken borçlanma, enflasyon ve işsizlik göstergeleri minimum değerler kapsamındadır. Veriler 2000-2023 döneminin ortalamasını kapsamaktadır. Modeldeki değişkenlere dair bilgiler Tablo 2'de gösterilmektedir.

Tablo 2. Değişkenlere İlişkin Tanımlayıcı Bilgiler

Değişkenler	Kısaltma	Tanımı	Kaynak
Maksimum			
Kişi Başı GSYH	KBGSYH	Kişi başı gayri safi yurtiçi hasıla (sabit fiyatlar, 2021 uluslararası dolar)	IMF (2025)
Yatırım	YAT	Yatırımların GSYH içindeki payı (%)	
Cari Denge	CD	Cari dengenin GSYH içindeki payı (%)	
Bütçe Dengesi	BD	Bütçe dengesinin GSYH içindeki payı (%)	
Minimum			
Enflasyon	ENF	Ortalama tüketici fiyatlarının değişimi (%)	
İşsizlik	İŞZ	İşsiz sayısının toplam işgücü içindeki payı (%)	
Borçlanma	BORÇ	Genel hükümet brüt borçlanma miktarının GSYH içindeki payı (%)	

YÖNTEM

LOPCOW Yöntemi

Analiz bölümünde ilk olarak, Ecer ve Pamucar (2022) tarafından geliştirilen LOPCOW (Logarithmic Percentage Change-driven Objective Weighting) yöntemi ile kriterlerin yani değişkenlerin objektif ağırlıklandırmaları hesaplanmaktadır. LOPCOW yöntemi, standart sapmaların yüzdelik değerleri ve ortalama kare hesaplamasıyla verilerin boyut farklılıklarını ortadan kaldırmaktadır. Tablo 3, dört aşamadan oluşan LOPCOW yönteminin metodolojisini göstermektedir (Ecer ve Pamucar, 2022):

FULL TEXTS BOOK

Tablo 3. LOPCOW Yönteminin Aşamaları

1. Aşama: Karar Matrisinin Oluşturulması	$x_{ij} = \begin{bmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & \cdots & x_{mn} \end{bmatrix}$ $i = 1,2,\dots,m; j = 1,2,\dots,n$	(1)
2. Aşama: Normalize Karar Matrisinin Elde Edilmesi	$r_{ij} = \frac{x_{ij} - x_{j_{min}}}{x_{j_{max}} - x_{j_{min}}}$ Fayda yönlü	(2)
	$r_{ij} = \frac{x_{j_{max}} - x_{ij}}{x_{j_{max}} - x_{j_{min}}}$ Maliyet yönlü	(3)
	$R = \begin{bmatrix} r_{11} & r_{12} & \cdots & r_{1n} \\ r_{21} & r_{22} & \cdots & r_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ r_{m1} & r_{m2} & \cdots & r_{mn} \end{bmatrix}$	(4)
3. Aşama: Her Kriter İçin Yüzdelik Değerlerin PV_{ij} Matrisinin Oluşturulması	$PV_{ij} = \left\lvert ln\left(\frac{\sqrt{\frac{\sum_{i=1}^{m} r_{ij}^2}{m}}}{\sigma} \right) * 100 \right\rvert$	(5)
4. Aşama: Objektif Ağırlıkların (W_j) Elde Edilmesi	$W_j = \frac{PV_{ij}}{\sum_{i=1}^{n} PV_{ij}}$	(6)

Yukarıdaki aşamalar yardımıyla kriterlere dair W_j kriter ağırlık değerleri hesaplanmaktadır. Bu değerler büyükten küçüğe doğru sıralanarak kriterlerin önem derecesini ortaya koymaktadır.

PIV Yöntemi

Analizin ikinci aşamasında, ağırlandırma değerleri elde edilen değişkenler üzerinden Muzaffal ve Muzakkir (2018) tarafından geliştirilen PIV (Proximity Indexed Value) yöntemi ile ülkelerin ekonomik performansları sıralanmaktadır. Bu yöntemde pozitif ideal çözüm hesaplanarak optimal çözüme en yakın olan alternatif belirlenmektedir. Yöntemin metodolojik aşamaları Tablo 4'te gösterilmektedir (Mufazzal ve Muzakkir, 2018: 430-431; Goswami vd., 2022: 1156):

FULL TEXTS BOOK

Tablo 4. PIV Yönteminin Aşamaları


1. Aşama: Karar Matrisinin Oluşturulması	$x_{ij} = \begin{bmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & \cdots & x_{mn} \end{bmatrix}$ $i = 1, 2, \ldots, m; j = 1, 2, \ldots, n$	(7)
	$r_{ij} = \frac{x_{ij}}{\sqrt{\sum_{i=1}^{m} x_{ij}^2}}$	(8)
2. Aşama: Normalize Karar Matrisi	$R = \begin{bmatrix} r_{11} & r_{12} & \cdots & r_{1n} \\ r_{21} & r_{22} & \cdots & r_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ r_{m1} & r_{m2} & \cdots & r_{mn} \end{bmatrix}$	(9)
3. Aşama: Ağırlıklandırılmış Normalize Karar Matrisi	$V_{ij} = w_{ij} * R_{ij}$	(10)
4. Aşama: Ağırlıklı Yakınlık Endeksinin (Z_{ij}) Hesaplanması	$Z_{ij} = \begin{cases} V_{max} - V_{ij} \ \ (fayda\ kriteri) \\ V_{ij} - V_{min} \ (maliyet\ kriteri) \end{cases}$	(11)
5. Aşama: Genel Yakınlık Değerlerinin (d_i) Belirlenmesi	$d_i = \sum_{j=1}^{n} V_j$	(12)

Elde edilen d_i değerleri küçükten büyüğe doğru sıralanmakta ve en küçük değere sahip olan alternatif en yüksek performansı göstermektedir.

BULGULAR

Uygulama aşamasının başlangıcında LOPCOW yöntemi ile değişkenlere dair ağırlık değerleri (W_j) hesaplanmaktadır. Grafik 1'e göre, analiz kapsamında en önemli değişkenler sırasıyla enflasyon, borç ve işsizlik göstergeleridir. Önem derecesi en düşük olan değişkenin ise kişi başına düşen GSYH olduğu tespit edilmektedir. Bu sonuçlar aynı zamanda incelenen dönemde OECD ülkeleri için enflasyon, borç ve işsizliğin başlıca ekonomik sorunlar olduğuna işaret etmektedir.

Grafik 1. LOPCOW Ağırlıklandırma Yönteminin Sonuçları



FULL TEXTS BOOK

Ağırlıklandırma değerlerinin elde edilmesinin ardından yedi gösterge üzerinden OECD ülkelerinin ekonomik performans sıralamaları PIV yöntemi ile belirlenmektedir. Bu kapsamda ilk olarak analizde kullanılan değişkenlere ait karar matrisi Tablo 5'te gösterilmektedir.

Tablo 5. LOPCOW ve PIV Yöntemlerinde Kullanılan Karar Matrisi

Ülkeler	KBGSYH	YAT	CD	BD	BORÇ	ENF	İŞZ
Avustralya	54541.06	25.71	-3.19	-2.10	28.82	2.86	5.37
Avusturya	60665.89	24.84	1.88	-2.34	75.85	2.43	5.28
Belçika	57842.12	23.76	1.17	-2.52	101.83	2.32	7.29
Kanada	53203.45	23.04	-0.85	-0.81	86.53	2.22	7.00
Şili	24696.52	24.35	-1.94	-1.12	17.25	3.78	8.23
Kolombiya	15112.34	21.00	-2.98	-2.47	44.76	5.33	11.84
Kosta Rika	19618.86	19.23	-3.81	-4.12	42.11	6.33	8.91
Çekya	41752.89	28.01	-1.87	-6.72	33.53	3.23	5.49
Danimarka	63161.72	21.69	5.82	0.61	42.83	1.88	4.46
Estonya	34891.99	29.28	-4.01	-0.97	9.46	4.25	8.56
Finlandiya	54364.86	23.74	1.45	-0.05	54.78	1.97	8.18
Fransa	51542.74	22.17	-0.27	-3.68	85.91	1.90	8.84
Almanya	57561.42	20.64	5.50	-1.29	67.43	2.04	6.12
Yunanistan	34879.49	18.41	-6.75	-2.68	150.73	2.21	15.32
Macaristan	30825.14	25.25	-2.84	-4.88	69.66	5.26	6.49
İzlanda	59485.23	21.97	-3.67	-1.23	90.01	4.92	4.50
İrlanda	74607.50	27.05	-0.63	-2.30	58.13	2.03	7.69
İsrail	39114.13	23.82	2.42	-3.51	70.70	1.67	7.60
İtalya	50543.01	20.30	0.11	-2.84	122.69	2.18	9.30
Japonya	42393.40	25.42	2.95	-5.62	205.86	0.32	3.87
Kore	40814.13	31.82	2.86	1.14	32.42	2.50	3.54
Letonya	26655.38	27.84	-4.96	-4.11	32.00	4.27	10.69
Litvanya	32083.02	21.76	-2.88	-2.31	31.24	3.42	10.11
Lüksemburg	132319.16	19.18	6.98	1.66	16.63	2.49	4.96
Meksika	20826.26	22.65	-1.10	-2.39	45.20	4.76	3.90
Hollanda	63403.64	20.81	6.41	-1.24	53.97	2.40	5.56
Yeni Zelanda	42325.24	23.21	-4.08	-0.92	30.93	2.55	4.84
Norveç	84739.10	24.96	12.43	-4.25	38.93	2.42	3.80
Polonya	29711.23	20.97	-2.93	-3.82	48.68	3.59	9.94
Portekiz	37003.32	20.76	-4.57	-3.96	99.30	2.18	9.21
Slovak Cumhuriyeti	29916.68	25.40	-3.72	-3.28	46.73	4.01	12.17
Slovenya	39160.02	24.14	1.38	-2.96	51.21	3.31	6.48
İspanya	43104.39	23.38	-1.88	-3.84	76.74	2.36	15.57
İsveç	56483.23	23.23	5.13	0.19	42.02	2.01	7.36
İsviçre	75733.18	26.52	7.62	0.14	45.25	0.61	2.92
Türkiye	22897.23	27.49	-3.36	-3.74	41.06	20.33	10.06
Birleşik Krallık	50354.40	17.54	-3.01	-3.24	70.26	2.52	5.49
ABD	62733.14	21.34	-3.42	-5.72	92.33	2.56	5.76

PIV yönteminin ikinci aşamasında, değişkenlerin karar matrisi üzerinden normalize karar matrisi hesaplanmakta ve Tablo 6'da raporlanmaktadır. Tablo 7 ve Tablo 8'de sırasıyla ağırlıklı normalize karar matrisi (LOPCOW yöntemi katsayıları ile) ve ağırlıklı yakınlık endeksi gösterilmektedir.

FULL TEXTS BOOK

Tablo 6. Normalize Karar Matrisi

Ülkeler	KBGSYH	YAT	CD	BD	BORÇ	ENF	İŞZ
Avustralya	0.1692	0.1760	-0.1224	-0.1094	0.0644	0.1023	0.1085
Avusturya	0.1882	0.1701	0.0722	-0.1220	0.1693	0.0868	0.1067
Belçika	0.1795	0.1627	0.0451	-0.1317	0.2273	0.0829	0.1474
Kanada	0.1651	0.1578	-0.0327	-0.0421	0.1932	0.0794	0.1414
Şili	0.0766	0.1667	-0.0744	-0.0586	0.0385	0.1351	0.1663
Kolombiya	0.0469	0.1438	-0.1144	-0.1290	0.0999	0.1904	0.2393
Kosta Rika	0.0609	0.1317	-0.1461	-0.2150	0.0940	0.2260	0.1800
Çekya	0.1295	0.1918	-0.0718	-0.3507	0.0748	0.1154	0.1109
Danimarka	0.1960	0.1485	0.2233	0.0317	0.0956	0.0673	0.0901
Estonya	0.1083	0.2005	-0.1540	-0.0505	0.0211	0.1519	0.1729
Finlandiya	0.1687	0.1626	0.0557	-0.0028	0.1223	0.0702	0.1654
Fransa	0.1599	0.1518	-0.0102	-0.1922	0.1918	0.0680	0.1787
Almanya	0.1786	0.1413	0.2110	-0.0676	0.1506	0.0728	0.1236
Yunanistan	0.1082	0.1260	-0.2589	-0.1401	0.3365	0.0789	0.3096
Macaristan	0.0956	0.1729	-0.1090	-0.2545	0.1555	0.1880	0.1313
İzlanda	0.1846	0.1504	-0.1409	-0.0643	0.2010	0.1756	0.0909
İrlanda	0.2315	0.1852	-0.0243	-0.1199	0.1298	0.0724	0.1555
İsrail	0.1214	0.1631	0.0929	-0.1832	0.1578	0.0597	0.1537
İtalya	0.1568	0.1390	0.0041	-0.1484	0.2739	0.0779	0.1879
Japonya	0.1315	0.1740	0.1133	-0.2934	0.4596	0.0114	0.0783
Kore	0.1266	0.2179	0.1098	0.0596	0.0724	0.0894	0.0715
Letonya	0.0827	0.1906	-0.1901	-0.2144	0.0714	0.1525	0.2160
Litvanya	0.0995	0.1490	-0.1105	-0.1208	0.0698	0.1223	0.2044
Lüksemburg	0.4105	0.1313	0.2679	0.0868	0.0371	0.0889	0.1003
Meksika	0.0646	0.1551	-0.0424	-0.1246	0.1009	0.1701	0.0788
Hollanda	0.1967	0.1425	0.2457	-0.0649	0.1205	0.0856	0.1124
Yeni Zelanda	0.1313	0.1589	-0.1565	-0.0480	0.0691	0.0910	0.0977
Norveç	0.2629	0.1709	0.4770	-0.2221	0.0869	0.0863	0.0768
Polonya	0.0922	0.1436	-0.1123	-0.1994	0.1087	0.1283	0.2008
Portekiz	0.1148	0.1421	-0.1754	-0.2067	0.2217	0.0780	0.1861
Slovak Cumhuriyeti	0.0928	0.1739	-0.1428	-0.1713	0.1043	0.1432	0.2459
Slovenya	0.1215	0.1653	0.0530	-0.1545	0.1143	0.1184	0.1309
İspanya	0.1337	0.1600	-0.0722	-0.2004	0.1713	0.0842	0.3147
İsveç	0.1752	0.1591	0.1967	0.0098	0.0938	0.0717	0.1487
İsviçre	0.2350	0.1816	0.2924	0.0074	0.1010	0.0218	0.0590
Türkiye	0.0710	0.1882	-0.1287	-0.1954	0.0917	0.7264	0.2033
Birleşik Krallık	0.1562	0.1201	-0.1154	-0.1692	0.1569	0.0901	0.1109
ABD	0.1946	0.1461	-0.1314	-0.2988	0.2061	0.0915	0.1165

FULL TEXTS BOOK

Tablo 7. Ağırlıklı Normalize Karar Matrisi

Ülkeler	KBGSYH	YAT	CD	BD	BORÇ	ENF	İŞZ
Avustralya	0.0142	0.0194	-0.0107	-0.0139	0.0125	0.0253	0.0163
Avusturya	0.0158	0.0187	0.0063	-0.0155	0.0330	0.0214	0.0160
Belçika	0.0150	0.0179	0.0039	-0.0167	0.0442	0.0205	0.0221
Kanada	0.0138	0.0174	-0.0029	-0.0053	0.0376	0.0196	0.0212
Şili	0.0064	0.0183	-0.0065	-0.0074	0.0075	0.0334	0.0250
Kolombiya	0.0039	0.0158	-0.0100	-0.0164	0.0194	0.0470	0.0359
Kosta Rika	0.0051	0.0145	-0.0128	-0.0273	0.0183	0.0558	0.0270
Çekya	0.0109	0.0211	-0.0063	-0.0446	0.0146	0.0285	0.0167
Danimarka	0.0164	0.0163	0.0195	0.0040	0.0186	0.0166	0.0135
Estonya	0.0091	0.0221	-0.0135	-0.0064	0.0041	0.0375	0.0260
Finlandiya	0.0141	0.0179	0.0049	-0.0004	0.0238	0.0173	0.0248
Fransa	0.0134	0.0167	-0.0009	-0.0244	0.0373	0.0168	0.0268
Almanya	0.0150	0.0156	0.0184	-0.0086	0.0293	0.0180	0.0186
Yunanistan	0.0091	0.0139	-0.0226	-0.0178	0.0655	0.0195	0.0465
Macaristan	0.0080	0.0190	-0.0095	-0.0323	0.0303	0.0464	0.0197
İzlanda	0.0155	0.0166	-0.0123	-0.0082	0.0391	0.0434	0.0137
İrlanda	0.0194	0.0204	-0.0021	-0.0152	0.0253	0.0179	0.0234
İsrail	0.0102	0.0179	0.0081	-0.0233	0.0307	0.0147	0.0231
İtalya	0.0131	0.0153	0.0004	-0.0189	0.0533	0.0192	0.0282
Japonya	0.0110	0.0191	0.0099	-0.0373	0.0895	0.0028	0.0118
Kore	0.0106	0.0240	0.0096	0.0076	0.0141	0.0221	0.0107
Letonya	0.0069	0.0210	-0.0166	-0.0273	0.0139	0.0376	0.0324
Litvanya	0.0083	0.0164	-0.0096	-0.0154	0.0136	0.0302	0.0307
Lüksemburg	0.0344	0.0145	0.0234	0.0110	0.0072	0.0220	0.0151
Meksika	0.0054	0.0171	-0.0037	-0.0158	0.0196	0.0420	0.0118
Hollanda	0.0165	0.0157	0.0215	-0.0082	0.0235	0.0211	0.0169
Yeni Zelanda	0.0110	0.0175	-0.0137	-0.0061	0.0134	0.0225	0.0147
Norveç	0.0220	0.0188	0.0417	-0.0282	0.0169	0.0213	0.0115
Polonya	0.0077	0.0158	-0.0098	-0.0253	0.0212	0.0317	0.0302
Portekiz	0.0096	0.0156	-0.0153	-0.0263	0.0431	0.0192	0.0279
Slovak Cumhuriyeti	0.0078	0.0191	-0.0125	-0.0218	0.0203	0.0353	0.0369
Slovenya	0.0102	0.0182	0.0046	-0.0196	0.0223	0.0292	0.0197
İspanya	0.0112	0.0176	-0.0063	-0.0255	0.0333	0.0208	0.0473
İsveç	0.0147	0.0175	0.0172	0.0012	0.0183	0.0177	0.0223
İsviçre	0.0197	0.0200	0.0255	0.0009	0.0197	0.0054	0.0089
Türkiye	0.0060	0.0207	-0.0112	-0.0248	0.0178	0.1794	0.0305
Birleşik Krallık	0.0131	0.0132	-0.0101	-0.0215	0.0305	0.0223	0.0167
ABD	0.0163	0.0161	-0.0115	-0.0380	0.0401	0.0226	0.0175

FULL TEXTS BOOK

Tablo 8. Ağırlıklı Yakınlık Endeksi

Ülkeler	KBGSYH	YAT	CD	BD	BORÇ	ENF	İŞZ
Avustralya	0.0202	0.0046	0.0524	0.0249	0.0084	0.0224	0.0074
Avusturya	0.0186	0.0053	0.0354	0.0265	0.0288	0.0186	0.0072
Belçika	0.0194	0.0061	0.0377	0.0278	0.0401	0.0177	0.0133
Kanada	0.0206	0.0066	0.0445	0.0164	0.0335	0.0168	0.0124
Şili	0.0280	0.0056	0.0482	0.0185	0.0034	0.0305	0.0161
Kolombiya	0.0305	0.0082	0.0517	0.0274	0.0153	0.0442	0.0271
Kosta Rika	0.0293	0.0095	0.0544	0.0384	0.0142	0.0530	0.0182
Çekya	0.0235	0.0029	0.0479	0.0556	0.0105	0.0257	0.0078
Danimarka	0.0180	0.0076	0.0222	0.0070	0.0145	0.0138	0.0047
Estonya	0.0253	0.0019	0.0551	0.0174	0.0000	0.0347	0.0171
Finlandiya	0.0203	0.0061	0.0368	0.0114	0.0197	0.0145	0.0160
Fransa	0.0210	0.0073	0.0426	0.0355	0.0332	0.0140	0.0180
Almanya	0.0194	0.0084	0.0232	0.0196	0.0252	0.0152	0.0097
Yunanistan	0.0253	0.0101	0.0643	0.0288	0.0614	0.0167	0.0376
Macaristan	0.0264	0.0050	0.0512	0.0434	0.0262	0.0436	0.0109
İzlanda	0.0189	0.0074	0.0540	0.0192	0.0350	0.0405	0.0048
İrlanda	0.0150	0.0036	0.0438	0.0263	0.0211	0.0151	0.0145
İsrail	0.0242	0.0060	0.0336	0.0343	0.0266	0.0119	0.0142
İtalya	0.0213	0.0087	0.0413	0.0299	0.0492	0.0164	0.0194
Japonya	0.0234	0.0048	0.0318	0.0483	0.0853	0.0000	0.0029
Kore	0.0238	0.0000	0.0321	0.0035	0.0100	0.0193	0.0019
Letonya	0.0275	0.0030	0.0583	0.0383	0.0098	0.0348	0.0236
Litvanya	0.0261	0.0076	0.0513	0.0264	0.0095	0.0274	0.0218
Lüksemburg	0.0000	0.0095	0.0183	0.0000	0.0031	0.0191	0.0062
Meksika	0.0290	0.0069	0.0454	0.0269	0.0155	0.0392	0.0030
Hollanda	0.0179	0.0083	0.0202	0.0193	0.0193	0.0183	0.0080
Yeni Zelanda	0.0234	0.0065	0.0553	0.0171	0.0093	0.0197	0.0058
Norveç	0.0124	0.0052	0.0000	0.0393	0.0128	0.0185	0.0027
Polonya	0.0267	0.0082	0.0515	0.0364	0.0170	0.0289	0.0213
Portekiz	0.0248	0.0083	0.0570	0.0373	0.0390	0.0164	0.0191
Slovak Cumhuriyeti	0.0266	0.0048	0.0541	0.0328	0.0162	0.0325	0.0281
Slovenya	0.0242	0.0058	0.0370	0.0307	0.0181	0.0264	0.0108
İspanya	0.0232	0.0064	0.0480	0.0365	0.0292	0.0180	0.0384
İsveç	0.0197	0.0065	0.0245	0.0098	0.0141	0.0149	0.0135
İsviçre	0.0147	0.0040	0.0161	0.0101	0.0156	0.0026	0.0000
Türkiye	0.0284	0.0033	0.0529	0.0359	0.0137	0.1765	0.0217
Birleşik Krallık	0.0213	0.0108	0.0517	0.0325	0.0264	0.0194	0.0078
ABD	0.0181	0.0079	0.0531	0.0490	0.0360	0.0198	0.0086

FULL TEXTS BOOK

Tablo 9. PIV Yöntemi Genel Yakınlık Endeksi (d_i) ve Sıralamalar


Ülkeler	d_i	Sıralama	Ülkeler	d_i	Sıralama
Lüksemburg	0.0563	1	Meksika	0.1658	20
İsviçre	0.0630	2	Birleşik Krallık	0.1700	21
Danimarka	0.0878	3	Litvanya	0.1700	22
Kore	0.0905	4	Fransa	0.1714	23
Norveç	0.0908	5	Çekya	0.1739	24
İsveç	0.1030	6	İzlanda	0.1799	25
Hollanda	0.1114	7	İtalya	0.1861	26
Almanya	0.1208	8	Polonya	0.1899	27
Finlandiya	0.1247	9	ABD	0.1926	28
Yeni Zelanda	0.1372	10	Slovak Cumhuriyeti	0.1952	29
İrlanda	0.1394	11	Letonya	0.1952	30
Avustralya	0.1404	12	Japonya	0.1966	31
Avusturya	0.1404	13	İspanya	0.1997	32
Şili	0.1503	14	Portekiz	0.2020	33
Kanada	0.1507	15	Kolombiya	0.2043	34
İsrail	0.1509	16	Macaristan	0.2065	35
Estonya	0.1516	17	Kosta Rika	0.2169	36
Slovenya	0.1531	18	Yunanistan	0.2443	37
Belçika	0.1620	19	Türkiye	0.3324	38

PIV yönteminin son aşamasında d_i değerleri hesaplanmakta ve ülkelerin ekonomik performansları bu değerlerin küçükten büyüğe doğru sıralanması ile belirlenmektedir. Genel yakınlık endeksi sonuçlarına göre, en düşük değere sahip olan Lüksemburg en iyi ekonomik performansı sergileyen ülke konumundadır. Lüksemburg'u İsviçre ve Danimarka takip etmektedir. Diğer taraftan en büyük d_i değerine sahip olan Türkiye, Yunanistan ve Kosta Rika en düşük ekonomik performans sergileyen ülkelerdir.

Analizin üçüncü aşamasında PIV yönteminden elde edilen performans değerleri, vergi yükü bağlamında incelenmektedir. Grafik 2, OECD ülkelerinin analiz dönemi ortalamasına göre vergi yükü oranlarını göstermektedir. İlgili dönemde OECD ülkelerinin genel vergi yükü ortalaması %32.8'dir. Ülke bazlı istatistiklere göre Danimarka (%45.8), İsveç (%44.2), Fransa (%44.1), Belçika (%43.4) ve Fransa'nın (%42.5) en yüksek vergi yüküne sahip iken, Meksika, Kolombiya, Şili, Kosta Rika ve Türkiye en düşük vergi yükü oranına sahiptir.

FULL TEXTS BOOK

Grafik 2. OECD Ülkelerinde Vergi Yükü (%) (2000-2023 Ortalaması)



Kaynak: OECD (2025)


Tablo 10'da OECD ülkelerinin PIV performans değerleri, %10'luk vergi dilimlerine göre sınıflandırılmaktadır. Buna göre, vergi yükü %10-%20 arasında 2 ülke, %20-%30 arasında 11 ülke, %30-%40 arasında 17 ülke ve %40-%50 arasında 8 ülke yer almaktadır. Vergi yükü sınıflandırmalarında yer alan ülkelerin PIV değerleri ortalaması Grafik 3 yardımıyla görselleştirilmektedir. Grafik 3'e göre, vergi yükü sınıflandırmalarında oranlar arttıkça, PIV değeri düşmektedir. Başka deyişle, vergi yükü oranlarının yüksek olduğu ülke sınıflandırmalarında ekonomik performans olumlu yönde etkilenmektedir.

Tablo 10. Vergi Yüküne Göre OECD Ülkelerinin Sınıflandırılması (2000-2023 Ortalaması)

%10-%20	PIV Skoru	%20-%30	PIV Skoru	%30-%40	PIV Skoru	%40-%50	PIV Skoru
Meksika	0.165812	**İsviçre**	0.063036	**Lüksemburg**	0.05625	**Danimarka**	0.087756
Kolombiya	0.204332	**Kore**	0.090457	**Hollanda**	0.11137	**Norveç**	0.090777
		İrlanda	0.13937	**Almanya**	0.120779	**İsveç**	0.10297
		Avustralya	0.14042	**Yeni Zelanda**	0.137158	**Finlandiya**	0.124747
		Şili	0.150303	**Kanada**	0.150744	**Avusturya**	0.140425
		Litvanya	0.170025	**İsrail**	0.150891	**Belçika**	0.162018
		ABD	0.192556	**Estonya**	0.151621	**Fransa**	0.171447
		Letonya	0.19524	**Slovenya**	0.153064	**İtalya**	0.186137
		Japonya	0.19655	**Birleşik Krallık**	0.170014		
		Kosta Rika	0.216918	**Çekya**	0.173893		
		Türkiye	0.332428	**İzlanda**	0.179869		
				Polonya	0.189916		
				Slovak Cumhuriyeti	0.195222		
				İspanya	0.19966		
				Portekiz	0.20197		
				Macaristan	0.206521		
				Yunanistan	0.244261		
Ortalama	0.185072	**Ortalama**	0.171573	**Ortalama**	0.164306	**Ortalama**	0.133285

FULL TEXTS BOOK

Grafik 3. Vergi Yükü Sınıflamasına Göre OECD Ülkelerinin Ortalama PIV Skorları



SONUÇ VE POLİTİKA ÖNERİLERİ

Çalışma, 38 OECD ülkesinin 2000-2023 dönemi ekonomik performansını kişi başı GSYH, işsizlik, enflasyon, yatırım, cari denge, bütçe dengesi ve borçlanma değişkenleri üzerinden PIV yöntemi ile analiz etmektedir. Analiz sonucuna göre, ekonomik performansı en yüksek ülkeler Lüksemburg, İsviçre, Danimarka, G. Kore ve Norveç'tir. Ekonomik performansı en düşük olan ülkeler ise Türkiye, Yunanistan, Kosta Rika, Macaristan ve Kolombiya şeklinde sıralanmaktadır. Ekonomik performans açısından listenin başındaki ülkelerde kişi başı GSYH'nin çok yüksek olması dikkat çekmektedir. Ayrıca ilk dört sıralamadaki ülkelerin cari açık veya bütçe açığı sorunu yaşamadığı tespit edilmektedir. En alt sıralamada olan ülkelerde ise cari ve bütçe dengelerinin negatif olduğu görülmektedir. Ayrıca bu ülkelerde işsizlik, enflasyon ve borçlanma rasyolarının da göreceli olarak yüksek olması, ekonomik performansı baskılamaktadır.

Çalışmanın bir diğer çıktısı, ülke sınıflandırmalarında vergi yükü oranı ile ekonomik performans arasında pozitif ilişkinin keşfedilmesidir. Hükümetlerin vergi gibi mali araçlar ile ekonomiye müdahalesi, birçok iktisadi görüşün temel tartışmaları arasında yer almaktadır. Örneğin minimal devlet müdahalesini savunan Klasik yaklaşım, vergi veya harcama gibi mali araçlar ile ekonomiye müdahalenin sınırlı olması gerektiğini ileri sürmektedir. Buna karşılık 1929 Buhranı sonrası önem kazanan Keynesyen yaklaşım, para ve maliye politikası araçları ile ekonomiye müdahale edilmesini olağan kabul etmektedir. Çalışmada vergi yükü ile ekonomik performans arasında pozitif bağlantının bulunması, OECD ülkelerinde Keynesyen ve benzeri iktisadi görüşlerin geçerli olduğuna işaret etmektedir. Bu çerçevede, genel olarak ekonomik performansı yüksek olan OECD ülkelerinde vergi politikalarının etkin kullanıldığı ve piyasa aktörlerinin (tüketiciler ve üreticiler) davranışları üzerinde bozucu etkilerinin olmadığı ifade edilebilir.

Gelecekteki araştırmalar farklı örneklemleri ve/veya dönemleri yeni geliştirilen ÇKKV yöntemleri ile inceleyebilirler. Ayrıca araştırmacılar ekonomik, sosyal, kültürel vb. alanlarda ülkelerin performansını ölçebilir ve çeşitli göstergeler ile ilişkilendirebilirler.

KAYNAKLAR

Apan, M., & Tiyek, R. (2023). Makroekonomik performans ve Entegre CRITIC tabanlı MABAC karar verme yaklaşımı: Türkiye ekonomisinin 2008-2021 dönem verisinden kanıtlar. *Süleyman Demirel Üniversitesi Vizyoner Dergisi,* 14(100. Yıl Özel Sayısı), 44-65.

FULL TEXTS BOOK

Avcı, T., & Yavuz, E. (2022). Türkiye'nin maliye politikası performansının SAW ve MAUT yöntemiyle analizi. Kübra İrday Demir, Pervin Zülfigarova (eds.), 8th Cukurova International Scientific Researches Conference (pp. 1100-1113), IKSAD Publishing.

Bektaş, S., & Baykuş, O. (2023). CRITIC ve MAIRCA yöntemleriyle Türk Dünyası ülkeleri, Türkiye ve Rusya'nın 2010-2020 dönemi için makroekonomik performanslarının analizi. *Uluslararası İktisadi ve İdari İncelemeler Dergisi,* (39), 107-122.

Demir, A. & Bakırcı F. (2014). OECD üyesi ülkelerin ekonomik etkinliklerinin veri zarflama analiziyle ölçümü. *Atatürk Üniversitesi İktisadi ve İdari Bilimler Dergisi,* 28(2), 109-132.

Ecer, F., & Pamucar, D. (2022). A novel LOPCOW-DOBI multi-criteria sustainability performance assessment methodology: An application in developing country banking sector. *Omega*, 112, 102690.

Ersoy, N. (2023). MEREC-MULTIMOOSRAL modeli ile OECD ülkelerinin makroekonomik performanslarının değerlendirilmesi. *Gümüşhane Üniversitesi Sosyal Bilimler Dergisi,* 14(2), 471-491.

Goswami, S. S., Mohanty, S. K., & Behera, D. K. (2022). Selection of a green renewable energy source in India with the help of MEREC integrated PIV MCDM tool. *Materials Today: Proceedings*, 52, 1153-1160.

IMF (2025). International Monetary Fund, The World Economic Outlook database, https://www.imf.org/en/Publications/WEO/weo-database/2025/april (Erişim Tarihi: 18.06.2025).

Kahreman, Y. (2024). D8 ülkelerinin ekonomik performanslarının CRITIC/LOPCOW-COCOSO modeli ile değerlendirilmesi. *Anadolu Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi,* 25(1), 534-559.

Koca, G., Ekinci, F., & Şimşek, M. (2019). Vergi gelirleri bakımından OECD ülkelerinin performansının Bütünleşik ENTROPİ-ARAS yöntemi ile değerlendirilmesi. *Elektronik Sosyal Bilimler Dergisi*, 18(70), 964-985.

Koşaroğlu, Ş. M. (2021). E7 ülkelerinin makroekonomik performanslarının ENTROPİ ve ARAS yöntemleriyle karşılaştırılması. *Dumlupınar Üniversitesi Sosyal Bilimler Dergisi,* (68), 203-221.

Künç, G. Y., & Yaşa, A. A. (2019). Türkiye-OECD ülkelerine ait bütçe göstergelerinin TOPSIS ve VIKOR yöntemleri ile karşılaştırılması. *Bitlis Eren Üniversitesi Sosyal Bilimler Dergisi,* 8(2), 366-384.

Mendoza, G. A., & Prabhu, R. (2000). Multiple criteria decision making approaches to assessing forest sustainability using criteria and indicators: a case study. *Forest Ecology and Management*, 131(1-3), 107-126.

Mufazzal, S., & Muzakkir, S. M. (2018). A new multi-criterion decision making (MCDM) method based on proximity indexed value for minimizing rank reversals. *Computers & Industrial Engineering*, 119, 427–438.

OECD (2025). OECD Data Explorer, https://data-explorer.oecd.org/?lc=en (Erişim Tarihi: 23.06.2025).

Şahin, C., & Öztel, A. (2017). Ülkelerin yaşanabilirlik düzeylerinin COPRAS yöntemiyle karşılaştırmalı analizi: BRICS ülkeleri ve Türkiye, *Uluslararası Batı Karadeniz Sosyal ve Beşeri Bilimler Dergisi*, 1(1), 75-84.

Topçu, B. A., & Oralhan, B. (2017). Türkiye ve OECD ülkeleri'nin temel makroekonomik göstergeler açısından çok kriterli karar verme yöntemleri ile karşılaştırılması. *Journal of Academic Value Studies*, 3(14), 260-277.

Urfalıoğlu, F., & Genç, T. (2013). Çok kriterli karar verme teknikleri ile Türkiye'nin ekonomik performansının Avrupa Birliği Üye Ülkeleri ile karşılaştırılması. *Marmara Üniversitesi İktisadi ve İdari Bilimler Dergisi*, 35(2), 329-360.

WB (2025). World Bank, World Economic Outlook database: April 2025, https://www.imf.org/en/Publications/WEO/weo-database/2025/april (Erişim Tarihi: 23.06.2025).

FULL TEXTS BOOK

SOSYO-EKONOMİK VE MALİ PERFORMANSIN BÖLGESEL ANALİZİ: TÜRKİYE İBBS DÜZEY-2 ÜZERİNE BİR UYGULAMA

REGIONAL ANALYSIS OF SOCIO-ECONOMIC AND FISCAL PERFORMANCE: AN IMPLEMENTATION ON TÜRKİYE NUTS LEVEL-2

Doç. Dr. Ersin YAVUZ

Pamukkale Üniversitesi, İktisadi ve İdari Bilimler Fakültesi, Maliye Bölümü, Denizli, Türkiye

Öğr. Gör. Tahsin AVCI

Pamukkale Üniversitesi, Uygulamalı Bilimler Fakültesi, Uluslararası Ticaret ve Lojistik Bölümü, Denizli, Türkiye

ÖZET

Çalışmanın amacı, Türkiye'de İBBS Düzey-2 alt bölgelerinin sosyo-ekonomik ve mali göstergeler açısından performansını analiz etmektir. Bu kapsamda, 26 bölgenin performansı çok kriterli karar verme yaklaşımlarından COBRA (Comprehensive Distance-Based Ranking) yöntemi ile ölçülmektedir. Modelde ekonomik (kişi başına GSYH, işgücüne katılım, işsizlik, enflasyon, ithalat ve ihracat), sosyal (yoksulluk ve gelir dağılımı) ve mali (vergi gayreti ve vergi gelirlerinin toplam tahsilat içindeki payı) alanlarda olmak üzere toplam 10 gösterge kullanılmaktadır. Analizin ilk aşamasında değişkenler ENTROPİ yöntemi ile ağırlıklandırılmaktadır. İkinci aşamada ise COBRA yaklaşımı ile bölgelerin performans endeksleri hesaplanmakta ve sıralama yapılmaktadır. Analiz sonucuna göre, TR10 (İstanbul), TR51 (Ankara), TR42 (Kocaeli, Sakarya, Düzce, Bolu, Yalova), TR31 (İzmir) ve TR21 (Tekirdağ, Edirne, Kırklareli) bölgeleri en yüksek endeks değeri alırken, TRC2 (Şanlıurfa, Diyarbakır), TRB2 (Van, Muş, Bitlis, Hakkari), TRA2 (Ağrı, Kars, Iğdır, Ardahan), TRC3 (Mardin, Batman, Şırnak, Siirt) ve TRC1 (Gaziantep, Adıyaman, Kilis) bölgeleri en düşük endekse sahiptir. Bulgular, Ankara ve İzmir'in yanı sıra Marmara bölgesindeki illerin yüksek performans gösterdiğini ortaya koymaktadır. Buna karşılık, Doğu Anadolu ve Güneydoğu Anadolu bölgelerindeki illerin performans düzeylerinin görece düşük olduğu tespit edilmektedir. Bu durum, bölgesel kalkınma farklarının devam ettiğine işaret etmektedir. Dolayısıyla hükümetlerin özellikle performansı düşük bölgelere yönelik teşvik mekanizmaları uygulaması, bölgesel eşitsizliklerin azaltılmasına ve ülke genelinde dengeli bir kalkınmanın sağlanmasına katkı sağlayacaktır.

Anahtar kelimeler: COBRA Yöntemi, Çok Kriterli Karar Verme, İBBS Düzey-2, Performans Analizi.

ABSTRACT

The aim of paper is to analyze the performance of the Nomenclature of Territorial Units for Statistics (NUTS) Level-2 sub-regions in Türkiye in terms of socio-economic and fiscal indicators. Within this scope, the performance of 26 regions is measured by COBRA (Comprehensive Distance-Based Ranking) method, one of the multi-criteria decision-making approaches. In the model, a total of 10 indicators are used in economic (GDP per capita, labor force participation, unemployment, inflation, imports and exports), social (poverty and income distribution), and fiscal (tax effort and share of tax revenues in total collection) fields. In the first stage of the analysis, variables are weighted with the ENTROPY method. In the second stage, the performance indexes of the regions are calculated and ranked with the COBRA approach. According to the analysis, TR10 (İstanbul), TR51 (Ankara), TR42 (Kocaeli, Sakarya, Düzce, Bolu, Yalova), TR31 (İzmir) and TR21 (Tekirdağ, Edirne, Kırklareli) regions obtain the highest index values, while TRC2 (Şanlıurfa, Diyarbakır), TRB2 (Van, Muş, Bitlis, Hakkari), TRA2 (Ağrı, Kars, Iğdır, Ardahan), TRC3 (Mardin, Batman, Şırnak, Siirt) and TRC1 (Gaziantep, Adıyaman, Kilis) regions have the lowest index values. The findings reveal that Ankara and İzmir, as well as provinces in the Marmara region have high performance. Conversely, provinces in Eastern Anatolia and Southeastern Anatolia have relatively low performance levels. This points to the existence of regional development disparities. Hence, the implementation of incentive mechanisms by

FULL TEXTS BOOK

governments, especially for low-performing regions, will contribute to reducing regional inequalities and ensuring a balanced development across the country.

Keywords: COBRA Method, Multi-Criteria Decision Making, NUTS Level-2, Performance Analysis.

GİRİŞ

Bölge kavramının tanımı üzerine görüş birliği bulunmamaktadır. Kullanıldığı alana bağlı olarak kavramın ifade ettiği anlam değişebilmektedir. Bu kapsamda, bölge kavramı bazen ekonomik, sosyal ve kültürel açıdan homojen olan bir mekan parçasını ifade ederken, bazen de işlevsel açıdan bütünlük gösteren birimleri belirtmektedir. Tüm bunlardan bağımsız olarak bölge yani mekan kavramı, kalkınmanın kritik bileşenlerinden biri olarak kabul edilmektedir. Bölgesel organizasyonlar tarafından oluşturulan vizyon, ulusal ve uluslararası rekabette etkin rol oynayabilmektedir (DPT, 2000: 7-8).

Kalkınma süreçlerinin coğrafi olarak farklılık göstermesi, ekonomik faaliyetlerin belirli mekanlarda yoğunlaşması, olanakları ve/veya doğal kaynakları nedeniyle belirli kent merkezlerinde daha fazla üretim yapılması gibi faktörler bölgelerin homojen ve eşit gelişmesini engellemiştir (Köse vd., 2012: 78). Bu durum, iktisadın temel tartışma konularından biri olan bölgelerarası eşitsizlik sorunsalına neden olmuştur. Küresel düzeyde ülkelerin gelişmiş, gelişmekte olan veya az gelişmiş olarak sınıflandırılmasının temelinde mekansal eşitsizliklerin olduğu ifade edilebilir (IMF, 2025). Benzer şekilde, ülkeleri çeşitli göstergeler üzerinden inceleyerek düşük, orta, yüksek ve çok yüksek olarak sınıflandıran İnsani Gelişme Endeksi, birçok faktörün yanı sıra mekansal düzeydeki dengesizliklerin bir sonucu olarak görülebilir (UNDP, 2025).

Küresel düzeye benzer şekilde ulusal sınırlar içinde de bölgelerarası gelişmişlik farklılıkları bulunmaktadır. Bu kapsamda Türkiye'de bölgesel kalkınma ve bölgelerarası dengesizliklerin giderilmesine ilişkin çabalar Cumhuriyetin ilk yıllarından itibaren görülmektedir. İlk dönemlerde kamu yatırımlarının ve sanayi tesislerinin Anadolu'ya yaygınlaştırılması yönünde adımlar atılmıştır. 1960 yılında Devlet Planlama Teşkilatı'nın (DPT) kurulmasının ardından bölgesel farklılıklar ile daha yoğun mücadele edilmiştir. Bu kapsamda Doğu Marmara Bölgesi, Çukurova Bölgesi, Antalya ve Zonguldak illerine ilişkin bölgesel kalkınma projeleri, Güneydoğu Anadolu Projesi, Çorum-Çankırı kırsal kalkınma projesi, Doğu Karadeniz ve Yeşilırmak Havzası projeleri vb. örnekler verilebilir. 2000'li yılların başında Avrupa Birliği uyum süreci kapsamında DPT ve Türkiye İstatistik Kurumu (TÜİK) koordinasyonunda İstatistiki Bölge Birimleri Sınıflandırılması (İBBS) tanımlanmıştır. İBBS, 28.08.2002 tarihli Bakanlar Kurulu Kararı ile yürürlüğe girmiştir. Sınıflandırma kapsamında Düzey 1 için 12 bölge, Düzey 2 için 26 bölge ve Düzey 3 için 81 il belirlenmiştir (Kalkınma Ajansları, 2025).

Çalışma, örneklem olarak Düzey 2 kapsamında yer alan 26 bölgeye odaklanmaktadır (Tablo 1). Bu sınıfta yer alan bölgeler arasında önemli farklılıklar bulunmaktadır. Örneğin 2023 yılında kişi başı GSYH miktarı incelendiğinde, en üst sırada yer alan TR10 (İstanbul) bölgesi, en alt sırada yer alan TRB2 (Van, Muş, Bitlis, Hakkari) bölgesinden 4.14 kat daha yüksek miktara sahiptir. Gelişmişlik farkı nedeniyle 2024 yılında TR51 (Ankara) bölgesinin net göç hızı %8.9 olarak gerçekleşirken, bu oran TRB2 (Van, Muş, Bitlis, Hakkari) bölgesinde %-20 şeklindedir (TÜİK, 2025). Bölgesel eşitsizliklere dair benzer birçok istatistiki örnek verilebilir. Dolayısıyla Türkiye'de bölgeler arasındaki dengesizliklere ilişkin analizler yapmak ve elde edilen ampirik bulgular temelinde politika önerileri geliştirmek büyük öneme sahiptir.

FULL TEXTS BOOK

Tablo 1. İBBS Düzey-2 Bölgeleri

BÖLGE KODU	BÖLGE ADI	BÖLGE KODU	BÖLGE ADI	BÖLGE KODU	BÖLGE ADI
TR10	İstanbul	TR52	Konya, Karaman	TR90	Trabzon, Ordu, Giresun, Rize, Artvin, Gümüşhane
TR21	Tekirdağ, Edirne, Kırklareli	TR61	Antalya, Isparta, Burdur	TRA1	Erzurum, Erzincan, Bayburt
TR22	Balıkesir, Çanakkale	TR62	Adana, Mersin	TRA2	Ağrı, Kars, Iğdır, Ardahan
TR31	İzmir	TR63	Hatay, Kahramanmaraş, Osmaniye	TRB1	Malatya, Elazığ, Bingöl, Tunceli
TR32	Aydın, Denizli, Muğla	TR71	Kırıkkale, Aksaray, Niğde, Nevşehir, Kırşehir	TRB2	Van, Muş, Bitlis, Hakkari
TR33	Manisa, Afyon, Kütahya, Uşak	TR72	Kayseri, Sivas, Yozgat	TRC1	Gaziantep, Adıyaman, Kilis
TR41	Bursa, Eskişehir, Bilecik	TR81	Zonguldak, Karabük, Bartın	TRC2	Şanlıurfa, Diyarbakır
TR42	Kocaeli, Sakarya, Düzce, Bolu, Yalova	TR82	Kastamonu, Çankırı, Sinop	TRC3	Mardin, Batman, Şırnak, Siirt
TR51	Ankara	TR83	Samsun, Tokat, Çorum, Amasya		

Kaynak: TÜİK (2025)

Çalışma, çok kriterli karar verme yöntemlerinden yararlanarak İBBS Düzey-2 bölgelerinde sosyo-ekonomik ve mali performans analizi yapmaktadır. Bu kapsamda çalışmanın ilgili literatüre iki temel katkısı bulunmaktadır. İlk olarak, çalışma bölgesel performans analizi yaparken sosyo-ekonomik göstergelere ek olarak mali göstergeleri modele dahil etmektedir. Bölgesel performansa yönelik literatürde çeşitli sosyo-ekonomik göstergeler olmasına karşın, mali göstergelere dair ampirik kanıtlar sınırlıdır. İkinci olarak, çalışma çok kriterli karar verme yaklaşımı olarak COBRA yöntemini uygulamaktadır. Çalışma, COBRA yöntemi ile ilgili yazına metodolojik yenilik sunmaktadır.

Çalışmanın diğer bölümleri şu şekilde düzenlenmektedir: İkinci bölüm, çok kriterli karar verme yöntemleri, bölgesel performans analizleri ve COBRA yöntemine dair literatürü açıklamaktadır. Üçüncü bölüm, modelde yer alan değişkenlere dair tanımlayıcı bilgileri sunmaktadır. Dördüncü bölüm, COBRA yönteminin metodolojisine yer vermektedir. Beşinci bölüm uygulamaya dair sonuçları, son bölüm ise bulgulara ilişkin tartışmaları ve politika önerilerini içermektedir.

LİTERATÜR İNCELEMESİ

Genellikle mikro alanlarda uygulanan çok kriterli karar verme yöntemleri, son yıllarda makro göstergeler düzeyinde ilgi görmektedir. Araştırmacılar bu yöntemleri kullanarak belirlenen göstergeler üzerinden ülkelerin performanslarını analiz etmektedir. Örneğin, Urfalıoğlu ve Genç (2013) Avrupa Birliği ve Türkiye'nin ekonomik performansını kişi başı GSYH, ekonomik büyüme, istihdam, ihracat, ithalat ve enflasyon göstergeleri üzerinden ELECTRE (Elemination and Choice Translating Reality English), PROMETHEE (The Preference Ranking Organization Method for Enrichment Evaluation) ve TOPSIS (Technique for Order Preference by Similarity to Ideal Solution) yöntemleriyle tespit etmektedir. Demir ve Bakırcı (2014) OECD ülkelerinin performansını Veri Zarflama Analizi ile kişi başı GSYH, işsizlik, ortalama çalışma süresi, doğrudan yabancı yatırım, ihracat, ithalat, vergi geliri, satınalma gücü paritesi, kişi başı karbon emisyonu, gıda üretim, gelir ve fiyat endeksleri üzerinden araştırmaktadır. Yalçın ve Yavuz (2016) İslam İşbirliği Teşkilatı'na üye ülkelerin ekonomik performanslarını istihdam oranı, insani gelişme endeksi, kişi başı GSYH, doğrudan yabancı yatırım, işsizlik oranı ve enflasyon oranı üzerinden ARAS (Additive Ratio Assesement) yöntemi ile analiz etmektedir. Şahin ve Öztel (2017) GSYH, ortalama yaşam süresi, insani gelişme endeksi, işsizlik ve nüfus yoğunluğu yardımıyla BRICS-T ülkelerinde COPRAS (Complex Proportional Assessment) yöntemiyle performans değerlendirmesi yapmaktadır. Yaraşır Tülümce ve Yavuz (2019) Avrupa Birliği ve Türkiye'de performans ölçümünü büyüme oranı, insani gelişme endeksi, yönetişim rasyosu, enflasyon oranı, işsizlik oranı ve gini katsayısı göstergeleri yardımıyla ARAS yöntemi ile gerçekleştirmektir. Samut (2024) Türkiye'de makroekonomik performansı ekonomik büyüme, enflasyon, istihdam, ihracat ve ithalat göstergeleri üzerinden TOPSIS yöntemiyle araştırmaktadır.

Literatürde çok kriterli karar verme yöntemleriyle ülkelerin yanı sıra Türkiye özelinde il veya bölge düzeyinde de performans analizleri yapılmaktadır. Gazibey vd. (2014) 52 gösterge üzerinden illerin

FULL TEXTS BOOK

sürdürülebilirlik performansını TOPSIS yöntemiyle araştırmaktadır. Genç (2016) Doğu ve Güneydoğu Anadolu Bölgesi'nden seçilen illerde sosyo-ekonomik performansı 9 değişken üzerinden VIKOR yöntemi ile incelemektedir. Yavuz vd. (2017) sosyal güvenlik kapsamının il nüfusuna oranı, sosyal güvenlik kapsamındaki aktif çalışanların il nüfusuna oranı, iş yeri sayısı, sosyal güvenlik kapsamındaki bakmakla yükümlü tutulanların il nüfusuna oranı ve sosyal güvenlik kapsamındaki emeklilerin il nüfusuna oranı üzerinden 81 ilin sosyal güvenlik alanındaki performansını MOORA (Multi-Objective Optimization by Ratio Analysis) oran ve referans yöntemleriyle ölçmektedir. Belgin ve Avşar (2019) Düzey-2 ve iller düzeyinde ar-ge ve yenilik performanslarını 27 değişken üzerinden Gri İlişkisel Analiz Yöntemi ile değerlendirmektedir. Erikli ve Türkoğlu (2020) Düzey-2 bölgelerinde işgücü piyasası performansını 5 gösterge üzerinden EDAS (Evaluation Based on Distance from Average Solution) yöntemi ile ölçmektedir. Çetin ve Kuvat (2022) Düzey-2 bölgelerinde ekonomik performansı 9 gösterge üzerinden COPRAS yöntemi ile analiz etmektedir. Örtlek (2024) Düzey-2 bölgelerinde sosyo-ekonomik performansı ekonomi, turizm, eğitim, istihdam, gelir dağılımı ve yoksulluk alanlarındaki göstergeler üzerinden COPRAS yöntemi ile incelemektedir.

Bu çalışmada kullanılan COBRA yöntemi, görece yeni geliştirilen yöntemlerden biri olduğu için çalışma sayısı sınırlıdır. Bu yöntemi kullanarak Çıtak ve Ünlü (2024) BIST kapsamında işlem gören bankaların sürdürülebilirlik raporlarını; Şimşek (2024) Türkiye'deki üniversitelerin performansını; Yürüyen ve Topal (2025) G7 ülkelerinin lojistik performansını; Saraç ve Karamaşa (2025) OECD ülkelerinde yaşam kalitesi performansını analiz etmektedir. Literatür genel olarak incelendiğinde, bu çalışma Düzey-2 bölgeleri için sosyo-ekonomik göstergelere mali göstergeleri dahil ederek ve güncel yöntemler arasında yer alan COBRA metodunu uygulayarak ilgili yazına katkı sunmaktadır.

VERİ

Çalışma, Düzey-2 kapsamında Türkiye'de bölgesel sosyo-ekonomik ve mali performansı araştırmaktadır. Analiz dönemi olarak verilerin erişilebilirliği dikkate alınarak 2021 yılı belirlenmiştir. Modelde performans sıralaması 5 maksimum ve 5 minimum olmak üzere toplam 10 değişken üzerinden yapılmaktadır. Tablo 2, analizde kullanılan değişkenlere dair tanımlayıcı bilgiler ve kaynakları hakkında detaylı bilgi sunmaktadır.

Tablo 2. Değişkenlere İlişkin Tanımlayıcı Bilgiler

	Değişken	Türü	Kısaltma	Tanım	Kaynak
Maksimum	Kişi Başı GSYH	Ekonomik	KBGSYH	Kişi Başına Gayrisafi Yurtiçi Hasıla (2009 Bazlı) ($)	TÜİK (2025)
	İşgücüne Katılma Oranı	Ekonomik	İKO	Yaş Gruplarına Göre İşgücüne Katılma Oranı (15 Yaş ve Üzeri) (%)	TÜİK (2025)
	İhracat	Ekonomik	İH	Ekonomik Faaliyetlere Göre İhracat (1000 ABD Doları) (Genel Toplam)	TÜİK (2025)
	Vergi Gayreti	Mali	VG	Vergi Tahsilinin Vergi Tahakkuku İçindeki Payı (%)	GİB (2025)
	Vergi Geliri Payı	Mali	VGP	Vergi Gelirinin Toplam Vergi Gelirleri İçindeki Payı (%)	GİB (2025)
Minimum	İşsizlik	Ekonomik	İŞZ	Yaş Gruplarına Göre İşsizlik Oranı (15 Yaş ve Üzeri) (%)	TÜİK (2025)
	Enflasyon	Ekonomik	ENF	Yıl Sonu İtibariyle Bir Önceki Yılın Aynı Ayına Göre Tüketici Fiyatları Endeksi Değişim Oranı (2003=100) (%)	TÜİK (2025)
	İthalat	Ekonomik	İT	Ekonomik Faaliyetlere Göre İthalat (1000 ABD Doları) (Genel Toplam)	TÜİK (2025)
	Yoksulluk	Sosyal	YOK	Eşdeğer Hanehalkı Kullanılabilir Fert Gelirine Göre Yoksul Sayıları ve Yoksulluk Oranları (Medyan Gelirin %60'ına Göre / Yoksulluk Oranı) (%)	TÜİK (2025)
	Gini Katsayısı	Sosyal	GİNİ	Eşdeğer Hanehalkı Kullanılabilir Fert Gelirine Göre Gini Katsayısı	TÜİK (2025)

FULL TEXTS BOOK

YÖNTEM

ENTROPİ Yöntemi

Analizde uygulanan değişkenlere ilişkin ağırlıklandırma işlemi, 1865 yılında Rudolph Clausius tarafından literatüre kazandırılan ENTROPİ yöntemi ile yapılmaktadır (Zhang, 2011: 444). Mühendislik, matematik, fizik ve bilgi teorisi alanlarında sıklıkla kullanılmasından dolayı Claude E. Shannon (1948) bu yöntemi enformasyon teorisiyle açıklamaktadır. ENTROPİ metodu kriterlerin önem derecelerinin belirlenmesinde kullanılan objektif ağırlıklandırma yöntemidir. ENTROPİ yönteminin matematiksel gösterimini ifade eden aşamalar Tablo 3'te gösterilmektedir (Wu vd., 2011: 5163-5165; Li vd., 2011: 2085-2091).

Tablo 3. ENTROPİ Yönteminin Metodolojik Aşamaları

Adım 1: Başlangıç Karar Matrisinin (D) Düzenlenmesi	$\mathbf{D} = \begin{bmatrix} \boldsymbol{x_{11}} & \boldsymbol{x_{12}} & \cdots & \boldsymbol{x_{1n}} \\ \boldsymbol{x_{21}} & \boldsymbol{x_{22}} & \cdots & \boldsymbol{x_{2n}} \\ \vdots & \vdots & \ddots & \vdots \\ \boldsymbol{x_{m1}} & \boldsymbol{x_{m2}} & \cdots & \boldsymbol{x_{mn}} \end{bmatrix} \quad \boldsymbol{i = 1, 2, \dots, m; j = 1, 2, \dots, n}$	(1)
Adım 2: Karar Matrisinin Normalizasyonu (R)	$r_{ij} = \frac{x_{ij}}{\sqrt{\sum_{i=1}^{m}(x_{ij})^2}}$ Fayda maksimizasyonu	(2)
	$r_{ij} = 1 - \frac{x_{ij}}{\sqrt{\sum_{i=1}^{m}(x_{ij})^2}}$ Maliyet minimizasyonu	(3)
	$R = \begin{bmatrix} r_{11} & r_{12} & \cdots & r_{1n} \\ r_{21} & r_{22} & \cdots & r_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ r_{m1} & r_{m2} & \cdots & r_{mn} \end{bmatrix}$	(4)
Adım 3: Tüm Kriterlerin ENTROPİ Değerlerinin Hesaplanması	$f_{ij} = \frac{r_{ij}}{\sum_{i=1}^{m} r_{ij}}$	(5)
	$e_j = -\frac{\sum_{i=1}^{m} f_{ij} * lnf_{ij}}{\ln(m)}$	(6)
Adım 4: Kriterlerin Ağırlık Değerlerinin Belirlenmesi	$W_j = \frac{1 - e_j}{n - \sum_{i=1}^{m} e_j}$	(7)
	$\sum_{i=1}^{n} W_j = 1$	(8)

Yöntem kapsamında dört aşama sonucunda kriterlere dair W_j değerleri elde edilmektedir. İlgili değerler büyükten küçüğe doğru sıralanarak, kriterlerin önem sıralaması tespit edilmektedir.

COBRA Yöntemi

Analizde performans sıralaması için Krstic vd. (2022) tarafından geliştirilen COBRA (Comprehensive Distance Based Ranking) yöntemi uygulanmaktadır. Bu yöntemde pozitif ideal, negatif ideal ve ortalama çözüme dayalı hesaplama yapılmaktadır. Üç yöntemin uzaklıklarının belirlenmesinde Öklid ve Taxicab uzaklıklarından faydalanılmaktadır. Yönteme ilişkin matematiksel denklemlerin aşamaları Tablo 4'te gösterilmektedir (Krstic vd., 2022; Şimşek 2024; Saraç ve Karamaşa 2025).

FULL TEXTS BOOK

Tablo 4. COBRA Yönteminin Metodolojik Aşamaları

Aşama 1: Karar Matrisinin Oluşturulması (i. alternatifin j. kriterini göstermektedir.)	$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$ $i = 1,2, \dots, n$ ve $j = 1,2, \dots, m$	(9)
Aşama 2: Normalize Karar Matrisinin Elde Edilmesi (Δ)	$\Delta = [a_{ij}]$	(10)
	$a_{ij} = \frac{a_{ij}}{max\ a_{ij}}$	(11)
Aşama 3: Ağırlıklandırılmış Normalize Karar Matrisi	$\Delta_W = [w_j * a_{ij}]$	(12)
Aşama 4: Pozitif İdeal ($\boldsymbol{PIS_j}$), Negatif İdeal ($\boldsymbol{NIS_j}$) ve Ortalama Çözüm ($\boldsymbol{AS_j}$) Değerlerinin Hesaplanması	$PIS_j = max_i(w_j * a_{ij}) \quad \forall_j = 1, \dots, m$ için $j \in$ fayda yönlü kriter	(13)
	$PIS_j = min_i(w_j * a_{ij}) \quad \forall_j = 1, \dots, m$ için $j \in$ maliyet yönlü kriter	(14)
	$NIS_j = min_i(w_j * a_{ij}) \quad \forall_j = 1, \dots, m$ için $j \in$ fayda yönlü kriter	(15)
	$NIS_j = max_i(w_j * a_{ij}) \quad \forall_j = 1, \dots, m$ için $j \in$ maliyet yönlü kriter	(16)
	$AS_j = \frac{\sum_{i=1}^{n} w_j * a_{ij}}{n} \quad \forall_j = 1, \dots, m$ için $j \in$ fayda ve maliyet yönlü kriter	(17)
Aşama 5: Alternatiflerin Her Biri İçin Pozitif İdeal ($d(PISj)$) ve Negatif İdeal ($d(NISj)$) Çözümlere Olan Uzaklıkları İle Ortalama Çözüme Olan Pozitif ve Negatif Uzaklık Değerlerinin Hesaplanması	$d(S_j) = dE(S_j) + (\sigma * dE(S_j) * dT(S_j) \quad j = 1, \dots, m$	(18)
	$dE(S_j)$: öklid uzaklığı, $dT(S_j)$: taxicab uzaklığı σ: düzeltme katsayısı $\sigma = max_i\ dE(S_j)_i - min_i dE(S_j)_i$	(19)
	$dE(PIS_j)_i = \sqrt{\sum_{j=1}^{m}(PIS_j - w_j * a_{ij})^2}$	(20)
	$dT(PIS_j)_i = \sum_{j=1}^{m}\lvert PIS_j - w_j * a_{ij}\rvert$	(21)
	$d(PIS_j) = dE(PIS_j) + (\sigma * dE(PIS_j) * dT(PIS_j) \quad j = 1, \dots, m$	(22)
	$\sigma = max_i\ dE(PIS_j)_i - min_i dE(PIS_j)_i$	(23)
	$dE(NIS_j)_i = \sqrt{\sum_{j=1}^{m}(NIS_j - w_j * a_{ij})^2}$	(24)
	$dT(PIS_j)_i = \sum_{j=1}^{m}\lvert PIS_j - w_j * a_{ij}\rvert$	(25)
	$d(NIS_j) = dE(NIS_j) + (\sigma * dE(NIS_j) * dT(NIS_j) \quad j = 1, \dots, m$	(26)
	$\sigma = max_i\ dE(NIS_j)_i - min_i dE(NIS_j)_i$	(27)

FULL TEXTS BOOK

	Ortalama çözümden pozitif ve negatif uzaklık için Öklidyen ve Taksicap mesafelerinin hesaplanması;	
	$dE(AS_j)_i^{+} = \sqrt{\sum_{j=1}^{m} \tau^{+} (AS_j - w_j * a_{ij})^2}$	(28)
	$dT(AS_j)_i^{+} = \sum_{j=1}^{m} \tau^{+} \lvert AS_j - w_j * a_{ij} \rvert$	(29)
	Burada $\tau^{+} = \begin{cases} 1 & eğer\ AS_j < w_j * a_{ij} \\ 0 & eğer\ AS_j > w_j * a_{ij} \end{cases}$	(30)
	$d(AS_j)_i^{+} = dE(AS_j)^{+} + (\sigma * dE(AS_j)^{+} * dT(AS_j)^{+} \quad j = 1, \dots, m$	(31)
	$\sigma = max_i\, dE(AS_j)^{+} - min_i dE(AS_j)^{+}$	(32)
	$dE(AS_j)_i^{-} = \sqrt{\sum_{j=1}^{m} \tau^{-} (AS_j - w_j * a_{ij})^2}$	(33)
	$dT(AS_j)_i^{-} = \sum_{j=1}^{m} \tau^{-} \lvert AS_j - w_j * a_{ij} \rvert$	(34)
	Burada $\tau^{-} = \begin{cases} 1 & eğer\ AS_j > w_j * a_{ij} \\ 0 & eğer\ AS_j < w_j * a_{ij} \end{cases}$	(35)
	$d(AS_j)_i^{-} = dE(AS_j)^{-} + (\sigma * dE(AS_j)^{-} * dT(AS_j)^{-} \quad j = 1, \dots, m$	(36)
	$\sigma = max_i\, dE(AS_j)^{-} - min_i dE(AS_j)^{-}$	(37)
Aşama 6: Alternatiflerin Kapsamlı Mesafelerinin Hesaplanması	$dC_i = \frac{d(PIS_j)_i - d(NIS_j)_i + d(AS_j)_i^{+} + d(AS_j)_i^{-}}{4}$	(38)


Yukarıdaki aşamaların uygulanması sonucunda dC_i skorları elde edilerek küçükten büyüğe doğru sıralanmaktadır. En küçük değere sahip alternatifin en iyi performans sergilediği kabul edilmektedir.

BULGULAR

Analizde performans sıralaması öncesinde değişkenlerin ağırlıklandırmaları ENTROPİ yöntemi yardımıyla elde edilmektedir. Grafik 1'de yer alan W_j skorlarına göre, değişkenler arasında enflasyon, işgücüne katılma oranı ve GİNİ katsayısı göstergelerinin diğerlerine kıyasla önem dereceleri daha yüksektir. Vergi gelirlerinin payı, ihracat ve ithalat göstergeleri ise en az önem derecesine sahiptir.

FULL TEXTS BOOK

Grafik 1. ENTROPİ Ağırlıklandırma Yöntemi Sonuçları



Değişkenlerin önem derecelerinin belirlenmesinin ardından performans analizi için COBRA yöntemi uygulanmaktadır. Analizin veri setini ifade eden karar matrisi Tablo 5'te gösterilmektedir.

Tablo 5. ENTROPİ ve COBRA Yöntemlerinde Kullanılan Karar Matrisi

BÖLGE KODU	İKO	İH	KBG	VG	VGP	İT	YOK	İŞZ	GİNİ	ENF
TR10	54.00	108666008	15684	83.38	43.99	138122531	19.60	12.30	0.434	32.75
TR21	57.30	3092417	12539	70.78	0.49	2819437	19.20	9.30	0.356	37.30
TR22	49.40	964377	8627	79.09	0.39	622697	17.30	7.90	0.367	35.90
TR31	52.10	14737269	11647	83.55	10.38	10956663	19.50	14.70	0.373	34.00
TR32	53.10	6043118	8125	71.81	0.51	2694891	18.10	10.20	0.346	40.51
TR33	51.30	3945052	8115	71.02	0.30	3710095	15.20	6.90	0.339	39.82
TR41	52.40	12948626	10694	70.95	1.19	9601602	13.90	9.40	0.298	35.34
TR42	52.50	18975637	13141	74.99	1.91	19567414	13.20	9.60	0.320	38.42
TR51	52.40	9496357	13084	75.36	11.29	14601816	17.40	13.40	0.350	35.69
TR52	50.20	3299207	7397	75.73	0.51	1269054	16.10	7.90	0.370	39.21
TR61	55.10	3020626	8612	71.77	0.48	1639479	19.00	11.80	0.361	38.98
TR62	50.30	6790746	7649	79.57	1.67	7512501	22.40	13.20	0.391	35.89
TR63	48.10	6151463	6467	78.79	0.79	8697389	19.10	17.10	0.377	37.07
TR71	50.20	661215	6527	70.37	0.09	523495	18.00	13.00	0.353	39.41
TR72	48.40	3711414	7205	65.94	0.34	1842365	16.70	11.70	0.375	38.89
TR81	49.10	1563088	7484	77.47	0.29	3104999	12.70	10.90	0.292	41.20
TR82	51.60	688595	6915	75.69	0.06	351902	18.80	5.80	0.335	41.03
TR83	53.20	4098699	5587	73.43	0.23	2094599	18.20	7.90	0.374	41.97
TR90	55.10	2157388	5787	76.21	0.15	260105	15.50	9.90	0.329	42.08
TRA1	47.90	51230	5978	77.00	0.08	112440	18.50	10.00	0.351	39.16
TRA2	50.70	143205	3902	61.70	0.05	104799	21.60	13.70	0.401	42.64
TRB1	48.30	718733	5703	79.76	0.11	128870	10.40	10.20	0.294	41.49
TRB2	46.60	208842	3543	65.06	0.08	179593	17.60	20.40	0.355	36.37
TRC1	50.00	10475436	6937	57.60	0.37	7513806	12.90	10.10	0.350	37.49
TRC2	40.60	599646	3489	59.73	0.33	449457	17.10	15.00	0.372	38.47
TRC3	42.70	2002973	4852	59.75	0.09	479173	19.30	29.80	0.377	37.06

Karar matrisindeki veriler üzerinden Tablo 6'da normalize karar matrisi ve Tablo 7'de ağırlıklı normalize karar matrisleri hesaplanmaktadır. Ağırlıklı karar matrisi hesaplanırken normalize karar matrisi değerleri, ENTROPİ yöntemiyle elde edilen katsayılar ile çarpılmaktadır.

FULL TEXTS BOOK

Tablo 6. Normalize Karar Matrisi

BÖLGE KODU	İKO	İH	KBG	VG	VGP	İT	YOK	İŞZ	GİNİ	ENF
TR10	0.9424	1.0000	1.0000	0.9980	1.0000	1.0000	0.8750	0.4128	1.0000	0.7681
TR21	1.0000	0.0285	0.7995	0.8471	0.0112	0.0204	0.8571	0.3121	0.8203	0.8748
TR22	0.8621	0.0089	0.5501	0.9466	0.0089	0.0045	0.7723	0.2651	0.8456	0.8419
TR31	0.9092	0.1356	0.7426	1.0000	0.2360	0.0793	0.8705	0.4933	0.8594	0.7974
TR32	0.9267	0.0556	0.5180	0.8594	0.0117	0.0195	0.8080	0.3423	0.7972	0.9500
TR33	0.8953	0.0363	0.5174	0.8500	0.0067	0.0269	0.6786	0.2315	0.7811	0.9339
TR41	0.9145	0.1192	0.6818	0.8492	0.0271	0.0695	0.6205	0.3154	0.6866	0.8288
TR42	0.9162	0.1746	0.8379	0.8976	0.0433	0.1417	0.5893	0.3221	0.7373	0.9010
TR51	0.9145	0.0874	0.8342	0.9020	0.2566	0.1057	0.7768	0.4497	0.8065	0.8370
TR52	0.8761	0.0304	0.4716	0.9063	0.0116	0.0092	0.7188	0.2651	0.8525	0.9196
TR61	0.9616	0.0278	0.5491	0.8590	0.0110	0.0119	0.8482	0.3960	0.8318	0.9142
TR62	0.8778	0.0625	0.4877	0.9524	0.0380	0.0544	1.0000	0.4430	0.9009	0.8417
TR63	0.8394	0.0566	0.4123	0.9430	0.0180	0.0630	0.8527	0.5738	0.8687	0.8694
TR71	0.8761	0.0061	0.4162	0.8423	0.0020	0.0038	0.8036	0.4362	0.8134	0.9242
TR72	0.8447	0.0342	0.4594	0.7892	0.0077	0.0133	0.7455	0.3926	0.8641	0.9121
TR81	0.8569	0.0144	0.4772	0.9273	0.0066	0.0225	0.5670	0.3658	0.6728	0.9662
TR82	0.9005	0.0063	0.4409	0.9059	0.0014	0.0025	0.8393	0.1946	0.7719	0.9622
TR83	0.9284	0.0377	0.3562	0.8789	0.0051	0.0152	0.8125	0.2651	0.8618	0.9843
TR90	0.9616	0.0199	0.3690	0.9121	0.0033	0.0019	0.6920	0.3322	0.7581	0.9869
TRA1	0.8360	0.0005	0.3812	0.9216	0.0019	0.0008	0.8259	0.3356	0.8088	0.9184
TRA2	0.8848	0.0013	0.2488	0.7385	0.0011	0.0008	0.9643	0.4597	0.9240	1.0000
TRB1	0.8429	0.0066	0.3636	0.9546	0.0024	0.0009	0.4643	0.3423	0.6774	0.9730
TRB2	0.8133	0.0019	0.2259	0.7787	0.0018	0.0013	0.7857	0.6846	0.8180	0.8530
TRC1	0.8726	0.0964	0.4423	0.6894	0.0085	0.0544	0.5759	0.3389	0.8065	0.8792
TRC2	0.7086	0.0055	0.2225	0.7149	0.0074	0.0033	0.7634	0.5034	0.8571	0.9022
TRC3	0.7452	0.0184	0.3094	0.7151	0.0020	0.0035	0.8616	1.0000	0.8687	0.8691

Tablo 7. Ağırlıklı Normalize Karar Matrisi

BÖLGE KODU	İKO	İH	KBG	VG	VGP	İT	YOK	İŞZ	GİNİ	ENF
TR10	0.100	0.087	0.105	0.106	0.078	0.093	0.093	0.044	0.106	0.082
TR21	0.106	0.002	0.084	0.090	0.001	0.002	0.091	0.033	0.087	0.093
TR22	0.092	0.001	0.058	0.101	0.001	0.000	0.082	0.028	0.090	0.090
TR31	0.097	0.012	0.078	0.106	0.018	0.007	0.092	0.052	0.091	0.085
TR32	0.099	0.005	0.054	0.091	0.001	0.002	0.086	0.036	0.085	0.101
TR33	0.095	0.003	0.054	0.090	0.001	0.003	0.072	0.024	0.083	0.099
TR41	0.097	0.010	0.072	0.090	0.002	0.006	0.066	0.033	0.073	0.088
TR42	0.097	0.015	0.088	0.095	0.003	0.013	0.063	0.034	0.078	0.096
TR51	0.097	0.008	0.088	0.096	0.020	0.010	0.082	0.047	0.086	0.089
TR52	0.093	0.003	0.050	0.096	0.001	0.001	0.076	0.028	0.091	0.098
TR61	0.102	0.002	0.058	0.091	0.001	0.001	0.090	0.042	0.088	0.097
TR62	0.093	0.005	0.051	0.101	0.003	0.005	0.106	0.047	0.096	0.089
TR63	0.089	0.005	0.043	0.100	0.001	0.006	0.090	0.061	0.092	0.092
TR71	0.093	0.001	0.044	0.090	0.000	0.000	0.085	0.046	0.086	0.098
TR72	0.090	0.003	0.048	0.084	0.001	0.001	0.079	0.041	0.092	0.097
TR81	0.091	0.001	0.050	0.099	0.001	0.002	0.060	0.039	0.072	0.103
TR82	0.096	0.001	0.046	0.096	0.000	0.000	0.089	0.021	0.082	0.102
TR83	0.099	0.003	0.037	0.093	0.000	0.001	0.086	0.028	0.092	0.105
TR90	0.102	0.002	0.039	0.097	0.000	0.000	0.073	0.035	0.081	0.105
TRA1	0.089	0.000	0.040	0.098	0.000	0.000	0.088	0.035	0.086	0.098
TRA2	0.094	0.000	0.026	0.078	0.000	0.000	0.102	0.049	0.098	0.106
TRB1	0.090	0.001	0.038	0.101	0.000	0.000	0.049	0.036	0.072	0.103
TRB2	0.086	0.000	0.024	0.083	0.000	0.000	0.083	0.072	0.087	0.091
TRC1	0.093	0.008	0.047	0.073	0.001	0.005	0.061	0.036	0.086	0.093
TRC2	0.075	0.000	0.023	0.076	0.001	0.000	0.081	0.053	0.091	0.096

FULL TEXTS BOOK

TRC3	0.079	0.002	0.033	0.076	0.000	0.000	0.091	0.106	0.092	0.092
PIS	0.106	0.087	0.105	0.106	0.078	0.000	0.049	0.021	0.072	0.082
NIS	0.075	0.000	0.023	0.073	0.000	0.093	0.106	0.106	0.106	0.106
AS	0.094	0.007	0.053	0.092	0.005	0.006	0.081	0.043	0.087	0.096

Tablo 8. COBRA Yöntemine Göre Nihai Sıralama

BÖLGE KODU	d(PIS)	d(NIS)	d(AS)+	d(AS)-	dCi	Sıralama
TR10	0.11227	0.16776	0.15846	0.01407	-0.04997	1
TR51	0.11313	0.13389	0.03893	0.00686	-0.01321	2
TR42	0.11131	0.14363	0.03733	0.02262	-0.01176	3
TR31	0.11768	0.12951	0.03594	0.01095	-0.00921	4
TR21	0.12920	0.14175	0.03509	0.01271	-0.00873	5
TR41	0.11869	0.14093	0.01938	0.02451	-0.00428	6
TR62	0.14457	0.11812	0.02810	0.00711	0.00136	7
TR61	0.13795	0.12637	0.01315	0.00817	0.00165	8
TR22	0.13464	0.13655	0.01004	0.01861	0.00167	9
TR32	0.13576	0.12841	0.00848	0.00943	0.00208	10
TR33	0.13287	0.13753	0.00411	0.02209	0.00333	11
TR52	0.13631	0.13388	0.00577	0.01780	0.00361	12
TR90	0.14162	0.13221	0.01344	0.02132	0.00432	13
TR81	0.13515	0.13568	0.00941	0.02829	0.00459	14
TR82	0.14196	0.13744	0.01101	0.02562	0.00478	15
TR83	0.14528	0.12939	0.01226	0.02268	0.00658	16
TR72	0.14112	0.12245	0.00497	0.01330	0.00675	17
TR63	0.14706	0.10934	0.02243	0.01195	0.00681	18
TRB1	0.14062	0.14097	0.01201	0.04066	0.00707	19
TRA1	0.14601	0.12643	0.00858	0.01882	0.00745	20
TR71	0.14544	0.12032	0.00576	0.01398	0.00834	21
TRC1	0.13552	0.12933	0.00146	0.02997	0.00868	22
TRC3	0.17776	0.09733	0.06476	0.03160	0.01182	23
TRA2	0.16467	0.11246	0.02677	0.03205	0.01437	24
TRB2	0.16332	0.10664	0.02993	0.03376	0.01513	25
TRC2	0.16133	0.11237	0.01139	0.03985	0.01935	26

COBRA yönteminin tüm aşamalarının uygulanmasının ardından Tablo 8'deki nihai sıralama elde edilmektedir. Buna göre, en düşük dCi skorlarına sahip TR10, TR51, TR42, TR31 ve TR21 bölgeleri en yüksek performans sergileyen beş bölgeyi oluşturmaktadır. Listenin en altında olan TRC2, TRB2, TRA2, TRC3 ve TRC1 bölgeleri ise en yüksek dCi skorları ile en düşük performansı ortaya koymaktadır.

SONUÇ VE POLİTİKA ÖNERİLERİ

Çalışmada, Türkiye'nin alt bölgelerinde sosyo-ekonomik ve mali performans düzeyi Düzey-2 sınıflaması esas alınarak ölçülmektedir. COBRA yönteminin uygulandığı analiz bölümünde kişi başına GSYH, işgücüne katılım, işsizlik, enflasyon, ithalat, ihracat, yoksulluk, gini katsayısı, vergi gayreti ve vergi gelirleri payı değişkenlerinin 2021 yılı verileri incelenmektedir. Elde edilen performans değerlerine göre, TR10 (İstanbul) bölgesi en başarılı bölge olarak sıralamanın en üstünde yer almaktadır. Nüfus, sanayileşme, kentleşme, ekonomik göstergeler gibi birçok açıdan Türkiye'nin en önemli ili olan İstanbul'un, ilk sırada yer alması beklentilerle uyumludur. Sıralamada TR10 bölgesini TR51 (Ankara), TR42 (Kocaeli, Sakarya, Düzce, Bolu, Yalova), TR31 (İzmir) ve TR21 (Tekirdağ, Edirne, Kırklareli) bölgeleri takip etmektedir. Ankara ve İzmir'in yanı sıra en başarılı illerin genel olarak Marmara Bölgesi kapsamında olduğu tespit edilmektedir. Diğer yandan performans sıralamasının en sonunda yer alan bölgeler TRC2 (Şanlıurfa, Diyarbakır), TRB2 (Van, Muş, Bitlis, Hakkari), TRA2 (Ağrı, Kars, Iğdır, Ardahan), TRC3 (Mardin, Batman, Şırnak, Siirt) ve TRC1 (Gaziantep, Adıyaman, Kilis) şeklindedir. Son sırada yer alan bölge ve illerin ise Doğu Anadolu ve Güneydoğu Anadolu bölgelerinde yer aldığı gözlenmektedir.

Analiz sonuçları genel olarak değerlendirildiğinde, ülkenin batı bölgelerinden doğu bölgelerine doğru sosyo-ekonomik ve mali performansın azaldığı sonucuna ulaşılmaktadır. Bu nedenle, bölgesel teşvik mekanizmalarının kapsamı genişletilerek uygulanması gerekmektedir. Maliye politikasının temel amaçlarından biri olan bölgelerarası dengesizliklerin giderilmesi, hükümetlerin öncelikli politikaları

FULL TEXTS BOOK

arasında olmalıdır. Bu kapsamda gelişmişlik seviyesi düşük bölgelere yönelik eğitim, sağlık, kültür vb. alanlarda kamu yatırımlarının artırılması, kamu personeli ücretlerinin teşvik edici şekilde düzenlenmesi, vergi teşviklerinin kapsamının genişletilmesi (Avantajlı gelir vergisi tarifelerinin uygulanması, indirimli kurumlar vergisinin uygulanması, bazı bölgelerde emlak vergisinin indirimli alınması veya alınmaması vb.), tarım ve hayvancılık faaliyetlerine yönelik teşviklerin kapsamının genişletilmesi, bölgesel dinamikleri dikkate alan farklı asgari ücret seviyelerinin belirlenmesi ve doğal kaynaklar analiz edilerek bazı bölgelerin belirli mal ve hizmetlerde ihtisaslaşması önerileri sunulabilir. Ayrıca kentleşme oranının ve nüfus yoğunluğunun yüksek olduğu bölgelerden kırsal kesime dönüşü teşvik edecek politikaların uygulanması da bölgesel eşitsizliklerin minimize edilmesine katkı sağlayacaktır.

Çalışma, bölgesel performans düzeyi hakkında çeşitli bilgiler sunmasına karşın bazı sınırlılıklar içermektedir. Çalışmada sosyo-ekonomik ve mali alanda 10 gösterge incelenmektedir. Gelecekteki araştırmalar, veri erişilebilirliğini dikkate alarak daha yakın dönemleri daha fazla sayıda gösterge ile inceleyebilir. Çalışmada İBBS Düzey-2 kapsamında 26 bölge örneklem olarak belirlenmiştir. Araştırmacılar veri çeşitliliğini gözeterek benzer çalışmayı Düzey-3 yani il düzeyinde gerçekleştirebilir. Son olarak analiz bölümde ağırlıklandırma için ENTROPİ ve sıralama için COBRA yöntemleri uygulanmaktadır. Gelecek çalışmalarda, metodolojik açıdan yenilik sunan farklı çok kriterli karar verme yaklaşımlarından yararlanılabilir.

KAYNAKLAR

Belgin, Ö., & Avşar, B. A. (2019). Türkiye'de bölgeler ve iller düzeyinde ar-ge ve yenilik performansının Gri İlişkisel Analiz Yöntemi ile ölçülmesi. *Verimlilik Dergisi*, (2), 27-48.

Çetin, B., & Kuvat, Ö. (2022). Türkiye'de ekonomik göstergeler açısından Düzey 2 bölgelerinin geliştirilmiş ENTROPİ ve CRITIC temelli COPRAS yöntemi ile sıralanması. *Aksaray Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi,* 14(1), 11-36.

Çıtak, Ö., & Ünlü, U. (2024). BİST'te işlem gören bankaların sürdürülebilirlik raporlarının ÇKKV yöntemi ile değerlendirilmesi. *Alanya Akademik Bakış,* 8(3), 840-857.

Demir, A. & Bakırcı F. (2014). OECD üyesi ülkelerin ekonomik etkinliklerinin veri zarflama analiziyle ölçümü. *Atatürk Üniversitesi İktisadi ve İdari Bilimler Dergisi,* 28(2), 109-132.

DPT (2000). Bölgesel Gelişme Özel İhtisas Komisyonu Raporu. Sekizinci Beş Yıllık Kalkınma Planı, Devlet Planlama Teşkilatı, DPT: 2502 – ÖİK: 523, Ankara.

Erikli, S., & Türkoğlu, S. P. (2020). Türkiye'de Düzey-2 kapsamında yer alan bölgelerin işgücü piyasası performans analizi. *Beykoz Akademi Dergisi*, 8(2), 229-250.

Gazibey, Y., Keser, A., & Gökmen, Y. (2014). Türkiye'de illerin sürdürülebilirlik boyutları açısından değerlendirilmesi. *Ankara Üniversitesi SBF Dergisi,* 69(3), 511-544.

Genç, T. (2016). Doğu ve Güneydoğu Anadolu'dan seçilen illerin sosyo-ekonomik değerlendirilmesine ilişkin bir hibrit çok kriterli karar verme yaklaşımı. *Çukurova Üniversitesi Sosyal Bilimler Enstitüsü Dergisi*, 25(2), 71-88.

GİB (2025). Gelir İdaresi Başkanlığı, İstatistikler, Bütçe Gelirleri, https://www.gib.gov.tr/sites/default/files/fileadmin/user_upload/VI/GBG/Tablo_78.xls.htm (Erişim Tarihi: 25.06.2025).

IMF (2025). International Monetary Fund, https://www.imf.org/en/Publications/WEO/weo-database/2025/april (Erişim Tarihi: 05.07.2025).

Kalkınma Ajansları (2025). Kalkınma Planlamasında İstatistiki Bölge Birimleri Sınıflandırması, https://ka.gov.tr/sayfalar/kalkinma-planlamasinda-istatistiki-bolge-birimleri-siniflandirmasi--24 (Erişim Tarihi: 05.07.2025).

Köse, S., Eser, U., & Konur, F. (2012). Türkiye'de bölgesel gelişmişlik farkları: Bir veri zarflama analizi (Düzey-2 bölgeleri). *Manisa Celal Bayar Üniversitesi Sosyal Bilimler Dergisi,* 10(2), 77-97.

FULL TEXTS BOOK

Krstić, M., Agnusdei, G. P., Miglietta, P. P., Tadić, S., & Roso, V. (2022). Applicability of industry 4.0 technologies in the reverse logistics: a circular economy approach based on comprehensive distance based ranking (COBRA) method. *Sustainability*, 14(9), 5632, 1-30.

Li, X., Wang, K., Liu, L., Xin, J., Yang, H., & Gao, C. (2011). Application of the entropy weight and TOPSIS method in safety evaluation of coal mines. *Procedia Engineering*, 26, 2085-2091.

Örtlek, Z. (2024). Sosyoekonomik performansın analizi: Türkiye Düzey 2 bölgesi. *Nevşehir Hacı Bektaş Veli Üniversitesi SBE Dergisi,* 14(4), 1934-1954.

Samut, P. K. (2024). Türkiye Cumhuriyeti'nin 100 yılının ekonomik değerlendirmesi ve çok kriterli karar verme ile ekonomik performans analizi. *Optimum Ekonomi ve Yönetim Bilimleri Dergisi,* 11(2), 265-286.

Saraç, B., & Karamaşa, Ç. (2025). OECD ülkelerinin yaşam kalitesi düzeyleri açısından LOPCOW-COBRA Entegre Yöntemi ile incelenmesi. Gülay Demir (ed.), Sosyal Bilimlerde Stratejik Karar Verme: Çok Kriterli Karar Verme Yöntemleri ile Uygulamalar (ss. 89-112), Özgür Yayınları, Gaziantep.

Shannon, C. E. (1948). A mathematical theory of communication. *The Bell System Technical Journal*, 27(3), 379-423.

Şahin, C., & Öztel A. (2017). Ülkelerin yaşanabilirlik düzeylerinin COPRAS yöntemiyle karşılaştırmalı analizi: BRICS ülkeleri ve Türkiye. *Uluslararası Batı Karadeniz Sosyal ve Beşeri Bilimler Dergisi,* 1(1), 75-84.

Şimşek, A. (2024). URAP verilerine dayalı üniversite sıralamalarında çok kriterli karar verme yaklaşımı: LOPCOW Temelli COBRA yöntemi. *Premium e-Journal of Social Sciences (PEJOSS),* 8(49), 1766-1779.

TÜİK (2025). Türkiye İstatistik Kurumu, Bölgesel İstatistikler, https://biruni.tuik.gov.tr/bolgeselistatistik/degiskenlerUzerindenSorgula.do# (Erişim Tarihi: 25.06.2025).

UNDP (2025). United Nations Development Programme, Human Development Report, https://hdr.undp.org/data-center/documentation-and-downloads (Erişim Tarihi: 05.07.2025).

Urfalıoğlu, F., & Genç, T. (2013). Çok kriterli karar verme teknikleri ile Türkiye'nin ekonomik performansının Avrupa Birliği Üye ülkeleri ile karşılaştırılması. *Marmara Üniversitesi İktisadi ve İdari Bilimler Dergisi,* 35(2), 329-360.

Wu, J., Sun, J., Liang, L., & Zha, Y. (2011). Determination of weights for ultimate cross efficiency using Shannon entropy. *Expert Systems with Applications*, 38(5), 5162-5165.

Yalçın, E., & Yavuz, E. (2016). İslam İşbirliği Teşkilatı'na üye ülkelerin ekonomik göstergelerinin ARAS (Additive Ratio Assesment) yöntemi ile analizi. XVII. Uluslararası Ekonometri, Yöneylem Araştırması ve İstatistik Sempozyumu (ss. 846-857), Tam Metin Kitabı.

Yaraşır Tülümce, S., & Yavuz, E. (2019). Kamu kesiminin büyüklüğü ve performansının analizi: Avrupa Birliği Ülkeleri ve Türkiye Uygulaması. Konya: Çizgi Kitabevi.

Yavuz, E., Avcı, T., & Çağlar, A. E. (2017). Sosyal Güvenlik Kurumları açısından illerin analizi: MOORA yöntemi. *Adnan Menderes Üniversitesi Sosyal Bilimler Enstitüsü Dergisi,* 4(4), 1-17.

Yürüyen, A. A., & Topal, B. (2025) G7 ülkelerinin lojistik performanslarının PSI Tabanlı COBRA yöntemi ile incelenmesi. *Dicle Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi,* 15(29), 39-57.

Zhang, H., Gu, C. L., Gu, L. W., & Zhang, Y. (2011). The evaluation of tourism destination competitiveness by TOPSIS & information entropy–A case in the Yangtze River Delta of China. *Tourism management*, 32(2), 443-451.

FULL TEXTS BOOK

İlgili makama;

8. Uluslararası Akdeniz Bilimsel Araştırmalar Kongresi 13-15 Ağustos 2025 tarihleri arasında Antalya, Türkiye'de 27 farklı ülkenin akademisyen/araştırmacılarının katılımıyla gerçekleşmiştir. Kongre kapsamında sunumu yapılan 420 bildirinin 206 adeti Türkiye'den katılımcılar tarafından; 214 bildiri ise 27 ülkeden katılımcılar tarafından sunulmuştur. Kongre 16 Ocak 2020 Akademik Teşvik Ödeneği Yönetmeliğine getirilen "*Tebliğlerin sunulduğu yurt içinde veya yurt dışındaki etkinliğin uluslararası olarak nitelendirilebilmesi için Türkiye dışında en az beş farklı ülkeden sözlü tebliğ sunan konuşmacının katılım sağlaması ve tebliğlerin yarıdan fazlasının Türkiye dışından katılımcılar tarafından sunulması esastır.*" değişikliğine uygun düzenlenmiştir.

Bilgilerinize arz edilir,
Saygılarımla,



Dr. Nurlan AKHMETOV
DÜZENLEME KURULU ÜYESİ
Hoca Ahmet Yesevi Uluslararası Kazak-Türk Üniversitesi

T.C.
RECEP TAYYİP ERDOĞAN ÜNİVERSİTESİ
Pazar Meslek Yüksekokulu Müdürlüğü

2025 AİLE YILI

Sayı : E-98028430-900-2040 14.07.2025
Konu : Kongre Düzenleme ve Bilim Kurulu Görevlendirmesi

Sayın Dr. Öğr. Üyesi Mehmet TATOĞLU

İlgi : 13.07.2025 tarihli dilekçeniz.

İlgide kayıtlı dilekçenize istinaden 13-15 Ağustos 2025 tarihleri arasında Antalya'da düzenlenecek olan **"8. Uluslararası Akdeniz Bilimsel Araştırmalar Kongresi"** adlı uluslararası kongrenin Düzenleme ve Bilim Kurulunda görev alma isteğiniz Müdürlüğümüzce uygun görülmüştür.

Bilgilerinize rica ederim.

Dr. Öğr. Üyesi Seda OKUR
Müdür V.

Bu belge, güvenli elektronik imza ile imzalanmıştır.
Belge Doğrulama Kodu : ce0830daf904 Belge Takip Adresi: http://ebys.erdogan.edu.tr/EBYS/eimzadogrulama

Katori Mevkii 53300 Pazar/RİZE
Telefon No: (0 464) 612 73 17 Faks No: (0 464) 612 73 16
e-Posta: pazarmyo@erdogan.edu.tr İnternet Adresi: www.erdogan.edu.tr
Kep Adresi: erdogan@hs01.kep.tr

Bilgi için: Meryem TEMELCİ
Bilgisayar İşletmeni
Telefon No: 2902